Major chemical and mineral components of profundal surface sediments in Minnesota lakes¹

Walter E. Dean²

Department of Geology, Syracuse University, Syracuse, New York 13210

Eville Gorham

Department of Ecology and Behavioral Biology, University of Minnesota, St. Paul 55108

Abstract

Minnesota lakes, characterized primarily on the basis of water chemistry, can also be classified according to the proportions of organic matter, CaCO_3 , and clastic minerals in their profundal sediments. The northeastern (group 1) lakes are characterized by shallow to very deep basins in Precambrian crystalline rocks, active outlets, and dilute surface waters. Profundal sediments of these lakes contain no sedimentary calcium carbonate and can be subdivided on the basis of organic content. Highly organic sediments of group 1 lakes contain more than 30% organic matter (as loss on ignition), but most group 1 lake sediments contain less than this.

The central group 2 lakes occupy shallow to moderately deep basins in calcareous glacial till, have inactive outlets as a result of net evaporation, contain more concentrated salts than group 1 lakes, and commonly precipitate CaCO_3 during the summer. These lakes can be subdivided on the basis of sedimentary CaCO_3 . Sediments of group 2 lakes with more than 30% organic material contain little or no CaCO_3 . Southern group 2 lake sediments are characterized by 10–30% CaCO_3 and 10–30% organic matter. Sediments of western group 2 lakes contain 30–50% CaCO_3 and about twice as much clastic as organic materials. Group 3 lakes occupy shallow depressions in calcareous, sulfur-rich glacial drift of the arid southwestern prairie region and contain even more concentrated salts than group 2 lakes. Sediments in these lakes are mainly clastic, averaging about 12% organic matter and 20% CaCO_3 .

Variations in water and sediment chemistry, and in sediment mineralogy of these lakes are clearly related to an increase in climatic aridity and aquatic productivity from northeast to southwest. Differences in nature of the underlying glacial drift and in degree of erosion also affect the composition of the lake sediments.

Minnesota has more than 15,000 lakes in a landscape characterized by diverse bedrock and surficial geology (Fig. 1). It also exhibits marked climatic gradients. Annual precipitation is 70–80 cm in the eastern half of the state and 50–70 cm in the western half (Ackroyd et al. 1967). Runoff exceeds 25 cm in northeastern Minnesota, 20 cm in the southeast, and falls to less than 5 cm in the west (Minn. Dep. Conserv. 1962). Mean annual temperatures range from 2°C in the northeast to 8°C in the south (Baker and Strub 1965). According to Bright

(1968), the balance of precipitation minus evaporation in the state is positive (0–8 cm) only in the northeast and is distinctly negative (–15 to –30 cm) in the west.

The chemistry of surface waters reflects the combined influences of these environmental variables, and on this basis Gorham et al. (in prep.) have divided Minnesota lakes into four distinct geographic groups (Fig. 2). The northeastern (group 1) lakes contain dilute bicarbonate waters (conductivity $<160 \mu\text{mhos cm}^{-1}$ at 25°C) in shallow to relatively deep basins (up to 64 m) in Precambrian crystalline rocks. In this heavily forested region precipitation exceeds evaporation, so that the northeastern lakes have active outlets throughout the year. A few group 1 lakes are also found elsewhere on coarse, noncalcareous drift. The central (group 2) lakes contain bicarbonate waters of intermediate concentration (conductiv-

¹ Contribution No. 135, Limnological Research Center, University of Minnesota. This work was supported by National Science Foundation grants GB-6018 and GB 18800 and the Graduate School, University of Minnesota.

² Present address: U.S. Geological Survey, Branch of Regional Geochemistry, Bldg. 53, Federal Center, Lakewood, Colorado 80225.

Fig. 1. Maps of bedrock geology (A) and major surficial deposits (B) (modified from Ackroyd et al. 1967 and Winter 1974).

ity 160–500 $\mu\text{mhos cm}^{-1}$ at 25°C) in shallow to moderately deep basins in calcareous drift. The central lakes are found in both forest and prairie regions, generally have inactive outlets, and many (or most) of them precipitate carbonate biogenically during summer (Megard 1968). Group 3 lakes have high concentrations of bicarbonate and sulfate waters (conductivity >500 $\mu\text{mhos cm}^{-1}$ at 25°C) and occupy shallow depressions in sulfur-bearing glacial drift (cm. Winter 1974) in the southwestern prairie region. Group 4 consists of a very few lakes along the western border with high concentrations of sodium and sulfate (conductivity >7,000 $\mu\text{mhos cm}^{-1}$ at 25°C); there are many more such lakes in the Dakotas (e.g. Mitten et al. 1968). Within Minnesota, concentrations of dissolved salts in lake water increases more than 1,000-fold from the northeastern lakes to those of the southwest—this gradient reflecting mainly the change from positive to negative balance between precipitation and evaporation along the same gradient (Bright 1968). The boundaries between groups are relatively sharp, however, suggesting that surficial or bedrock geology,

or both, are more important than climate in determining actual group boundaries.

Here we analyze the major components of profundal sediments from 46 lakes (14 group 1, 27 group 2, 4 group 3, and 1 group 4) in relation to environmental factors. The top 10 cm of profundal sediments from the deepest portions of the lakes were chosen for analysis in the belief that they would provide the best integration of allochthonous materials from the drainage basin and autochthonous materials produced within the lake. Analyses of sediments along transects in several lakes demonstrated that excessive solution of sedimentary carbonate minerals in the deeper portions of some lakes is the most serious error in this assumption. This problem is discussed in a later section.

We are grateful to H. E. Wright, Jr., G. J. Brunskill, and R. Y. Anderson for helpful reviews of the manuscript. Additional suggestions, particularly regarding clay mineralogy, were provided by W. Parham.

Methods

Undisturbed samples of profundal sediments were collected with a Jenkin sampler

Fig. 2. Map of lake location and profundal sediment composition for all 46 lakes. Lakes are numbered in order of increasing salinity, used for reference in tables. Symbols reflect surface water salinity (groups 1-4), further modified to separate lakes with profundal sediments rich in carbonate and those rich in organic matter.

(Mortimer 1941, 1942). Depth of penetration ranged from 10-15 cm in clayey sediments with little organic matter to more than 30 cm in flocculent highly organic sediments. The top 10 cm of core used does not represent the same time interval in all cores. Samples were stored in polyethylene containers precleaned with hydrochloric acid and rinsed with distilled water.

In the laboratory, samples were air-dried at 90°C and ground to powder. Percent organic matter and carbonate were determined by loss on ignition at 550°C and 1,000°C (Dean 1974). Ignition loss at 550°C was assumed to represent ignition of organic matter; this was checked chromatographically with a Hewlett-Packard CHN analyzer (Fig. 3). Ignition loss between 550° and 1,000°C was assumed to represent loss of CO₂ from carbonate minerals and percent calcium carbonate calculated by multiplying the percent ignition loss between 550° and 1,000°C by 2.27, the factor for converting CO₂ to CaCO₃.

Fig. 3. Scatter plot of organic matter (by loss on ignition at 550°C) and organic carbon in profundal sediments of all 46 lakes.

The accuracy of this method has been checked by comparison with several standard methods (Dean 1974). Lake sediments with considerable clay but no carbonate usually have a 1.0 to 1.8% weight loss between 550° and 1,000°C, presumably due to loss of water from clay lattices; this converts to 2.3 to 4.1% CaCO₃ if we assume that all of the loss is due to CO₂ evolved from carbonate. However, standard additions of CaCO₃ to noncalcareous lake sediment show nearly 100% recovery by the ignition loss method for CaCO₃ concentrations above 10% (Dean 1974). The difference between 100% and the sum of ignition loss organic matter and carbonate was taken as the inorganic, noncarbonate clastic fraction. Ignition loss determinations for organic, carbonate, and clastic components are given in Table 1 as percent dry weight.

Total and soluble Ca and Mg were determined with an atomic absorption attachment for a Beckman DU spectrophotometer. Samples (500 mg) for analysis of total Ca and Mg were digested and evaporated to dryness successively in HNO₃, HF, and HCl. The final dried residue was taken up in 1.0 N HCl. For samples high in Ca and Mg, 100-fold dilutions were necessary. The final solutions contained 1% lanthanum to suppress interference by silica and alumina. Samples for soluble Ca and Mg were leached with 1.0 N ammonium acetate at pH 5.2; according to Wangersky and Joensuu (1967) and confirmed here, this

Table 1. Location, morphometry, and composition of profundal sediments from all 46 lakes. Lakes listed in order of increasing total cation concentration in surface waters.

			1	2	3	4	5	6	7	8	9	10
LAKE NO.	LAKE NAME	COUNTY	AREA (SQ.KM.)	MAX. DEPTH (M)	L.O.I. 550-C (%)	TOTAL CARBON (%)	ORGANIC CARBON (%)	H (%)	N (%)	L.O.I. 1000-C (%)	TOTAL CA (%)	SOLUBLE CA (%)
GROUP 1	1 BIG	ST. LOUIS	6.88	4.57	38.10	18.04	17.60	2.75	1.78	3.10	0.93	0.70
	2 IRON	COOK	0.42	5.79	40.20	25.31	24.90	3.55	1.79	3.70	0.84	0.70
	3 MOUNTAIN	COOK	8.45	64.01	19.20	10.03	9.60	1.69	0.86	3.30	0.81	0.50
	4 TROUT	COOK	1.05	21.34	21.00	9.00	8.50	1.40	1.00	4.50	1.03	0.50
	5 CLEARWATER	COOK	5.36	39.62	15.80	6.11	5.80	1.03	0.50	2.80	0.84	0.50
	6 WILSON	LAKE	2.47	14.02	17.40	8.00	7.70	1.70	0.60	2.40	0.79	0.40
	7 SAND POINT	ST. LOUIS	7.33	53.34	17.90	7.59	7.10	1.73	0.63	4.20	0.78	0.60
	8 KIMBALL	COOK	0.32	5.49	46.90	23.94	23.50	3.58	2.85	3.60	0.88	0.60
	9 CRANE	ST. LOUIS	12.50	24.38	16.90	7.06	6.60	1.22	0.82	4.00	0.96	0.70
	10 O'LEARY	ST. LOUIS	0.80	17.07	19.20	9.04	8.70	1.38	0.97	2.20	1.21	0.60
	11 JOSEPHINE	HUBBARD	0.01	11.55	40.20	20.92	20.60	2.89	2.09	2.50	0.87	0.90
	12 ARCO	HUBBARD	0.01	7.60	41.60	21.65	21.30	2.85	2.36	3.60	0.91	1.00
	13 DEMING	HUBBARD	0.01	17.00	47.90	25.05	24.60	3.31	2.74	3.80	0.63	0.80
	14 FAROUAR	DAKOTA	0.27	1.83	16.30	6.98	6.70	1.05	0.92	2.60	0.80	0.60
GROUP 2	15 GEORGE	ANOKA	2.19	9.75	39.50	17.67	17.30	3.47	2.16	3.10	1.15	1.00
	16 LINWOOD	ANOKA	1.98	10.67	35.20	17.16	15.90	3.24	1.97	10.40	5.80	5.10
	17 CEDAR BOG	ANOKA	0.00	1.22	60.00	28.98	28.60	3.95	3.47	4.20	1.14	0.90
	18 SPECTACLE	ISANTI	0.97	14.33	43.00	21.47	21.10	3.11	2.66	3.50	1.11	1.10
	19 GLADSTONE	CROW WING	1.95	9.45	43.40	21.17	20.90	3.12	2.61	3.80	1.61	1.50
	20 HAM	ANOKA	0.53	6.71	27.20	17.19	12.70	1.78	1.39	37.30	16.40	16.50
	21 GREEN	ISANTI	3.25	8.53	20.90	14.27	10.60	1.36	1.26	30.90	12.70	12.90
	22 REEDS	WASECA	0.76	17.68	24.30	13.71	12.50	1.83	1.45	10.30	3.44	1.50
	23 FRANCIS	LESUEUR	3.63	17.68	26.20	14.06	12.80	1.87	1.60	11.10	3.51	2.50
	24 NOKAY	CROW WING	2.64	13.11	38.40	21.41	19.50	2.76	2.48	15.70	7.60	6.80
	25 BEAVER	STEELE	0.40	7.32	16.60	12.16	9.00	1.27	0.99	26.60	10.30	9.70
	26 MOOSE	ITASCA	4.86	21.34	6.80	11.81	8.40	1.19	0.90	28.30	11.00	10.70
	27 BALL CLUB	ITASCA	15.93	25.91	30.60	17.83	17.10	2.50	1.85	5.60	1.94	1.70
	28 ST. OLAF	WASECA	0.41	9.14	16.10	10.44	8.20	1.22	0.96	18.00	6.80	6.90
4 GR. 3	29 CHRISTMAS	HENNEPIN	1.05	27.43	18.80	9.53	8.30	1.25	0.99	10.30	3.61	1.50
	30 LOTUS	CARVER	0.98	10.67	16.40	9.06	7.00	0.58	0.87	17.10	6.90	6.60
	31 CLEAR	LESUEUR	1.27	6.10	16.50	10.28	8.80	1.48	0.97	12.10	4.80	4.20
	32 LONG	CLEARWATER	0.68	24.38	12.50	10.82	6.00	0.82	0.64	40.30	15.20	16.20
	33 ELK	CLEARWATER	1.21	29.57	24.20	11.78	9.50	1.80	0.99	19.10	6.10	7.80
	34 ITASCA	CLEARWATER	4.36	13.72	22.30	15.00	10.80	1.52	1.26	34.60	14.30	12.60
	35 LITTLE PINE	OTTERTAIL	8.24	19.20	17.80	14.14	8.50	1.17	1.11	46.30	19.90	18.10
	36 GROVE	POPE	1.53	9.45	21.90	15.53	11.20	1.43	1.25	35.60	14.10	14.50
	37 FISH	COTTONWOOD	1.21	7.92	16.00	9.95	7.50	1.23	0.92	21.00	7.10	6.10
	38 SALLIE	BECKER	5.11	14.33	18.80	15.05	9.50	1.51	1.20	46.40	16.50	17.10
	39 COWDRY	DOUGLAS	0.96	15.85	21.50	14.58	10.80	1.59	1.35	31.30	12.00	12.70
	40 MAPLE	DOUGLAS	3.38	21.95	18.40	14.66	9.60	1.22	1.13	42.60	15.90	16.30
	41 MINA	DOUGLAS	1.72	37.49	17.10	13.48	8.80	1.49	1.04	39.40	15.30	17.20
	42 SHETEK	MURRAY	17.40	5.49	15.40	12.04	8.90	1.26	0.91	22.50	9.00	8.10
43 BIG KANDIYOH	KANDIYOH	15.38	3.05	13.30	10.34	7.60	1.13	0.72	22.50	9.30	8.30	
44 ELK	GRANT	0.77	7.92	15.60	10.13	7.80	1.35	0.81	19.00	6.80	5.30	
45 DEAD COON	LINCOLN	2.35	3.35	6.40	5.20	2.90	0.51	0.34	19.10	7.50	6.40	
46 SALT	LACQUI PARLE	1.00	0.30	11.30	5.38	3.30	0.89	0.41	17.10	6.00	5.70	
	mean		3.20	15.82	24.59	14.02	12.06	1.85	1.36	16.33	6.24	5.93
	standard dev.		4.36	12.92	12.24	5.79	6.28	0.90	0.73	14.02	5.69	5.86
	low organic	mean	4.78	29.45	17.96	7.98	7.59	1.40	0.79	3.25	0.90	0.55
	Group 1	standard dev.	4.39	21.09	1.74	1.30	1.27	0.29	0.19	0.88	0.15	0.09
	high organic	mean	2.56	11.23	41.92	21.58	20.99	3.16	2.37	5.12	1.95	1.75
	Group 1 and 2	standard dev.	4.44	6.38	7.09	3.52	3.66	0.41	0.50	3.75	2.17	1.92
	intermediate	mean	1.58	15.49	18.19	11.28	9.20	1.37	1.06	17.39	6.35	5.75
	Group 2	standard dev.	1.47	8.57	5.63	1.71	1.95	0.39	0.25	6.57	2.68	3.26
	high carb.	mean	2.98	17.16	19.84	14.47	9.85	1.39	1.16	38.47	15.23	15.41
	Group 2	standard dev.	2.44	9.17	3.90	1.62	1.83	0.27	0.21	5.56	2.21	2.10
	Group 3	mean	7.38	4.02	12.40	8.62	6.10	1.23	0.64	20.05	7.72	6.76
		standard dev.	8.28	2.85	3.78	3.12	2.79	0.34	0.25	2.38	1.41	1.37

reagent dissolves calcite and aragonite but not dolomite. Results are given in Table 1 as percent dry weight of sample.

Sediments for size analyses were pre-treated with 1.0 N ammonium acetate at pH 5.2 to remove calcium carbonate and 30% H₂O₂ to remove most of the organic fraction. They were then washed through a 44- μ nylon sieve. The sand-size fraction

retained on the sieve was washed, dried, weighed, and mounted on slides for mineralogical analyses. Material passing the sieve (silt + clay) was centrifuged to separate silt and clay fractions (Jackson 1967). After centrifugation, the supernatant containing the clay was filtered through a 0.45- μ Millipore filter and saved for X-ray analysis. The silt-size fraction remaining

Table 1. Continued.

			11	12	13	14	15	16	17	18	19
LAKE NO.	LAKE NAME	COUNTY	X-RAY CALCITE (CM)	X-RAY ARAGON (CM)	X-RAY DOLOM. (CM)	TOTAL MG (%)	SOLUBLE MG (%)	CLASTIC (%)	SAND (%)	SILT (%)	CLAY (%)
GROUP 1	1 BIG	ST. LOUIS	0.0	0.0	0.0	0.29	0.10	58.80	0.70	19.60	38.00
	2 IRON	COOK	0.0	0.0	0.0	0.44	0.07	56.10	1.30	18.70	21.40
	3 MOUNTAIN	COOK	0.0	0.0	0.0	0.74	0.05	77.50	0.60	31.20	31.80
	4 TROUT	COOK	0.0	0.0	0.0	0.44	0.12	74.50	0.30	30.10	37.00
	5 CLEARWATER	COOK	0.0	0.0	0.0	0.88	0.13	81.40	2.60	33.80	32.80
	6 WILSON	LAKE	0.0	0.0	0.0	0.82	0.10	80.20	23.50	21.70	34.40
	7 SAND POINT	ST. LOUIS	0.0	0.0	0.0	0.56	0.04	77.90	13.80	46.60	16.40
	8 KIMBALL	COOK	0.0	0.0	0.0	0.53	0.08	49.50	0.30	34.90	22.20
	9 CRANE	ST. LOUIS	0.0	0.0	0.0	0.87	0.07	79.10	0.10	22.80	43.20
	10 O'LEARY	ST. LOUIS	0.0	0.0	0.0	0.93	0.08	78.60	2.60	32.30	24.40
	11 JOSEPHINE	HUBBARD	0.0	0.0	0.0	0.51	0.08	57.30	1.50	23.50	21.80
	12 ARCO	HUBBARD	0.0	0.0	0.0	0.51	0.10	54.80	0.60	25.60	18.40
	13 DEMING	HUBBARD	0.0	0.0	0.0	0.43	0.08	48.30	0.70	18.20	16.00
	14 FARQUAR	DAKOTA	0.0	0.0	0.0	0.62	0.08	81.10	0.40	15.40	24.60
GROUP 2	15 GEORGE	ANOKA	0.0	0.0	0.0	0.52	0.13	57.40	2.10	25.50	25.20
	16 LINWOOD	ANOKA	10.16	0.0	2.79	0.32	0.15	54.40	1.20	21.40	19.00
	17 CEDAR BOG	ANOKA	0.0	0.0	0.0	0.24	0.08	35.80	0.30	10.40	12.80
	18 SPECTACLE	ISANTI	0.0	0.0	0.0	0.44	0.13	53.50	2.20	23.90	16.60
	19 GLADSTONE	CROW WING	0.0	0.0	0.0	0.45	0.15	52.80	6.90	29.90	15.20
	20 HAM	ANOKA	25.15	0.0	0.0	0.48	0.25	35.50	2.20	9.90	12.80
	21 GREEN	ISANTI	20.32	0.0	0.76	0.48	0.23	48.20	1.80	15.70	15.20
	22 REEDS	WASECA	4.32	0.0	2.03	0.98	0.38	65.40	1.50	25.10	24.60
	23 FRANCIS	LESUEUR	2.54	0.0	3.56	1.20	0.64	62.70	2.10	24.70	24.40
	24 NOKAY	CROW WING	8.89	0.0	0.0	0.46	0.16	45.90	2.00	25.00	25.00
	25 BEAVER	STEELE	13.97	0.0	1.78	1.13	0.44	56.80	6.60	30.00	25.40
	26 MOOSE	ITASCA	16.51	0.0	2.29	1.45	0.77	54.90	5.80	17.50	24.20
	27 BALL CLUB	ITASCA	1.27	0.0	0.0	0.55	0.28	63.80	11.50	28.70	16.80
	28 ST. OLAF	WASECA	11.43	0.0	1.78	0.98	0.42	65.90	1.40	34.30	24.60
GROUP 3	29 CHRISTMAS	HENNEPIN	5.08	0.0	2.54	0.84	0.46	70.90	1.00	31.70	24.00
	30 LOTUS	CARVER	4.57	0.0	1.02	1.00	0.46	66.50	1.30	27.10	40.40
	31 CLEAR	LESUEUR	5.08	0.0	1.02	1.09	0.53	71.40	0.70	16.70	32.40
	32 LONG	CLEARWATER	25.15	0.0	1.02	1.08	0.66	47.20	2.30	24.20	11.20
	33 ELK	CLEARWATER	0.0	0.0	0.0	0.45	0.30	56.70	0.30	18.00	7.20
	34 ITASCA	CLEARWATER	25.15	0.51	0.0	0.57	0.34	43.10	0.20	9.40	9.80
	35 LITTLE PINE	OTTERTAIL	25.15	0.0	0.51	0.71	0.45	35.90	1.50	11.60	13.20
	36 GROVE	POPE	21.59	0.0	0.0	0.93	0.52	42.50	0.30	16.70	18.00
	37 FISH	COTTONWOOD	11.43	1.02	1.02	1.11	0.58	63.00	2.40	26.70	27.60
	38 SALLIE	BECKER	25.15	0.0	1.02	1.23	0.72	34.80	0.90	9.90	14.60
	39 COWDRY	DOUGLAS	19.05	1.27	1.02	0.94	0.67	47.20	0.30	18.70	16.40
	40 MAPLE	DOUGLAS	16.51	1.27	1.02	1.16	0.79	39.00	1.50	11.40	12.40
	41 MINA	DOUGLAS	22.86	0.76	0.0	0.90	0.68	43.50	5.90	19.50	15.40
	42 SHETEK	MURRAY	17.78	0.0	1.78	1.66	0.93	62.10	2.10	22.90	17.60
4	43 BIG KANDIYOH	KANDIYOH	16.51	1.02	2.29	1.14	0.39	64.20	1.60	36.40	22.20
	44 ELK	GRANT	25.15	1.52	5.59	2.79	0.78	65.40	6.10	28.30	21.00
	45 DEAD COON	LINCOLN	7.62	0.0	4.32	1.28	0.44	74.50	5.40	29.10	29.00
	46 SALT	LACUI PARLE	6.35	0.0	1.52	1.46	0.59	71.60	1.90	21.30	29.00
	mean		8.59	0.15	0.88	0.84	0.34	58.86	2.88	23.39	22.30
	standard dev.		9.58	0.40	1.29	0.45	0.26	13.69	4.22	8.08	8.39
	low organic	mean	0.00	0.00	0.00	0.73	0.08	78.78	5.49	29.24	30.58
	Group 1	standard dev.	0.00	0.00	0.00	0.18	0.03	2.24	8.59	9.46	8.42
	high organic	mean	1.56	0.00	0.21	0.43	0.12	52.95	2.40	23.48	20.65
	Group 1 and 2	standard dev.	3.56	0.00	0.77	0.10	0.06	6.97	3.22	6.13	6.43
	intermediate	mean	7.49	0.10	1.70	1.02	0.50	63.42	2.39	25.18	25.48
	Group 2	standard dev.	5.43	0.32	1.00	0.26	0.14	5.79	2.10	6.12	8.27
	high carb.	mean	22.61	0.38	0.54	0.85	0.53	41.69	1.69	14.70	13.90
	Group 2	standard dev.	3.14	0.54	0.49	0.28	0.20	5.12	1.67	5.04	2.48
	Group 3	mean	14.68	0.51	3.10	1.67	0.63	67.56	3.42	27.60	23.76
		standard dev.	7.78	0.72	1.77	0.66	0.23	5.25	2.15	5.96	5.07

in the centrifuge tube was dried and mounted on glass slides for X-ray analysis. Results of size analyses are given in Table 1.

Sand grains were mounted on a glass slide with epoxy cement and ground with 400-mesh carborundum grit to give a flat surface, then etched with HF fumes, and stained with sodium cobaltinitrite and amaranth (Boone and Wheeler 1968). This

procedure stains plagioclase red, orthoclase yellow, and leaves quartz grains clear. Percentages of quartz, plagioclase, orthoclase, and accessory minerals were determined by point-counting at least 100 grains under a petrographic microscope. Only 23 samples contained enough sand grains to be counted (Table 2).

Mineralogy of the silt-size fraction was

Table 1. Continued.

	LAKE NO.	LAKE NAME	COUNTY	20	21	22	23	24	25	26	27
				X-RAY 4.26Å QUARTZ (CM)	PEAK HTS. 4.20Å F'SPAR (CM)	2-44J 4.18Å F'SPAR (CM)	RELATIVE PEAK HEIGHTS (GLYCOL)				X-RAY TOTAL QUARTZ (CM)
							ILLITE (CLAY)	CHLOR. (CLAY)	QUARTZ (CLAY)	F'SPAR (CLAY)	
GROUP 1	1	BIG	ST. LOUIS	1.27	1.27	2.54	1.00	1.00	3.00	2.00	3.81
	2	IRON	COOK	1.52	1.02	4.06	0.0	0.0	1.00	0.0	6.35
	3	MOUNTAIN	COOK	1.27	0.25	2.03	2.00	2.00	2.00	1.00	7.62
	4	TROUT	COOK	1.27	0.25	1.02	0.0	1.00	1.00	0.0	7.62
	5	CLEARWATER	COOK	4.57	0.0	0.0	2.00	2.00	2.00	0.0	8.89
	6	WILSON	LAKE	1.52	1.02	3.56	1.00	1.00	1.00	1.00	15.24
	7	SAND POINT	ST. LOUIS	1.27	1.27	2.54	0.0	1.00	1.00	1.00	5.00
	8	KIMBALL	COOK	2.54	3.05	5.84	1.00	0.0	1.00	1.00	8.89
	9	CRANE	ST. LOUIS	4.57	3.05	5.84	1.00	2.00	2.00	1.00	11.43
	10	O'LEARY	ST. LOUIS	3.81	1.78	4.57	2.00	2.00	2.00	2.00	25.15
	11	JOSEPHINE	HUBBARD	5.08	4.57	8.64	1.00	2.00	2.00	2.00	17.78
	12	ARCO	HUBBARD	6.10	2.54	3.81	1.00	2.00	2.00	2.00	16.51
	13	DEMING	HUBBARD	5.08	2.54	5.84	1.00	2.00	2.00	2.00	17.78
	14	FARQUAR	DAKOTA	4.57	1.78	2.54	2.00	2.00	2.00	1.00	25.15
	15	GEORGE	ANOKA	3.56	0.25	2.54	1.00	0.0	1.00	1.00	10.16
GROUP 2	16	LINWOOD	ANOKA	1.78	1.27	2.79	0.0	0.0	1.00	0.0	6.35
	17	CEDAR ROG	ANOKA	4.83	3.56	4.83	0.0	0.0	0.0	0.0	0.0
	18	SPECTACLE	ISANTI	4.32	2.54	3.05	0.0	0.0	1.00	0.0	19.05
	19	GLADSTONE	CROW WING	0.0	0.0	0.0	0.0	0.0	1.00	1.00	0.0
	20	HAM	ANOKA	3.05	2.54	4.57	0.0	0.0	0.0	0.0	10.16
	21	GREEN	ISANTI	2.03	1.78	2.29	0.0	1.00	1.00	0.0	8.89
	22	REEDS	WASECA	3.81	0.0	0.0	2.00	2.00	2.00	1.00	17.78
	23	FRANCIS	LESUEUR	6.35	3.56	5.33	2.00	2.00	2.00	2.00	10.16
	24	NOKAY	CROW WING	2.29	3.05	3.30	1.00	1.00	1.00	0.0	6.35
	25	BEAVER	STEELE	5.33	0.0	2.79	4.00	2.00	2.00	1.00	13.97
	26	MOOSE	ITASCA	6.60	3.81	8.38	2.00	2.00	2.00	1.00	8.89
	27	BALL CLUB	ITASCA	0.76	0.0	0.25	0.0	0.0	2.00	0.0	8.89
	28	ST. OLAF	WASECA	6.35	0.0	0.0	2.00	2.00	2.00	2.00	13.97
	29	CHRISTMAS	HENNEPIN	3.56	0.0	2.54	2.00	2.00	2.00	2.00	19.05
	30	LOTUS	CARVER	5.08	2.54	3.56	2.00	2.00	2.00	2.00	7.62
GROUP 3	31	CLEAR	LESUEUR	7.62	4.06	5.33	2.00	2.00	2.00	2.00	16.51
	32	LONG	CLEARWATER	2.54	0.51	2.29	2.00	2.00	2.00	2.00	8.89
	33	ELK	CLEARWATER	4.06	1.27	2.79	0.0	0.0	1.00	0.0	0.0
	34	ITASCA	CLEARWATER	3.56	5.59	6.86	0.0	0.0	1.00	0.0	3.05
	35	LITTLE PINE	OTTERTAIL	4.83	6.60	6.35	1.00	0.0	2.00	2.00	2.54
	36	GROVE	POPE	5.33	2.54	4.06	2.00	2.00	2.00	1.00	8.89
	37	FISH	BECKER	6.86	2.03	3.81	2.00	2.00	2.00	0.0	13.97
	38	SALLIE	DOUGLAS	6.35	4.57	6.86	2.00	2.00	2.00	2.00	5.08
	39	COWDRY	DOUGLAS	2.54	2.03	3.30	1.00	1.00	2.00	1.00	6.35
	40	MAPLE	DOUGLAS	6.60	4.06	5.33	2.00	2.00	2.00	2.00	5.08
	41	MINA	DOUGLAS	3.30	3.30	4.06	1.00	1.00	2.00	1.00	5.08
	42	SHETEK	MURRAY	5.33	1.52	2.29	2.00	2.00	2.00	1.00	15.24
	43	BIG KANDIYOH	KANDIYOHI	5.59	2.03	2.03	2.00	3.00	2.00	1.00	15.24
	44	ELK	GRANT	5.59	1.52	2.54	3.00	3.00	2.00	1.00	13.97
	45	DEAD COON	LINCOLN	7.11	0.0	2.29	3.00	3.00	2.00	2.00	22.86
	46	SALT	LACQUA PARLE	7.37	2.79	4.06	2.00	2.00	2.00	2.00	16.51
		mean		4.10	2.04	3.55	1.30	1.36	1.65	1.07	10.81
		standard dev.		2.00	1.63	2.08	0.99	0.96	0.60	0.80	6.38
	low organic	mean		2.86	1.18	2.76	1.25	1.63	1.63	0.88	13.27
	Group 1	standard dev.		1.65	1.03	1.88	0.89	0.52	0.52	0.64	7.93
	high organic	mean		3.01	1.97	3.65	0.54	0.62	1.38	0.85	9.38
	Group 1 and 2	standard dev.		1.94	1.45	2.32	0.52	0.87	0.77	0.90	6.59
	intermediate	mean		5.56	1.73	3.45	2.00	1.80	1.90	1.30	12.19
	Group 2	standard dev.		1.41	1.70	2.52	0.94	0.63	0.32	0.82	5.69
	high carb.	mean		4.01	3.35	4.60	1.10	1.10	1.60	1.10	6.40
	Group 2	standard dev.		1.65	1.86	1.72	0.88	0.88	0.70	0.88	2.67
	Group 3	mean		6.20	1.57	2.64	2.40	2.60	2.00	1.40	16.76
		standard dev.		0.96	1.02	0.81	0.55	0.55	0.00	0.55	3.52

determined by X-ray diffraction with a General Electric XRD-5 X-ray unit with Ni-filtered Cu-K α radiation, 35 kV, 24 mA, 1° divergence slit, 0.1° receiving slit, 4-s time constant, full-scale deflection of 4,000 cps, scan rate of 2° 2 θ per minute, and

chart speed of 1 inch (2.54 cm) per minute. Representative patterns are shown in Fig. 4. Results of these analyses, expressed as relative peak heights in centimeters in Table 1, provide a semiquantitative basis for comparing the mineralogy of silt from dif-

Table 2. Mineralogy of sand-size fraction from profundal sediments of 23 lakes.

LAKE	SAND (% clastic)	QUARTZ (%)	POTASH FELDSPAR (%)	PLAGIOCLASE FELDSPAR (%)	OPAQUE MINERALS (%)	OTHER (%)
BIG	2.1	46	18	31	4	1
IRON	3.1	29	15	45	1	9
TROUT	0.5	39	35	23		
WILSON	29.5	26	26	30	4	13
O'LEARY	4.4	58	14	26	<1	2
ARCO	1.3	47	23	19	9	1
LINWOOD	3.0	52	14	33	1	<1
SPECTACLE	5.1	53	16	28		
GLADSTONE	13.3	45	24	27	2	2
HAM	8.8	55	17	22	3	3
REEDS	2.9	46	11	41	<1	1
ST. OLAF	2.3	73	12	13	<1	2
CHRISTMAS	3.2	48	18	33	1	<1
LOTUS	2.0	51	12	31	1	1
LONG	6.2	56	5	32	1	5
LITTLE PINE	5.8	47	20	29	2	3
GROVE	0.9	34	30	33	2	2
FISH	4.3	38	19	42	2	0
SALLIE	3.5	47	18	32	1	2
COWDRY	1.0	46	23	23	7	1
SHETEK	1.4	50	26	23	<1	<1
DEAD COON	8.4	51	15	31	1	2
SALT	3.6	41	16	38	4	2

ferent lake sediments. In most cases two distinct feldspar peaks were recognized in the 3.2Å region—one at 3.20Å and one at 3.18Å. They probably represent orthoclase and plagioclase feldspar respectively.

Semiquantitative analyses for mineralogy of the clay-size fraction ($<2\ \mu$) were made on oriented glass-slide mounts by X-ray diffraction with Ni-filtered Cu-K α radiation, as for silt-size mineralogy. The clay minerals were oriented with c-axes perpendicular to the glass slide by pipetting a heavy slurry of clay in water onto the slide, allowing the clay minerals to settle out with a preferred orientation, and evaporating the water. X-ray diffractograms were made on air-dried clay mounts and on the same mounts after saturation with ethylene glycol to test for expansion of montmorillonite. No attempt was made to separate the 7Å kaolinite and chlorite peaks. Representative patterns are presented in Fig. 5. Quantification of peak intensity is difficult owing to the poor crystallinity of most samples. Some samples (e.g. Cedar Bog Pond, Fig. 5) appear almost entirely amorphous to X-radiation and produce no peaks at the in-

strument settings used. All samples were run at the same settings to produce comparable diffraction patterns. Relative intensities of mineral peaks in the clay-size fraction are given in Table 3. The degree of crystallinity, based on peak sharpness and intensity (Carroll 1970), is also given in relative terms.

Mineralogy of untreated samples was determined by X-ray diffraction of randomly oriented powdered samples mounted on glass slides: representative patterns are presented in Fig. 6. Semiquantitative results, expressed as peak height in centimeters for calcite, aragonite, dolomite, and quartz, are given in Table 1.

Results

The geographic patterns of organic, carbonate, and clastic materials in profundal sediments of Minnesota Lakes are shown in Fig. 7.

Organic matter—Organic matter as loss on ignition in profundal sediments is greatest, though not invariably high, in lakes of the northern and eastern forested regions

Fig. 4. Representative X-ray diffraction patterns of silt-size fractions of profundal sediments of six lakes. Q—Quartz; F—feldspar; D—dolomite; I—illite; Chl—chlorite.

(Fig. 7A). In all the lakes it ranges from a high of 60% (Cedar Bog, east-central) to a low of 6.4% (Dead Coon, southwestern prairie) (Table 1; Figs. 7A and 8). The mean is 24.6% (SD 12.2). This corresponds to a mean organic-C (total-C minus carbonate-C) of $12.1 \pm 6.3\%$, and a mean *ignition loss : organic-C* ratio of 2.13 ± 0.40 (Fig. 3). When organic content is measured as percent total N, most of these sediments have an *organic-C : N* ratio of less than 10 on a weight percent basis, with a mean ratio of 8.9 (Fig. 9).

Carbonate—Profundal sediments in the central lakes may contain up to 46% carbonate as calculated from loss on ignition (Fig. 7B). Mean sedimentary carbonate in

all the lakes is 16.3% (SD 14). Mean carbonate content in profundal sediments from the 27 carbonate lakes (those with greater than 10% sedimentary carbonate) is 26.3%. Sedimentary carbonate is restricted to those lakes with total cation concentrations in surface waters greater than 2 epm, which corresponds to Ca^{2+} concentrations >1 epm (20 ppm) (Fig. 10). Theoretical saturation values of 22–24 ppm Ca^{2+} can be expected from the solubility of CaCO_3 at 25°C and CO_2 concentrations in the range 0.033–0.044% (Hutchinson 1957).

The zero-carbonate baseline in Figs. 8 and 10, at about 3% carbonate, represents sediments with no carbonate and is presumably due to loss of water from clay lattices

Fig. 5. Representative X-ray diffraction patterns of clay-size fractions, solvated with ethylene glycol, of profundal sediments of eight lakes. Q—Quartz; F—feldspar; I—illite; M—montmorillonite; Chl—chlorite; K—kaolinite.

(Dean 1974). The value of about 3% applies only to those samples lacking carbonate. If we assume that those sediments with 550°–1,000°C ignition losses <5% contain no carbonate, then the mean carbonate content of sediments in all 46 lakes would be lowered slightly—to about 15%. The lakes with profundal sediments lacking carbonate include all group 1 lakes, and four of the more dilute group 2 lakes with relatively

high organic sediments, all located on highly leached outwash of the Anoka Sand Plain (Sims and Morey 1972).

Most of the carbonate is in the form of low-Mg calcite, although minor amounts of high-Mg calcite, dolomite, and aragonite were detected by X-ray diffraction (Table 1, Figs. 6 and 11). The dolomite occurs in lake sediments in southern and western Minnesota (Table 1). In addition, X-ray

Table 3. Mineralogy of clay-size ($<2 \mu$) fraction from profundal sediments of all 46 lakes.

LAKE	Degree of crystall- inity *	ILLITE (10 Å)		KAOLINITE + CHLORITE (7 Å)		QUARTZ (3.35 Å)		FELDSPAR (3.2 Å)		12 - 14 Å peak	
		untr.†	gly.‡	untr.	gly.	untr.	gly.	untr.	gly.	untr.	gly.
Big Iron	poor	tr	tr	tr	tr	++	++	+	+	A	A
Mountain Trout	poor	0	0	0	0	tr	0	0	0	A	A
Clearwater	fair	+	+	+	+	+	+	tr	tr	A	A
Wilson	poor	0	0	tr	tr	tr	tr	0	0	P	A
Sand Point	poor	tr	0	tr	0	tr	tr	tr	tr	P	A
Kimball	poor	tr	0	0	0	tr	tr	tr	tr	P	A
Crane	fair	tr	tr	+	+	+	+	tr	tr	P	A
O'Leary	good	+	+	+	+	+	+	+	+	P ⁺	17 Å
Josephine	fair	tr	tr	+	+	+	+	+	+	P	P
Arco	fair	tr	tr	+	+	+	+	+	+	P	A
Deming	fair	tr	tr	+	+	+	+	+	+	P	A
Farguar	poor	+	+	+	+	+	+	tr	tr	A	A
George	poor	tr	tr	0	0	tr	tr	tr	tr	P	A
Linwood	poor	0	0	0	0	tr	tr	0	0	P	A
Cedar Bog	poor	0	0	0	0	0	0	0	0	0	0
Spectacle	poor	tr	tr	tr	0	+	+	tr	tr	P	A
Gladstone	poor	0	0	tr	0	tr	tr	0	0	P	A
Ham	poor	0	0	0	0	0	tr	0	0	A	A
Green	poor	0	0	tr	tr	tr	tr	0	0	P	17 Å
Reeds	good	+	+	+	+	+	+	tr	tr	P	P
Francis	good	+	+	+	+	+	+	+	+	P	17 Å
Nokay	poor	tr	0	tr	tr	tr	tr	0	0	A	A
Beaver	excel.	++	+	+	+	+	+	tr	tr	P ⁺	P
Moose	good	+	0	+	+	+	+	tr	tr	P	P
Ball Club	poor	0	0	0	0	+	tr	0	0	A	A
St. Olaf	good	+	+	+	+	+	+	+	tr	P	P
Christmas	fair	+	tr	+	tr	+	+	+	tr	P	A
Lotus	fair	+	+	+	+	+	+	+	+	P ⁺	17 Å
Clear	good	+	tr	+	tr	+	+	+	tr	P	A
Long	fair	+	+	+	+	+	+	+	+	P	P
Elk	poor	0	0	0	0	tr	tr	0	0	A	A
Itasca	poor	0	0	0	0	tr	tr	0	0	A	A
Little Pine	fair	tr	tr	0	tr	+	+	+	tr	P	P
Grove	fair	+	tr	+	0	+	+	tr	0	A	A
Fish	fair	+	tr	+	tr	+	+	0	0	A	A
Sallie	fair	+	+	+	+	+	+	+	tr	P	P
Cowdry	poor	tr	0	tr	0	+	tr	tr	0	A	A
Maple	poor	+	+	+	+	+	+	+	+	A	A
Mina	poor	tr	tr	tr	tr	+	+	tr	0	A	A
Shetek	excel.	+	+	++	+	+	+	tr	tr	A	P
Big Kandiyo	excel.	+	+	+	+	+	+	tr	tr	P	P
Elk (Grant Co.)	excel.	++	++	++	++	+	+	tr	tr	P	P
Dead Coon	excel.	++	++	++	++	+	+	+	+	P	P
Salt	excel.	+	+	+	+	+	+	+	+	P	P

* Relative degree of crystallinity for these samples only.

† Untreated sample, dry on slide.

‡ Sample after treatment with ethylene glycol.

§ Peak intensities: ++ = very well defined peak; + = well defined peak; tr = trace or peak very poorly defined; 0 = no peak above background; P = well defined 12-14 Å; A = only low angle background slope at 12-14 Å.

Fig. 6. Representative X-ray diffraction patterns of untreated samples of profundal sediments of eight lakes. Q—Quartz; F—feldspar; C—calcite; D—dolomite; A—aragonite.

diffraction patterns of sediments from several of these lakes (e.g. Francis, Beaver, Fish, Cowdry, Sallic, Dead Coon, and Salt) exhibit shoulders at about $29.7^\circ 2\theta$ on the main calcite peak (29.4°), suggesting that some high-Mg calcite may be present. Some dolomite in group 3 and high-carbonate group 2 lakes may therefore be a diagenetic product of high-Mg calcite, as suggested by Müller et al. (1972), or it may be detrital, derived from Paleozoic dolostones. Müller et al. (1972) found primary aragonite only in lakes with a $Mg : Ca$ ratio in the water of >12 . The small amounts of aragonite in Minnesota lakes are probably derived from mollusc shells because the $Mg : Ca$ in waters from these lakes is generally <2 .

The lake with the most instructive carbonate mineralogy is Elk Lake, Grant Co. (No. 44). Its sediments contain more do-

lomite than any of the other lakes (Table 1, Fig. 6). Although Elk is not the most saline of the lakes studied, it is the only one with a high $Mg : Ca$ ratio (7.7) in the water. X-ray diffraction patterns of its sediments commonly show two distinct calcite peaks, one at $29.4^\circ 2\theta$ (low-Mg calcite) and one at about 29.7° (high-Mg calcite) (Fig. 11). Figure 12 shows that Elk Lake sediments (point marked 44) contain much more total Mg than any other lake sediments, but it is largely in dolomite which is not soluble in 1.0 N ammonium acetate at pH 5.2 (Wangersky and Joensuu 1967). According to Müller et al. (1972), diagenetic dolomite is found in lakes with high-Mg calcite as a primary carbonate mineral and a $Mg : Ca$ ratio in the water of >7 .

For most sediments high in carbonate (i.e. $>10\%$), about half of the Mg is soluble in acid ammonium acetate, presumably in

Fig. 7. Maps showing distribution of percent organic matter (A), percent carbonate (B), and percent clastic material (C) in profundal sediments of all 46 lakes. All values calculated from loss on ignition at 550° and 1,000°C (*see text*). Note that minimum values for percent clastic and percent organic matter are 35% and 6%.

solid solution for Ca in the calcite structure (Fig. 12). The total soluble Mg, expressed as percent dry weight, increases with increasing $Mg:Ca$ ratio of the water (Fig. 13). In those lakes which precipitate the most calcium carbonate, i.e. the group 3 and

high-carbonate group 2 lakes, the proportion of Mg coprecipitated with calcite, expressed as percent calcium carbonate, increases with increasing $Mg:Ca$ ratio in the lake water (Fig. 14). In other words, in lakes of western and southwestern Minnesota, the $Mg:$

Fig. 8. Trilinear plot of carbonate, organic, and clastic fractions in profundal sediments of all 46 lakes. The 0-carbonate refers to ignition loss between 550–1,000°C of sediments containing no carbonate minerals. Numbers adjacent to points refer to lakes (Fig. 2) specifically discussed in text.

Ca ratio of sedimentary calcium carbonate increases with increasing *Mg* : *Ca* in the water. Figures 13 and 14 also show that for low-carbonate group 2 lakes (sedimentary carbonate <30%; *Mg* : *Ca* ratio in water generally <1.0), there is no apparent rela-

Fig. 9. Scatter plot of weight percent nitrogen and weight percent organic carbon in profundal sediments from all 46 lakes.

Fig. 10. Semilogarithmic plot of carbonate by ignition loss in profundal sediments and total cations in surface waters from all 46 lakes.

tionship between soluble *Mg* and the *Mg* : *Ca* ratio of the water. However, some of these lakes contain considerable soluble *Mg*. If this *Mg* is present in solid solution for *Ca* in calcite, then these lakes must have sedimentary carbonates with a higher *Mg* : *Ca* ratio than those of group 3 and high-carbonate group 2. This is unlikely because the *Mg* : *Ca* ratios in these waters are much lower than in the western and southwestern lakes. Presumably the soluble *Mg* of the low-carbonate group 2 lakes is adsorbed on

Fig. 11. X-ray diffraction patterns from slow ($0.4^\circ 2\theta \text{ min}^{-1}$) scans between 29° and $32^\circ 2\theta$ of sediment samples from two depths in short core from Elk Lake, Grant Co. HMC, LMC, and D refer to peaks for high-Mg calcite, low-Mg calcite, and dolomite.

Fig. 12. Scatter plot of Mg soluble in 1.0 N ammonium acetate (pH 5.2) and total Mg in profundal sediments from all 46 lakes.

clays, because soluble Mg is not high in lakes with high sedimentary organic matter (Table 1).

Clastic material—The clastic fraction is defined as the noncarbonate, inorganic mineral material in a sample. Lake sediments with the largest clastic fractions occur in northeastern and southwestern Minnesota (Fig. 7C). Lake sediments from central Minnesota contain relatively low clastic fractions, as a result of dilution by carbonate and organic matter (Figs. 7A and 7B). The range of clastic material is from 35–81%, with a mean of 58.9% (SD 13.7).

After destruction of the organic and carbonate components, the clastic fraction was further subdivided into sand, silt, and clay

Fig. 13. Scatter plot of sedimentary Mg soluble in 1.0 N ammonium acetate (pH 5.2), expressed as percent dry weight, and Mg:Ca ratio in surface waters from all 46 lakes. (Source of data—Gorham et al. in prep.)

Fig. 14. Scatter plot of sedimentary Mg soluble in 1.0 N ammonium acetate (pH 5.2), expressed as percent CaCO_3 , and Mg:Ca ratio in surface waters from all 46 lakes.

fractions. Significant proportions of sand occur in very few profundal sediments and exceed 10% dry weight only in three northern lakes. Silt and clay are present in about equal proportions in most samples. Mean values and standard deviations for sand, silt, and clay, expressed as percent dry weight, are 2.9% (± 4.2), 23.4% (± 8.1), and

Fig. 15. Map showing ratio of 4.26 Å quartz peak height to 3.18 Å feldspar peak height from X-ray diffraction traces of silt-size materials in profundal sediments from 38 lakes.

22.3% (± 8.4). Expressed as percent clastic material, the respective means are 7.3%, 47.6%, and 46.8%.

The sand-size fraction (Table 2) is characterized by a predominance of quartz (46.9%), with lesser amounts of plagioclase (29.8%) and orthoclase (18.6%). No regional variations exist in the proportions of these minerals.

The silt-size fraction is also dominated by quartz and, to a lesser extent, plagioclase and orthoclase feldspar (Table 1 and Fig. 4). Most X-ray diffractograms exhibit trace or moderate low angle peaks for clay minerals as well; most common are illite (10Å) and kaolinite + chlorite (7Å). Figure 15 illustrates a marked northeast to southwest change in the mineralogy of the silt-size fraction. On the basis of peak-height ratios, the abundance of quartz relative to feldspar is greatest in group 3 sediments of the arid southwest, and lowest in group 1 sediments of the humid northeast.

Most X-ray diffractograms of clay-size material are characterized by a 3.2Å feldspar peak, a 3.35Å quartz peak, 3.5Å and 7Å kaolinite + chlorite peaks, and a 10Å illite peak (Table 3; Fig. 5). For some samples it was possible to distinguish between the 3.57Å kaolinite peak and the 3.52Å chlorite peak, but for most the 3.5Å region yielded a broad single peak; no attempt was made to separate the two peaks by heating. In some samples, the presence of montmorillonite is indicated by the development of a 17Å peak from 12–14Å mixed-layer clays after treatment with ethylene glycol.

The values in Table 3 show no well defined regional differences in clay mineralogy, although there is an apparent increase in all clay minerals in sediments from group 3 lakes in southwestern Minnesota. However, values for percent clay in Table 1 show no regional variation in the amount of clay-size material. Therefore, the apparent increase in clay minerals in the southwestern lake sediments is due not to an increase in actual clay mineral content, but to a higher degree of crystallinity of the clay-size fraction, resulting in stronger X-ray diffraction peaks. Most of the clay-size

material in sediments from group 1 and group 2 lakes in northeastern and central Minnesota appears to be X-ray amorphous. Ross and Kerr (1931) have suggested that the name allophane be applied to all X-ray amorphous clay minerals, regardless of composition. Most of this allophane is probably opaline silica derived from diatom remains. However, the fact that the clay-crystallinity gradient corresponds to the regional climatic gradient of the state suggests that climate has caused a greater alteration of clays in the northeast than in the southwest.

Discussion

Lake grouping—Figures 2 and 8 show that the geographic groups of the lakes defined initially on the basis of water chemistry (Gorham et al. in prep.) can also be differentiated on the basis of profundal sediment chemistry and mineralogy. The sediments of dilute group 1 lakes on non-calcareous drift in northeastern Minnesota are characterized by an absence of carbonate minerals. Profundal sediments of the deep (>10 m) group 1 lakes contain much less organic material (16–21% loss on ignition at 550°C) than those of the shallow group 1 lakes (38–47% loss on ignition). Three very small outlying group 1 lakes in Hubbard Co. (Fig. 2; Table 1) are also rich in organic matter despite depths of 7.6, 11.6, and 17.0 m. All three have unusually high ratios of depth to area and are sheltered from wind mixing; as a result, they do not undergo normal mixing and may remain stratified for several years.

Group 2 lakes, of intermediate ionic concentration, have the most diverse sediments, due mainly to variation in the amount of precipitated calcium carbonate. They are distinguished from group 1 lakes by the presence of carbonate minerals in profundal sediments, except in four of the most dilute group 2 lakes with highly organic sediments (>39% loss on ignition at 550°C). Waters of these high-organic group 2 lakes commonly have Ca^{2+} concentrations greater than 1.0 cpm (20 ppm), a level sufficient

to precipitate CaCO_3 in other dilute, low-organic group 2 lakes. For example, George, Cedar Bog, Spocket, Ham, and Green are five relatively dilute group 2 lakes on the Anoka Sand Plain with very similar water chemistries (Fig. 2). George, Cedar Bog, and Spocket contain >39% organic matter and no carbonate, whereas Ham and Green contain 28 and 21% organic matter and 37 and 30% carbonate. Decomposition of organic matter in the profundal zone of the high-organic group 2 lakes apparently generates enough CO_2 to redissolve any calcium carbonate that may have formed. Two other dilute group 2 lakes, Nokay and Linwood (Fig. 2), with slightly less organic matter (38% and 35%), contain 16% and 10% carbonate in their profundal sediments. These two lakes are transitional between the carbonate-free, high-organic group 2 lakes and those intermediate in both components.

Sediments of intermediate group 2 lakes, containing 10–30% organic matter and 10–30% carbonate, also tend to be intermediate in dissolved ionic concentration within group 2 lakes. With two exceptions, Moose and Elk (Clearwater Co.) (Fig. 2), intermediate group 2 lakes are in south-central and southwestern Minnesota, where evaporation exceeds precipitation, so that dissolved salts become concentrated particularly during summer. Calcium and alkalinity concentrations in these lakes are generally greater than 1.0 and 2.0 epm and are greater than 2.0 and 3.0 epm during summer (Gorham et al. in prep.; Bright 1968). Elk Lake (Clearwater Co., No. 33 in Fig. 2) is in west-central Minnesota, an area dominated by high-carbonate group 2 lakes, although it classifies with the south-central and southeastern intermediate group 2 lakes. This lake has a maximum depth of nearly 30 m; it is the second deepest carbonate lake studied. The profundal sediments at this depth have a carbonate content of only 19%, which would classify Elk as an intermediate group 2 lake. However, sediments from five other stations in Elk Lake, ranging from 11–24 m in depth, contain 24–70% CaCO_3 , with CaCO_3 con-

tent roughly proportional to depth. Most sediments in this lake would place it as a high-carbonate (>30%) group 2 lake, and indeed the surface waters commonly have Ca^{2+} and alkalinity concentrations of about 2 and 3 epm (Megard 1968; Gorham et al. in prep.). Elk Lake, like nearby Itasca and Long Lakes (Megard 1968), undoubtedly precipitates large quantities of calcium carbonate during summer, but dissolution of carbonate in the deeper waters of the lake has greatly reduced the amount of carbonate being incorporated into deeper profundal sediments. Moose Lake, the other geographic exception in the intermediate group 2 category, has 28.3% carbonate and therefore also comes close to being classified with high-carbonate group 2 lakes.

The high carbonate (>30%) group 2 lakes are all in west-central Minnesota (Fig. 2). They generally contain less than 30% organic matter and are also characterized by inactive outlets and concentrations of dissolved Ca^{2+} and alkalinity in excess of 2.0 and 3.0 epm. Detailed studies by Megard (1968) on the precipitation of CaCO_3 in two typical high-carbonate group 2 lakes (Long and Itasca) showed that their surface waters are supersaturated with respect to calcite during summer and precipitate large quantities of calcite. Megard also found a close correlation between dissolved Ca^{2+} and alkalinity depletion (presumably as precipitated CaCO_3) and photosynthetic carbon assimilation in six typical high-carbonate group 2 lakes. For the six lakes, an average of 1 mole of Ca^{2+} was removed for every 4 moles of carbon fixed by phytoplankton photosynthesis. Because of the high rate of calcite precipitation, the high-carbonate group 2 lakes generally have ratios of dissolved $\text{Mg} : \text{Ca} > 1.0$ (Figs. 13 and 14).

Group 3 lakes occupy shallow depressions in thin, gypsum-bearing glacial drift in the western and southwestern prairie regions. Dissolved salts in the waters of these arid regions are more concentrated than in those of group 1 or group 2 lakes; they are rich in dissolved magnesium, calcium, sulfate, and bicarbonate. Profundal sediments from

group 3 lakes resemble those of intermediate group 2 lakes (Fig. 8), but are distinguished from them by the excellent degree of crystallinity of the clay-size fraction (Table 3), by their low organic content (<16% loss on ignition at 550°C: Fig. 7A), and by a strong predominance of the clastic fraction (62–75%: Fig. 7C). The high Mg^{2+} concentration in group 3 lake waters is reflected by relatively high Mg concentrations in the sediments, presumably due to coprecipitation with calcite (Figs. 12–14). The percentage of carbonate in group 3 lake sediments (17–26%) is less than in high-carbonate group 2 lake sediments (Figs. 8 and 10), despite higher concentrations of Ca^{2+} and alkalinity in the waters of group 3 lakes. This anomaly presumably reflects dilution by high silting in these shallow prairie lakes.

A rare fourth group of lakes, characterized by very saline waters (conductivity >7,000 $\mu mhos\ cm^{-1}$ at 25°C) dominated by sodium and sulfate ions, occurs along the boundary between Minnesota and the Dakotas. Group 4 lakes are represented in this study by Salt Lake. The profundal sediment of Salt Lake is similar to those of group 3 lakes and has been included with them in this discussion. If we treat profundal sediments as a three-component system (carbonate, organic, and clastic), two trends are followed by Minnesota lakes from an inorganic clastic base (Fig. 8). The first trend, illustrated by group 1 lakes and most high-organic group 2 lakes, involves addition of organic matter. The second trend, illustrated by the remaining group 2 lakes and those of group 3, involves addition of carbonate to a predominantly inorganic clastic fraction plus 10–20% organic matter. As mentioned above, two high-organic group 2 lakes (Nokay and Linwood) appear to be transitional.

To test the grouping of lakes, based primarily on water chemistry but subdivided in relation to major sediment components, we ran a Q-mode factor analysis on the 25 organic-, carbonate-, and clastic-related variables listed in Table 1, plus lake area and maximum depth. By this method, similarities among lakes are examined based

on the measured variables and coefficients computed for each lake. These coefficients, called factor loadings or factor scores, are measures of the degree of similarity among lakes (e.g. Harman 1967; Morrison 1967; Sneath and Sokal 1973). So that we could compare percentages, X-ray diffraction peak heights, areas, and depths in the same analysis, all values for each variable were divided by the maximum value of that variable occurring in all 46 lakes (i.e. $x_{i,j}/x_{jmax}$; $i = 1$ to 46 lakes; $j = 1$ to 27 variables). The input matrix, therefore, consisted of 27 converted variables, each with a range of 0 to 1.0, and 46 observations (lakes). The actual analysis was done using a FORTRAN IV program developed by Ondrick and Srivastava (1970). A total of 10 factors was extracted, although 83% of the variance in the data was accounted for about equally by the first three factors, after a varimax rotation (Table 4).

Plots of factor loadings for the first three rotated factors are shown in Fig. 16. Lakes with the highest loadings for factor 1 are all high-carbonate group 2 lakes, whereas lakes with low loadings for this factor are those with little or no sedimentary carbonate. Lakes with high loadings for factor 2 are the high-organic group 1 and group 2 lakes, and lakes with high loadings for factor 3 are those lakes of all groups with high proportions of clastic material.

Figure 16A illustrates the degree to which the three lake groups can be segregated by the factor analysis. Group 3 lakes are segregated by their low loadings for factor 2 (organic fraction) coupled with intermediate loadings for factor 1 (carbonate fraction). Group 2 lakes exhibit a wide range in loadings for both factors, and a few high-organic group 2 lakes with no sedimentary carbonate overlap the high-organic group 1 lakes, which are clearly segregated from the group 3 lakes.

Figure 16B shows the same organic- and carbonate-enrichment trends as Fig. 8. Lakes of all groups with high loadings for factor 3 (clastic fraction) form a base from which we can see diverging trends toward high and low loadings for factor 1 (car-

Table 4. Loadings for first three rotated factors of *Q*-mode factor analysis, 27 variables, 46 lakes.

LAKE NO.	LAKE NAME	FACTOR 1	FACTOR 2	FACTOR 3
1	BIG	0.24	0.63	0.54
2	IRON	0.24	0.90	0.21
3	MOUNTAIN	0.21	0.39	0.59
4	TROUT	0.19	0.55	0.47
5	CLEARWATER	0.20	0.32	0.65
6	WILSON	0.19	0.44	0.59
7	SAND POINT	0.23	0.42	0.43
8	KIMBALL	0.28	0.86	0.35
9	CRANE	0.29	0.39	0.66
10	O'LEARY	0.22	0.42	0.81
11	JOSEPHINE	0.30	0.67	0.61
12	ARCO	0.28	0.68	0.64
13	DEMING	0.28	0.73	0.57
14	FARQUAR	0.20	0.40	0.82
15	GEORGE	0.24	0.80	0.45
16	LINWOOD	0.48	0.74	0.24
17	CEDAR BOG	0.28	0.93	0.10
18	SPECTACLE	0.26	0.86	0.32
19	GLADSTONE	0.23	0.81	0.23
20	HAM	0.86	0.41	0.08
21	GREEN	0.81	0.38	0.24
22	REEDS	0.33	0.46	0.74
23	FRANCIS	0.44	0.44	0.73
24	NOKAY	0.53	0.73	0.28
25	BEAVER	0.58	0.27	0.68
26	MOOSE	0.72	0.25	0.57
27	BALL CLUB	0.28	0.61	0.24
28	ST. OLAF	0.48	0.28	0.76
29	CHRISTMAS	0.34	0.33	0.80
30	LOTUS	0.48	0.29	0.75
31	CLEAR	0.45	0.37	0.78
32	LONG	0.78	0.11	0.51
33	ELK	0.58	0.57	0.24
34	ITASCA	0.88	0.38	0.10
35	LITTLE PINE	0.87	0.23	0.29
36	GROVE	0.80	0.30	0.48
37	FISH	0.58	0.31	0.62
38	SALLIE	0.84	0.22	0.46
39	COWDRY	0.79	0.31	0.40
40	MAPLE	0.81	0.20	0.49
41	MINA	0.87	0.24	0.33
42	SHETCK	0.63	0.22	0.59
43	BIG KANDIYOH	0.59	0.20	0.64
44	ELK	0.57	0.16	0.66
45	DEAD COON	0.41	0.13	0.86
46	SALT	0.48	0.23	0.82
SUM OF SQUARES		12.65	11.57	14.11
% VARIANCE EXPL.		27.50	25.16	30.67

bonate fraction). The lakes with high loadings for factor 1 are the western high-carbonate lakes of group 2, those with low loadings are the group 1 lakes together with a few group 2 lakes lacking sedimentary carbonate. Three lakes (Linwood, Nokay, and Elk) appear intermediate with regard to these trends (cf. Fig. 8). Linwood and Nokay (No. 16 and 24) form a high-organic, intermediate carbonate subgroup of the group 2 lakes, and Elk Lake (Clearwater Co., No. 33) should probably be placed

MINNESOTA LAKE SEDIMENTS Q-MODE FACTOR ANALYSIS

27 Variables
46 Lakes

Fig. 16. Scatter plots of rotated factor loadings, *Q*-mode factor analysis of 27 variables in profundal sediments from all 46 lakes. A—Rotated factor 1 vs. rotated factor 2; B—rotated factor 1 vs. rotated factor 3. Numbers adjacent to points refer to lake numbers in Fig. 2 and Table 4.

with the high-carbonate group 2 lakes, for reasons discussed above.

Associations of variables—Relations among the 27 variables listed in Table 1 are expressed as a correlation matrix in Table 5. To simplify analysis of these relationships, the correlation matrix was used as input for *R*-mode factor analysis using the FORTRAN IV program developed by Ondrick and Srivastava (1970). Results of the *R*-mode factor analysis from the Ondrick and Srivastava program were checked using another factor analysis program (BMD X72)

(Dixon 1970). Results of both factor analyses were almost identical. Loadings for 10 factors from the Ondrick and Srivastava program, after varimax rotation, are given in Table 6. Each of the first two factors accounts for over 20% of the variance in the data. The highest loadings for factor 1 are for organic-related variables: loss on ignition at 550°C, total carbon, organic carbon, hydrogen, and nitrogen. The highest loadings for factor 2 are for carbonate-related variables: calcium carbonate calculated from loss on ignition between 550° and 1,000°C, soluble Ca, X-ray calcite peak height, and, to a lesser extent, soluble Mg.

The clastic variables contribute variance to both factor 1 (organic) and factor 2 (carbonate), but are inversely related to the principal variables in each factor. This reflects the negative correlation between percent clastic material and both percent carbonate and percent organic matter (Table 5) and also the two trends (Fig. 8, Fig. 16B) toward either organic or carbonate enrichment of a clastic base. Tables 5 and 6 also show that the inverse relationship between carbonate and clastic material is closer than the inverse relationship between organic matter and clastic material.

The clay-mineral variables of the clastic fraction, illite, chlorite + kaolinite, and clay-size quartz, are negatively correlated with organic matter but not with carbonate. On the other hand, the total percentage of clastic material, and its major component variables, percent silt and percent clay, do show a negative correlation with the carbonate fraction. We speculate that this anomaly can be explained by considering the geographic distribution of lakes exhibiting these trends. Most high-organic lakes (groups 1 and 2) are located in north-central and northeastern Minnesota (Fig. 2). In these forested regions, a relatively thick, moist, acid humus has developed, so that most of the clastic material transported to the lakes is highly weathered and low in total volume. Dilution of this clastic base with organic matter, both accumulating at a relatively slow rate, will result in an inverse relationship between the organic frac-

tion and the highly altered clays of the clastic fraction. The carbonate-rich lakes, on the other hand, occupy shallow basins in the arid western and southwestern prairie regions. There, the soil cover is not well developed, and erosion is therefore likely to be more rapid. As a result these lakes receive a greater volume of clastic material which, because of increased rate of erosion and low available moisture, is probably less altered by decomposition. Dilution of this clastic base with CaCO_3 at a relatively rapid rate results in an inverse relationship between carbonate and clastic material, most of which is relatively unaltered silt and clay.

The difference in degree of decomposition of clastic materials in the northeastern and central lakes, relative to the southwestern lakes, is suggested by the degree of crystallinity of the clay minerals. The crystallinity index given in Table 3 is much higher for clays from the southwestern group 3 lakes than for the northeastern and central lake sediments. This increase in degree of crystallinity is not related to the amount of clay in the sediments or to the total amount of clastic material in the sediments. Gorham et al. (in prep.), in their study of water chemistry of Minnesota lakes, found that the boundary between group 2 and group 3 lakes is not gradual, as might be expected by increased aridity alone. They conclude that the sharp boundary is the result of differences in glacial drift composition superimposed on the regional climatic gradient. The sharp increase in clay crystallinity in group 3 lake sediments is probably related, therefore, to drift composition as well as to climate.

Factor 3, accounting for 10% of the variance in the data, is essentially a clay mineralogy factor. The fact that no other clastic variables have high loadings in this factor suggests that the mineralogy of the clay-size material is independent of the total amount of clastic material being introduced into the lakes.

One might expect pronounced regional differences in the clay mineralogy of the profundal sediments, because bedrock types

Table 5. Correlation coefficient matrix for 27 variables, 46 lakes. 99% significance level = ± 0.37 for $N = 46$; correlations significant at the 99% level are underlined.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
2	0.23														
3	-0.25	-0.20													
4	-0.20	-0.26	<u>0.91</u>												
5	-0.21	-0.22	<u>0.97</u>	<u>0.96</u>											
6	-0.19	-0.15	<u>0.96</u>	<u>0.89</u>	<u>0.95</u>										
7	-0.23	-0.26	<u>0.96</u>	<u>0.94</u>	<u>0.96</u>	<u>0.92</u>									
8	0.05	-0.06	<u>-0.46</u>	-0.14	-0.40	-0.50	-0.34								
9	0.06	-0.09	<u>-0.44</u>	-0.13	-0.38	-0.48	-0.32	<u>0.99</u>							
10	0.04	-0.06	<u>-0.41</u>	-0.10	-0.36	-0.45	-0.30	<u>0.99</u>	<u>0.99</u>						
11	0.09	-0.14	<u>-0.44</u>	-0.13	-0.37	-0.45	-0.33	<u>0.93</u>	<u>0.94</u>	<u>0.91</u>					
12	0.03	-0.04	-0.24	-0.12	-0.21	-0.23	-0.19	<u>0.37</u>	<u>0.34</u>	<u>0.33</u>	<u>0.45</u>				
13	0.01	-0.23	-0.42	-0.35	-0.38	-0.37	-0.34	0.20	0.19	0.12	0.33	0.34			
14	0.13	-0.10	-0.64	-0.52	-0.57	-0.59	-0.56	0.34	0.30	0.26	0.46	0.49	<u>0.74</u>		
15	0.12	-0.12	-0.57	-0.32	-0.49	-0.56	-0.44	<u>0.72</u>	<u>0.67</u>	<u>0.65</u>	<u>0.70</u>	<u>0.48</u>	<u>0.58</u>	<u>0.76</u>	
16	0.16	0.24	-0.40	-0.66	-0.45	-0.34	-0.50	-0.63	-0.64	-0.66	-0.57	-0.16	0.15	0.20	-0.25
17	0.15	0.20	-0.18	-0.22	-0.17	-0.07	-0.22	-0.14	-0.14	-0.14	-0.10	-0.01	0.03	0.12	-0.05
18	0.15	0.33	-0.12	-0.30	-0.15	-0.07	-0.17	-0.49	-0.50	-0.51	-0.41	-0.05	0.17	0.11	-0.23
19	0.08	-0.01	-0.24	-0.44	-0.28	-0.23	-0.30	-0.48	-0.47	-0.50	-0.45	-0.19	0.07	0.12	-0.21
20	-0.10	-0.32	-0.35	-0.25	-0.31	-0.39	-0.25	0.29	0.26	0.23	0.23	0.19	<u>0.44</u>	<u>0.56</u>	<u>0.54</u>
21	0.01	-0.15	0.07	0.24	0.11	0.02	0.16	<u>0.43</u>	<u>0.44</u>	<u>0.42</u>	0.36	0.15	-0.14	0.02	0.22
22	-0.12	-0.12	0.09	0.25	0.15	0.09	0.15	0.27	0.27	0.27	0.20	0.03	-0.10	0.03	0.14
23	-0.03	-0.10	-0.58	-0.51	-0.53	-0.58	-0.51	0.21	0.18	0.14	0.22	0.20	<u>0.58</u>	<u>0.74</u>	<u>0.53</u>
24	0.07	0.00	-0.58	-0.55	-0.54	-0.61	-0.54	0.09	0.05	0.02	0.13	0.24	<u>0.53</u>	<u>0.68</u>	<u>0.42</u>
25	0.25	0.09	-0.46	-0.41	-0.41	-0.47	-0.42	0.11	0.07	0.06	0.09	0.19	0.30	<u>0.47</u>	<u>0.42</u>
26	-0.08	-0.05	-0.23	-0.21	-0.23	-0.28	-0.21	0.11	0.08	0.07	0.02	-0.05	0.26	0.33	0.30
27	-0.13	-0.25	-0.24	-0.33	-0.22	-0.24	-0.22	-0.30	-0.30	-0.33	-0.21	-0.08	0.28	0.29	-0.02

Table 5. Continued.

	16	17	18	19	20	21	22	23	24	25
17	0.29									
18	<u>0.62</u>	0.29								
19	<u>0.70</u>	0.07	0.35							
20	-0.01	-0.29	-0.19	0.06						
21	-0.52	-0.27	-0.60	-0.26	0.40					
22	-0.40	-0.17	-0.52	-0.14	0.38	0.85				
23	0.29	-0.03	0.21	0.34	<u>0.62</u>	-0.12	-0.02	0.83		
24	<u>0.42</u>	0.06	0.26	<u>0.38</u>	<u>0.61</u>	-0.10	-0.01	<u>0.67</u>	<u>0.69</u>	
25	0.29	-0.12	0.15	0.35	<u>0.39</u>	-0.02	-0.00	<u>0.54</u>	<u>0.55</u>	0.65
26	0.10	-0.07	0.07	0.18	<u>0.44</u>	0.18	0.25	<u>0.49</u>	<u>0.57</u>	0.34
27	<u>0.53</u>	0.04	0.26	0.31	<u>0.40</u>	-0.21	-0.05			

and associated glacial drift vary considerably across the state (Fig. 1). Precambrian metamorphic and igneous rocks predominate in the northeastern and central parts, Cretaceous shale dominates the lithology in the west, and Paleozoic sandstone and dolomite are dominant in the southeast. The absence of marked regional differences suggests that weathering, as controlled by temperature, available moisture, soil pH, etc., has a greater influence on clay mineralogy than bedrock type. Pluth et al. (1970) also report a general similarity in mineralogy of the $<5\text{-}\mu$ size fraction of 16 soils throughout Minnesota.

However, a hint of bedrock control on clay mineralogy is suggested by factor 10, which accounts for 10.3% of the variance in the data. Factors 10 and 2 suggest three contributions of Mg to lake sediments. In factor 2, soluble Mg is related to the carbonate variables, indicating that most Mg soluble in 1.0 N ammonium acetate is apparently derived from Mg coprecipitated with calcite. In factor 10, however, the highest loadings are for total Mg and X-ray dolomite. This association indicates that in the western and southwestern lakes, which are high in sedimentary Mg, much of it is tied up as dolomite which is not dissolved by ammonium acetate. A third contribution of Mg indicated by factor 10 is from clay minerals. Table 5 shows that total Mg has high positive correlations with both illite and chlorite. Illite is derived mainly from the alteration of potassium feldspar and muscovite, both abundant minerals in acid igneous and metamorphic rocks. Most chlorite is derived from pre-existing metamorphic rocks (e.g. chlorite schist) and some sedimentary rocks. It is a common detrital mineral especially in areas where metamorphic rocks have been glaciated (Grim 1953). Consequently chlorite would be expected to be a major clay mineral in Minnesota soils and lake sediments and a major source of Mg in the sediments. Data from Pluth et al. (1970) show that chlorite is common in soils throughout Minnesota, but is most abundant in the northeastern and east-central parts underlain by meta-

Table 6. Loadings for 10 rotated factors of R-mode factor analysis, 27 variables, 46 lakes.

VARIABLE	THE ROTATED MATRIX OF FACTOR LOADINGS									
	FACTOR 1	FACTOR 2	FACTOR 3	FACTOR 4	FACTOR 5	FACTOR 6	FACTOR 7	FACTOR 8	FACTOR 9	FACTOR 10
AREA	0.16	-0.01	-0.02	-0.03	-0.97	0.00	-0.06	0.09	0.06	-0.00
MAXIMUM DEPTH	0.24	0.09	-0.01	-0.07	-0.10	0.02	-0.07	0.88	0.16	0.22
L.O.I. 500° C	-0.91	0.23	0.16	0.03	0.10	-0.05	0.07	-0.05	0.05	0.22
TOTAL CARBON	-0.95	-0.09	0.11	0.11	0.04	-0.01	0.06	-0.08	0.05	0.15
ORGANIC CARBON	-0.95	0.17	0.12	0.06	0.04	-0.03	0.04	0.06	0.03	0.17
HYDROGEN	-0.90	0.27	0.21	0.02	0.04	-0.04	-0.04	-0.02	0.08	0.16
NITROGEN	-0.94	0.12	0.13	0.07	0.07	-0.05	0.09	-0.08	0.02	0.13
CARBONATE	0.25	-0.94	-0.06	0.16	0.00	0.07	0.05	-0.01	0.08	-0.10
TOTAL Ca	0.24	-0.94	-0.02	0.16	-0.02	0.05	0.04	-0.05	0.07	-0.07
SOLUBLE Ca	0.22	-0.95	-0.03	0.15	0.01	0.05	0.04	-0.02	0.08	-0.02
CALCITE (X-RAY)	0.21	-0.88	0.01	0.08	-0.06	0.18	0.00	-0.09	0.01	-0.21
ARGONITE (X-RAY)	0.08	-0.27	-0.01	0.03	0.00	0.90	-0.01	0.02	0.01	-0.25
DOLOMITE (X-RAY)	0.19	-0.09	-0.12	-0.14	0.02	0.08	-0.03	-0.11	-0.03	-0.87
TOTAL Mg	0.41	-0.16	-0.26	0.04	-0.07	0.27	-0.10	-0.05	-0.06	-0.74
SOLUBLE Mg	0.28	-0.59	-0.25	0.09	-0.09	0.17	0.01	-0.05	0.08	-0.57
CLASTIC	0.55	0.76	-0.09	-0.23	-0.08	-0.02	-0.11	0.06	-0.14	-0.07
% SAND	0.16	0.12	0.08	-0.11	-0.06	0.01	-0.95	0.07	-0.01	-0.01
% SILT	0.12	0.50	-0.12	-0.51	-0.08	0.03	-0.16	-0.38	-0.10	-0.19
% CLAY	0.43	0.69	-0.28	-0.02	-0.02	-0.02	0.06	-0.29	0.30	0.00
4.26 Å QUARTZ (SILT)	0.22	-0.10	-0.30	0.45	0.08	-0.01	0.30	-0.16	-0.32	-0.53
4.20 Å F'SPAR (SILT)	-0.12	-0.32	-0.02	0.89	-0.07	0.07	0.10	-0.03	0.04	0.05
4.18 Å F'SPAR (SILT)	-0.13	-0.15	-0.10	0.93	0.08	-0.02	-0.01	0.00	-0.01	-0.01
ILLITE (CLAY)	0.41	-0.03	-0.60	-0.07	0.09	0.05	0.08	-0.09	-0.15	-0.52
CHLORITE (CLAY)	0.45	0.11	-0.58	-0.00	-0.01	0.16	0.14	0.01	-0.29	-0.43
QUARTZ (CLAY)	0.30	0.03	-0.84	-0.03	-0.23	0.17	0.01	0.01	-0.02	-0.12
FELDSPAR (CLAY)	0.08	-0.02	-0.86	0.20	0.13	-0.18	-0.07	0.04	-0.07	-0.16
QUARTZ (TOTAL)	0.24	0.35	-0.30	-0.04	0.10	-0.02	-0.02	-0.24	-0.76	-0.16
SUM OF SQUARES	6.04	5.66	2.70	2.36	1.11	1.08	1.13	1.17	0.98	2.78
% VARIANCE EXPL.	22.38	20.98	9.99	8.75	4.12	4.00	4.19	4.32	3.61	10.31

morphic rocks. Group 1 sediments contain up to 1.0% total Mg, most of which is not soluble in 1.0 N ammonium acetate (Fig. 12). Most of the Mg in these sediments is probably tied up in clay mineral structures, particularly in chlorite. This is in agreement with Callender's (1969) conclusion that most of the Mg in sediments from Lake Superior and Lake Michigan not accounted for by calcite and dolomite is tied up in chlorite. Brunskill et al. (1971), from their study of sediments in lakes in the Experimental Lakes Area, northwestern Ontario, also conclude that Mg is not very mobile and tends to remain in sediments.

Another potentially useful index of the degree of weathering is the ratio of quartz to feldspar in a particular size fraction. Because all feldspars are considerably less durable, chemically and physically, than quartz, increased weathering should increase quartz relative to feldspar. With an increased rate of erosion and a decreased degree of weathering, one might expect both the amount of feldspar and the percentage of silt, relative to quartz, to increase with a consequent positive correlation between percent silt and feldspar peak height. However, from Tables 5 and 6 it appears that this is not so for the silt-size fraction of these lake sediments. The loadings for factor 4, which accounts for 8.7% of the variance in the data, reflect an inverse relationship between the percentage of silt and feldspar peak heights in X-ray diffractograms of silt-size material. This apparent contradiction may be explained by observing the geographic pattern of silt-size mineralogy. The ratio of the 4.26Å quartz peak to the 3.18Å feldspar peak in silt-size material is mapped in Fig. 15. In the northeastern group 1 lakes, the *quartz* : *feldspar* peak height ratio is generally less than 1.0. However, in the southwestern group 3 lakes, this ratio is generally greater than 1.0. The Precambrian crystalline igneous and metamorphic rocks of northeastern Minnesota would be expected to contain relatively high proportions of both plagioclase and orthoclase. The Cretaceous shales of southwestern Minnesota are themselves weathering

products and should therefore contain much lower proportions of feldspars relative to quartz and clay minerals. Therefore, even though chemical weathering is more intense in the northeast, decomposing feldspars at a faster rate, the weathering products that accumulate as silt in the lakes are enriched in feldspars relative to the less decomposed clastic debris in the southwest. Whereas weathering has more control than bedrock lithology on clay-size mineralogy, bedrock type appears to be the dominant control on silt-size mineralogy. Presumably this difference is a result of the much larger ratio of surface area to volume in clay-size material, which would make it more vulnerable to decomposition than the silt fractions.

The five rotated factors discussed above account for 72.4% of the total variance in the data and incorporate most of the relationships among variables. Of the remaining five factors, each explains only about 4% of the total variance and is the result of a single variable. The main contributing variables in each of these five remaining factors are: lake area for factor 5, X-ray aragonite for factor 6, percent sand for factor 7, maximum depth for factor 8, and X-ray quartz for factor 9. It is surprising that lake area and depth do not correlate appreciably with any sedimentary variables (Table 5). As discussed above, the main control exerted by depth on mineralogy is its effect on the dissolution of calcite. However, most calcite-precipitating lakes in Minnesota are sufficiently shallow that dissolution is probably minor. Table 5 shows a slight negative correlation between maximum depth and organic matter but it is not significant at the 99% level. A hint of depth control on amount of sedimentary organic matter is given by the fact that all deep lakes (>30 m) contain less than 20% organic matter in their profundal sediments. However, for lakes shallower than 30 m, there is no relationship between depth and sedimentary organic matter.

Comparison with other lakes—The high-organic group 1 and group 2 lakes are the only Minnesota lakes that have organic con-

Table 7. Loss on ignition, organic carbon, total nitrogen, *organic-C : N* ratio, and carbonate in Minnesota lake sediments compared with other lake groups. All values are based on weight percent dry weight.

	Loss on Ignition	Organic Carbon	Total Nitrogen	Organic-C Nitrogen	Carbonate	Reference
<u>MINNESOTA LAKES:</u>						
8 Low-Org. G 1	18 ± 2*	8 ± 1	0.8 ± 0.2	9.6	none	
13 High-Org. G 1&2	42 ± 7	21 ± 4	2.4 ± 0.5	8.9	none	
10 Intermed. G 2	18 ± 6	9 ± 2	1.1 ± 0.3	8.7	17 ± 7	
10 High-Carb. G 2	20 ± 4	10 ± 2	1.2 ± 0.2	8.5	39 ± 6	
5 Group 3	12 ± 4	6 ± 3	0.6 ± 0.3	9.5	20 ± 2	
46 Total	25 ± 12	12 ± 6	1.4 ± 0.7	8.9		
16 Precambrian Shield Lakes, Experimental Lakes Area, N.W. Ontario	44 ± 14	20 ± 7	2.1 ± 0.8	9.7	none	Brunskill et al. 1971
<u>GREAT LAKES:</u>						
Ontario [†]		3.5	0.39	8.2	5.0 [‡]	Kemp 1971
Erie [†]		2.5	0.33	8.5	5.9 [‡]	"
Huron		2.7	0.39	7.7	5.8	"
South. Michigan		1.6±0.7	0.20±0.09	7.6	8.5 ± 4.2	Callender 1969
Central Michigan		1.0±0.9	0.18±0.11	5.5	6.4 ± 4.6	"
North. Michigan		1.7±0.8	0.24±0.11	7.1	8.8 ± 4.6	"
Green Bay		2.6±1.4	0.31±0.20	8.4	4.7 ± 3.3	"
Superior		2.4	0.23	10.6	1.14	"
<u>WISCONSIN LAKES:</u>						
Mendota		7.3	0.93	7.8	35	Konrad et al. 1970
La Belle		7.3	0.65	11.1	45	Compiled by Brunskill et al. 1971
23 Wisc. Lakes	46	20±10	1.8±0.7§	11.0	not given	
<u>ENGLISH LAKES:</u>						
5 Group 1	14.6	6.1	0.49	12.0	none	Gorham et al. 1974
5 Group 2	16.2	7.1	0.57	13.0	none	
6 Group 3	17.0	7.8	0.65	12.0	none	
16 Total	16.0	7.0	0.58	12.3	none	

* All ± values represent one standard deviation.

† Means of basin samples only; 50 samples for Ontario and 59 samples for Erie.

‡ Calculated from values of carbonate-carbon.

§ Kjeldahl Nitrogen.

|| Group 1 (least productive) includes Wastwater, Thirlmere, Buttermere, Ennerdale, and Crummock; Group 2 (intermediate productivity) includes Haweswater, Coniston, Rydal, Windermere North Basin, and Derwentwater; Group 3 (most productive) includes Loweswater, Windermere South Basin, Blelham Tarn, Ullswater, Bassenthwaite, and Esthwaite.

tents in their profundal sediments even approaching those of lakes in Wisconsin and the Experimental Lakes Area (ELA) of Ontario (Table 7). Most of those lakes resemble group 1 lakes in their water chemistry. As Brunskill et al. (1971) point out, the high organic content of profundal sediments in the ELA lakes results not from

high primary production, but from lack of dilution by clastic material and an abundance of organic matter in soils of the drainage basin. Lack of correlation between organic productivity and sedimentary organic matter is also noted by Kleerekoper and Grenier (1952) for Lake Lauzon, a deep lake on the Precambrian shield of Quebec.

Although Lake Lauzon is oligotrophic, profundal sediments contain up to 75% organic matter (average 45.6%), attributed to inhibition of organic decomposition. Gorham (1958) found that sediments from 54 dilute and generally unproductive Scottish lakes had ignition loss values ranging from 4–70% with a median of only 20%. Those lakes, however, are in a mountainous region favorable to erosion of clastic material.

Brunskill et al. (1971) concluded that most organic material in the ELA lakes is derived from their drainage basins. A similar conclusion was reached by Mackereith (1966) for organic matter in profundal sediments from the English lakes. Data on sedimentary pigments suggest that an appreciable part of the sedimentary organic matter in deep unproductive group 1 lakes in northeastern Minnesota was probably derived from soils in the drainage basins (Sanger and Gorham 1970; Gorham and Sanger 1975). However, in the more productive of the English lakes, as in the more productive Minnesota lakes, pigment data (Gorham 1960; Gorham and Sanger 1967, 1975; Gorham et al. 1974; Sanger and Gorham 1970) demonstrate clearly that most organic matter is derived from primary production within the lakes.

The profundal sediments of the low-organic group 1 lakes are most like those of the English lakes insofar as they generally contain less than 20% organic matter. They are not nearly so low in organic content as profundal sediments from the Great Lakes. The main difference between sediments from the low-organic group 1 Minnesota lakes and the English lakes is the higher C : N ratio of the English sediments. Because both the English and Minnesota samples were analyzed in the same laboratory at about the same time, there is little chance that this difference can be attributed to analytical technique. The higher C : N ratio in the English lake sediments may mean that on average these lakes receive a greater input of more resistant soil organic matter, rich in carbon, than the Minnesota lakes. In general, differences of C : N ratio in pro-

fundal sediments are not very great, with an overall range of 5.5–13.0 (Table 7).

The high-carbonate profundal sediments from the western Minnesota lakes are clearly distinguishable from those of most other lake groups listed in Table 7, except the high-carbonate Wisconsin lakes, Mendota and LaBelle.

References

- ACKROYD, E. A., W. C. WALTON, AND D. L. HILLS. 1967. Ground-water contribution to stream flow and its relation to basin characteristics in Minnesota. Minn. Geol. Surv. Rep. Invest. 6. 36 p.
- BAKER, D. G., AND J. H. STRUB. 1965. Climate of Minnesota, Part 3. Univ. Minn. Agr. Exp. Sta. Tech. Bull. 248. 12 p.
- BOONE, G. M., AND E. P. WHEELER. 1968. Staining for cordierite in thin section. Am. Mineral. 53: 327–331.
- BRIGHT, R. C. 1968. Surface water chemistry of some Minnesota lakes, with preliminary notes on diatoms. Univ. Minn. Limnol. Res. Center, Interim Rep. 3. 58 p.
- BRUNSKILL, G. J., D. POVOLEDO, B. W. GRAHAM, AND M. P. STANTON. 1971. Chemistry of surface sediments of sixteen lakes in the Experimental Lakes Area, northwestern Ontario. J. Fish. Res. Bd. Can. 28: 277–294.
- CALLENDER, E. 1969. Geochemical characteristics of Lakes Michigan and Superior sediments. Proc. 12th Conf. Great Lakes Res. 1969: 124–160.
- CARROLL, D. 1970. Clay minerals: A guide to their X-ray identification. Geol. Soc. Am. Spec. Pap. 126. 80 p.
- DEAN, W. E., JR. 1974. Determination of carbonate and organic matter in calcareous sediments and sedimentary rocks by loss on ignition: Comparison with other methods. J. Sediment. Petrol. 44: 242–248.
- DIXON, W. J. [Ed.]. 1970. BMD Biomedical computer programs, X-series supplement. Univ. Calif. Publ. Automatic Computation 2 (2nd ed.). 260 p.
- GORHAM, E. 1958. The physical limnology of northern Britain: An epitome of the bathymetrical survey of the Scottish freshwater lochs, 1897–1909. Limnol. Oceanogr. 3: 40–50.
- . 1960. Chlorophyll derivatives in surface muds from the English lakes. Limnol. Oceanogr. 5: 29–33.
- , J. W. G. LUND, J. E. SANGER, AND W. E. DEAN, JR. 1974. Some relationships between algal standing crop, water chemistry, and sediment chemistry in the English lakes. Limnol. Oceanogr. 19: 601–617.

- , AND J. E. SANGER. 1967. Plant pigments in woodland soils. *Ecology* **48**: 306–308.
- , AND ———. 1975. Fossil pigments in Minnesota lake sediments, and their bearing upon balance between terrestrial and aquatic inputs to sedimentary organic matter. *Int. Ver. Theor. Angew. Limnol. Verh.* **19**: in press.
- GRIM, R. E. 1953. *Clay mineralogy*. McGraw-Hill.
- HARMAN, H. H. 1967. *Modern factor analysis*, 2nd ed. Chicago.
- HUTCHINSON, G. E. 1957. *Treatise on limnology*, v. 1. Wiley.
- JACKSON, M. L. 1967. *Soil chemical analysis—advanced course*. M. L. Jackson, Dep. Soil. Sci., Univ. Wisconsin, Madison. 894 p.
- KEMP, A. L. W. 1971. Organic carbon and nitrogen in the surface sediments of Lake Ontario, Erie, and Huron. *J. Sediment. Petrol.* **41**: 537–548.
- KLEEREKOPER, H., AND F. GRENIER. 1952. The bottom sediments of Lake Lauzon, Montcalm County, Province of Quebec, I. Chemical and physical investigations. *Can. J. Zool.* **30**: 219–242.
- KONRAD, J. G., D. R. KEENEY, G. CHESTERS, AND K. CHEN. 1970. Nitrogen and carbon distribution in sediment cores of selected Wisconsin lakes. *J. Water Pollut. Control Fed.* **42**: 2094–2101.
- MACKERETH, F. J. H. 1966. Some chemical observations on post-glacial lake sediments. *Phil. Trans. R. Soc. Lond. Ser. B* **250**: 165–213.
- MEGARD, R. P. 1968. Planktonic photosynthesis and the environment of carbonate deposition in lakes. Univ. Minn. Limnol. Res. Center, Interim Rep. 2. 47 p.
- MINNESOTA DEPARTMENT OF CONSERVATION. 1962. *Water resources of Minnesota, a study guide*. Minn. Dep. Conserv. Div. Waters Bull. 16. 28 p.
- MITTEN, H. T., C. H. SCOTT, AND P. G. ROSENE. 1968. Chemical quality of surface waters in Devils' Lake Basin, North Dakota, 1952–60. U.S. Geol. Surv. Water Supply Pap. 1859–8. 42 p.
- MORRISON, D. F. 1967. *Multivariate statistical methods*. McGraw-Hill.
- MORTIMER, C. H. 1941. The exchange of dissolved substances between mud and water in lakes. *J. Ecol.* **29**: 280–329.
- . 1942. The exchange of dissolved substances between mud and water in lakes. *J. Ecol.* **30**: 147–201.
- MÜLLER, G., G. IRION, AND U. FÖRSTNER. 1972. Formation and diagenesis of inorganic Ca-Mg carbonates in the lacustrine environment. *Naturwissenschaften* **59**: 158–164.
- ONDRICK, C. W., AND G. S. SRIVASTAVA. 1970. CORFAN-FORTRAN IV computer program for correlation, factor analysis (R- and Q-mode) and varimax rotation. *Kansas Geol. Surv. Computer Contrib.* 42.
- PLUTH, D. J., R. S. ADAMS, JR., R. H. RUST, AND J. R. PETERSON. 1970. Characteristics of selected horizons from 16 soil series in Minnesota. *Univ. Minn. Agr. Exp. Sta. Tech. Bull.* 272. 34 p.
- ROSS, C. S., AND P. F. KERR. 1931. The clay minerals and their identification. *J. Sediment. Petrol.* **1**: 55–65.
- SANGER, J. E., AND E. GORHAM. 1970. The diversity of pigments in lake sediments and its ecological significance. *Limnol. Oceanogr.* **15**: 59–69.
- SIMS, P. K., AND G. B. MOREY. 1972. *Geology of Minnesota: A centennial volume*. Minn. Geol. Surv. 632 p.
- SNEATH, P. H. A., AND R. R. SOKAL. 1973. *Numerical taxonomy, the principles and practice of numerical classification*. Freeman.
- WANGERSKY, P. J., AND O. I. JOENSUU. 1967. The fractionation of carbonate in deep-sea cores. *J. Geol.* **75**: 148–177.
- WINTER, T. C. 1974. The natural quality of ground water in Minnesota. Minn. Dep. Nat. Resour., Div. Waters, Soils, Minerals. Bull. 26. 25 p.

Submitted: 13 January 1975

Accepted: 24 November 1975