**EIGENVALUES OF ALMOST SKEW SYMMETRIC MATRICES
AND TOURNAMENT MATRICES**

By

Shmuel Friedland

IMA Preprint Series # 911

December 1991

Eigenvalues of almost skew symmetric matrices and tournament matrices

Shmuel Friedland

University of Illinois at Chicago and Institute for Mathematics and its Applications

November 20, 1991

ABSTRACT

A real square matrix C is called almost skew symmetric if $C = S + A$ where S is a rank one real symmetric matrix and A is a real skew symmetric matrix. We shall show that the real eigenvalues of almost skew symmetric matrices satisfy remarkable inequalities. We shall apply these and other inequalities to estimate the spectral radii of the tournament matrices.

§1. Introduction

Let C be a real $n \times n$ matrix. Set

$$C = S + A, \quad S = \frac{C + C^T}{2}, \quad A = \frac{C - C^T}{2}. \quad (1.1)$$

Thus, S and A are the symmetric and the skew (anti) symmetric components of C . We call C almost skew symmetric if S is a rank one matrix. In the terminology of [M-P] $C - \frac{I}{2}$ is a pseudo-tournament matrix. In §2 we discuss remarkable properties of the spectrum of almost skew symmetric matrices. Under certain conditions we estimate the largest real eigenvalue of an almost skew symmetric C from below and above when the nonzero eigenvalue of S is positive. Our estimate are quadratic in the entries of A . We also give conditions on A which ensure the existence of real eigenvalues for C in the case where the dimension of C is even. Our results are almost unrelated to the results discussed in [M-P]. A tournament matrix is a square $(0,1)$ matrix T such that $T + T^T = J - I$ where J is the all ones matrix. In §3 we note that a tournament matrix is an almost skew symmetric matrix minus $\frac{I}{2}$. We then apply our results to get some estimates on the spectral radius of tournament matrices. Our results can be viewed as a natural extension of the basic inequality due to Kirkland [Kir]. In §4 we bring a generalization of Ostrowski's inequality [Ost] for the spectral radius of nonnegative matrices which we apply to generalized tournament matrices. (A nonnegative matrix T is called a generalized tournament matrix if $T + T^T = J - I$.) In the last section we discuss the extremal tournament matrices of an even order n which maximize the spectral radius of the tournament matrices and the corresponding values of their spectral radii as a function of n . We were not able to prove or disprove the Brualdi - Li conjecture [B-L]

on these extremal matrices. We did show that the Brualdi-Li matrix maximizes a function which is closely related to the spectral radius of a tournament matrix.

§2. Perturbation of a rank one symmetric matrix by a skew symmetric matrix

As usual, denote by $M_n(\mathbf{F})$, $S_n(\mathbf{F})$, $A_n(\mathbf{F})$ be the algebra of $n \times n$ matrices, the space of $n \times n$ symmetric matrices and the space of $n \times n$ skew symmetric matrices with the entries in a field $\mathbf{F}$. Let $C \in M_n(\mathbf{R})$ and assume the decomposition (1.1). Let $\text{spec}(C) = \{\lambda_1(C), \dots, \lambda_n(C)\}$ be the spectrum of C . Then $\text{spec}(S), \sqrt{-1}\text{spec}(A) \subset \mathbf{R}$. Arrange the eigenvalues of S in the decreasing order:

$$\lambda_1(S) \geq \lambda_2(S) \geq \dots \geq \lambda_n(S).$$

As $x^T C x = x^T S x$, $x \in \mathbf{R}^n$ it follows that

$$\text{spec}(C) \cap \mathbf{R} \subset [\lambda_n(S), \lambda_1(S)].$$

Thus, the real eigenvalues of C can be viewed as a perturbation of the eigenvalues of S . More precisely, consider the coefficients of the characteristic polynomial of C as polynomials in the entries of S and A . As

$$\det(\lambda I - C) = \det(\lambda I - (S + A)) = \det(\lambda I - A^T) = \det(\lambda I - (S - A))$$

we deduce that the coefficients of the characteristic polynomial of C involve only even degree monomials in the entries of A . That is, the characteristic polynomial of C is a quadratic perturbation of the characteristic polynomial of S by the entries of A . The following theorem exhibits more precisely what happens to the spectrum of a real symmetric matrix S under the perturbation by a real skew symmetric matrix A :

Theorem 2.1. *Let*

$$C(z) = S + zA, \quad S \in S_n(\mathbf{R}), \quad A \in A_n(\mathbf{R}), \quad z \in \mathbf{C}. \quad (2.2)$$

Then for a small enough disk $|z| < r$ the matrix $C(z)$ has n analytic eigenvalues $\lambda_1(z), \dots, \lambda_n(z)$ with corresponding n independent analytic eigenvectors. Let $\lambda \in \text{spec}(S)$ be an eigenvalue of multiplicity k . Assume that the eigenvalues $\lambda_i(z), \dots, \lambda_{i+k-1}(z)$ correspond to λ , i.e. $\lambda_j(0) = \lambda$, $j = i, \dots, i+k-1$. Suppose furthermore that k is odd. Then at least one of the eigenvalues $\lambda_j(z)$, $i \leq j \leq i+k-1$ is real analytic. Furthermore, any real analytic eigenvalue $\lambda_l(z)$ is a function of z^2 .

Proof. Let $z = \sqrt{-1}w$. Then $C(\sqrt{-1}w) = S + wA_1$, $A_1 = \sqrt{-1}A$. Thus, A_1 is a hermitian matrix. It then follows that $C(\sqrt{-1}w)$ is hermitian for real w . Rellich's theorem, e.g. [Kat], yields that for a small enough disk $|w| < r$ the matrix $C(\sqrt{-1}w)$

has n real analytic eigenvalues $\lambda_1(w), \dots, \lambda_n(w)$ and corresponding n independent analytic eigenvectors. Replace $w = -\sqrt{-1}z$ to deduce the corresponding result for $C(z)$.

Assume that $\lambda_j(z)$ is real analytic. As $\lambda_j(\sqrt{-1}w)$ is real analytic we deduce that the power series of $\lambda_j(z)$ contain only even powers of z . Suppose finally that $\lambda \in \text{spec}(A)$ is of an odd multiplicity k . Assume that z is real. Then the eigenvalues of $C(z)$ are either real or come as conjugate complex pairs. As k is odd one of the eigenvalues $\lambda_i(z), \dots, \lambda_{i+k-1}(z)$ is real analytic. $\diamond$

In the space $\mathbf{R}^n$ we consider the standard inner product $(x, y) = x^T y$ and the l_2 norm $\|x\| = (x, x)^{\frac{1}{2}}$. The following lemma is well known, e.g. [G-V, §9.1.1, p'483] but we bring its proof for completeness:

Lemma 2.3. *Let $A \in A_n(\mathbf{R})$. Assume that $e \in \mathbf{R}^n$, $\|e\| = 1$. Let $U = \text{span}\{e, Ae, \dots, A^{k-1}e\} \subset \mathbf{R}^n$ be a cyclic subspace of A , that is $e, Ae, \dots, A^{k-1}e$ are linearly independent and $A^k e \in U$. Assume that $k > 1$. Let $e^1 = e, \dots, e^k$ be the orthonormal vectors obtained by the Gram-Schmidt process from the vectors $e, Ae, \dots, A^{k-1}e$:*

$$e^1 = e, e^2 = \frac{Ae}{\|Ae\|}, e^j = \frac{Ae^{j-1} - ((e^{j-2})^T Ae^{j-1})e^{j-2}}{\|Ae^{j-1} - ((e^{j-2})^T Ae^{j-1})e^{j-2}\|}, j = 3, \dots, k. \quad (2.4)$$

Then in the basis $e^1, \dots, e^k$ the matrix A is represented by a skew symmetric $k \times k$ tridiagonal matrix $\hat{A} = (\hat{a}_{ij})_1^k$ whose lower nontrivial diagonal is given by:

$$\hat{a}_{21} = \alpha_1 = \|Ae\|, a_{(j+1)j} = \alpha_j = \|Ae^j + \alpha_{j-1}e^{j-1}\| = (e^{j+1})^T Ae^j, j = 2, \dots, k-1. \quad (2.5)$$

Proof. As $x^T Ax = 0$ we deduce that Ae is orthogonal to e . Since $k > 1$ $Ae \neq 0$ hence $e^2 = \frac{Ae}{\|Ae\|}$. Assume by the induction hypothesis that (2.4) holds for $j = 1, \dots, m-1$. Let e^m be given by (2.4). As $(e^{m-1})^T Ae^{m-1} = (e^{m-1})^T e^{m-2} = 0$ it follows that $(e^m)^T e^{m-1} = 0$. The formula for e^m yields that $(e^m)^T e^{m-2} = 0$. Consider $(e^m)^T e^j$ for $j < m-2$. By the induction hypothesis $(e^{m-2})^T e^j = 0$. As $Ae^j \in \text{span}\{e^1, \dots, e^{j+1}\}$ we deduce that $(e^{m-1})^T Ae^j = 0$. Hence, $(e^m)^T e^j = 0$. This shows that $e^1, \dots, e^k$ form an orthonormal basis of U which is constructed by the Gramm-Schmidt process. The formula (2.4) yields that A is represented by a tridiagonal matrix $\hat{A}$ in the orthonormal basis $e^1, \dots, e^k$. As A is skew symmetric so is $\hat{A}$. The equalities (2.5) follow straight forward from (2.4). $\diamond$

Let $\bar{\mathbf{R}} = \mathbf{R} \cup \{-\infty\}$. We adopt the natural convention $-\infty < x, \forall x \in \mathbf{R}$. Let $x, y \in \bar{\mathbf{R}}$. We then let $x \prec y$ ($y \succ x$) if either $x = y = -\infty$ or $y \in \mathbf{R}$ and $x < y$.

Theorem 2.6. *Let $S \in S_n(\mathbf{R})$, $A \in A_n(\mathbf{R})$. Assume furthermore that S is a rank one matrix with positive eigenvalue $\delta > 0$ and the corresponding eigenvector $Se = \delta e$, $\|e\| = 1$. If $Ae = 0$ then the $\text{spec}(S + A)$ is obtained from $\text{spec}(A)$ by removing one zero eigenvalue and replacing it by δ . Assume next that $Ae \neq 0$. Let $e^1 = e, \dots, e^k$ be the orthonormal basis of the cyclic subspace of $U = \text{span}\{e, Ae, \dots, A^{k-1}e\} \subset \mathbf{R}^n$ as in Lemma 2.3. Then in the basis $e^1, \dots, e^k$ S is represented by the $k \times k$ diagonal matrix $D = (d_{ij})_1^k = \text{diag}\{\delta, 0, \dots, 0\}$*

and A by the $k \times k$ tridiagonal skew symmetric matrix $\hat{A} = (\hat{a}_{ij})_1^k$ given by (2.5). Furthermore, any real eigenvalue of $D + \hat{A}$ is nonzero. Moreover, all nonzero real eigenvalues of $S + A$ are the eigenvalues of $D + \hat{A}$. Set:

$$q_l(\lambda) = \det(\lambda I - (d_{ij} + \hat{a}_{ij})_1^l), \quad l = 1, \dots, k. \quad (2.7)$$

Denote by ρ_l the maximal real root of $q_l(\lambda)$ for $l = 1, \dots, k$. If l is even and $q_l(\lambda)$ does not have a real root we then let $\rho_l = -\infty$. Then

$$\rho_i > \rho_j, \quad j = i + 1, \dots, k; \quad i = 1, 3, \quad \rho_j \succ \rho_2, \quad j = 3, \dots, k. \quad (2.8)$$

Suppose that $q_{2i-1}(\lambda)$, $i = 1, \dots, \lfloor \frac{k+1}{2} \rfloor$ have exactly one simple real root ρ_{2i-1} , $i = 1, \dots, \lfloor \frac{k+1}{2} \rfloor$. Then

$$\begin{aligned} \rho_{2i-1} &> \rho_{2i+1}, \quad i = 1, \dots, \lfloor \frac{k-1}{2} \rfloor, \\ \rho_{2i} &\prec \rho_{2i+2}, \quad i = 1, \dots, \lfloor \frac{k-2}{2} \rfloor, \\ \rho_{2i-1} &> \rho_{2j}, \quad i = 1, \dots, \lfloor \frac{k+1}{2} \rfloor, \quad j = 1, \dots, \lfloor \frac{k}{2} \rfloor. \end{aligned} \quad (2.9)$$

Proof. Let $U^\perp \subset \mathbf{R}^n$ be the orthogonal complement of U . As A is skew symmetric we deduce that $AU^\perp \subset U^\perp$. Clearly, $S\text{span}(e)^\perp = 0$. Hence $SU \subset U$, $SU^\perp = 0$ and

$$\text{spec}(S + A) = \text{spec}(S + A|_U) \cup \text{spec}(A|_{U^\perp}).$$

Assume first that $Ae = 0$. Then $U = \text{span}(e)$ and the above equality yield that the $\text{spec}(S + A)$ is obtained from the $\text{spec}(A)$ by replacing a zero eigenvalue of A by δ .

We now assume that $Ae \neq 0$, that is, $k = \dim(U) > 1$. Suppose that $(S + A)x = 0$, $0 \neq x \in U$. As $0 = x^T(S + A)x = x^T Sx$ we deduce that $Sx = 0$. Thus $Ax = 0$. As U is a cyclic subspace of A we deduce that $U = \text{span}(x)$ contrary to our assumption. Thus, the matrix $D + A$ does not have a zero eigenvalue. We now show the inequalities (2.8).

Let

$$p_l(\lambda) = \det(\lambda I - (d_{ij} + \hat{a}_{ij})_{k-l+1}^k), \quad l = 1, \dots, k. \quad (2.10)$$

Set $p_0(\lambda) = q_0(\lambda) = 1$. The following recursive formulas are established in a straightforward manner:

$$\begin{aligned} p_k(\lambda) &= q_k(\lambda) = q_i(\lambda)p_{k-i}(\lambda) + q_{i-1}(\lambda)\alpha_i^2 p_{k-i-1}, \quad i = 1, \dots, k-1, \\ p_i(\lambda) &= \lambda p_{i-1}(\lambda) + \alpha_{k-i+1}^2 p_{i-2}, \quad i = 2, \dots, k-1. \end{aligned} \quad (2.11)$$

Note that $p_i(\lambda) > 0$, $\lambda > 0$, $i = 1, \dots, k-1$. The formulas (2.11) yield

$$q_k(\lambda) = (\lambda - \delta)p_{k-1}(\lambda) + \alpha_1^2 p_{k-2}(\lambda).$$

Hence, $q_k(\lambda) \geq \alpha_1^2 p_{k-2}(\lambda) > 0$ for $\lambda \geq \delta (> 0)$. Therefore, $\rho_k < \rho_1 = \delta$. By considering the matrix $(d_{ij} + \hat{a}_{ij})_1^l$ we deduce that $\rho_l < \rho_1$, $l = 2, \dots, k$. Consider ρ_2 . If $\rho_2 = -\infty$ we trivially have that $\rho_j \succ \rho_2$, $j = 3, \dots, k$. Assume now that $\rho_2 \in \mathbf{R}$. Our arguments yield that $0 < \rho_2 < \delta$. The first set of the equalities of (2.11) imply:

$$q_k(\lambda) = q_2(\lambda)p_{k-2}(\lambda) + (\lambda - \delta)\alpha_2^2 p_{k-3}(\lambda).$$

Thus $q_k(\rho_2) = (\delta - \rho_2)\alpha_2^2 p_{k-3}(\rho_2) < 0$. Hence, q_k has a real root and $\rho_k > \rho_2$. The arguments above show that each q_l , $l = 3, \dots, k$ has a real root and $\rho_l > \rho_2$, $l = 3, \dots, k$. Assume finally that all q_{2i-1} have simple real roots. We prove the inequalities (2.9) by the induction on k . Suppose (2.9) holds for $k = l$. Let $k = l + 1$. The first set of equalities of (2.11) yield that

$$q_{l+1} = \lambda q_l + \alpha_l^2 q_{l-1}. \quad (2.12)$$

Suppose first that l is even. Then the induction hypothesis yield that $\rho_l < \rho_{l-1}$. It then follows that $q_{l+1}(\lambda) > 0$ for $\lambda \geq \rho_{l-1}$. Hence, $\rho_{l+1} < \rho_{l-1}$. If $\rho_l = -\infty$ we obviously have the inequality $\rho_l < \rho_{l+1}$. If $0 < \rho_l$ then (2.12) yields $q_{l+1}(\rho_l) = \alpha_l^2 q_{l-1}(\rho_l) < 0$. The last inequality follows from the inequality $\rho_l < \rho_{l-1}$ and the assumption that ρ_{l-1} is the only real root of q_{l-1} which is also simple. Therefore $\rho_{l+1} > \rho_l$. Similar arguments apply in the case l is odd. $\diamond$

The two roots of:

$$q_2(\lambda) = \lambda^2 - \delta\lambda + \alpha_1^2 \quad (2.13)$$

are given by

$$\tau_1 = \frac{\delta - \sqrt{\delta^2 - 4\alpha_1^2}}{2}, \quad \tau_2 = \frac{\delta + \sqrt{\delta^2 - 4\alpha_1^2}}{2}. \quad (2.14)$$

Thus, if $\delta^2 \geq 4\alpha_1^2$ then $\rho_2 = \tau_2 \geq \frac{\delta}{2}$. Then Theorem 2.6 yields:

$$\rho_i > \rho_2 = \frac{\delta + \sqrt{\delta^2 - 4\alpha_1^2}}{2}, \quad i = 3, \dots, k, \quad \delta^2 \geq 4\alpha_1^2. \quad (2.15)$$

In particular, the matrix $S + A$ has a positive real eigenvalue which is not less than ρ_2 . This last statement can be deduced from the recent inequality of [Kir] for hypertournament matrices. We adopt Kirkland's proof to state a more precise result about the spectrum of almost skew symmetric matrices in the case $\delta^2 \geq 4\alpha_1^2$.

Theorem 2.16. *Let $S \in S_n(\mathbf{R})$, $A \in A_n(\mathbf{R})$. Assume furthermore that S is a rank one matrix with positive eigenvalue $\delta > 0$ and the corresponding eigenvector $Se = \delta e$, $\|e\| = 1$. Set $\alpha_1 = \|Ae\|$. Assume that $\delta^2 \geq 4\alpha_1^2$. Then $S + A$ has one real eigenvalue λ satisfying the inequality:*

$$\lambda \geq \frac{\delta + \sqrt{\delta^2 - 4\alpha_1^2}}{2}. \quad (2.17)$$

Furthermore, all other $n - 1$ eigenvalues of $A + S$ satisfy the inequality:

$$\Re(\lambda) \leq \frac{\delta - \sqrt{\delta^2 - 4\alpha_1^2}}{2}. \quad (2.18)$$

Proof. Using a continuity argument it is enough to consider the case $\delta^2 > 4\alpha_1^2$. Let

$$H = S - \frac{\delta}{2}I, \quad M_t = S + A - tI, \quad P_t = M_t H + H M_t^T, \quad \frac{\delta - \sqrt{\delta^2 - 4\alpha_1^2}}{2} < t < \frac{\delta + \sqrt{\delta^2 - 4\alpha_1^2}}{2}. \quad (2.19)$$

Then $P_t = N_t \oplus \frac{t\delta}{2}I_{n-2}$. A straightforward calculation shows that the 2×2 matrix N_t is positive definite for the values of t prescribed in (2.19). Thus, P_t is positive definite. The result of Ostrowski and Schneider [O-S] claims that M_t and H have the same inertia. That is, M_t has exactly one eigenvalue with positive real part and all other $n - 1$ eigenvalues have negative real part. As M_t is a real matrix we deduce that the unique eigenvalue with a positive real part must be real. To deduce (2.17) let t tend to $\rho_2 = \tau_2$ from below and to deduce (2.18) let t tend to τ_1 from above. $\diamond$

Theorem 2.20 For $\delta > 0$, $\alpha_1^2, \alpha_2^2 \geq 0$ Let $\rho_3(\alpha_1^2, \alpha_2^2)$ be the largest real root of

$$q_3(\lambda) = \lambda^3 - \delta\lambda^2 + \lambda(\alpha_1^2 + \alpha_2^2) - \delta\alpha_2^2. \quad (2.21)$$

For a fixed value $\alpha_1^2 > 0$, $4\alpha_1^2 \leq \delta^2$, ρ_3 is a strictly increasing function of α_2^2 such that

$$\rho(\alpha_1^2, 0) = \rho_2(\alpha_1^2) = \frac{\delta + \sqrt{\delta^2 - 4\alpha_1^2}}{2}, \quad \rho_2(\alpha_2^2, \infty) = \delta, \quad \delta^2 \geq 4\alpha_1^2 > 0. \quad (2.22)$$

For a fixed value of $\alpha_2 > 0$ ρ_3 is a strictly decreasing function of α_1 on the interval $[0, \frac{\delta}{2}]$ such that $\rho_3(0, \alpha_2^2) = \delta$. For a fixed value $\alpha_2^2 \geq \frac{\delta^2}{3}$ ρ_3 is a strictly decreasing function of α_1^2 such that:

$$\rho_3(0, \alpha_2^2) = \delta, \quad \rho_3(\infty, \alpha_2^2) = 0, \quad (2.23)$$

Assume finally that $\alpha_1^2, \alpha_2^2 > 0$ and $4\alpha_1^2 \leq \delta^2$. Then

$$0 < \rho_3 - \rho_2 < \frac{\alpha_2^2(\delta - \rho_2)}{(2\rho_2 - \delta)\rho_2 + \alpha_2^2}. \quad (2.24)$$

Let $S \in S_n(\mathbf{R})$, $A \in A_n(\mathbf{R})$. Assume furthermore that S is a rank one matrix with positive eigenvalue $\delta > 0$ and the corresponding eigenvector $Se = \delta e$, $\|e\| = 1$. Assume that $A^2 e \neq 0$. Set

$$\alpha_1^2 = \|Ae\|^2, \quad \alpha_2^2 = \left\| \frac{A^2 e}{\alpha_1} + Ae \right\|^2 = \frac{\|A^2 e\|^2}{\|Ae\|^2} - \|Ae\|^2. \quad (2.25)$$

Then $S + A$ has exactly one positive simple eigenvalue λ satisfying

$$\rho_2(\alpha_1^2) < \lambda \leq \rho_3(\alpha_1^2, \alpha_2^2). \quad (2.26)$$

Proof. Using the definitions (2.13) and (2.2) we deduce in a straightforward manner the equality

$$q_3(\lambda) = \lambda q_2(\lambda) + \alpha_2^2(\lambda - \delta), \quad (2.27)$$

which is a special case of (2.12). Then ρ_3 is the maximal real solution of

$$\frac{\lambda q_2(\lambda)}{\delta - \lambda} = \alpha_2^2. \quad (2.28)$$

Note that $q_2(\lambda)$ is an increasing function of λ for $\lambda \geq \frac{\delta}{2}$. Also, for $\delta^2 \geq 4\alpha_1^2 > 0, \alpha_2^2 > 0$ the argument in the proof of Theorem 2.6 yields that

$$\delta > \rho_3(\alpha_1^2, \alpha_2^2) > \rho_2(\alpha_1^2) \geq \frac{\delta}{2}, \quad 0 < 4\alpha_1^2 \leq \delta^2, \quad 0 < \alpha_2^2. \quad (2.29)$$

It then follows that for a fixed α_1 , $\delta^2 \geq 4\alpha_1^2 > 0$ ρ_3 is an increasing function in $\alpha_2^2 > 0$. Thus, when $\alpha_2^2 = 0$ we obtain the first equality in (2.23). As $\rho_3 < \delta$ we deduce that if $\alpha_2^2 \rightarrow \infty$ then $\rho_3 \rightarrow \delta$. This proves the second equality of (2.23).

Fix the value of $\alpha_2 > 0$. Then for $\alpha_1 \in (0, \frac{\delta}{2})$ we know that $\rho_3 > \rho_2 > \frac{\delta}{2}$. Note that the equalities (2.27) and (2.13) directly yield that for a fixed $\lambda > 0$, q_3 is an increasing function of α_1^2 . Fix the values of α_1^2, α_2^2 . Then q_3 is a strictly increasing function of λ for $\lambda > \frac{\delta}{2}$. These arguments imply straightforward the $\rho_3(\alpha_1, \alpha_2)$ is a strictly decreasing function of $\alpha_1 \in (0, \frac{\delta}{2})$. Assume that $\alpha_2^2 \geq \frac{\delta^2}{3}$. A straightforward calculation shows that $q_3' \geq 0$ for all λ . The previous arguments yield that ρ_3 is a strictly decreasing function of $\alpha_1 > 0$. The equalities (2.23) are straightforward.

We now prove (2.24). Assume that $0 < \alpha_1 \leq \frac{\delta}{2}$, $0 < \alpha_2$. Then the inequality $\rho_2 < \rho_3$ is a consequence of (2.8). Clearly

$$q_2(\rho_3) = q_2'(\rho_2)(\rho_3 - \rho_2) + (\rho_3 - \rho_2)^2 > q_2'(\rho_2)(\rho_3 - \rho_2).$$

Since ρ_3 satisfies the equation (2.28) from the above inequality and the fact $q_2'(\rho_2) \geq 0$ we deduce:

$$\alpha_2^2 > \frac{\rho_3 q_2'(\rho_2)(\rho_3 - \rho_2)}{\delta - \rho_3} \geq \frac{\rho_2 q_2'(\rho_2)(\rho_3 - \rho_2)}{\delta - \rho_3}.$$

The inequality (2.24) is equivalent to the observation that α_2^2 is greater than the last term in the above inequality. $\diamond$

Theorem 2.30. *Let $S \in S_n(\mathbf{R})$, $A \in A_n(\mathbf{R})$. Assume furthermore that S is a rank one matrix with positive eigenvalue $\delta > 0$ and the corresponding eigenvector $Se = \delta e$, $\|e\| = 1$. Assume that $Ae \neq 0$. Let*

$$e^1 = e, \dots, e^k, D, \hat{A}, q_1, \dots, q_k, \rho_1, \dots, \rho_k$$

be defined as in Theorem 2.6. Assume that $\alpha_1 = \|Ae\| \leq \frac{\delta}{2}$. Then all $\rho_1, \dots, \rho_k$ are positive numbers which satisfy the strict inequalities:

$$\begin{aligned} \rho_{2i-1} &> \rho_{2i+1}, \quad i = 1, \dots, \lfloor \frac{k-1}{2} \rfloor, \\ \rho_{2i} &< \rho_{2i+2}, \quad i = 1, \dots, \lfloor \frac{k-2}{2} \rfloor, \\ \rho_{2i-1} &> \rho_{2j}, \quad i = 1, \dots, \lfloor \frac{k+1}{2} \rfloor, \quad j = 1, \dots, \lfloor \frac{k}{2} \rfloor. \end{aligned} \quad (2.31)$$

Proof. We prove (2.31) by the induction on k . For $k = 1, 2, 3$ we showed the validity of (2.31). Assume that (2.31) holds for $k = l \geq 3$. Let $k = l + 1$. Suppose first that l is even. The induction hypothesis yields that $\rho_l < \rho_{l-1}$. Use (2.12) to deduce that $q_{l+1}(\lambda) > 0$ for $\lambda \geq \rho_{l-1}$. Hence $\rho_{l+1} < \rho_{l-1}$. Combine Theorem 2.16, (2.12) and the induction hypothesis $\rho_{l-1} > \rho_l > \rho_2$ to deduce that $q_{l+1}(\rho_l) = \alpha_l^2 q_{l-1}(\rho_l) < 0$. Therefore $\rho_{l+1} > \rho_l$. Similar arguments apply in the case l is odd. $\diamond$

Note that under the assumptions of Theorem 2.30 ρ_k is the maximal real eigenvalue of the matrix $S + A$.

§3. Tournament matrices

A *tournament matrix* is a square $(0, 1)$ matrix T such that $T + T^T = J - I$ where J is the all ones matrix. These matrices have many interesting properties, see [Moo] and the references therein. The spectral properties of these matrices were studied by quite a few numbers of authors, see for example [M-P] and [Kir] and the references therein. We first note that any tournament matrix is an almost skew symmetric matrix minus $\frac{I}{2}$

$$T = C - \frac{I}{2}, \quad C = \frac{J}{2} + A, \quad A = \frac{T - T^T}{2}, \quad J = uu^T, \quad u = (1, \dots, 1)^T. \quad (3.1)$$

To estimate the spectral radius $\rho(T)$ of T it is enough to estimate the maximal real eigenvalue of an almost skew symmetric matrix C . In the notation of Section 2

$$\delta = \frac{n}{2}, \quad e = \frac{u}{\sqrt{n}} = \frac{(1, \dots, 1)^T}{\sqrt{n}}. \quad (3.2)$$

Set

$$b = (b_1, \dots, b_n)^T = Tu - T^T u \in \mathbb{Z}^n. \quad (3.3)$$

That is b_i is the number of wins minus the number of losses of the player number i . We call b the balance vector. Use (3.1), (2.4) and (2.25) to deduce:

$$\alpha_1^2 = \frac{b^T b}{4n}, \quad \alpha_2^2 = -\frac{b^T A^2 b}{b^T b} - \frac{b^T b}{4n}. \quad (3.4)$$

We assume that $\alpha_1 = \alpha_2 = 0$ if $b = 0$. Theorems 2.6 and 2.30 yield:

Theorem 3.5 *Let T be an $n \times n$ tournament matrix. Assume (3.1) – (3.5). Let $\rho_3(\alpha_1^2, \alpha_2^2)$ be the biggest real root of (2.21). Then*

$$\rho(T) \leq \rho_3(\alpha_1^2, \alpha_2^2) - \frac{1}{2}. \quad (3.6)$$

Suppose furthermore that $4b^Tb \leq n^3$. Let $\rho_2(\alpha_1^2)$ be the maximal root of (2.13). Then

$$\rho(T) \geq \rho_2(\alpha_1^2) - \frac{1}{2} \quad (3.7)$$

and equality holds iff either $b = 0$ or $b \neq 0, \alpha_2 = 0$.

The inequality (3.7) is due to Kirkland [Kir]. He also gives the condition for equality stated in a different form.

The difficulty in using the inequality (3.6) is due to the fact that ρ_3 is a solution to the cubic equation. We now bring an upper bound on the spectral tournament matrices using Ostrowski's theorem. Let $B = (b_{ij})_1^n$ be a nonnegative matrix. Set

$$r_i(B) = \sum_{j=1}^n b_{ij}, c_i(B) = r_i(B^T), \hat{r}_i(B) = r_i(B) - b_{ii}, \hat{c}_i(B) = \hat{r}_i(B^T) \quad i = 1, \dots, n. \quad (3.8)$$

Ostrowski's theorem [Ost], see also [Bru] for generalization of Ostrowski's theorem, claims that $\text{spec}(B)$ - the spectrum of B is contained in the following complex disks:

$$|z - b_{ii}| \leq \hat{r}_i(B)^t \hat{c}_i(B)^{1-t}, \quad i = 1, \dots, n$$

for any $0 \leq t \leq 1$. In particular:

$$\rho(B) \leq \mu_t(B), \quad \mu_t(B) = \max_{1 \leq i \leq n} b_{ii} + \hat{r}_i(B)^t \hat{c}_i(B)^{1-t}, \quad 0 \leq t \leq 1. \quad (3.9)$$

Corollary 3.10 *Let T be an $n \times n$ tournament matrix. Then*

$$\rho(T) \leq \mu_{\frac{1}{2}}(T). \quad (3.11)$$

Assume that n is even. Let ρ_n^* be the maximal spectral radius of all $n \times n$ tournament matrices. By abusing the notation we let $\rho^* = \rho_n^*$. A conjecture due to Brualdi-Li [B-L] gives a "natural" candidate for the tournament matrix T^* , $\rho(T^*) = \rho^*$.

$$\begin{aligned} Q &= (Q_{ij})_1^2, \quad Q_{11} = Q_{22} = V, \quad Q_{12} = I + V^T, \quad Q_{21} = V^T, \\ V &= (v_{ij})_1^{\frac{n}{2}}, \quad v_{ij} = 1 \text{ for } 1 \leq i < j \leq \frac{n}{2}, \quad v_{ij} = 0 \text{ for } 1 \leq j \leq i \leq \frac{n}{2}. \end{aligned} \quad (3.12)$$

In particular, Q is almost regular. Recall that a tournament matrix T is called almost regular if $|r_i(T) - \frac{n-1}{2}| = \frac{1}{2}$, $i = 1, \dots, n$. A weaker version of Brualdi-Li conjecture is that any extremal matrix T^* is almost regular. We now study these conjectures.

Theorem 3.13 *Let T be a tournament matrix of an even order n . Then*

$$\rho(T) \leq \sqrt{\frac{n}{2}(\frac{n}{2} - 1)}. \quad (3.14)$$

Assume furthermore that T satisfies the condition

$$|r_i(T) - \frac{n-1}{2}| > \frac{1}{2}, \quad i = 1, \dots, n. \quad (3.15)$$

Then

$$\rho(T) \leq \sqrt{(\frac{n}{2} + 1)(\frac{n}{2} - 2)}. \quad (3.16)$$

In particular, any tournament matrix which satisfies (3.15) satisfies the inequality

$$\rho(T) < \frac{n-1}{2} - \frac{1}{n + \sqrt{n^2 - 4}}. \quad (3.17)$$

Proof. Note that for any tournament matrix of an even order n we have the inequality $r_i(T)c_i(T) \leq \frac{n}{2}(\frac{n}{2} - 1)$. Hence, (3.11) yields (3.14). The assumption (3.15) implies that $r_i(T)c_i(T) \leq (\frac{n}{2} + 1)(\frac{n}{2} - 2), i = 1, \dots, n$. Thus, (3.11) implies (3.16). A straightforward calculation shows that the right-hand side of (3.16) is strictly less than the right-hand side of (3.17). $\diamond$

The following lemma is due to Kirkland:

Lemma 3.18. *Let $T = (T_{ij})_1^{2m}$, $T_{ij} \in M_m(\mathbf{R})$, $i, j = 1, 2$ be an almost regular tournament matrix of an even order. Assume furthermore that the row sum of each of the first m rows of T is m and $T_{11} = T_{22} = V$ where V is given by (3.12). Then T is equal to the Brualdi-Li matrix Q given by (3.12).*

Proof. As T is an almost regular tournament matrix with $r_i(T) = m$, $i = 1, \dots, m$, we deduce that $c_{m+i}(T) = 2m - 1 - r_{m+i}(T) = m$, $i = 1, \dots, m$. Clearly, $r_i(V) = c_{m-i+1}(V) = m - i$, $i = 1, \dots, m$. The equalities $r_i(T) = c_{m+i}(T) = m$, $i = 1, \dots, m$, yield

$$r_i(T_{12}) = m - r_i(V) = i, \quad c_i(T_{12}) = m - c_i(V) = m - i + 1, \quad i = 1, \dots, m. \quad (3.19)$$

We claim that the above equations have a unique 0-1 matrix solution $T_{12} = I + V^T$. For $m = 1, 2$ the claim is trivial. Assume that the claim holds for $m \leq k-1$. Let $m = k$. As T_{12} is a 0-1 matrix it follows that the first column and the last row of T_{12} consists of all ones. The equality $r_1(T_{12}) = 1$ yields that the only nonzero entry on the first row of T_{12} is the entry $(1, 1)$. As $c_k(T_{12}) = 1$ we deduce that the only nonzero entry on the last column of T_{12} is the entry (k, k) . Delete the first and the last rows and columns of T_{12} and use the induction hypothesis to show that the equalities (3.19) have a unique 0-1 matrix solution $T_{12} = I + V^T$. As T is a tournament we deduce that $T_{21} = V^T$. $\diamond$

Theorem 3.20. *Let T be an almost regular tournament matrix of an even order n . Assume (3.1) - (3.4). Then $\alpha_1 = \frac{1}{2}$. In particular, Kirkland's inequality holds [Kir]:*

$$\rho(T) \geq \rho_2\left(\frac{1}{4}\right) - \frac{1}{2} = \frac{n-1}{2} - \frac{1}{n + \sqrt{n^2 - 4}}. \quad (3.21)$$

Furthermore

$$\alpha_2^2 \leq \frac{n^2 - 4}{12}. \quad (3.22)$$

Equality holds iff T is permutationally similar to the Brualdi-Li matrix given by (3.12). In particular, any almost regular tournament matrix T of an even order satisfies the inequality

$$\rho(T) \leq \rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right) - \frac{1}{2} \quad (3.23)$$

which implies

$$\rho(T) < \frac{n - 1}{2} - \frac{3}{8n}. \quad (3.24)$$

Proof. Let $n = 2m$. Then the balance vector b has m entries equal to 1 and m entries equal to -1 . Use the first part of (3.4) to deduce the equality $\alpha_1^2 = \frac{1}{4}$. According to Theorem 2.20 we have that $\rho(\frac{1}{4}, \alpha_2^2)$ is an increasing function of α_2^2 . (Here $\delta = \frac{n}{2}$.) Use (3.4) to deduce that the maximal value of α_2^2 is achieved for the almost regular tournament matrices for which $\|Ab\|$ is maximal. Since $Jb = 0$ we get:

$$\begin{aligned} \|Ab\|^2 &= \|(A + \frac{J}{2})b\|^2 = \|(T + \frac{I}{2})b\|^2 = b^T T^T T b + \frac{b^T b}{4} + \frac{b^T (T + T^T) b}{2} = \\ &= b^T T^T T b + \frac{n}{4} + \frac{b^T (J - I) b}{2} = b^T T^T T b - \frac{n}{4}. \end{aligned} \quad (3.25)$$

Thus, it is enough to consider almost regular tournament matrices which maximize $\|Tb\|^2$. Rename the indices $1, \dots, 2m$ so that the first m coordinates of b are equal to 1 and the last m coordinates of b are equal to -1 . Let $T = (t_{ij})_1^n = (T_{ij})_1^m$, $T_{ij} \in M_m(\mathbf{R})$. Assume that s_i, t_i are the i -th row sum of T_{11}, T_{22} respectively, for $i = 1, \dots, m$. It then follows that

$$\begin{aligned} b^T T^T T b &= \sum_1^m (2s_i - m)^2 + (2t_i - (m - 1))^2, \\ s_i, t_i &\in \mathbf{Z}, \quad 0 \leq s_i, t_i \leq m - 1, \quad \sum_1^m s_i = \sum_1^m t_i = \frac{m(m - 1)}{2}. \end{aligned}$$

We next observe that if $x \leq y$ are two integers then $x^2 + y^2 < (x - 2)^2 + (y + 2)^2$. We now maximize the expression

$$\sum_1^m (2s_i - m)^2 + (2t_i - (m - 1))^2. \quad (3.26)$$

on all tournament matrices of an order $n = 2m$. Let $T = (t_{ij})_1^n$. Assume that $t_{ij} = 1$, $1 \leq i \neq j \leq m$. Then we may change the value of (3.26) by letting $t_{ij} = 0$. This is equivalent to replacing s_i, s_j by $s_i - 1$ and $s_j + 1$ and leaving all other s_k and t_k unchanged. Assume that T maximizes (3.26). We claim that after renaming the first m vertices if necessary

we have the equality: $T_{11} = V$. In particular $s_i = m - i$, $i = 1, \dots, m$. Assume that we renamed the indices so that $s_1 \geq s_2 \geq \dots \geq s_m$. Suppose to the contrary that $s_m > 0$, that is, $t_{mj} = 1$, $1 \leq j < m$. Then replace s_m, s_j by $s_m - 1, s_j + 1$ to contradict the optimality of T . Thus, the last row of T_{11} is a zero row and the first $m - 1$ entries of the last column of T_{11} are equal to zero. It then follows that $s_i - 1$, $i = 1, \dots, m - 1$ are the i -th row sum of $(t_{ij})_1^{m-1}$. It now follows that $s_{m-1} - 1 = 0$. Thus, the last row of the matrix $(t_{ij})_1^{m-1}$ is equal to zero. Continuing in the same manner we deduce that $T_{11} = V$. Assume that we have renamed the indices $m + 1, \dots, 2m$ so that $t_1 \geq \dots \geq t_m$. Our arguments yield that $T_{22} = V$. In particular, Q maximize the expression (3.26).

Assume now that T is an almost regular matrix with $r_i(T) = m$, $i = 1, \dots, m$, which maximizes (3.26). By renaming the indices in the sets $\{1, \dots, m\}$ and $\{m + 1, \dots, 2m\}$ if necessary we deduce from the arguments above that $T_{11} = T_{22} = V$. Lemma 3.18 yields that $T = Q$.

In what follows we assume that T is an almost regular tournament matrix. Using the standard formula

$$\sum_1^k i^2 = \frac{(2k+1)(k+1)k}{6}$$

we deduce that

$$\|Tb\|^2 \leq \frac{(2m^2 + 1)m}{3}.$$

From (3.25), (3.4) and the above inequality we obtain the inequality (3.22) and the equality holds iff T is permutationally similar to the Brualdi-Li matrix. Theorem 2.20 yields the inequality (3.23). To deduce (3.24) let $r_n = \frac{n}{2} - \frac{3}{8n}$. A straightforward calculation shows that

$$\frac{r_n q_2(r_n)}{\delta - r_n} > \frac{n^2 - 4}{12}, \quad \delta = \frac{n}{2}, \quad \alpha_1^2 = \frac{1}{4}.$$

The arguments of the proof of Theorem 2.20 yield that

$$\rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right) < \frac{n}{2} - \frac{3}{8n}. \quad (3.27)$$

§4. A generalization of Ostrowski's inequality and applications

In this section we let $h = (1, \dots, 1)^T$.

Theorem 4.1 *Let B be an $n \times n$ nonnegative irreducible matrix matrix. Set*

$$\begin{aligned} Bu &= \rho(B)u, \quad B^T v = \rho(B)v, \\ u &= (u_1, \dots, u_n)^T, \quad v = (v_1, \dots, v_n)^T, \quad 0 < u_i, v_i, \quad i = 1, \dots, n, \quad \sum_1^n u_i v_i = 1. \end{aligned} \quad (4.2)$$

Let $0 \leq s \leq \frac{1}{n}$, $0 \leq t \leq 1$ be given. Suppose that

$$s \leq u_i v_i, \quad i = 1, \dots, n. \quad (4.3)$$

Then

$$\log \rho(B) \leq s \left(\sum_1^n t \log r_i(B) + (1-t) \log c_i(B) \right) + (1-s) \log \left(\max_{1 \leq i \leq n} t \log r_i(B) + (1-t) \log c_i(B) \right). \quad (4.4)$$

Assume that $0 < t < 1$ and B is fully indecomposable. (PBQ is irreducible for any permutation matrices P, Q .) Then the equality sign holds iff $\frac{B}{r_1(B)}$ is doubly stochastic.

Proof. The characterization of Friedland-Karlin [F-K] claims that

$$\log \rho(B) = \min_{x=(x_1, \dots, x_n), 0 < x_i, i=1, \dots, n} \sum_1^n u_i v_i \log \frac{(Bx)_i}{x_i}. \quad (4.5)$$

By choosing $x = h$ we deduce that

$$\log \rho(B) \leq \sum_1^n u_i v_i \log r_i(B).$$

Replace B by B^T to obtain

$$\log \rho(B) \leq \sum_1^n u_i v_i \log c_i(B).$$

Multiply the first inequality by t the second inequality by $1-t$ and use (4.3) to get (4.4). Suppose that B is fully indecomposable. Then the minimum in (4.5) is achieved only for the eigenvector $x = \alpha u, \alpha > 0$. (Contrary to the claim in [F-K], if B is irreducible but not fully indecomposable then the equality sign in (4.5) can be achieved for vectors x which may not be the eigenvector of B . See [Fri].) As $0 < t < 1$, the equality sign in (4.4) implies that h is the left and the right eigenvector of B . That is $\frac{B}{r_1(B)}$ is a doubly stochastic matrix. In that case we obviously have the equality sign in (4.4). $\diamond$

Remark 4.6 We view (4.4) as a generalization of (3.9). Indeed, for any nonnegative irreducible matrix B (4.4) yields the inequality

$$\rho(B) \leq \nu_t(B), \quad \nu_t(B) = \max_{1 \leq i \leq n} r_i(B)^t c_i(B)^{1-t}, \quad 0 \leq t \leq 1. \quad (4.7)$$

Use a continuity argument to deduce that the above inequality holds for any nonnegative B . It is not hard to show that (3.9) implies (4.7). Of course, in the case B has a zero diagonal (3.9) is equivalent to (4.7).

Remark 4.8 Clearly, if B is reducible we might have the equality sign in the inequality $\rho(B) \leq \nu_{\frac{1}{2}}(B)$ without the requirement that $\frac{A}{r_1(B)}$ is a doubly stochastic matrix. This also may be the case for an irreducible B which is not completely irreducible. Indeed, assume that C is a nonnegative cyclic matrix of even order - $n = 2m$. Suppose that all odd row sums of C and all even column sums of C are equal to $\alpha > 0$ and all even row sums and odd column row sums are equal to $\beta > 0$. As $C^n = \nu_{\frac{1}{2}}(B)^n I$ we deduce that we have the equality sign in $\rho(B) = \nu_{\frac{1}{2}}(B)$. Clearly, for $\alpha \neq \beta$ the matrix $\frac{B}{\alpha}$ is not doubly stochastic.

A generalized tournament matrix T is a nonnegative square matrix satisfying

$$T + T^T = J - I \quad (4.9).$$

See [M-P]. We now apply (4.4) to give an upper bound for the spectral radius of generalized tournament matrices. According to (3.1) any generalized tournament matrix is an almost skew symmetric matrix minus $\frac{I}{2}$. Thus, we can use the inequalities (3.6) – (3.7) for $\rho(T)$. As we pointed out before there is an intrinsic difficulty in applying (3.6) as $\rho_3(\alpha_1^2, \alpha_2^2)$ is a solution of a cubic equation. In what follows we shall deduce upper bounds for $\rho(T)$ using (4.4) provided that $\rho(T)$ is sufficiently big. This condition can be verified by using (3.7).

Let T be a generalized tournament matrix and assume that

$$Tu = \rho(T)u, \quad u = (u_1, \dots, u_n)^T, \quad u_i \geq 0, \quad i = 1, \dots, n, \quad u^T u = 1, \quad \alpha = \frac{\sum_1^n u_i}{n}. \quad (4.10)$$

Combine (4.9) and (4.10) to deduce

$$T^T u = Ju - (\rho(T) + 1)u = n\alpha h - (\rho(T) + 1)u. \quad (4.11)$$

Multiply (4.11) by u^T to deduce:

$$n^2 \alpha^2 = 2\rho(T) + 1. \quad (4.12)$$

It then follows

$$\epsilon = (\epsilon_1, \dots, \epsilon_n)^T = u - \alpha h, \quad \epsilon^T \epsilon = u^T u - 2u^T \alpha h + \alpha^2 h^T h = 1 - n\alpha^2 = 1 - \frac{2\rho(T) + 1}{n}. \quad (4.13)$$

Theorem 4.14. *Let T be an $n \times n$ generalized tournament matrix. Assume that*

$$\rho(T) > \frac{n-2}{2} + \frac{1}{2(n+1)}. \quad (4.15)$$

Then T is an irreducible matrix. Furthermore

$$\begin{aligned} \log \rho(T) \leq & \left(\frac{\sqrt{2\rho(T) + 1}}{n} - \sqrt{1 - \frac{2\rho(T) + 1}{n}} \right)^2 \frac{\sum_1^n \log r_i(T) c_i(T)}{2} + \\ & (1 - n \left(\frac{\sqrt{2\rho(T) + 1}}{n} - \sqrt{1 - \frac{2\rho(T) + 1}{n}} \right)^2) \log \nu_{\frac{1}{2}}(T). \end{aligned} \quad (4.16)$$

Proof. In view of (4.12) and (4.13) we deduce that (4.15) is equivalent to the inequality $\alpha^2 > \epsilon^T \epsilon$. By the definition $\alpha = \frac{u^T h}{n} > 0$. Hence, the inequality (4.15) yields that $0 < u_i$, $i = 1, \dots, n$. Let

$$T^T w = \rho(T)w, \quad w = (w_1, \dots, w_n)^T, \quad 0 \leq w_i, \quad i = 1, \dots, n, \quad w^T w = 1. \quad (4.17)$$

Using the arguments for the eigenvector u we deduce:

$$\alpha = \frac{w^T h}{n}, \quad \delta = (\delta_1, \dots, \delta_n)^T = w - \alpha h, \quad \delta^T \delta = 1 - \frac{2\rho(T) + 1}{n}. \quad (4.18)$$

Thus, the assumption (4.15) yields that w is a vector with strictly positive coordinates. It is well known that if a nonnegative matrix with positive spectral radius has strictly positive left and right eigenvectors then T is a direct sum of irreducible matrices. The assumption (4.9) implies that this sum reduces only to one component, i.e. T is irreducible. We next note that $0 < w^T u \leq (w^T w u^T u)^{\frac{1}{2}} = 1$. Also, the vector v satisfying (4.2) is given by:

$$v = \frac{1}{w^T u} w. \quad (4.19)$$

Therefore

$$u_i v_i \geq (\alpha - (\epsilon^T \epsilon)^{\frac{1}{2}})(\alpha - (\delta^T \delta)^{\frac{1}{2}}) = \left(\frac{\sqrt{2\rho(T) + 1}}{n} - \sqrt{1 - \frac{2\rho(T) + 1}{n}} \right)^2 = s, \quad i = 1, \dots, n. \quad (4.20)$$

Use (4.4) with $t = \frac{1}{2}$ and the above s to deduce (4.16). $\diamond$

Theorem 4.21 *Let T be an $n \times n$ generalized tournament matrix. Assume that*

$$\sqrt{2\rho(T) + 1} = \sqrt{n} \left(1 - \frac{\tau^2}{8n} \right), \quad 0 \leq \tau \leq 1. \quad (4.22)$$

Then

$$\log \rho(T) \leq (1 - \tau) \log \left(\prod_{i=1}^n r_i(T) c_i(T) \right)^{\frac{1}{2n}} + \tau \log \nu_{\frac{1}{2}}(T). \quad (4.23)$$

Proof. The condition (4.22) implies:

$$\frac{2\rho(T) + 1}{n} = 1 - \frac{\tau^2}{4n} + \frac{\tau^4}{64n^2}. \quad (4.24)$$

Hence

$$\sqrt{1 - \frac{2\rho(T) + 1}{n}} \leq \frac{\tau}{2\sqrt{n}}. \quad (4.25)$$

Therefore:

$$\left(\frac{\sqrt{2\rho(T) + 1}}{n} - \sqrt{1 - \frac{2\rho(T) + 1}{n}} \right)^2 \geq \frac{1}{n} \left(1 - \frac{\tau}{2} - \frac{\tau^2}{8n} \right)^2 \geq \frac{1 - \tau}{n}. \quad (4.26)$$

Use the above inequality in (4.16) to obtain (4.23). $\diamond$

For an $n \times n$ generalized tournament matrix T let:

$$\begin{aligned} \bar{r}_i(T) &= r_i(T) - \frac{n-1}{2}, \quad \bar{c}_i(T) = c_i(T) - \frac{n-1}{2} = -\bar{r}_i(T), \\ |\bar{r}_i(T)| &= |\bar{c}_i(T)| \leq \frac{n-1}{2}, \quad i = 1, \dots, n. \end{aligned} \quad (4.27)$$

Then:

$$r_i(T)c_i(T) = \left(\frac{n-1}{2}\right)^2 - \bar{r}_i(T)^2 = \left(\frac{n-1}{2}\right)^2 - \bar{c}_i(T)^2. \quad (4.28)$$

Set:

$$\omega(T) = \left(\prod_{i=1}^n r_i(T)c_i(T)\right)^{\frac{1}{2n}}. \quad (4.29)$$

Combine the standard inequality $\log(1-x) \leq -x$, $x \leq 1$ with (4.28) to deduce

$$\log \omega(T) \leq \log \frac{n-1}{2} - \frac{2}{n(n-1)^2} \sum_{i=1}^n \bar{r}_i(T)^2. \quad (4.30)$$

Corollary 4.31. *Let the assumptions of Theorem 4.21 hold. Then*

$$\log \frac{2\rho(T)}{n-1} \leq -(1-\tau) \frac{2}{n(n-1)^2} \sum_{i=1}^n \bar{r}_i(T)^2. \quad (4.31)$$

We stress again that the inequalities (4.16), (4.23) and (4.31) should be viewed as a refined version of (3.11). (Note that for generalized tournament matrices T $\mu_{\frac{1}{2}}(T) = \nu_{\frac{1}{2}}(T)$.) To apply these inequalities one should first use the lower bound (3.7) which in particular results in an upper bound for τ .

§5. Remarks and open problems

Inequalities (3.17) and (3.21) yield that a matrix T which satisfy (3.15) also satisfy $\rho(T) < \rho^*$, i.e. T is not maximal. The inequality (3.21) implies

$$\frac{n-1}{2} - \frac{1}{2n-1} < \rho_n^*. \quad (5.1)$$

The inequality (3.14) imply

$$\rho_n^* < \frac{n-1}{2} - \frac{1}{4n}. \quad (5.2)$$

Note that for almost regular matrices (3.24) gives a better upper bound than (5.2). Brualdi-Li conjecture implies in particular that the inequality (3.23) holds for all tournament

matrices of an even order n . We conjecture that for any tournament matrix of an even order n we have the inequality:

$$\rho_3(\alpha_1^2, \alpha_2^2) \leq \rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right). \quad (5.3)$$

Here α_1^2, α_2^2 are defined by (3.4). Furthermore, the equality sign holds iff T is permutationally similar to Brualdi-Li matrix.

The inequalities (5.1) – (5.2) imply that

$$\rho_n^* = \frac{n-1}{2} - \frac{K_n}{n}. \quad (5.4)$$

Roughly speaking $\frac{1}{4} \leq K_n \leq \frac{1}{2}$. Conjecture (5.3) yields the inequality $\frac{3}{8} \leq K_n$. It would be interesting to find the asymptotic distribution of the spectral radius of Q_n given by (3.12). A straightforward computation shows that

$$\rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right) - \frac{1}{2} = \frac{n-1}{2} - \frac{3}{8n} + O\left(\frac{1}{n^2}\right). \quad (5.5)$$

Thus, it is possible to estimate the asymptotics of $\rho(Q_n)$ by finding the corresponding value of α_3^2 and evaluating the asymptotics of $\rho_4(\alpha_1^2, \alpha_2^2, \alpha_3^2)$ for the matrix Q_n . We decided to skip this computation here. We conjecture that

$$\lim_{n \rightarrow \infty} K_n = \frac{3}{8}. \quad (5.6)$$

However, $\rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right) - \frac{1}{2} > \rho(T)$ for any almost regular tournament matrix.

Lemma 5.7 *Let $n = 2m$ and assume that T is a generalized tournament matrix such that $r_i(T) = m$, $i = 1, \dots, m$, $r_i(T) = m - 1$, $i = m + 1, \dots, 2m$. Suppose furthermore that $T = (T_{ij})_1^2$, $T_{ij} \in M_m(\mathbf{R})$, $T_{11} = T_{22} = V$ where V is given by (3.12). Then $\rho(T) \leq \rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right) - \frac{1}{2}$. Furthermore, $\rho(T) = \rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right) - \frac{1}{2}$ iff the following conditon hold:*

$$(T - T^T)(w, -w)^T = \tau(v, -v)^T, \quad (5.8)$$

$$v = (1, \dots, 1), w = (w_1, \dots, w_m) \in \mathbf{Z}^m, w_i = m + 1 - 2i, i = 1, \dots, m, \tau = -\frac{2(m^2 - 1)}{3}.$$

There is no almost regular tournament matrix which satisfies $\rho(T) = \rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right) - \frac{1}{2}$.

Proof. Let $2A = T - T^T$, $u = (v, v)^T$ and consider the cyclic subspace V generated by $u, Au, \dots, A^{n-1}u$. As in the proof of Theorem (3.20), we deduce that for any generalized tournament matrix which satisfies the conditions of the theorem one has the equalities $\alpha_1^2 = \frac{1}{4}$, $\alpha_2^2 = \frac{n^2 - 4}{12}$. Theorem 2.30 implies the inequality $\rho(T) \leq \rho_3\left(\frac{1}{4}, \frac{n^2 - 4}{12}\right) - \frac{1}{2}$ and the equality sign holds iff V is three dimensional. Clearly, $u, b = (v, -v)^T \in V$. It is

straightforward to show that Tb and hence Ab in view of (3.25) is spanned by $u, b, x = (w, -w)^T$. Furthermore, $u^T w = b^T w = 0$. Thus V is three dimensional iff Ax is a linear combination of u, b, w . As $x^T Ax = u^T Ax = 0$ it follows that V is three dimensional iff $2Ax = \tau b$. Moreover, $\tau = -\frac{x^T 2Ab}{b^T b}$. A straightforward calculation shows that $\tau = -\frac{2(m^2-1)}{3}$.

Assume that T is an almost regular tournament matrix which satisfies $\rho(T) = \rho_3(\frac{1}{4}, \frac{n^2-4}{12}) - \frac{1}{2}$. From the proof of Theorem 2.6 we obtain that $\text{spec}(T) \supset \text{spec}(F - \frac{I}{2})$, $F \in A_{n-3}(\mathbf{R})$. As $n-3$ is odd we deduce that $0 \in \text{spec}(F)$. That is $-\frac{1}{2} \in \text{spec}(T)$. This is impossible, since all the eigenvalues of T are algebraic integers. $\diamond$

We close the paper by giving some numerical data. For $n = 4, 6, 8, 10, 12$ we computed the quantities

$$\rho_2(\frac{1}{4}) - \frac{1}{2} < \rho(Q_n) < \rho_3(\frac{1}{4}, \frac{n^2-4}{12}) - \frac{1}{2}, \quad \delta = \frac{n}{2}$$

using MATLAB. The matrix Q_n is given by (3.12). Here are the results:

$$\begin{aligned} n = 4 : & \quad 1.3660, \quad 1.3953, \quad 1.3969; \\ n = 6 : & \quad 2.4142, \quad 2.4340, \quad 2.4350; \\ n = 8 : & \quad 3.4365, \quad 3.4513, \quad 3.4521; \\ n = 10 : & \quad 4.4495, \quad 4.4614, \quad 4.4620; \\ n = 12 : & \quad 5.4580, \quad 5.4680, \quad 5.4684. \end{aligned}$$

References

- [Bru] R.A. Brualdi, Matrices, eigenvalues and directed graphs, *Linear and Multilinear Alg.* 11 (1982), 143-165.
- [B-L] R.A. Brualdi and Q. Li, Problem 31, *Discrete Math.* 43 (1983), 329-330.
- [Fri] S. Friedland, Convex spectral functions, *Linear and Multilinear Alg.* 9 (1981), 299-316.
- [F-K] S. Friedland and S. Karlin, Some inequalities for the spectral radius of nonnegative matrices and applications, *Duke Math. J.* 42 (1975), 459-490.
- [G-V] G.H. Golub and C.F. Van Loan, *Matrix computations*, John Hopkins Univ. Press, 2nd edition, 1989.
- [Kat] T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, 1966.
- [Kir] S. Kirkland, Hypertournament matrices, score vectors and eigenvalues, *Linear and Multilinear Alg.*, to appear.
- [M-P] J.S. Maybee and N.J. Pullman, Tournament matrices and their generalizations, I, *Linear and Multilinear Alg.* 28 (1990), 57-70.
- [Moo] J.W. Moon, *Topics on tournaments*, Holt, Rinehart and Winston, 1968.
- [Ost] A. Ostrowski, Über das nichtverschwinden einer klasse von determinanten und die lokalisierung der charakteristischen wurzeln von matrizen, *Compositio Math.* 9 (1951), 209-226.
- [O-S] A. Ostrowski and H. Schneider, Some theorems on the inertia of general matrices, *J. Math. Analysis and Appl.* 4 (1962), 72-84.

Aknowledgements: I would like to thank to R. Brualdi, S. Kirkland, H. Schneider and B. Shader for useful remarks.

Recent IMA Preprints

#	Author/s	Title
818	Avner Friedman and Bei Hu ,	The Stefan problem with kinetic condition at the free boundary
819	M.A. Grinfeld ,	The stress driven instabilities in crystals: mathematical models and physical manifestations
820	Bei Hu and Lihe Wang ,	A free boundary problem arising in electrophotography: solutions with connected toner region
821	Yongzhi Xu, T. Craig Poling, and Trent Brundage ,	Direct and inverse scattering of time harmonic acoustic waves in an inhomogeneous shallow ocean
822	Steven J. Altschuler ,	Singularities of the curve shrinking flow for space curves
823	Steven J. Altschuler and Matthew A. Grayson ,	Shortening space curves and flow through singularities
824	Tong Li ,	On the Riemann problem of a combustion model
825	L.A. Peletier & W.C. Troy ,	Self-similar solutions for diffusion in semiconductors
826	C.J. van Duijn, L.A. Peletier & R.J. Schotting ,	On the analysis of brine transport in porous media
827	Minkyu Kwak ,	Finite dimensional description of convective reaction-diffusion equations
828	Minkyu Kwak ,	Finite dimensional inertial forms for the 2D Navier–Stokes equations
829	Victor A. Galaktionov and Sergey A. Posashkov ,	On some monotonicity in time properties for a quasilinear parabolic equation with source
830	Victor A. Galaktionov ,	Remark on the fast diffusion equation in a ball
831	Hi Jun Choe and Lihe Wang ,	A regularity theory for degenerate vector valued variational inequalities
832	Vladimir I. Oliker and Nina N. Uraltseva ,	Evolution of nonparametric surfaces with speed depending on curvature, II. The mean curvature case.
833	S. Kamin and W. Liu ,	Large time behavior of a nonlinear diffusion equation with a source
834	Shoshana Kamin and Juan Luis Vazquez ,	Singular solutions of some nonlinear parabolic equations
835	Bernhard Kawohl and Robert Kersner ,	On degenerate diffusion with very strong absorption
836	Avner Friedman and Fernando Reitich ,	Parameter identification in reaction-diffusion models
837	E.G. Kalnins, H.L. Manocha and Willard Miller, Jr. ,	Models of q -algebra representations I. Tensor products of special unitary and oscillator algebras
838	Robert J. Sacker and George R. Sell ,	Dichotomies for linear evolutionary equations in Banach spaces
839	Oscar P. Bruno and Fernando Reitich ,	Numerical solution of diffraction problems: a method of variation of boundaries
840	Oscar P. Bruno and Fernando Reitich ,	Solution of a boundary value problem for Helmholtz equation via variation of the boundary into the complex domain
841	Victor A. Galaktionov and Juan L. Vazquez ,	Asymptotic behaviour for an equation of superslow diffusion. The Cauchy problem
842	Josephus Hulshof and Juan Luis Vazquez ,	The Dipole solution for the porous medium equation in several space dimensions
843	Shoshana Kamin and Juan Luis Vazquez ,	The propagation of turbulent bursts
844	Miguel Escobedo, Juan Luis Vazquez and Enrike Zuazua ,	Source-type solutions and asymptotic behaviour for a diffusion-convection equation
845	Marco Biroli and Umberto Mosco ,	Discontinuous media and Dirichlet forms of diffusion type
846	Stathis Filippas and Jong-Shenq Guo ,	Quenching profiles for one-dimensional semilinear heat equations
847	H. Scott Dumas ,	A Nekhoroshev-like theory of classical particle channeling in perfect crystals
848	R. Natalini and A. Tesei ,	On a class of perturbed conservation laws
849	Paul K. Newton and Shinya Watanabe ,	The geometry of nonlinear Schrödinger standing waves
850	S.S. Sritharan ,	On the nonsmooth verification technique for the dynamic programming of viscous flow
851	Mario Taboada and Yuncheng You ,	Global attractor, inertial manifolds and stabilization of nonlinear damped beam equations
852	Shigeru Sakaguchi ,	Critical points of solutions to the obstacle problem in the plane
853	F. Abergel, D. Hilhorst and F. Issard-Roch ,	On a dissolution-growth problem with surface tension in the neighborhood of a stationary solution
854	Erasmus Langer ,	Numerical simulation of MOS transistors
855	Haïm Brezis and Shoshana Kamin ,	Sublinear elliptic equations in $\mathbb{R}^n$
856	Johannes C.C. Nitsche ,	Boundary value problems for variational integrals involving surface curvatures
857	Chao–Nien Chen ,	Multiple solutions for a semilinear elliptic equation on $\mathbb{R}^N$ with nonlinear dependence on the gradient
858	D. Brochet, X. Chen and D. Hilhorst ,	Finite dimensional exponential attractor for the phase field model
859	Joseph D. Fehribach ,	Mullins-Sekerka stability analysis for melting-freezing waves in helium-4
860	Walter Schempp ,	Quantum holography and neurocomputer architectures
861	D.V. Anosov ,	An introduction to Hilbert’s 21st problem
862	Herbert E Huppert and M Grae Worster ,	Vigorous motions in magma chambers and lava lakes

- 863 **Robert L. Pego and Michael I. Weinstein**, A class of eigenvalue problems, with applications to
instability of solitary waves
- 864 **Mahmoud Affouf**, Numerical study of a singular system of conservation laws arising in enhanced oil
reservoirs
- 865 **Darin Beigie, Anthony Leonard and Stephen Wiggins**, The dynamics associated with the chaotic
of tangles two dimensional quasiperiodic vector fields: theory and applications
- 866 **Gui-Qiang Chen and Tai-Ping Liu**, Zero relaxation and dissipation limits for hyperbolic conservation laws
- 867 **Gui-Qiang Chen and Jian-Guo Liu**, Convergence of second-order schemes for isentropic gas dynamics
- 868 **Aleksander M. Simon and Zbigniew J. Grzywna**, On the Larché-Cahn theory for stress-induced diffusion
- 869 **Jerzy Łuczka, Adam Gadomski and Zbigniew J. Grzywna**, Growth driven by diffusion
- 870 **Mitchell Luskin and Tsorng-Whay Pan**, Nonplanar shear flows for nonaligning nematic liquid crystals
- 871 **Mahmoud Affouf**, Unique global solutions of initial-boundary value problems for thermodynamic phase transitions
- 872 **Richard A. Brualdi, Keith L. Chavey and Bryan L. Shader**, Rectangular L -matrices
- 873 **Xinfu Chen, Avner Friedman and Bei Hu**, The thermistor problem with zero-one conductivity II
- 874 **Raoul LePage**, Controlling a diffusion toward a large goal and the Kelly principle
- 875 **Raoul LePage**, Controlling for optimum growth with time dependent returns
- 876 **Marc Hallin and Madan L. Puri**, Rank tests for time series analysis a survey
- 877 **V.A. Solonnikov**, Solvability of an evolution problem of thermocapillary convection in an infinite time interval
- 878 **Horia I. Ene and Bogdan Vernescu**, Viscosity dependent behaviour of viscoelastic porous media
- 879 **Kaushik Bhattacharya**, Self-accommodation in martensite
- 880 **D. Lewis, T. Ratiu, J.C. Simo and J.E. Marsden**, The heavy top: a geometric treatment
- 881 **Leonid V. Kalachev**, Some applications of asymptotic methods in semiconductor device modeling
- 882 **David C. Dobson**, Phase reconstruction via nonlinear least-squares
- 883 **Patricio Aviles and Yoshikazu Giga**, Minimal currents, geodesics and relaxation of variational integrals on
mappings of bounded variation
- 884 **Patricio Aviles and Yoshikazu Giga**, Partial regularity of least gradient mappings
- 885 **Charles R. Johnson and Michael Lundquist**, Operator matrices with chordal inverse patterns
- 886 **B.J. Bayly**, Infinitely conducting dynamos and other horrible eigenproblems
- 887 **Charles M. Elliott and Stefan Luckhaus**, 'A generalised diffusion equation for phase separation of a
multi-component mixture with interfacial free energy'
- 888 **Christian Schmeiser and Andreas Unterreiter**, The derivation of analytic device models by asymptotic methods
- 889 **LeRoy B. Beasley and Norman J. Pullman**, Linear operators that strongly preserve the index of imprimitivity
- 890 **Jerry Donato**, The Boltzmann equation with lie and cartan
- 891 **Thomas R. Hoffend Jr., Peter Smereka and Roger J. Anderson**, A method for resolving the laser induced
local heating of moving magneto-optical recording media
- 892 **E.G. Kalnins, Willard Miller, Jr. and Sanchita Mukherjee**, Models of q -algebra representations:
the group of plane motions
- 893 **T.R. Hoffend Jr. and R.K. Kaul**, Relativistic theory of superpotentials for a nonhomogeneous, spatially
isotropic medium
- 894 **Reinhold von Schwerin**, Two metal deposition on a microdisk electrode
- 895 **Vladimir I. Oliker and Nina N. Uraltseva**, Evolution of nonparametric surfaces with speed depending on
curvature, III. Some remarks on mean curvature and anisotropic flows
- 896 **Wayne Barrett, Charles R. Johnson, Raphael Loewy and Tamir Shalom**, Rank incrementation via
diagonal perturbations
- 897 **Thierry Dombre, Alain Pumir, and Eric D. Siggia**, On the interface dynamics for convection in porous media
- 898 **Mingxiang Chen, Xu-Yan Chen and Jack K. Hale**, Structural stability for time-periodic one-dimensional
parabolic equations
- 899 **Hong-Ming Yin**, Global solutions of Maxwell's equations in an electromagnetic field with the temperature-
dependent electrical conductivity
- 900 **Robert Grone, Russell Merris and William Watkins**, Laplacian unimodular equivalence of graphs
- 901 **Miroslav Fiedler**, Structure-ranks of matrices
- 902 **Miroslav Fiedler**, An estimate for the nonstochastic eigenvalues of doubly stochastic matrices
- 903 **Miroslav Fiedler**, Remarks on eigenvalues of Hankel matrices
- 904 **Charles R. Johnson, D.D. Olesky, Michael Tsatsomeros and P. van den Driessche**, Spectra with positive
elementary symmetric functions
- 905 **Pierre-Alain Gremaud**, Thermal contraction as a free boundary problem
- 906 **K.L. Cooke, Janos Turi and Gregg Turner**, Stabilization of hybrid systems in the presence of feedback delays
- 907 **Robert P. Gilbert and Yongzhi Xu**, A numerical transmutation approach for underwater sound propagation
- 908 **LeRoy B. Beasley, Richard A. Brualdi and Bryan L. Shader**, Combinatorial orthogonality
- 909 **Richard A. Brualdi and Bryan L. Shader**, Strong hall matrices
- 910 **Håkan Wennerström and David M. Anderson**, Difference versus Gaussian curvature energies; monolayer versus
bilayer curvature energies applications to vesicle stability
- 911 **Shmuel Friedland**, Eigenvalues of almost skew symmetric matrices and tournament matrices