LARGE GAMES: FAIR AND STABLE OUTCOMES

BY

MYRNA HOLTZ WOODERS

AND

WILLIAM R. ZAME

IMA Preprint Series # 92

August 1984

INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
UNIVERSITY OF MINNESOTA
514 Vincent Hall
206 Church Street S.E.
Minneapolis, Minnesota 55455

#	Author(s)	Title	#	Author(s)	Title
1	Workshop Summaries from the September 1982 workshop on Statistical Mechanics, Dynamical Systems and Turbulence		25	Scott J. Spector,	On the Absence of Bifurcation for Elastic Bars in Uniaxial Tension
2	Raphael De laLlave,	A Simple Proof of C. Siegel's Center Theorem	26	W.A. Coppel,	Maps on an Interval
3	H. Simpson, S. Spector,	On Copositive Matrices and Strong Ellipticity for Isotropic Elastic Materials	27	James Kirkwood,	Phase Transitions in the Ising Model with Traverse Field
4	George R. Sell,	Vector Fields in the Vicinity of a Compact Invariant Manifold	28	Luis Magalhães,	The Asymptotics of Solutions of Singularly Perturbed Functional Differential Equations: and Concentrated Delays are Different
5	Milan Miklavcic,	Non-linear Stability of Asymptotic Suction	29	Charles Tresser,	Homoclinic Orbits for Flow in R^3
6	Hans Weinberger,	A Simple System with a Continuum of Stable Inhomogeneous Steady States	30	Charles Tresser,	About some Theorems by L.P. Sil'nikov
7	Bau-Sen Du,	Period 3 Bifurcation for the Logistic Mapping	31	Michael Alzenmann,	On the Renormalized Coupling Constant and the Susceptibility in ϕ^4 Field Theory and the Ising Model in Four Dimensions
8	Hans Weinberger,	Optimal Numerical Approximation of a Linear Operator	32	C. Eugene Wayne,	The KAM Theory of Systems with Short Range Interactions I
9	L.R. Angel, D.F. Evans, B. Minham,	Three Component Ionic Microemulsions	33	M. Slemrod, J. E. Marsden,	Spatial Chaos in a Van der Waals Fluid due to Periodic Thermal Fluctuations
10	D.F. Evans, D. Mitchell, S. Mukherjee, B. Minham,	Surfactant Diffusion; New Results and Interpretations	34	J. Kirkwood, C.E. Wayne,	Percolation in Continuous Systems
11	Leif Arkeryd,	A Remark about the Final Aperiodic Regime for Maps on the Interval	35	Luis Magalhães,	Invariant Manifolds for Functional Differential Equations Close to Ordinary Differential Equations
12	Luis Magalhães,	Manifolds of Global Solutions of Functional Differential Equations	36	C. Eugene Wayne,	The KAM Theory of Systems with Short Range Interactions II
13	Kenneth Meyer,	Tori in Resonance	37	Jean De Canniere,	Passive Quasi-Free States of the Noninteracting Fermi Gas
14	C. Eugene Wayne,	Surface Models with Nonlocal Potentials: Upper Bounds	38	Elias C. Alfántis,	Maxwell and van der Waals Revisited
15	K.A. Pericak-Spector,	On Stability and Uniqueness of Fluid Flow Through a Rigid Porous Medium	39	Elias C. Alfántis,	On the Mechanics of Modulated Structures
16	George R. Sell,	Smooth Linearization Near a Fixed Point	40	William Ruckle,	The Strong ϕ Topology on Symmetric Sequence Spaces
17	David Wolkind,	A Nonlinear Stability Analysis of a Model Equation for Alloy Solidification	41	Charles R. Johnson,	A Characterization of Borda's Rule Via Optimization
18	Pierre Collet,	Local C^∞ Conjugacy on the Julia Set for some Holomorphic Perturbations of $z \rightarrow z$	42	Hans Meinberger,	Kazuo Kishimoto, The Spatial Homogeneity of Stable Equilibria of Some Reaction-Diffusion Systems on Convex Domains
19	Henry C. Simpson,	Scott J. Spector, On the Modified Bessel Functions of the First Kind On Barrelling for a Material in Finite Elasticity	43	K.A. Pericak-Spector,	W.O. Williams, On Work and Constraints in Mixtures
20	George R. Sell,	Linearization and Global Dynamics	44	H. Rosenberg, E. Toubiana,	Some Remarks on Deformations of Minimal Surfaces
21	P. Constantin, C. Fofas,	Global Lyapunov Exponents, Kaplan-Yorke Formulas and the Dimension of the Attractors for 2D Navier-Stokes Equations	45	Stephan Pellikan,	The Duration of Transients
22	Milan Miklavcic,	Stability for Semilinear Parabolic Equations with Noninvertible Linear Operator	46	V. Capasso, K.L. Cooke, M. Witten,	Random Fluctuations of the Duration of Harvest
23	P. Collet, H. Epstein, G. Gallavotti,	Perturbations of Geodesic Flows on Surfaces of Constant Negative Curvature and their Mixing Properties	47	E. Fabes, D. Stookey,	The L^p -Intergrability of Green's Functions and Fundamental Solutions for Elliptic and Parabolic Equations
24	J.E. Dunn, J. Serrin,	On the Thermodynamics of Interstitial working	48	H. Brezis,	Semilinear Equations in R^N without conditions at infinity
			49	M. Slemrod,	Lax-Friedrichs and the Viscosity-Capillarity Criterion

Recent IMA Preprints (continued)		
#	Author(s)	Title
50	C. Johnson, W. Barrett,	Spanning Tree Extensions of the Hadamard-Fischer Inequalities
51	Andrew Postlewaite, David Schmiedler,	Revelation and Implementation under Differential Information
52	Paul Blanchard,	Complex Analytic Dynamics on the Riemann Sphere
53	G. Levitt, H. Rosenberg,	Topology and Differentiability of Labyrinths in the Disc and Annulus
54	G. Levitt, H. Rosenberg,	Symmetry of Constant Mean Curvature Hyper-surfaces in Hyperbolic Space
55	Ennio Stacchetti,	Analysis of a Dynamic, Decentralized Exchange Economy
56	Henry Simpson, Scott Spector,	On Failure of the Complementing Condition and Nonuniqueness in Linear Elastostatics
57	Craig Tracy,	Complete Integrability in Statistical Mechanics and the Yang-Baxter Equations
58	Tongren Ding,	Boundedness of Solutions of Duffing's Equation
59		Abstracts for the Workshop on Price Adjustment, Quantity Adjustment, and Business Cycles
60	Rafael Rob,	The Coase Theorem an Informational Perspective
61	Joseph Jerome,	Approximate Newton Methods and Homotopy for Stationary Operator Equations
62	Rafael Rob,	A Note on Competitive Bidding with Asymmetric Information
63	Rafael Rob,	Equilibrium Price Distributions
64	William Rucklë,	The Linearization Projection, Global Theories
65	Russell Johnson, Kenneth Palmer, George R. Sell,	Ergodic Properties of Linear Dynamical Systems
66	Stanley Reiter,	How a Network of Processors can Schedule Its Work
67	R.N. Goldman, D.C. Heath,	Linear Subdivision is Strictly a Polynomial Phenomenon
68	R. Giachetti, R. Johnson,	The Floquet Exponent for Two-dimensional Linear Systems with Bounded Coefficients
69	S. Williams,	Realization and Nash Implementation: Two Aspects of Mechanism Design
70	S. Williams,	Sufficient Conditions for Nash Implementation
71	Nicholas Yannelis, William R. Zame,	Equilibria in Banach Lattices Without Ordered Preferences
72	W. Harris, Y. Sibuya,	The Reciprocals of Solutions of Linear Ordinary Differential Equations
73	S. Pelikan,	A Dynamical Meaning of Fractal Dimension
74	D. Heath, W. Sudderth,	Continuous-Time Portfolio Management: Minimizing the Expected Time to Reach A G
75	J.S. Jordan,	Information Flows Intrinsic to the Stability Economic Equilibrium
76	J. Jerome,	An Adaptive Newton Algorithm Based on Numerical Inversion: Regularization Post Condition
77	David Schmiedler,	Integral Representation Without Additivity
78		Abstracts for the Workshop on Bayesian Analysis in Economics and Game Theory
79	G. Chichilnisky, G.M.Heal,	Existence of a Competitive Equilibrium in L_p and Sobolev Spaces
80	Thomas Seidman,	Time-dependent solutions of a Nonlinear system in Semiconductory Theory, II: Boundedness and Periodicity
81	Yakar Kannal,	Engaging in R&D and the Emergence of Expected Non-convex Technologies
82	Herve Moulin,	Choice Functions over a finite set: A Summary
83	Herve Moulin,	Choosing from a Tournament
84	David Schmiedler,	Subjective Probability and Expected Utility without Additivity
85	I.G. Kevrekidis, R. Aris, L.D. Schmidt, and S. Pelikan,	The Numerical Computation of Invariant Circles of Maps
86	F. William Lawvere,	State Categories, Closed Categories, and the Existence Semi-Continuous Entropy Functions
87	F. William Lawvere,	Functional Remarks on the General Concept of Chaos
88	Steven R. Williams,	Necessary and Sufficient Conditions for the Existence of a Locally Stable Message Process
89	Steven R. Williams,	Implementing a Generic Smooth Function
90	Dilip Abreu,	Infinitely Repeated Games with Discounting: A General Theory
91	J.S. Jordan,	Instability in the Implementation of Walrasian Allocations

 

LARGE GAMES: FAIR AND STABLE OUTCOMES

by

Myrna Holtz Wooders^{*}

and

William R. Zame^{**}

^{*} Department of Economics, University of Toronto

^{**} Department of Mathematics, State University of New York at Buffalo

ACKNOWLEDGMENTS

We are indebted to Robert Aumann, John Chapman, Yakar Kannai and Hans Weinberger for several helpful discussions.

This work was done at the Institute for Mathematics and its Applications, University of Minnesota. The hospitality and financial support of the I.M.A. are gratefully acknowledged.

We are also grateful for financial support to the Social Sciences and Humanities Research Council, Canada (Wooders) and the National Science Foundation, U.S.A. (Zame).

1. INTRODUCTION

Kuhn and Tucker (1950) posed, as one of a list of important problems for study, "to establish significant asymptotic properties of n -person games, for large n ." This paper takes a step in that direction.

We establish that for a broad class of large games, fair outcomes are nearly stable. Moreover, the larger the game, the more nearly stable is the fair outcome. In more precise terms, the Shapley value of a large game is in the ϵ -core, and ϵ is very small if the game is very large. The framework we use is that of games in characteristic function form with side payments. This framework can accommodate diverse economic models with money, including private goods exchange economies (with divisible and indivisible goods), coalition-production economies, economies with local public goods, and economies with club goods.

The convergence of the Shapley value to the competitive payoff (and hence to the core) for private goods exchange economies was suggested by Shubik, and first demonstrated by Shapley (1964) in the context of replication economies with money. It has subsequently been extended by Shapley and Shubik (1969), Champsaur (1975), Mas-Colell (1976), Cheng (1977) and others.* Some of these extensions treat the case of economies without money, but they are all restricted to the context of private goods exchange economies** with divisible goods and concave, monotone utility functions. Moreover, they all treat either replicated sequences of economies, or convergent sequences of economies.*** This kind of private goods framework, however, rules out many natural economic

*This list is not intended to be complete.

**Strictly speaking, Champsaur allows for production, but only of a restricted kind (additive production).

***The work of Aumann and Shapley (1974), Aumann (1975), Hart (1977) and others on values of economies with a continuum of agents is of a somewhat different nature since it begins with a limiting continuum economy.

phenomena, such as indivisible goods, nonmonotonic or nonconcave utility functions, general production, local public goods, or club goods. Our framework is broad enough to encompass all these situations (in the presence of money^{*}). Moreover, our framework does not require replicated economies or convergent sequences of economies.

To model large games we introduce the notion of a "technology" (to be understood in a broad sense). A technology encompasses all possible opportunities available to any conceivable group of agents in society. More specifically, a technology consists of a space Ω of possible attributes of agents, and a mapping Λ which specifies the maximum utility obtainable by any group of agents, given the attributes of its members. We make natural assumptions of compactness of the space Ω of attributes, and superadditivity, continuity and marginal boundedness of the mapping Λ . A game is determined from such a technology by specifying a finite set of players, and the attributes of each player. (In a private goods economy, for example, we would specify, for each agent, an initial endowment and a utility function.)

The stability concept we employ is the individually rational ϵ -core. A payoff is in this ϵ -core if it is feasible, Pareto optimal, individually rational, and has the property that no coalition can improve upon it by more than ϵ for each of its members.

Our main result shows that, given an $\epsilon > 0$, any game, derived from such a technology, which is sufficiently large (in the sense of having many players whose attributes are sufficiently close) has the property that the Shapley value (a fair outcome) is in the individually rational ϵ -core (the set of nearly stable outcomes).

In the special case of private goods exchange economies with divisible goods and money (but allowing for nonconcave and nonmonotone utility functions),

^{*} Without money, such economic settings may be modelled as games without side-payments. In that framework, the analog of the Shapley value is the NTU value (or λ -transfer value). We have some positive, but fragmentary results in this direction.

such ϵ -core payoffs can be "approximately decentralized" by prices (in the sense of Anderson (1978)). In this setting, therefore, our result implies that the Shapley value is an approximately competitive payoff (if the economy is sufficiently large). We stress, however, that this is a very special case of our result.

Section 2 gives some examples which motivate our work and indicate the generality of our framework. We collect general information about game theory in Section 3, and describe our notion of technologies in Section 4. In Section 5 we give a precise statement of our main result and some comments on its meaning, and describe the two main ideas of its proof. The proof itself is divided into four sections: Section 6 is devoted to the first main idea, Section 7 contains a probabilistic estimate, Section 8 is devoted to the second main idea, and the proof is completed in Section 9. Finally, Section 10 relates the ϵ -cores of a sequence of games to the core of the nonatomic limiting game.

This work is, in part, an outgrowth of an earlier paper (Wooders and Zame (1984)), which established the nonemptiness of the ϵ -cores of large games. There is an overlap of ideas between the two papers, but the thrust is quite different and we do not assume here any familiarity with Wooders and Zame (1984). In particular, the present paper may be read quite independently.

2. EXAMPLES

In order to motivate our framework and main result, and to indicate the wide variety of contexts to which they apply, we give several economic examples. Since our purpose is largely illustrative, we will be somewhat informal in our discussions and not go into detail in any of the examples; in particular, we will not give any numerical calculations.*

We begin with a simple economy with a local public good.

EXAMPLE 1: We consider an economy with two types of agents: entertainers and spectators. Within types, the agents are identical. The only activity of this economy is the presentation of performances by a group of entertainers for an audience of spectators. A group of entertainers can perform for an audience of any size, but the quality of such a performance will suffer if the audience is too large (because of crowding). We assume there are no interactions between entertainers performing for the same audience. Thus the value of a performance by a group of m entertainers to an audience of n spectators may reasonably be assumed to have an expression of the form $v(m,n) = mF(n)$, where F is a function which is initially increasing but eventually decreasing.** We assume that this value is expressed in monetary terms and money is freely transferable among all agents.

In this economic setting, our Theorem 1 guarantees that the Shapley value (of the associated game) is in the ϵ -core provided the economy has enough entertainers and enough spectators.

For our second example, we treat a simple coalition-production economy.

*A simple numerical example of the nonemptiness of the ϵ -core for a coalition-production economy is worked out in Wooders and Zame (1984).

**If $m \geq 2$ and n is large, it may be sensible for the group of m entertainers and n spectators to divide into smaller groups.

EXAMPLE 2: We consider an economy whose sole activity is the production of a single divisible good, call it rope. The rope produced is to be sold on world markets at a fixed price of p dollars per foot. Agents in the economy are characterized by their skill level, on a scale of 0 to 1. A group of n agents with skill levels $s_1, \dots, s_n$ can produce rope according to the cost function

$$c(q) = K + A(s, n)q^2,$$

where $c(q)$ is the cost of producing q feet of rope, K is a fixed cost, and $s = s_1 + \dots + s_n$ is the total skill level of the group. Evidently, A is an inverse measure of efficiency, so it seems reasonable to assume that A is continuous and decreasing as a function of s (groups which are more skilled are more efficient) and is eventually increasing as a function of n (due to congestion and management costs, for example). We assume that $A(s, n)$ is positive and bounded away from zero (no free production). It is routine to verify that the maximal profit obtainable by a coalition of n agents with total skill level s is

$$P(s, n) = \frac{p^2}{2A(s, n)} - K.$$

At issue in this economy is the efficient division of agents into productive coalitions and the division among all agents of the total profits. Theorem 1 tells us that the Shapley value, which is a fair division of the optimal profits, is nearly stable, in the sense that no group could do very much better, provided the number of agents is sufficiently large and the skill levels are sufficiently dispersed.

For our final example, we describe a rather general private goods exchange economy with money.

EXAMPLE 3: We consider a private goods exchange economy with m divisible goods, ℓ indivisible goods, and money. Divisible goods are available in arbitrary nonnegative amounts, while indivisible goods are available only in nonnegative integral amounts. An allocation for an agent is thus a vector (x, y) in $\mathbb{R}_+^m \times \mathbb{Z}_+^\ell$.

and we take $\mathbb{R}_+^m \times \mathbb{Z}_+^\ell$ as the consumption set for each agent. We assume that the initial endowments of all agents lie in some compact set $E \subset \mathbb{R}_+^m \times \mathbb{Z}_+^\ell$ (but we do not exclude the possibility that some agents have zero initial endowment of some or all goods).

We let $C_+^b(\mathbb{R}_+^m \times \mathbb{Z}_+^\ell)$ be the space of bounded, continuous functions from $\mathbb{R}_+^m \times \mathbb{Z}_+^\ell$ to $\mathbb{R}_+$. This is a complete metric space when equipped with the distance function

$$\text{dist}(u_1, u_2) = \sup \{ |u_1(x, y) - u_2(x, y)| : (x, y) \in \mathbb{R}_+^m \times \mathbb{Z}_+^\ell \} .$$

We assume that the utility functions of all agents lie in some compact set $\Phi \subset C_+^b(\mathbb{R}_+^m \times \mathbb{Z}_+^\ell)$. We assume in addition that Φ is a uniformly Lipschitz family, i. e., there is a constant C such that

$$|u(x_1, y_1) - u(x_2, y_2)| \leq C \text{dist}((x_1, y_1), (x_2, y_2)) ,$$

for each (x_1, y_1) and (x_2, y_2) in $\mathbb{R}_+^m \times \mathbb{Z}_+^\ell$ and each u in Φ .^{*}

Finally, we assume that utility is (linearly) transferable between agents, in the form of monetary payments.^{**}

In this framework, to specify completely an exchange economy with n agents we need to specify for each agent i an initial endowment $e_i \in E$ and a utility function $u_i \in \Phi$. Theorem 1 guarantees that, given an $\epsilon > 0$, if the number of agents is sufficiently large and their endowments and utility functions are sufficiently dispersed in $E \times \Phi$, then the Shapley value for the economy is in the ϵ -core.

A few remarks may help to put this example into perspective vis-a-vis the existing literature.

^{*}See Remark 1 following this example for comments on these assumptions.

^{**}More formally, we could, in the usual way, introduce money explicitly, and assume that each agent's utility for an allocation (x, y, ξ) of goods and money is of the form $u(x, y) + \xi$.

REMARK 1: We have assumed that utility functions are bounded and Lipschitz. These assumptions are parallel to, but weaker than, the assumptions made by Mas-Colell (1977). On the other hand, Shapley (1964) and Champsaur (1975) do not assume that utility functions are bounded or Lipschitz. However, these authors consider only replica economies and concave utility functions. In the replication framework, concave utility functions may always be replaced by suitable modifications which are bounded and Lipschitz without altering the Shapley value or the ϵ -core payoffs.

REMARK 2: In the framework we have described, competitive equilibria need not exist. However, at least in the absence of indivisible goods, the work of Anderson (1978) may be used to show that allocations corresponding to payoffs in the ϵ -core may be approximately decentralized by prices. Hence in this context, we can conclude that the Shapley value is a nearly competitive payoff, provided the number of agents is large and endowments and utility functions are sufficiently dispersed. We point out that no assumptions need be made about the relative distribution of endowments or utility functions. All the other value convergence theorems of which we are aware do make assumptions on the distributions of endowments and utility functions: Shapley (1964) and Champsaur (1975) treat replica economies, for example, while Mas-Colell (1977) treats convergent sequences of economies.

3. GAMES

By a game (in characteristic function form, with sidepayments) we mean a pair (N, v) where N is a finite set (the set of players) and v is a function (the characteristic function) from the set 2^N of subsets of N to the set $\mathbb{R}_+$ of non-negative real numbers, with the property that $v(\emptyset) = 0$. We usually refer to subsets S of N as coalitions; the number $v(S)$ is the worth of S . If the player set N is understood, we frequently refer to v itself as the game. We say that v is superadditive if for all disjoint subsets S, S' of N we have

$$v(S \cup S') \geq v(S) + v(S') .$$

By a payoff for (N, v) we mean a vector x in $\mathbb{R}^N$; it is convenient to use functional notation, so for $i \in N$, $x(i)$ is the i -th component of x . We say that x is feasible if $x(N) \leq v(N)$ (where $x(S) = \sum_{i \in S} x(i)$ for each $S \subset N$).

For $\epsilon \geq 0$, a feasible payoff x is in the ϵ -core of (N, v) if

- (a) $x(N) = v(N)$ (Pareto optimality);
- (b) $x(S) \geq v(S) - \epsilon |S|$ for all subsets S of N .

(We use $|S|$ to denote the number of elements of the set S .) We say x is in the individually rational ϵ -core if it is in the ϵ -core and in addition

- (c) $x(i) \geq v(\{i\})$ for all $i \in N$ (individual rationality).

When $\epsilon = 0$, the ϵ -core (which coincides with the individually rational ϵ -core) is simply the core.

Less formally, a feasible, Pareto optimal payoff x belongs to the ϵ -core of (N, v) if no group of players can guarantee for themselves a payoff which each of them finds better than x by more than ϵ . As Shapley and Shubik (1966) point out, such a payoff can be interpreted as stable if players are nearly optimizing, or satisficing; or if there is an organizational or communicational cost to formation of coalitions (proportional to the size of the coalition).

The Shapley value $Sh(v)$ of the game (N, v) is the payoff whose i -th component is given by

$$\text{Sh}(v, i) = \frac{1}{|N|} \sum_{J=0}^{|N|-1} \frac{1}{\binom{|N|-1}{J}} \sum_{\substack{S \subset N \setminus \{i\} \\ |S|=J}} [v(S \cup \{i\}) - v(S)]$$

In other words, $\text{Sh}(v, i)$ is player i 's average marginal contribution to coalitions in N . The Shapley value is a feasible, Pareto optimal and individually rational payoff. It is frequently interpreted as representing a "fair" payoff since it yields to each player his expected contribution. It can also be given various other interpretations (as a von Neumann-Morgenstern utility function for example, see Roth (1977)).

4. TECHNOLOGIES

We want to formalize the notion of a large game for which the worth of a coalition depends in a continuous fashion on the attributes of its members. To do this, we introduce the notion of a technology.*

Let Ω be a compact metric space. By a (positive) divisor** on Ω we mean a function f from Ω to the set $\mathbb{Z}_+$ of nonnegative integers for which the support of f ,

$$\text{support}(f) = \{\omega \in \Omega : f(\omega) \neq 0\} ,$$

is finite. We denote the set of divisors on Ω by $D(\Omega)$. Note that the sum of divisors (defined pointwise) is a divisor, and that the product of a divisor with a nonnegative integer is a divisor. We write 0 for the divisor which is identically zero. We write $f \leq g$ if $f(\omega) \leq g(\omega)$ for each $\omega \in \Omega$. For ω_0 a point of Ω , we write χ_{ω_0} for the divisor given by

$$\begin{aligned} \chi_{\omega_0}(\omega) &= 0 \quad \text{if } \omega \neq \omega_0 ; \\ \chi_{\omega_0}(\omega_0) &= 1 . \end{aligned}$$

By the norm of a divisor f we mean

$$\|f\| = \sum_{\omega \in \Omega} f(\omega) .$$

(Notice that this is a finite sum, since f has finite support.)

In essence, a divisor is simply an (unordered) list of elements of Ω , with each element ω appearing as many times as its multiplicity $f(\omega)$.

By a technology we mean a pair (Ω, Λ) where Ω is a compact metric space (the space of attributes) and $\Lambda : D(\Omega) \rightarrow \mathbb{R}_+$ is a function with the following properties:

*In Wooders and Zame (1984) the term "pre-game with attributes" was used; the term "technology" was suggested by Y. Kannai, and seems much more descriptive.

**This is the usual mathematical term.

- (i) $\Lambda(0) = 0$;
- (ii) $\Lambda(f+g) \geq \Lambda(f) + \Lambda(g)$ (superadditivity) ;
- (iii) there is a constant M such that $\Lambda(f+\chi_\omega) \leq \Lambda(f) + M$ for each $\omega \in \Omega$, $f \in D(\Omega)$ (we say M is an individual marginal bound);
- (iv) for every $\epsilon > 0$ there is a $\delta > 0$ such that

$$\left| \Lambda(f+\chi_{\omega_1}) - \Lambda(f+\chi_{\omega_2}) \right| < \epsilon$$

whenever $f \in D(\Omega)$ and $\omega_1, \omega_2 \in \Omega$ with $\text{dist}(\omega_1, \omega_2) < \delta$ (continuity).

The interpretation we have in mind is that a technology encompasses all the economic possibilities for every conceivable group of agents. A point of Ω represents a complete description of the relevant attributes of an agent (endowment, utility function, etc.). A divisor f represents a group of agents, of whom $f(\omega)$ are described by the attribute ω ; the total number of agents in the group is just $\|f\|$. The number $\Lambda(f)$ represents the maximal possible payoff the members of this group could achieve (using their own resources) by cooperation. The requirement that $\Lambda(0) = 0$ means that the group of no agents can achieve nothing. Superadditivity has its usual interpretation: one of the possibilities open to the group represented by $f+g$ is to split into the groups represented by f and by g , and share the proceeds. (Note that disjointness is not required here; if $f(\omega_0) = g(\omega_0) = 1$, for example, then the group represented by $f+g$ has $f(\omega_0) + g(\omega_0) = 2$ agents described by the attribute ω_0 .) The existence of an individual marginal bound simply means that there are no agents whose (potential) contributions to society are arbitrarily large. Continuity of Λ simply means that agents with similar attributes are good substitutes for each other.

If Ω is finite, we frequently refer to its elements as types; agents of the same type are exact substitutes. (Notice that continuity of Λ is automatic in this case.)

To derive a game from the technology (Ω, Λ) , we specify a finite set N and a function $\alpha : N \rightarrow \Omega$ (an attribute function). We associate with each subset S of

N a divisor $\text{div}(\alpha \mid S)$ given by

$$\text{div}(\alpha \mid S)(\omega) = |\alpha^{-1}(\omega) \cap S| \quad ;$$

i. e. , $\text{div}(\alpha \mid S)(\omega)$ is the number of players in S possessing the attribute ω . We then define the characteristic function $v_\alpha : 2^N \rightarrow \mathbb{R}_+$ by

$$v_\alpha(S) = \Lambda(\text{div}(\alpha \mid S)) \quad .$$

It is easily checked that (N, v_α) is a superadditive game.

5. THE SHAPLEY VALUE AND THE ϵ -CORE

Having described the framework, we can now state our main result. If the space Ω of attributes is finite (and so consists of a finite number of types), it seems natural to think of a game as large if it has many players of each type; thus each player has many exact substitutes. If Ω is not finite, it thus seems natural to think of a game as large if each player has many near substitutes.

THEOREM 1: Let (Ω, Λ) be a technology. For each $\epsilon > 0$ there are a $\delta(\epsilon) > 0$ and an integer $n(\epsilon)$ with the following property: If (N, v_α) is a game derived from the technology (Ω, Λ) , and for each player i in N there exist $n(\epsilon)$ distinct players $j_1, \dots, j_{n(\epsilon)}$ in N such that $\text{dist}(\alpha(i), \alpha(j_n)) < \delta(\epsilon)$ for each n , then the Shapley value of (N, v_α) is in the individually rational ϵ -core of (N, v_α) .

A number of remarks are in order. Most importantly, we stress that our result is valid for games and not simply for private goods exchange economies. The case of private goods exchange economies (with money) is covered by our framework, but is a special case.

Continuity of Λ assures us that players whose attributes are close together are close substitutes. The parameter $\delta(\epsilon)$ is thus a measure of how close these substitutes must be.

For the remainder of our remarks, we will, for ease of exposition, assume that the attribute space Ω is finite and speak of types.

We stress that our result asserts that the Shapley value is in the ϵ -core whenever there are enough players of each type; we do not need to know anything at all about the relative proportions of players of each type. By contrast, all the extant value convergence results for private goods exchange economies treat convergent sequences of economies, or sequences of replica economies, so that the relative proportions of players of each type play a crucial role.* Sequences of

* Of course, these value convergence theorems also assert the convergence of the value to the competitive equilibrium, which is meaningless in our framework.

games are also of interest to us (we discuss them in Section 10) but our main result is much more general.

It is important that the game have enough players of each type (or at least of each type which is represented at all). It will not suffice merely to have enough players in total; Example 2 of Wooders and Zame (1984) shows that this weaker condition does not guarantee that there is any payoff at all in the individually rational ϵ -core.

We note that the Shapley value may be in the ϵ -core and yet not be close to any point in the core. The most obvious reason for this is that the core may be empty. However, the Shapley value need not be close to the core even if the core is not empty, as the following simple example illustrates.

EXAMPLE (The market for gloves): The attribute space Ω consists of two points, $\Omega = \{L, R\}$. Every divisor f on Ω may then be written uniquely in the form $f = n_L \chi_L + n_R \chi_R$; define $\Lambda(n_L \chi_L + n_R \chi_R) = \min(n_L, n_R)$. (This is the market for (indivisible) gloves. Agents of type L own one left glove; agents of type R own one right glove. The value of a pair of gloves is one dollar; odd gloves and fractional combinations are valueless. Money is perfectly divisible and freely transferable.) If we consider a derived game (N, v_Q) with k players of type L and $k+1$ players of type R, we see that the core of (N, v_Q) consists of the unique payoff for which the players of type L each receive one dollar, while the players of type R receive nothing. The Shapley value, on the other hand, shares the total payoff of k dollars more evenly, with players of type L receiving only a slightly higher payoff than players of type R. (See Shapley and Shubik (1969) for a detailed numerical discussion of this example.)

Notice that in this example, if we let k tend to infinity and imagine the limiting continuum market, the imbalance between left and right gloves disappears. In the limit, the Shapley value divides the total payoff equally between players of type L and type R; this payoff is indeed in the core of the limiting continuum market. We will pursue this idea in Section 10.

Finally, we want to make some remarks about the proof. The usual proofs of value convergence theorems for private goods exchange economies depend of course on the special structure: divisible goods, concave utility functions, etc. At the heart of all such proofs is the idea that the marginal contribution of a player to a large coalition depends essentially only on the approximate distribution of types within the coalition. The law of large numbers guarantees that, for most large coalitions, the distribution of types within the coalition is approximately the same as in the economy as a whole. Thus, the Shapley value for each player is approximately the same as his marginal contribution to the entire economy, which is his competitive payoff.*

Such an approach cannot work for games in general, since a player's marginal contribution to a coalition may depend very critically on the precise make-up of the coalition. This is the case, for instance, in the glove market, and will commonly be the case whenever we are modelling indivisible goods, or coalition-production, or club goods, etc. Our approach is quite different.

At the heart of our proof are two ideas. The first of these ideas is that, even in large games, the blocking power is concentrated in small coalitions. A little more formally: if the feasible, Pareto optimal payoff x is not in the ϵ -core of a game derived from the given technology, so that there is a coalition S with $v(S) > x(S) + \epsilon|S|$, then there is a small coalition S' with $v(S') > x(S') + \frac{1}{2}\epsilon|S'|$. The second idea is that, in a large game, no small group can profit by forming a syndicate (i.e., an a priori coalition). More formally, the Shapley value which would accrue to a group if they agreed to act as a single unit is very nearly the sum of their individual Shapley values.

Our results on the blocking power of small coalitions are in Section 6. Section 7 contains a probabilistic estimate which is used in Section 8, where we discuss the power of syndicates. The proof of Theorem 1 is given in Section 9; it follows quite easily from the two basic ideas. We also discuss the question of estimating the parameters $\delta(\epsilon)$ and $n(\epsilon)$.

*This discussion is over-simplified, but we think it captures the essential idea.

6. THE POWER OF SMALL COALITIONS

Throughout this section, we fix a technology (Ω, Λ) . Our goal is to show that, for games derived from this technology, the blocking power is concentrated in small coalitions. We first isolate a combinatorial lemma which will be used several times.

LEMMA 1: Let $\omega_1, \dots, \omega_K$ be distinct points in Ω and let $\theta_1, \dots, \theta_K$ be strictly positive real numbers whose sum is one. Let $\{\phi_i\}$ and $\{\psi_j\}$ be sequences of divisors in $D(\Omega)$ such that:

- (a) $\phi_i(\omega) = 0$ for each i and each $\omega \notin \{\omega_1, \dots, \omega_K\}$ (no restriction is made for ψ);
- (b) $\lim_{i \rightarrow \infty} \|\phi_i\| = \lim_{j \rightarrow \infty} \|\psi_j\| = +\infty$;
- (c) $\lim_{i \rightarrow \infty} (\phi_i(\omega_k) / \|\phi_i\|) = \lim_{j \rightarrow \infty} (\psi_j(\omega_k) / \|\psi_j\|) = \theta_k$ for each k , $1 \leq k \leq K$.

For each i, j let r_{ij} be the largest integer such that $r_{ij}\phi_i \leq \psi_j$. Then

$$\lim_{i \rightarrow \infty} \limsup_{j \rightarrow \infty} (r_{ij} \|\phi_i\| / \|\psi_j\|) = \lim_{i \rightarrow \infty} \liminf_{j \rightarrow \infty} (r_{ij} \|\phi_i\| / \|\psi_j\|) = 1.$$

PROOF: For each i, j the integer r_{ij} is characterized by the inequalities

$$r_{ij}\phi_i(\omega_k) \leq \psi_j(\omega_k) \quad \text{for each } k;$$

$$(r_{ij} + 1)\phi_i(\omega_\ell) > \psi_j(\omega_\ell) \quad \text{for some } \ell.$$

We now fix an arbitrary real number ρ with $0 < \rho < 1$. By (c) above, and the fact that each θ_k is strictly positive, we can find integers i_0 and j_0 such that, for $i \geq i_0$ and $j \geq j_0$,

$$(1 - \rho)\theta_k \leq \phi_i(\omega_k) / \|\phi_i\| \leq (1 + \rho)\theta_k$$

$$(1 - \rho)\theta_k \leq \psi_j(\omega_k) / \|\psi_j\| \leq (1 + \rho)\theta_k$$

for each k . Combining these inequalities with those which characterize r_{ij} and simplifying yields

$$\frac{1-\rho}{1+\rho} \frac{\|\psi_j\|}{\|\phi_i\|} - 1 \leq r_{ij} \leq \frac{1+\rho}{1-\rho} \frac{\|\psi_j\|}{\|\phi_i\|}$$

or equivalently

$$\frac{1-\rho}{1+\rho} - \frac{\|\phi_i\|}{\|\psi_j\|} \leq r_{ij} \frac{\|\phi_i\|}{\|\psi_j\|} \leq \frac{1+\rho}{1-\rho}$$

provided that $i \geq i_0$ and $j \geq j_0$. If we let j tend to infinity (holding i fixed) and recall that $\|\psi_j\|$ tends to infinity, we obtain

$$\begin{aligned} \frac{1-\rho}{1+\rho} &\leq \liminf_{j \rightarrow \infty} r_{ij} \frac{\|\phi_i\|}{\|\psi_j\|} \\ &\leq \limsup_{j \rightarrow \infty} r_{ij} \frac{\|\phi_i\|}{\|\psi_j\|} \\ &\leq \frac{1+\rho}{1-\rho} \end{aligned}$$

for each $i \geq i_0$. Letting our arbitrary ρ tend to zero, so that i_0 tends to infinity, yields the desired result. $\square$

Our next task is to obtain some information about payoffs for sequences of divisors with a given limiting distribution. (Note carefully the difference in assumptions between Lemma 1 and Lemma 2.)

LEMMA 2: Let $\omega_1, \dots, \omega_T$ be distinct points in Ω and let $\theta_1, \dots, \theta_T$ be non-negative real numbers whose sum is one. Let $\{f_i\}$ be a sequence of divisors in $D(\Omega)$ such that $\lim_{i \rightarrow \infty} \|f_i\| = +\infty$ and $\lim_{i \rightarrow \infty} (f_i(\omega_t)/\|f_i\|) = \theta_t$ for each t . Then $\lim_{i \rightarrow \infty} (\Lambda(f_i)/\|f_i\|)$ exists. Moreover, this limit is independent of the sequence $\{f_i\}$ and depends only on the points $\omega_1, \dots, \omega_T$ and the numbers $\theta_1, \dots, \theta_T$.

PROOF: After renumbering if necessary, we may assume that $\theta_1, \dots, \theta_K$ are strictly positive and that $\theta_{K+1} = \dots = \theta_T = 0$ for some K , $1 \leq K \leq T$. Define

a new sequence of divisors $\{\phi_i\}$ by

$$\begin{aligned}\phi_i(\omega_k) &= f_i(\omega_k) \quad \text{for } 1 \leq k \leq K-1, \\ \phi_i(\omega_K) &= \|f_i\| - \sum_{k=1}^{K-1} f_i(\omega_k), \\ \phi_i(\omega) &= 0 \quad \text{if } \omega \notin \{\omega_1, \dots, \omega_K\}.\end{aligned}$$

It is evident that the points $\omega_1, \dots, \omega_K$, the numbers $\theta_1, \dots, \theta_K$ and the sequence $\{\phi_i\}$ satisfy all the conditions of Lemma 1 (i.e., take $\psi_i = \phi_i$). For each i, j let r_{ij} be the largest integer such that $r_{ij}\phi_i \leq \phi_j$. We are first going to show that $\lim(\Lambda(\phi_i)/\|\phi_i\|)$ exists.

Note that if M is an individual marginal bound for the technology (Ω, Λ) then

$$0 \leq \frac{\Lambda(\phi_i)}{\|\phi_i\|} \leq M \quad \text{for each } i.$$

Hence $\liminf(\Lambda(\phi_i)/\|\phi_i\|)$ and $\limsup(\Lambda(\phi_i)/\|\phi_i\|)$ both exist, and

$$\liminf(\Lambda(\phi_i)/\|\phi_i\|) \leq \limsup(\Lambda(\phi_i)/\|\phi_i\|).$$

To see that the reverse inequality holds we note that since $r_{ij}\phi_i \leq \phi_j$, the super-additivity of Λ yields

$$\Lambda(\phi_j) \geq \Lambda(r_{ij}\phi_i) \geq r_{ij}\Lambda(\phi_i)$$

and hence that

$$\frac{\Lambda(\phi_j)}{\|\phi_j\|} \geq r_{ij} \frac{\Lambda(\phi_i)}{\|\phi_i\|} = \left(\frac{r_{ij}\|\phi_i\|}{\|\phi_j\|} \right) \left(\frac{\Lambda(\phi_i)}{\|\phi_i\|} \right)$$

for each i, j . Let ϵ be a small positive number. In view of Lemma 1 and the definition of $\limsup$, we can find an index i_0 such that

 

$$\left(\frac{r_{i_0 j} \|\phi_{i_0}\|}{\|\phi_j\|} \right) \left(\frac{\Lambda(\phi_{i_0})}{\|\phi_{i_0}\|} \right) \geq \limsup \frac{\Lambda(\phi_i)}{\|\phi_i\|} - \epsilon$$

for every index j greater than some j_0 (which depends on i_0). On the other hand, we can, by the definition of $\liminf$, find an index $j_1 \geq j_0$ for which

$$\frac{\Lambda(\phi_{j_1})}{\|\phi_{j_1}\|} \leq \liminf \frac{\Lambda(\phi_j)}{\|\phi_j\|} + \epsilon.$$

Combining these inequalities yields that

$$\liminf \frac{\Lambda(\phi_j)}{\|\phi_j\|} + \epsilon \geq \limsup \frac{\Lambda(\phi_i)}{\|\phi_i\|} - \epsilon.$$

Since ϵ was arbitrary, we conclude that $\liminf(\Lambda(\phi_j)/\|\phi_j\|) \geq \limsup(\Lambda(\phi_i)/\|\phi_i\|)$ and hence that $L = \lim(\Lambda(\phi_i)/\|\phi_i\|)$ exists.

Returning to our original sequence $\{f_i\}$ of divisors, we want to show that $L = \lim(\Lambda(f_i)/\|f_i\|)$. To see this, note that $\|f_i\| = \|\phi_i\|$ and that our individual marginal bound and the definition of ϕ_i yield:

$$\begin{aligned} \left| \frac{\Lambda(f_i)}{\|f_i\|} - \frac{\Lambda(\phi_i)}{\|\phi_i\|} \right| &= \frac{|\Lambda(f_i) - \Lambda(\phi_i)|}{\|f_i\|} \\ &\leq 2M \frac{\sum_{k=1}^K f_i(\omega_k) - \|f_i\|}{\|f_i\|}. \end{aligned}$$

Since $(\sum f_i(\omega_k) - \|f_i\|)/\|f_i\|$ tends to zero (as i tends to infinity), we conclude that

$$\lim_{i \rightarrow \infty} (\Lambda(f_i)/\|f_i\|) = \lim_{i \rightarrow \infty} (\Lambda(\phi_i)/\|\phi_i\|) = L.$$

Finally, to show that this limit L is independent of the sequence $\{f_i\}$, let $\{g_i\}$ be another such sequence. Define a sequence $\{h_i\}$ by $h_{2i} = f_i$ and $h_{2i+1} = g_i$. By the above argument, $\lim(\Lambda(h_i)/\|h_i\|)$ exists. Since all subsequences of a convergent sequence converge to the same limit, we conclude that $L = \lim(\Lambda(f_i)/\|f_i\|) = \lim(\Lambda(g_i)/\|g_i\|)$, as asserted. This completes the proof. $\blacksquare$

 

We can now give the main result of this section.

LEMMA 3: For each $\epsilon > 0$ there is an integer $\ell(\epsilon)$ with the following property:

If (N, v_α) is any game derived from the technology (Ω, Λ) and $x \in \mathbb{R}^N$ is a feasible, Pareto optimal payoff which is not in the ϵ -core of (N, v_α) , then there is a coalition $S \subset N$ such that $|S| \leq \ell(\epsilon)$ and $v_\alpha(S) > x(S) + \frac{\epsilon}{2}|S|$.

PROOF: Suppose this were not so. Then we could find, for each integer ℓ , a finite set N_ℓ , an attribute function $\alpha_\ell : N_\ell \rightarrow \Omega$, an vector $x_\ell \in \mathbb{R}^{N_\ell}$, and a subset B_ℓ of N_ℓ with the following properties:

- (a) $x_\ell(N_\ell) = v_{\alpha_\ell}(N_\ell)$;
- (b) $v_{\alpha_\ell}(B_\ell) > x_\ell(B_\ell) + \epsilon|B_\ell|$,
- (c) if $v_{\alpha_\ell}(S) > x_\ell(S) + \frac{\epsilon}{2}|S|$, then $|S| > \ell$.

In order to obtain a contradiction, we are going to construct small representative subcoalitions S_ℓ of B_ℓ for which $v_{\alpha_\ell}(S_\ell)/|S_\ell|$ is approximately $v_{\alpha_\ell}(B_\ell)/|B_\ell|$ and $x_\ell(S_\ell)/|S_\ell|$ is (almost) less than $x_\ell(B_\ell)/|B_\ell|$.

We begin by choosing a $\delta > 0$ such that if $\omega_1, \omega_2 \in \Omega$ and $\text{dist}(\omega_1, \omega_2) < \delta$ then $|\Lambda(f + \chi_{\omega_1}) - \Lambda(f + \chi_{\omega_2})| < \epsilon/10$ for each divisor f (continuity of Λ). We then use compactness of Ω to write Ω as the disjoint union of a finite number of non-empty subsets $\Omega_1, \dots, \Omega_T$, each of diameter less than δ . For each t , choose and fix a point $\omega_t \in \Omega_t$.

For each ℓ and t , set

$$\theta_t^\ell = |\alpha_\ell^{-1}(\Omega_t) \cap B_\ell|/|B_\ell|$$

so that θ_t^ℓ is the relative proportion of players in B_ℓ with attributes in the set Ω_t . Notice that $0 \leq \theta_t^\ell \leq 1$ and that $\sum_{t=1}^T \theta_t^\ell = 1$ for each ℓ . Passing to a subsequence if necessary, we may assume that $\theta_t = \lim_{\ell \rightarrow \infty} \theta_t^\ell$ exists for each t . Renumbering if necessary, we may also assume that $\theta_1, \dots, \theta_K$ are strictly positive and $\theta_{K+1} = \dots = \theta_T = 0$ for some K , $1 \leq K \leq T$.

We now define divisors ϕ_ℓ and ψ_ℓ by

$$\begin{aligned}\phi_\ell(\omega_t) &= |\alpha_\ell^{-1}(\Omega_t) \cap B_\ell| & \text{for } 1 \leq t \leq K; \\ \phi_\ell(\omega) &= 0 & \text{for } \omega \notin \{\omega_1, \dots, \omega_K\}; \\ \psi_\ell(\omega_t) &= |\alpha_\ell^{-1}(\Omega_t) \cap B_\ell| & \text{for } 1 \leq t \leq T; \\ \psi_\ell(\omega) &= 0 & \text{for } \omega \notin \{\omega_1, \dots, \omega_T\}.\end{aligned}$$

(Notice that $\phi_\ell \leq \psi_\ell$.) It is easily checked that the points $\omega_1, \dots, \omega_T$, the numbers $\theta_1, \dots, \theta_T$ and the sequences $\{\phi_\ell\}$ and $\{\psi_\ell\}$ satisfy the conditions of Lemmas 1 and 2. Hence,

$$\lim_{\ell \rightarrow \infty} \frac{\Lambda(\phi_\ell)}{\|\phi_\ell\|} = \lim_{\ell \rightarrow \infty} \frac{\Lambda(\psi_\ell)}{\|\psi_\ell\|}$$

and both these limits exist; call this common limit L . Fix an index ℓ_0 so that $|L - \Lambda(\phi_\ell)/\|\phi_\ell\|| < \epsilon/10$ and $|L - \Lambda(\psi_\ell)/\|\psi_\ell\|| < \epsilon/10$ for every $\ell \geq \ell_0$.

For each i, j let r_{ij} be the largest integer such that $r_{ij}\phi_i \leq \psi_j$. In view of Lemma 1, we may choose an index $I \geq \ell_0$ and an index $J_0 \geq \ell_0$ such that (for M the individual marginal bound for the technology):

$$\frac{r_{IJ}\|\phi_I\|}{\|\psi_J\|} > \left(1 + \frac{\epsilon}{10M}\right)^{-1}$$

whenever $J \geq J_0$ (keeping I fixed).

For each $\ell \geq \max(I, J_0)$ we now define a coalition $S_\ell \subset N_\ell$ in the following way. For each t , $1 \leq t \leq K$, we consider all subsets A of $\alpha_\ell^{-1}(\Omega_t) \cap B_\ell$ with $|A| = \phi_I(\omega_t) = |\alpha_I^{-1}(\Omega_t) \cap B_I|$; there are of course only finitely many such sets. Among all such sets, we let S_ℓ^t be any one for which the aggregate payoff $x_\ell(A)$ is as small as possible. Finally, we set $S_\ell = \sum_{t=1}^K S_\ell^t$. Notice that $|S_\ell| = \|\phi_I\| \leq |B_I|$, so if we can show that

$$v_{\alpha_\ell}(S_\ell) > x_\ell(S_\ell) + \frac{\epsilon}{2}|S_\ell|,$$

 

or equivalently that

$$\frac{v_{\alpha_\ell}(S_\ell)}{|S_\ell|} > \frac{x_\ell(S_\ell)}{|S_\ell|} + \frac{\epsilon}{2} ,$$

we will have obtained a contradiction.

To see this, let us first estimate $v_{\alpha_\ell}(S_\ell)/|S_\ell|$. Our choice of the sets Ω_t and our construction of S_ℓ , together with the facts that $|S_\ell| = \|\phi_I\|$ and $\ell \geq I \geq \ell_0$ imply that

$$\begin{aligned} \frac{v_{\alpha_\ell}(S_\ell)}{|S_\ell|} &\geq \frac{\Lambda(\phi_I)}{\|\phi_I\|} - \frac{\epsilon}{10} \\ &> L - \frac{\epsilon}{10} - \frac{\epsilon}{10} \\ &> \frac{\Lambda(\psi_\ell)}{\|\psi_\ell\|} - \frac{\epsilon}{10} - \frac{\epsilon}{10} - \frac{\epsilon}{10} \\ &\geq \frac{v_{\alpha_\ell}(B_\ell)}{|B_\ell|} - \frac{\epsilon}{10} - \frac{\epsilon}{10} - \frac{\epsilon}{10} - \frac{\epsilon}{10} . \end{aligned}$$

By assumption, $v_{\alpha_\ell}(B_\ell) > x_\ell(B_\ell) + \epsilon|B_\ell|$, so we obtain

$$\frac{v_{\alpha_\ell}(S_\ell)}{|S_\ell|} > \frac{x_\ell(B_\ell)}{|B_\ell|} + \frac{6\epsilon}{10} .$$

We now estimate $x_\ell(S_\ell)/|S_\ell|$. By construction $r_{I\ell}\phi_I \leq \psi_\ell$ so that $r_{I\ell}\phi_I(\omega_t) \leq \psi_\ell(\omega_t)$ for $1 \leq t \leq K$. But $\phi_I(\omega_t) = |S_\ell^t| = |S_\ell \cap \alpha_\ell^{-1}(\Omega_t)|$, and $\psi_\ell(\omega_t) = |\alpha_\ell^{-1}(\Omega_t) \cap B_\ell|$, so $r_{I\ell}|S_\ell^t| = r_{I\ell}|S_\ell \cap \alpha_\ell^{-1}(\Omega_t)| \leq |\alpha_\ell^{-1}(\Omega_t) \cap B_\ell|$ for $1 \leq t \leq K$. Using the fact that S_ℓ^t minimizes $x_\ell(A)$ over all subsets A of $\alpha_\ell^{-\ell}(\Omega_t)$ with $|S_\ell^t|$ members, we obtain

$$\begin{aligned}
x_\ell(B_\ell) &\geq \sum_{t=1}^K x_\ell(B_\ell \cap \alpha_\ell^{-1}(\Omega_t)) \\
&\geq \sum_{t=1}^K r_{I\ell} x_\ell(S_\ell^t) \\
&= r_{I\ell} x_\ell(S_\ell) .
\end{aligned}$$

Recalling that $|S_\ell| = \|\phi_I\|$ and $|B_\ell| = \|\psi_\ell\|$, that $\ell \geq J_0$ (which gives an estimate on $r_{I\ell}$) and that $x_{\alpha_\ell}(B_\ell) < v_{\alpha_\ell}(B_\ell) + \epsilon|B_\ell|$, and using our individual marginal bound, we obtain:

$$\begin{aligned}
\frac{x_\ell(S_\ell)}{|S_\ell|} &\leq \frac{x_\ell(B_\ell)}{r_{I\ell} |S_\ell|} \\
&= \left(\frac{x_\ell(B_\ell)}{r_{I\ell} \|\phi_I\|} \right) \left(\frac{\|\psi_\ell\|}{|B_\ell|} \right) \\
&\leq \left(\frac{x_\ell(B_\ell)}{|B_\ell|} \right) \left(1 + \frac{\epsilon}{10M} \right) \\
&< \frac{v_{\alpha_\ell}(B_\ell)}{|B_\ell|} + \frac{\epsilon}{10} .
\end{aligned}$$

Combining this estimate with our estimate on $v_{\alpha_\ell}(S_\ell)/|S_\ell|$ yields

$$\begin{aligned}
\frac{x_\ell(S_\ell)}{|S_\ell|} &< \frac{v_{\alpha_\ell}(S_\ell)}{|S_\ell|} - \frac{6\epsilon}{10} + \frac{\epsilon}{10} \\
&= \frac{v_{\alpha_\ell}(S_\ell)}{|S_\ell|} - \frac{\epsilon}{2} ,
\end{aligned}$$

which is the contradiction we desire. This completes the proof. $\square$

7. A PROBABILISTIC ESTIMATE

As with most calculations involving the Shapley value, ours depends on a probabilistic estimate. The estimate we use is very simple, so we derive it here.

We consider an urn which contains a total of t balls, of which g are green. We draw, at random and without replacement, a sample of size k ($k \leq t$), and write h for the number of green balls in the sample. We want to estimate the probability that

$$\left| \frac{h}{g} - \frac{k}{t} \right| < \eta ,$$

where η is some predetermined parameter.

To obtain this estimate, we simply follow Feller (1968, p. 233). The number h is a random variable whose expected value is kg/t and whose variance is

$$\text{Var}(h) = \frac{kg(t-g)}{t^2} \left[1 - \frac{k-1}{t-1} \right] < \frac{kg(t-g)}{t^2}$$

Hence Chebyshev's inequality yields

$$\begin{aligned} \text{prob} \left(\left| \frac{h}{g} - \frac{k}{t} \right| \geq \eta \right) &= \text{prob} \left(\left| h - \frac{gk}{t} \right| \geq \eta g \right) \\ &< \frac{\text{Var}(h)}{(\eta g)^2} \\ &< \frac{kg(t-g)}{t^2 \eta^2 g^2} \\ &< \frac{1}{\eta^2 g} . \end{aligned}$$

Equivalently

$$\text{prob} \left(\left| \frac{h}{g} - \frac{k}{t} \right| < \eta \right) > 1 - \frac{1}{\eta^2 g} .$$

This is the estimate we shall need.

Note that this estimate is independent of t (the total number of balls) and of k (the sample size). Note too that for η fixed, this probability tends to one as g (the number of green balls) tends to infinity.

8. THE POWER OF SMALL SYNDICATES

The purpose of this Section is to study the way the Shapley value of a game changes if we allow a group of players to form a syndicate. We will show that, if the syndicate is small (in the sense that there are many near substitutes for members of the syndicate), then the change in the Shapley value is also small. Since this phenomenon seems of interest outside the context of technologies, we will formulate and prove a general result.

We consider games (N, v) which are not required to be superadditive. If S is a nonempty subset of N , then by the syndicate game (N_S, v_S) we mean the game whose player set is $N_S = (N \setminus S) \cup \{S\}$ and whose characteristic function is given by

$$\begin{aligned} v_S(W) &= v(W) \quad \text{if } \{S\} \notin W, \\ v_S(W) &= v([W \cap (N \setminus S)] \cup S) \quad \text{if } \{S\} \in W. \end{aligned}$$

That is, (N_S, v_S) is the game which results if we treat the syndicate S as an indivisible unit. We want to compare $\text{Sh}(v_S, \{S\})$ with $\sum_{i \in S} \text{Sh}(v, i)$; the difference between these two numbers might be called the power of the syndicate, i.e., the gain or loss resulting from formation of the syndicate.

We need some terminology. We will say that the positive number M is an individual marginal bound for the game (N, v) if

$$|v(W \cup \{j\}) - v(W)| \leq M$$

for each $j \in N$ and each $W \subset N$. For $\gamma > 0$, we will say that players $i, j \in N$ are γ -substitutes if

$$|v(W \cup \{i\}) - v(W \cup \{j\})| < \gamma$$

for every $W \subset N \setminus \{i, j\}$. Our result on syndication is the following.

LEMMA 4: Let M and γ be positive numbers and let s be a positive integer. Then there is an integer $r(M, \gamma, s)$ with the following property: If (N, v) is a game with an individual marginal bound of M , S is a subset of N with $|S| \leq s$, and for each

player i in S there are at least $r(M, \gamma, s)$ players not in S who are γ -substitutes for i , then

$$\left| \text{Sh}(v_S, \{S\}) - \sum_{i \in S} \text{Sh}(v, i) \right| \leq \frac{3|S|(|S|+1)\gamma}{2}.$$

PROOF: We are going to obtain an estimate for $\left| \text{Sh}(v_S, \{S\}) - \sum_{i \in S} \text{Sh}(v, i) \right|$ in terms of the number of γ -substitutes for members of S and some other parameters. We will then show that this estimate can be reduced to the one we want by judicious choice of these other parameters provided the number of γ -substitutes is large enough. If $s = 1$ there is nothing to prove (since $v_S = v$); we consider first the case $s = 2$.

We fix a game (N, v) and a coalition $S \subset N$ with $|S| = 2$. Write $S = \{a, b\}$, $N^* = N \setminus S$ and $n = |N| - 1 = |N^*| + 1 = |N_S|$. Let $A \subset N^*$ be a set of γ -substitutes for a , and let $B \subset N^*$ be a set of γ -substitutes for b . We may assume that $|A| = |B| = r$ (simply ignoring some γ -substitutes if necessary). Finally, we fix parameters ρ, τ and ξ with $0 < \rho < 1/2$, $0 < \tau$, and $0 < \xi < 1$.

For the remainder of the proof it is convenient to write

$$\Delta(W, i) = v(W \cup \{i\}) - v(W)$$

for $W \subset N$ and $i \in N$.

By definition,

$$\begin{aligned} \text{Sh}(v_S, \{S\}) &= \frac{1}{n} \sum_{J=0}^{n-1} \frac{1}{\binom{n-1}{J}} \sum_{\substack{W \subset N^* \\ |W|=J}} [v_S(W \cup \{S\}) - v_S(W)] \\ &= \frac{1}{n} \sum_{J=0}^{n-1} \frac{1}{\binom{n-1}{J}} \sum_{\substack{W \subset N^* \\ |W|=J}} [v(W \cup S) - v(W)] \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{n} \sum_{J=0}^{n-1} \frac{1}{\binom{n-1}{J}} \sum_{\substack{W \subset N^* \\ |W|=J}} \left[v(W \cup \{a, b\}) - v(W \cup \{a\}) \right] \\
&\quad + \frac{1}{n} \sum_{J=0}^{n-1} \frac{1}{\binom{n-1}{J}} \sum_{\substack{W \subset N^* \\ |W|=J}} \left[v(W \cup \{a\}) - v(W) \right] \\
&= \frac{1}{n} \sum_{J=0}^{n-1} \frac{1}{\binom{n-1}{J}} \sum_{\substack{W \subset N^* \\ |W|=J}} \Delta(W \cup \{a\}, b) \\
&\quad + \frac{1}{n} \sum_{J=0}^{n-1} \frac{1}{\binom{n-1}{J}} \sum_{\substack{W \subset N^* \\ |W|=J}} \Delta(W, a) .
\end{aligned}$$

Let us write Q for this last double sum and P for the next to last. We are going to estimate $|\text{Sh}(v, a) - Q|$ and $|\text{Sh}(v, b) - P|$.

As a guide to the intuition, we point out that Q is itself the Shapley value for player a in the game $(\bar{N}, \bar{v})$, where $\bar{N} = N \setminus \{b\}$ and $\bar{v}(W) = v(W)$ for each $W \subset \bar{N} = N \setminus \{b\}$. Since B consists of γ -substitutes for b , it is natural to expect that $Q = \text{Sh}(\bar{v}, a)$ should be close to $\text{Sh}(v, a)$ if $|B| = r$ is large. Verifying that this is so, however, takes quite a lot of work.

By definition,

$$\text{Sh}(v, a) = \frac{1}{n+1} \sum_{J=0}^n \frac{1}{\binom{n}{J}} \sum_{\substack{W \subset N \setminus \{a\} \\ |W|=J}} \Delta(W, a) .$$

Write

$$E = \frac{1}{n} \sum_{J=0}^{n-1} \frac{1}{\binom{n}{J}} \sum_{\substack{W \subset N \setminus \{a\} \\ |W|=J}} \Delta(W, a)$$

Using the individual marginal bound of M , we see that

$$(1) \quad |E - \text{Sh}(v, a)| \leq \frac{2M}{n} < \frac{2M}{r} .$$

Now, we want to estimate $|E - Q|$. Note two main differences between the expression for E and the expression for Q : different coefficients, and the presence of coalitions which contain b . We are going to approximate terms involving coalitions which contain b by terms not involving coalitions which contain b ; this will also have the effect of "correcting" the discrepancy in the coefficients.

We first introduce some sets of coalitions. For each J and k , we let $\Gamma(J, k)$ be the set of coalitions $W \subset N \setminus \{a\}$ for which $|W| = J$, $|W \cap B| = k$ and $b \notin J$; we let $\Gamma_b(J, k)$ be the set of coalitions $W \subset N \setminus \{a\}$ for which $|W| = J$, $|W \cap B| = k$ and $b \in W$.

We now use these sets to break up the expressions for E and Q . Set

$$\begin{aligned} E(J, k) &= \sum_{W \in \Gamma(J, k)} \Delta(W, a) , \\ E_b(J, k) &= \sum_{W \in \Gamma_b(J, k)} \Delta(W, a) , \\ E(J) &= \frac{1}{\binom{n}{J}} \sum_{k=0}^r [E(J, k) + E_b(J, k)] , \end{aligned}$$

and

$$Q(J) = \frac{1}{\binom{n-1}{J}} \sum_{k=0}^r E(J, k) .$$

Notice that

$$E = \frac{1}{n} \sum_{J=0}^{n-1} E(J)$$

and that

$$Q = \frac{1}{n} \sum_{J=0}^{n-1} Q(J) .$$

Thus, we want to estimate $|E(J) - Q(J)|$ for each J .

The estimate we use depends on the size of J . If $(J-1) \leq \rho(n-1)$ or $(J-1) \geq (1-\rho)(n-1)$ we use the obvious estimate given to us by the individual marginal bound:

$$(2) \quad |E(J) - Q(J)| \leq 2M .$$

If $\rho(n-1) < (J-1) < (1-\rho)(n-1)$, we need to be more careful. Let us fix such a J , and let I be the set of indices k for which

$$\left| \frac{k+1}{r-k} - \frac{J+1}{n-1-J} \right| < \tau$$

(recall that τ was one of our initial fixed parameters); let I' be the complementary set of indices.

Fix an index $k \in I$. For $W \in \Gamma_b(J, k)$ and $c \in B \setminus (W \cap B)$, write $W_c = (W \setminus \{b\}) \cup \{c\}$. The fact that b and c are γ -substitutes yields that

$$|\Delta(W, a) - \Delta(W_c, a)| \leq 2\gamma .$$

Hence if we average over the $r-k$ elements of $B \setminus (W \cap B)$, we obtain

$$\left| \Delta(W, a) - \frac{1}{r-k} \sum_{c \in B \setminus (W \cap B)} \Delta(W_c, a) \right| \leq 2\gamma .$$

Notice that W was chosen to be in $\Gamma_b(J, k)$ but that each W_c belongs to $\Gamma(J, k+1)$. Now, as W runs over all elements of $\Gamma_b(J, k)$, the sets W_c run over all elements of $\Gamma(J, k+1)$, but each element of $\Gamma(J, k+1)$ occurs exactly $k+1$ times (because there are $k+1$ elements which might have occurred as replacements for b). Hence if we sum over $\Gamma_b(J, k)$, we obtain

$$\left| E_b(J, k) - \frac{k+1}{r-k} E(J, k+1) \right| \leq 2\gamma |\Gamma_b(J, k)| .$$

Combining this with the fact that k is in I yields

$$\begin{aligned}
(3) \quad \left| E_b(J, k) - \frac{J}{n-1-J} E(J, k+1) \right| &\leq 2\gamma |\Gamma_b(J, k)| + \tau |E(J, k+1)| \\
&\leq 2\gamma |\Gamma_b(J, k)| + 2M\tau |\Gamma(J, k+1)| .
\end{aligned}$$

On the other hand, for indices $k \notin I$ we certainly have

$$(4) \quad |E_b(J, k)| \leq 2M |\Gamma_b(J, k)|$$

and

$$(5) \quad |E(J, k)| \leq 2M |\Gamma(J, k)| .$$

The next step is to show that most of the coalitions in $\bigcup_{k=0}^r \Gamma_b(J, k)$ actually belong to $\bigcup_{k \in I} \Gamma_b(J, k)$. Note first that if $k/r = (J-1)/(n-1)$, then direct calculation gives

$$(6) \quad \frac{k+1}{r-k} = \frac{J-1}{n-J} + \frac{1}{r \left(1 - \left(\frac{J-1}{n-1} \right) \right)} .$$

Since we have taken $J-1$ in the range $\rho(n-1) < (J-1) < (1-\rho)(n-1)$ and ρ is fixed, the second term on the right-hand side of equation (6) is certainly less than $\tau/3$ if r is sufficiently large. Since $n \geq r$, the first term on the right-hand side of (6) differs from $(J+1)/(n-1-J)$ by less than $\tau/3$ if r (and hence n) is large. That is,

$$\left| \frac{k+1}{r-k} - \frac{J+1}{n-1-J} \right| < 2\tau/3$$

provided that r is large and $k/r = (J-1)/(n-1)$. Hence

$$(7) \quad \left| \frac{k+1}{r-k} - \frac{J+1}{n-1-J} \right| < \tau$$

provided that r is large and that

$$\left| \frac{k}{r} - \frac{J-1}{n-1} \right|$$

is sufficiently small; if (7) holds, then $k \in I$.

We can now use our probabilistic estimate from Section 7 in the following way. Choosing at random a coalition W in $\Gamma_b(J)$ is the same thing as choosing at

random a set of $J-1$ elements from $N \setminus \{a, b\}$, and then adding the element b to this set. If $k = |W \cap B|$, the estimate of Section 7 tells us that with probability at least $1 - \zeta$ (recall that ζ is the third of our fixed parameters), the quantity

$$\left| \frac{k}{r} - \frac{J-1}{n-1} \right|$$

will be very small, provided only that r is sufficiently large. In other words,

$$(8) \quad \left| \bigcup_{k \in I'} \Gamma_b(J, k) \right| \leq \zeta \left| \bigcup_{k=0}^r \Gamma_b(J, k) \right|$$

provided that r is large.

We now begin to estimate $|E(J) - Q(J)|$ for $\rho(n-1) < (J-1) < (1-\rho)(n-1)$.

We write

$$(9) \quad E(J) = \frac{1}{\binom{n}{J}} \sum_{k \in I} [E(J, k) + E_b(J, k)] + \frac{1}{\binom{n}{J}} \sum_{k \in I'} [E(J, k) + E_b(J, k)] .$$

Combining the inequalities (4), (5) and (8), we see that the second of these sums is small; to be precise,

$$(10) \quad \left| \frac{1}{\binom{n}{J}} \sum_{k \in I'} [E(J, k) + E_b(J, k)] \right| \leq 4M\zeta .$$

We can use (3) to obtain an approximate expression for the first sum in (9),

$$(11) \quad \frac{1}{\binom{n}{J}} \sum_{k \in I} [E(J, k) + E_b(J, k)] \simeq \frac{1}{\binom{n}{J}} \sum_{k \in I} \left[E(J, k) + \frac{J}{n-1-J} E(J, k+1) \right] ,$$

with an error not exceeding $2\gamma + 2M\tau$. Notice that in the right-hand sum of (11), every one of the terms $E(J, k+1)$ appears twice, with the exception of the term corresponding to the smallest value k_0 of k which is in I : the term $E(J, k+1)$ appears the first time with a coefficient of $J/(n-1-J)$ and a second time with a coefficient of 1. Hence we can rewrite the right-hand side of (11) as

$$\begin{aligned}
(12) \quad & \frac{1}{\binom{n}{J}} \sum_{k \in I} \left[E(J, k) + \frac{J}{n-1-J} E(J, k+1) \right] \\
&= \frac{1}{\binom{n}{J}} \sum_{\substack{k \in I \\ k \neq k_0}} \left(\frac{J}{n-1-J} + 1 \right) E(J, k+1) + \frac{1}{\binom{n}{J}} E(J, k_0) .
\end{aligned}$$

Combining (5) with the same probabilistic estimate we used before, we can see that the last of these terms can be made small:

$$(13) \quad \left| \frac{1}{\binom{n}{J}} E(J, k_0) \right| \leq 2M\xi ,$$

if r is large.

We now point out a very convenient identity:

$$(14) \quad \frac{1}{\binom{n}{J}} \left(\frac{J}{n-1-J} + 1 \right) = \frac{1}{\binom{n-1}{J}} .$$

If we plug this identity into (12), and combine (9) with the estimates (10), (11) and (13), we conclude that:

$$(15) \quad E(J) \simeq \frac{1}{\binom{n-1}{J}} \sum_{\substack{k \in I \\ k \neq k_0}} E(J, k+1) ,$$

with an error not exceeding $4M\xi + 2\gamma + 2M\tau + 2M\xi$. But the right-hand side of (15) is just part of the expression for $Q(J)$; the terms which are missing are just the ones for which $k \in I'$, and just as before we can see that the sum of the missing terms does not exceed $4M\xi$, for large r .

To summarize, we have shown that

$$(16) \quad |E(J) - Q(J)| \leq 10M\xi + 2M\tau + 2\gamma ,$$

provided that $\rho(n-1) < (J-1) < (1-\rho)(n-1)$ and r is large. Recall from (2) that

$$|E(J) - Q(J)| \leq 2M$$

if $(J - 1) \leq \rho(n - 1)$ or $(J - 1) \geq (1 - \rho)(n - 1)$.

In order to estimate $|E - Q|$, we need only add all these estimates and divide by n . This yields

$$\begin{aligned} |E - Q| &\leq \frac{1}{n} [2\rho nM + n(10M\xi + 2M\tau + 2\gamma)] \\ &= M(2\rho + 10\xi + 2\tau) + 2\gamma . \end{aligned}$$

Combining this with (1) gives us the estimate we are after:

$$|Sh(v, a) - Q| \leq M(2\rho + 10\xi + 2\tau) + 2\gamma + \frac{2M}{r} ,$$

for r sufficiently large. It is now clear that we need not be very careful in our choice of the parameters ρ , τ and ξ . If we simply choose each of them very small, we obtain

$$|Sh(v, a) - Q| \leq 3\gamma ,$$

provided that r is sufficiently large. This is the desired estimate for Q .

The intuition underlying our estimate for P is similar to the intuition underlying our estimate for Q . We want to see that P is nearly the Shapley value for player b in the game $(\bar{N}, \bar{v})$, where $\bar{N} = N \setminus \{a\}$ and $\bar{v}(W) = v(W)$ for each $W \subset N \setminus \{a\}$. Since the preceding argument, with the roles of a and b reversed, shows that $|Sh(\bar{v}, b) - Sh(v, b)| \leq 3\gamma$ if r is large, this will give us the estimate we need.

Our previous argument was based on the idea of systematically replacing every term which involved a coalition containing b by terms which do not involve coalitions containing b . This time, we want to systematically replace every term involving a coalition containing a by terms which do not involve coalitions containing a . The twist is that all these terms occur in the expression for P and not in the expression for $Sh(\bar{v}, b)$. The terms we seek to replace are of the form $\Delta(W \cup \{a\}, b)$, and following the same procedure as before, we obtain the approximation

$$\Delta(W \cup \{a\}, b) \simeq \Delta(W \cup \{d\}, b) ,$$

for $d \in A \setminus (W \cap A)$. Notice that $W \cup \{d\}$ is a coalition in $\bar{N} \setminus \{b\} = N \setminus \{a, b\}$, and $|W \cup \{d\}| = |W| + 1$. Thus if we follow the same averaging procedure as before, the terms in the expression for P corresponding to coalitions W of size J become terms in the expression for $\text{Sh}(\bar{v}, b)$ corresponding to coalitions $W \cup \{d\}$ of size $J+1$. The averaging procedure also gives us an extra factor which (for most of the terms) is nearly equal to $(J+1)/(n-1-J)$. This extra factor is just what we need, since we can then use, instead of the identity (14), the identity

$$\frac{1}{\binom{n-1}{J}} \left(\frac{J+1}{n-1-J} \right) = \frac{1}{\binom{n-1}{J+1}} .$$

After carrying out the same sort of approximations as before, we obtain that

$$|P - \text{Sh}(\bar{v}, b)| \leq 3\gamma ,$$

provided that r is sufficiently large. Since, as we have already noted, the first argument (with the roles of a and b reversed) gives $|\text{Sh}(\bar{v}, b) - \text{Sh}(v, b)| \leq 3\gamma$, we conclude that

$$|P - \text{Sh}(v, b)| \leq 6\gamma ,$$

if r is sufficiently large. Combining our estimates for P and Q gives

$$\begin{aligned} |\text{Sh}(v_S, \{S\}) - \text{Sh}(v, a) - \text{Sh}(v, b)| &= |P+Q - \text{Sh}(v, a) - \text{Sh}(v, b)| \\ &\leq 9\gamma , \end{aligned}$$

provided that r is sufficiently large. This is the estimate we want, so the proof in the case $|S| = s = 2$ is complete.

Finally, we come to the general case. Let $|S| \leq s$ and write $S = \{a_1, \dots, a_{|S|}\}$. We write, as before,

$$\text{Sh}(v_S, \{S\}) = P_1 + \dots + P_{|S|} .$$

Just as above, we find that $|P_{|S|} - \text{Sh}(v, a_S)| \leq 3\gamma$, that $|P_{|S|-1} - \text{Sh}(v, a_{S-1})| \leq 6\gamma$, and so forth. Summing yields

$$\left| \text{Sh}(v_S, \{S\}) - \sum_{i=1}^{|S|} \text{Sh}(v, a_i) \right| \leq \frac{3|S|(|S|+1)\gamma}{2} ,$$

provided that r (the number of γ -substitutes for members of S) is sufficiently large. This is the estimate we needed, so the proof is complete. $\square$

Although we shall have no need of it, we point out that a similar argument can be used to show that for $i \notin S$ we also have

$$|\text{Sh}(v_S, i) - \text{Sh}(v, i)| < \frac{3|S|(|S|+1)\gamma}{2} ,$$

provided that the number of γ -substitutes for members of S is sufficiently large.

9. PROOF OF THEOREM 1

It is now very easy to combine Lemma 3 of Section 6 and Lemma 4 of Section 8 to give a proof of Theorem 1.

We are given a technology (Ω, Λ) and a positive number ϵ . Let $\ell(\epsilon)$ be the integer given by Lemma 3. Set

$$\gamma = \frac{\epsilon}{3(\ell(\epsilon) + 1)} \quad ,$$

and choose a positive number $\delta(\epsilon) > 0$ such that, for each $f \in D(\Omega)$,

$$|\Lambda(f + \chi_{\omega_1}) - \Lambda(f + \chi_{\omega_2})| < \gamma$$

whenever ω_1 and ω_2 belong to Ω and $\text{dist}(\omega_1, \omega_2) < \delta(\epsilon)$. Let $r(M, \gamma, \ell(\epsilon))$ be the integer given by Lemma 4. Finally set $n(\epsilon) = r(M, \gamma, \ell(\epsilon)) + \ell(\epsilon)$.

Now suppose we are given a game (N, v_α) derived from (Ω, Λ) and that for each player $i \in N$ there are at least $n(\epsilon)$ distinct players $j_1, \dots, j_{n(\epsilon)}$ such that $\text{dist}(\alpha(i), \alpha(j_k)) < \delta(\epsilon)$ for $1 \leq k \leq n(\epsilon)$. We must show that $\text{Sh}(v_\alpha)$ belongs to the individually rational ϵ -core of (N, v_α) . Certainly, $\text{Sh}(v_\alpha)$ is feasible, Pareto optimal and individually rational. If $\text{Sh}(v_\alpha)$ were not in the ϵ -core, we could, by Lemma 3, find a coalition $S \subset N$ such that $|S| \leq \ell(\epsilon)$ and

$$\sum_{i \in S} \text{Sh}(v_\alpha, i) < v_\alpha(S) - \frac{\epsilon}{2} |S| \quad .$$

On the other hand, our choice of $\delta(\epsilon)$ guarantees that each member of S has at least $n(\epsilon)$ γ -substitutes in N ; our choice of $n(\epsilon)$ guarantees that at least $r(M, \gamma, \ell(\epsilon))$ of these γ -substitutes do not belong to S . Hence Lemma 4 tells us that

$$\sum_{i \in S} \text{Sh}(v_\alpha, i) \geq \text{Sh}((v_\alpha)_S, \{S\}) - \frac{3|S|(|S| + 1)\gamma}{2} \quad .$$

On the other hand, individual rationality of the Shapley value for the game $(v_\alpha)_S$ guarantees that $\text{Sh}((v_\alpha)_S, \{S\}) \geq (v_\alpha)_S(\{S\}) = v_\alpha(S)$. Combining this fact with the

 

previous inequality and the definition of γ yields

$$\sum_{i \in S} \text{Sh}(v_\alpha, i) \geq v_\alpha(S) - \frac{\epsilon}{2} |S| .$$

which is a contradiction. We conclude that $\text{Sh}(v_\alpha)$ indeed belongs to the individually rational ϵ -core of (N, v_α) , as asserted. $\square$

It is of interest to obtain good estimates on $\delta(\epsilon)$ and $n(\epsilon)$, in terms of certain natural parameters of the technology (Ω, Λ) . The chief obstacle to doing so is that the integer $\ell(\epsilon)$ given by Lemma 3 depends on the technology in a way that seems difficult to quantify. If we think of $\ell(\epsilon)$ as known, however,^{*} it is not hard to obtain an estimate for $n(\epsilon)$. Simply keeping track of the various parameters in the proofs of Lemma 4 and Theorem 1 gives an estimate for $n(\epsilon)$ on the order of a constant times $(M/\epsilon)^3 (\ell(\epsilon))^3$.^{**} The number $\delta(\epsilon)$ of course, depends only on the modulus of continuity of Λ .

^{*} That is, if we regard $\ell(\epsilon)$ as a known parameter of the technology.

^{**} This may be compared with the estimate obtained by Mas-Colell, which is of the same order.

10. LIMITING BEHAVIOR

The example presented in Section 5 shows that the Shapley value of a large game, while in the individually rational ϵ -core, need not be close to any point in the core (even if the core is not empty). The same example also suggests that the Shapley value should be close, not to the core of the large finite game, but to the core of a limiting nonatomic game. The purpose of this section is to show that this is indeed the case. In fact, we show quite a lot more: the individually rational ϵ -cores of a sequence of games converge to the core of the limit game. To make this precise we require some preliminary discussion.

We will restrict our attention to the case where Ω is a finite set, say $\Omega = \{\omega_1, \dots, \omega_T\}$ so that we will speak of types. (Similar results could be established in more general contexts, but would be much more complicated, and lose much of their intuitive flavor.)

To define a nonatomic limiting game, we fix strictly positive* real numbers $\theta_1, \dots, \theta_T$ with $\sum \theta_t = 1$, and disjoint intervals $I_1, \dots, I_T$ on the real line for which $\text{length}(I_t) = \theta_t$. Set $I = \cup I_t$, let $\mathcal{B}$ be the family of Borel subsets of I , and let μ be the restriction to I of Lebesgue measure. (The interpretation we have in mind is that I represents a continuum of players of which the fraction $\theta_t = \mu(I_t)$ are of type t . The family $\mathcal{B}$ of all Borel subsets of I is the family of admissible coalitions.)

To define a nonatomic game on I , in the sense of Aumann and Shapley (1974), we must define a set function λ on $\mathcal{B}$ which is of bounded variation. To this end, let B be an element of $\mathcal{B}$. If $\mu(B) = 0$, we define $\lambda(B) = 0$. Otherwise, we write $\beta_t = \mu(B \cap I_t) / \mu(B)$ for each t ; note that $\beta_t \geq 0$ and that $\sum \beta_t = 1$. Choose a sequence $\{f_k\}$ of divisors on Ω such that $\|f_k\| \rightarrow \infty$ and $f_k(\omega_t) / \|f_k\| \rightarrow \beta_t$ for each t . We then define

* The case in which some of the θ_t 's are zero can be treated by restricting attention to a subset of Ω .

$$\lambda(B) = \left(\lim_{k \rightarrow \infty} \frac{\Lambda(f_k)}{\|f_k\|} \right) \cdot \mu(B) .$$

(In view of Lemma 2, this limit exists and is independent of the particular sequence $\{f_k\}$ of divisors we choose.) To see that λ is of bounded variation, we note that $0 \leq \Lambda(f)/\|f\| \leq M$ for any divisor f (where M is the individual marginal bound for the technology (Ω, Λ)) so that $0 \leq \lambda(B) \leq M\mu(B)$. (The construction we have given corresponds to the usual "fractionating process" for constructing nonatomic economies. The limit $\lim_{k \rightarrow \infty} \Lambda(f_k)/\|f_k\|$ is to be interpreted as the limiting per-capita payoff to a coalition with a given distribution of types, so $\lambda(B)$ is the limiting payoff, normalized relative to the number of players.)

Recall that the core of λ consists of all nonnegative, finitely-additive set functions $\sigma : \mathcal{B} \rightarrow \mathbb{R}$ such that $\sigma(I) = \lambda(I)$ and $\sigma(B) \geq \lambda(B)$ for each $B \in \mathcal{B}$. (If σ is in the core, we may interpret $\sigma(I_t)$ as the total (normalized) payoff to the set of agents of type t .) It is a useful fact that every element of the core treats agents of the same type equally.

LEMMA 5: Let σ belong to the core of λ . Then for each t and each Borel subset A of I_t ,

$$\sigma(A) = \sigma(I_t)\mu(A)/\mu(I_t) .$$

PROOF: Let us first perform a preliminary calculation. Suppose that B is a Borel subset of I such that $\mu(B \cap I_t)/\mu(B) = \mu(I_t) = \theta_t$ for each t (so that B has the same relative distribution as I); write $B' = I \setminus B$. The definition of λ implies that

$$\lambda(B) = \lambda(I)\mu(B) ,$$

$$\lambda(B') = \lambda(I)\mu(B') .$$

Since σ is in the core of λ , we obtain

$$\sigma(B) \geq \lambda(B) = \lambda(I)\mu(B) ,$$

$$\sigma(B') \geq \lambda(B') = \lambda(I)\mu(B') .$$

Additivity of σ and μ imply that

$$\sigma(I) = \sigma(B) + \sigma(B') \geq \lambda(I) .$$

Since $\sigma(I) = \lambda(I)$, we conclude that $\sigma(B) = \lambda(B)$ and $\sigma(B') = \lambda(B')$.

Now suppose that A_1, A_2 are subsets of I_t with $\mu(A_1) = \mu(A_2)$; we claim that $\sigma(A_1) = \sigma(A_2)$. To see this, we write $\rho = \mu(A_1)/\theta_t = \mu(A_2)/\theta_t$, and choose a subset C of $I \setminus I_t$ such that $\mu(C \cap I_s)/\theta_s = \rho$ for each $s \neq t$. Our preliminary calculation implies that

$$\sigma(A_1 \cup C) = \lambda(A_1 \cup C) = \lambda(I)\mu(A_1 \cup C) ,$$

$$\sigma(A_2 \cup C) = \lambda(A_2 \cup C) = \lambda(I)\mu(A_2 \cup C) .$$

Since $\mu(A_1 \cup C) = \mu(A_2 \cup C)$, we conclude that $\sigma(A_1 \cup C) = \sigma(A_2 \cup C)$, and additivity of σ implies that $\sigma(A_1) = \sigma(A_2)$.

Finally, let A be a Borel subset of I_t , and let m be a positive integer. There is a unique integer r such that

$$\frac{r}{m} \leq \frac{\mu(A)}{\theta_t} < \frac{r+1}{m} .$$

We may then choose disjoint Borel subsets $E_1, \dots, E_m$ of I_t such that $\bigcup E_i = I_t$, $\mu(E_i) = \theta_t/m$ for each i , $\bigcup_{i=1}^r E_i \subset A \subset \bigcup_{i=1}^{r+1} E_i$. Our previous calculations, together with additivity of σ yield that $\sigma(E_i) = \sigma(I_t)/m$ for each i , and hence that

$$r\sigma(I_t)/m \leq \sigma(A) \leq (r+1)\sigma(I_t)/m .$$

Letting m tend to infinity now yields the desired result. $\blacksquare$

For each σ in the core of λ , define $\bar{\sigma} \in \mathbb{R}^T$ by $\bar{\sigma}(t) = \sigma(I_t)/\mu(I_t)$. In view of Lemma 5, we can unambiguously interpret $\bar{\sigma}(t)$ as the per-capita payoff to players of type t . Let

$$\bar{C}(\lambda) = \{ \bar{\sigma} : \sigma \text{ is in the core of } \lambda \} ,$$

so that $\bar{C}(\lambda)$ is a subset of $\mathbb{R}^T$.

We now fix a sequence $\{(N_k, v_{\alpha_k})\}$ of games derived from the technology (Ω, Λ) . We assume that $|N_k| \rightarrow \infty$ and that $\alpha_k^{-1}(\omega_t)/|N_k| \rightarrow \theta_t$ for each t ; there is also no loss of generality in assume that $\alpha_k^{-1}(\omega_t) > 0$ for each k and t . We view the nonatomic game λ as the (normalized) limit of the games (N_k, v_{α_k}) . For $\epsilon > 0$ we will say that a payoff x in the individually rational ϵ -core of (N_k, v_k) is an equal-treatment payoff if $x(i) = x(j)$ whenever $\alpha_k(i) = \alpha_k(j)$ (so that players of the same type receive the same payoff). For such an x , we define $\bar{x} \in \mathbb{R}^T$ by $\bar{x}(t) = x(i)$ for any $i \in N_k$ with $\alpha(i) = t$; we write $\bar{C}_\epsilon(N_k, v_{\alpha_k})$ (for the set of such vectors, so that $\bar{C}_\epsilon(N_k, v_k)$ is a subset of $\mathbb{R}^T$). Of course, for $\bar{x} \in \bar{C}_\epsilon(N_k, v_k)$, we may interpret $\bar{x}(t)$ as the per-capita payoff to players of type t . Evidently, then, to show that the sets $\bar{C}_\epsilon(N_k, v_{\alpha_k})$ and $\bar{C}(\lambda)$ are close is to show that, in a natural sense, the individually rational ϵ -core of (N_k, v_{α_k}) (and the Shapley value in particular) is close to the core of λ ; this is what we are going to do. We use the Hausdorff distance between sets as the measure of closeness; for information about the Hausdorff distance and the $\limsup$ and $\liminf$ of a sequence of sets, we refer to Hildenbrand (1974).

THEOREM 2: Given $\delta_0 > 0$ and $\epsilon_0 > 0$ there is an ϵ_1 with $0 < \epsilon_1 < \epsilon_0$ and an integer k_0 such that

$$\text{dist}(\bar{C}_{\epsilon_1}(N_k, v_{\alpha_k}), \bar{C}(\lambda)) < \delta_0$$

for every $k \geq k_0$. Equivalently,

$$\begin{aligned} \bar{C}(\lambda) &= \bigcap_{\epsilon > 0} \limsup_{k \rightarrow \infty} \bar{C}_\epsilon(N_k, v_{\alpha_k}) \\ &= \bigcap_{\epsilon > 0} \liminf_{k \rightarrow \infty} \bar{C}_\epsilon(N_k, v_{\alpha_k}) . \end{aligned}$$

In particular, if $\xi_k = \text{Sh}(N_k, v_{\alpha_k})$, then

$$\lim_{k \rightarrow \infty} \text{dist}(\xi_k, \bar{C}(\lambda)) = 0 .$$

PROOF: We write $C_\epsilon^k = \bar{C}_\epsilon(N_k, v_{\alpha_k})$. Note that each C_ϵ^k is a compact set. Moreover, if $\bar{x} \in C_\epsilon^k$, then for each t ,

$$\bar{x}(t) \leq v_{\alpha_k}(N_k)/|\alpha_k^{-1}(\omega_t)| \leq \frac{M|N_k|}{|\alpha_k^{-1}(\omega_t)|}.$$

Since $|N_k|/|\alpha_k^{-1}(\omega_t)| \rightarrow \theta_t$, we conclude that the sets C_ϵ^k are in fact uniformly bounded. We have already noted that, given $\epsilon > 0$, the sets C_ϵ^k are nonempty for sufficiently large k . It follows then that $\limsup_{k \rightarrow \infty} C_\epsilon^k \neq \emptyset$ for each $\epsilon > 0$. Since $C_{\epsilon'}^k \subset C_\epsilon^k$ whenever $\epsilon' < \epsilon$, we also have that $\limsup C_{\epsilon'}^k \subset \limsup C_\epsilon^k$ whenever $\epsilon' < \epsilon$. Thus $\{\limsup C_\epsilon^k\}$ is a nested family of compact sets, and in particular, $\bigcap_{\epsilon > 0} \limsup C_\epsilon^k \neq \emptyset$.

The next step is to show that $\bigcap_{\epsilon > 0} \limsup C_\epsilon^k \subset \bar{C}(\lambda)$. Let $\bar{x} \in \bigcap_{\epsilon > 0} \limsup C_\epsilon^k$; then $\bar{x} \in \limsup C_\epsilon^k$ for each $\epsilon > 0$, so there is an increasing sequence $\{k_n\}$ of positive integers, a decreasing sequence $\{\epsilon_n\}$ of positive numbers tending to zero, and a sequence $\{\bar{x}_n\}$ converging to $\bar{x}$ with $\bar{x}_n \in C_{\epsilon_n}^{k_n}$ for each n . We claim that $\bar{x}$ is in $\bar{C}(\lambda)$.

To see this, set $\sigma = \sum_{t=1}^T \bar{x}(t)\mu_t$; we need to show that σ belongs to the core of λ . First of all, note that

$$\begin{aligned} \sigma(I) &= \sum_{t=1}^T \bar{x}(t)\mu_t(I) \\ &= \sum_{t=1}^T \bar{x}(t)\theta_t \\ &= \sum_{t=1}^T \left(\lim_{n \rightarrow \infty} \bar{x}_n(t) \right) \left(\lim_{n \rightarrow \infty} \frac{|\alpha_{k_n}^{-1}(\omega_t)|}{|N_{k_n}|} \right) \\ &= \sum_{t=1}^T \lim_{n \rightarrow \infty} \left(\frac{\bar{x}_n(t) |\alpha_{k_n}^{-1}(\omega_t)|}{|N_{k_n}|} \right) \end{aligned}$$

$$\begin{aligned}
&= \lim_{n \rightarrow \infty} \frac{\sum_{t=1}^T \bar{x}_n(t) |\alpha_{k_n}^{-1}(\omega_t)|}{|N_{k_n}|} \\
&= \lim_{n \rightarrow \infty} \frac{v_{\alpha_{k_n}}(N_{k_n})}{|N_{k_n}|} \\
&= \lambda(I) .
\end{aligned}$$

Thus σ is feasible. If σ were not in the core we could find a Borel set B for which $\lambda(B) > \sigma(B)$; hence there would be a positive number ρ such that

$$\frac{\lambda(B)}{\mu(B)} > \frac{\sigma(B)}{\mu(B)} + \rho .$$

Choose coalitions $S_n \subset N_{k_n}$ such that $|S_n| \rightarrow \infty$ and

$$\frac{|\alpha_{k_n}^{-1}(\omega_t) \cap S_n|}{|S_n|} \rightarrow \frac{\mu(B \cap I_t)}{\mu(B)}$$

for each t . The definition of λ yields that

$$\frac{\lambda(B)}{\mu(B)} = \lim_{n \rightarrow \infty} \frac{v_{\alpha_{k_n}}(S_n)}{|S_n|}$$

so we have

$$\frac{v_{\alpha_{k_n}}(S_n)}{|S_n|} > \sum_{t=1}^T \frac{\bar{x}_n(t) |\alpha_{k_n}^{-1}(\omega_t) \cap S_n|}{|S_n|} + \frac{\rho}{4}$$

for n large. If we write x_n for the equal-treatment payoff in the ϵ_n -core of $(N_{k_n}, v_{\alpha_{k_n}})$ which gives rise to $\bar{x}_n$, then our last inequality implies that

$$v_{\alpha_{k_n}}(S_n) > x_n(S_n) + \frac{\rho}{4} |S_n|$$

for large n . However, since $\epsilon_n \rightarrow 0$, this contradicts the fact that x_n is in the ϵ_n -core of (N_k, v_{α_k}) for large n . We conclude that σ is in the core of λ and that $\bar{x}$ belongs to $\bar{C}(\lambda)$, as asserted. Thus $\bigcap_{\epsilon > 0} \limsup C_\epsilon^k \subset \bar{C}(\lambda)$; note that in particular, $\bar{C}(\lambda) \neq \emptyset$.

The next step is to show that $\bar{C}(\lambda) \subset \bigcap_{\epsilon > 0} \liminf C_\epsilon^k$. Given $\bar{\sigma} \in \bar{C}(\lambda)$ and an $\epsilon > 0$, we must therefore construct, for each large k , a vector $\bar{x}_k \in C_\epsilon^k$ such that $\bar{x}_k \rightarrow \bar{\sigma}$. To this end, define, for each k , a vector $y_k \in \mathbb{R}^{N_k}$ by $y_k(i) = \bar{\sigma}(\alpha_k(i))$. We will show that a small perturbation of y_k is in the individually rational ϵ -core of (N_k, v_{α_k}) .

It is convenient and involves no loss of generality to assume that our games are zero-normalized, so that $v_{\alpha_k}(\{i\}) = 0$ for each k and i .

Write $\sigma = \sum \bar{\sigma}(t) \mu_t$, so that σ is in the core of λ . By definition,

$$\sigma(I) = \lim \frac{v_{\alpha_k}(N_k)}{|N_k|}.$$

Since

$$y_k(N_k) = \sum y_k(i) = \sum_{t=1}^T \bar{\sigma}(t) |\alpha_k^{-1}(\omega_t)|$$

and
$$|\alpha_k^{-1}(\omega_t)| / |N_k| \rightarrow \theta_t = \mu(I_t),$$

we conclude that

$$\frac{y_k(N_k)}{|N_k|} \rightarrow \sigma(I),$$

and

$$\frac{v_{\alpha_k}(N_k)}{y_k(N_k)} \rightarrow 1.$$

Thus if we set

$$z_k = \frac{v_{\alpha_k}(N_k)}{y_{\alpha_k}(N_k)} y_k ,$$

we obtain a feasible, Pareto optimal, individually rational equal-treatment payoff for the game (N_k, v_{α_k}) , and clearly $\bar{z}_k \rightarrow \bar{\sigma}$. We need to see that z_k is in the ϵ -core of (N_k, v_{α_k}) for k large. If this were not so then for each k we could find a coalition $S_k \subset N_k$ such that

$$\frac{v_{\alpha_k}(S_k)}{|S_k|} > \frac{z_k(S_k)}{|S_k|} + \epsilon .$$

Passing to a subsequence if necessary, we may assume that

$\beta_t = \lim_{k \rightarrow \infty} |\alpha_k^{-1}(\omega_t) \cap S_k|/|S_k|$ exists for each t . Let B be any Borel subset of I such that $\mu(B \cap I_t)/\mu(B) = \beta_t$ for each t . Then

$$\frac{\lambda(B)}{\mu(B)} = \lim_{k \rightarrow \infty} \frac{v_{\alpha_k}(S_k)}{|S_k|} .$$

On the other hand, the equal-treatment nature of z_k and the fact that $\bar{z}_k \rightarrow \bar{\sigma}$ imply that

$$\frac{\sigma(B)}{\mu(B)} = \lim_{k \rightarrow \infty} \frac{z_k(S_k)}{|S_k|} .$$

Hence

$$\lambda(B) > \sigma(B) + \epsilon \mu(B)$$

which contradicts the fact that σ is in the core of λ . We conclude that z_k is in the ϵ -core of (N_k, v_k) for k sufficiently large. Hence $\bar{C}(\lambda) \subset \bigcap_{\epsilon > 0} \liminf_{k \rightarrow \infty} C_{\epsilon}^k$.

We have now shown that

$$\bigcap_{\epsilon > 0} \limsup_{k \rightarrow \infty} C_{\epsilon}^k \subset \bar{C}(\lambda) \subset \bigcap_{\epsilon > 0} \liminf_{k \rightarrow \infty} C_{\epsilon}^k ;$$

Since $\limsup_{k \rightarrow \infty} C_{\epsilon}^k \supset \liminf_{k \rightarrow \infty} C_{\epsilon}^k$ for every ϵ , it follows that

$$\bigcap \limsup C_\epsilon^k = \bar{C}(\lambda) = \bigcap \liminf C_\epsilon^k.$$

The first statement of the Theorem is an easy consequence of this fact. To see this fix $\delta_0 > 0$, $\epsilon_0 > 0$, and let U be the δ_0 -neighborhood of $\bar{C}(\lambda)$; i.e.,

$$U = \{\bar{y} \in \mathbb{R}^T : \text{dist}(\bar{y}, \bar{C}(\lambda)) < \delta_0\}.$$

Since $\bar{C}(\lambda) = \bigcap \limsup C_\epsilon^k$, we can find an ϵ_1 , $0 < \epsilon_1 < \epsilon_0$, such that $U \supset \limsup C_{\epsilon_1}^k$. Hence $U \supset C_{\epsilon_1}^k$ for all sufficiently large k . Writing U_k for the δ_0 -neighborhood of $C_{\epsilon_1}^k$, we need to show that $\bar{C}(\lambda) \subset U_k$ for all sufficiently large k . If this were not so we could find a sequence of integers $\{k_n\}$ tending to infinity and a sequence $\{\bar{\sigma}_n\}$ in $\bar{C}(\lambda)$ such that $\text{dist}(\bar{\sigma}_n, C_{\epsilon_1}^{k_n}) \geq \delta_0$ for each n . Passing to a subsequence if necessary, we may assume that $\bar{\sigma}_n \rightarrow \bar{\sigma}$. But then $\text{dist}(\bar{\sigma}, C_{\epsilon_1}^{k_n}) \geq \frac{1}{2} \delta_0$ for k_n large, so that $\bar{\sigma}$ would be a point of $\bar{C}(\lambda)$ which was not in $\liminf C_{\epsilon_1}^k$. This is a contradiction, so we must have $\bar{C}(\lambda) \subset U_k$ for all sufficiently large k . In other words

$$\text{dist}(\bar{C}_{\epsilon_1}^k, \bar{C}(\lambda)) < \delta_0$$

for all sufficiently large k , as desired.

The final assertion, about the Shapley value, is an immediate consequence of this distance estimate and Theorem 1, so the proof is complete. $\square$

We remark that a similar argument may be used to show that, for every $\epsilon > 0$, $\liminf C_\epsilon^k = \limsup C_\epsilon^k$; this set may be interpreted as representing the set of equal-treatment payoffs in the ϵ -core of λ . (Of course, this is not true if $\epsilon = 0$, as the example in Section 5 shows.) We shall not go into the details here.

It is tempting to suppose that the (per-capita normalized) Shapley values of the games (N_k, v_{α_k}) converge to the asymptotic Shapley value of the nonatomic game λ . Unfortunately, λ need not have an asymptotic Shapley value (see the "three-handed glove" market in Aumann and Shapley (1984), for instance). Perhaps such a result could be proved using the more general value of Mertens (1983).

REFERENCES

- Anderson, R.M. (1978), An elementary core equivalence theorem, *Econometrica* 46, 1483-1487.
- Aumann, R.J. (1975), Values of markets with a continuum of traders, *Econometrica* 43, 611-646.
- Aumann, R.J. and L. Shapley (1974), "Values of Non-Atomic Games," Princeton Univ. Press, Princeton, NJ.
- Champsaur, P. (1975), Cooperation vs. competition, *J. Econ. Theory* 11, 394-417.
- Cheng, H.-C. (1981), On dual regularity and value convergence theorems, *J. Math. Econ.* 8, 37-57.
- Feller, W. (1968), "An Introduction to Probability Theory and its Applications," vol. I, John Wiley and Sons, New York.
- Hart, S. (1977), Values of non-differentiable markets with a continuum of traders, *J. Math. Econ.* 4, 103-116.
- Hildenbrand, W. (1974), "Core and Equilibria of a Large Economy," Princeton Univ. Press, Princeton, NJ.
- Kuhn, H.W. and A.W. Tucker (1950), Preface, in "Contributions to the Theory of Games," *Annals of Math. Studies* 24, v-xiii.
- Mas-Colell, A. (1977), Competitive and value allocations of large exchange economies, *J. Econ. Theory* 14, 419-438.
- Mertens, J.F. (1983), The Shapley value in the nondifferentiable case, Technical Report No. 417, The Economics Series, Institute for Mathematical Studies in the Social Sciences, Stanford University.
- Roth, A.E. (1977), The Shapley value as a vonNeumann-Morgenstern utility, *Econometrica* 45
- Shapley, L. (1964), Values of large games - VII: A general exchange economy with money, Rand Memorandum RM-4248-PR.
- Shapley, L.S. and M. Shubik (1966), "Quasi-cores in a monetary economy with nonconvex preferences, *Econometrica* 34, 805-827.
- Shapley, L. and M. Shubik (1969), Pure competition, coalitional power, and fair division, *Internat. Econ. Rev.* 10, 337-362.
- Wooders, M.H. and W.R. Zame (1984), Approximate cores of large games, *Econometrica*, to appear.