**ASYMPTOTICS FOR A PARABOLIC DOUBLE
OBSTACLE PROBLEM**

By

Xinfu Chen

and

Charles M. Elliott

IMA Preprint Series # 922

February 1992

ASYMPTOTICS FOR A PARABOLIC DOUBLE OBSTACLE PROBLEM

XINFU CHEN¹ AND CHARLES M. ELLIOTT²

Abstract. We consider a parabolic double obstacle problem which is a version of the Allen–Cahn equation $u_t = \Delta u - \varepsilon^{-2}\psi'(u)$ in $\Omega \times (0, \infty)$, where Ω is a bounded domain, ε is a small constant, and ψ is a double well potential; here we take ψ such that $\psi(u) = (1 - u^2)$ when $|u| \leq 1$ and $\psi(u) = \infty$ when $|u| > 1$. We study the asymptotic behavior, as $\varepsilon \rightarrow 0$, of the solution of the double obstacle problem. Under some natural restrictions on the initial data, we show that after a short time (of order $\varepsilon^2 |\ln \varepsilon|$), the solution takes value 1 in a region Ω_t^+ and value -1 in Ω_t^- , where the region $\Omega \setminus (\Omega_t^+ \cup \Omega_t^-)$ is a thin strip and is contained in a $O(\varepsilon |\ln \varepsilon|)$ neighborhood of a hypersurface Γ_t which moves with normal velocity equal to its mean curvature. We also study the asymptotic behavior, as $t \rightarrow \infty$, of the solution in the one-dimensional case. In particular, we prove that the ω -limit set consists of a singleton.

1 Introduction.

The Allen–Cahn equation

$$u_t = 2\alpha k \Delta u - \alpha \psi'(u) \quad \text{in } \mathcal{R}^3 \times (0, \infty) \quad (1.1)$$

was introduced by Allen–Cahn [1979] in order to describe the motion of an ‘anti-phase’ boundary separating two phases of a polycrystalline material. Each phase is characterized by the value taken by the order parameter u at the minima of the double equal well potential $\psi : \mathcal{R} \rightarrow \mathcal{R}$. With the scaling

$$\alpha = \frac{1}{\varepsilon^2}, \quad k = \frac{\varepsilon^2}{2}, \quad \varepsilon \rightarrow 0, \quad (1.2)$$

it was proposed that the interface evolves according to the mean curvature flow

$$V = -K^m \quad (1.3)$$

where V is the velocity of the interface and K^m is the sum of its principle curvatures. Formal asymptotics showing this were carried out by Rubinstein–Sternberg–Keller [1989]. Rigorous results have been obtained by Fife–Hsiao [1988], Carr–Pego

¹Department of Mathematics, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA. The author was supported by the Sloan Doctoral Dissertation Fellowship during the 1990/91 academic year.

²School of Mathematical and Physical Sciences, University of Sussex, Brighton, BN1 9QH, UK. The author was supported by the Institute for Mathematics and its Applications (Minneapolis) with funds provided by the National Science Foundation while he was visiting the IMA during the 1990/91 program on ‘Phase transitions and Free Boundary problems’.

[1989, 1990], Fusco–Hale [1989], and Fusco [1990] in the one dimensional case and by Bronsard–Kohn [1991], DeMottoli–Schatzman [1989], Chen [1990], and Evans–Soner–Souganides [1990] in the higher dimensional case.

The purpose of this paper is to study the version of (1.1) obtained by replacing the smooth double well potential ψ by

$$\psi(u) = \begin{cases} \frac{1}{2}(1 - u^2) & |u| \leq 1, \\ +\infty & |u| > 1. \end{cases} \quad (1.4)$$

With the scaling (1.2) it follows that we study the parabolic double obstacle problem: find $u(x, t)$, $(x, t) \in \Omega \times (0, T)$, such that for a.e. (x, t) ,

$$\begin{cases} (u_t - \Delta u - \frac{1}{\varepsilon^2}u) \text{Sign } u \leq 0, \\ |u| \leq 1, \\ (u_t - \Delta u - \frac{1}{\varepsilon^2}u)(|u| - 1) = 0. \end{cases} \quad (1.5)$$

In particular we impose the boundary condition

$$\frac{\partial u}{\partial n} = 0, \quad (x, t) \in \partial\Omega \times (0, T) \quad (1.6)$$

where Ω is a bounded domain in $\mathcal{R}^N$ with a sufficient smooth boundary $\partial\Omega$. (Either Ω is convex and $\partial\Omega$ is Lipschitz or $\partial\Omega$ is smooth.)

We note that (1.5) and (1.6) is a gradient flow for the energy functional

$$E(v) := \int_{\Omega} \left\{ |\nabla v|^2 + \frac{1}{\varepsilon^2}(1 - v^2) \right\} dx, \quad (1.7)$$

$$v \in \mathcal{K} := \left\{ \eta \in H^1(\Omega) : |\eta(x)| \leq 1 \text{ a.e. } x \in \Omega \right\}. \quad (1.8)$$

The existence theory for (1.5) and (1.6) together with asymptotic long time behavior and a comparison lemma is given in §2. Using the method of Chen [1990], the connection with smooth mean curvature flow is rigorously established in §3. Finally in §4 detailed and precise results concerning the asymptotic long time behavior in one space dimension are proved.

For other connections between double obstacle problems and curvature dependent phase boundary motion we refer to Blowey and Elliott [1991a,b,c].

2 The Double Obstacle Problem.

In this section we briefly expose the existence and regularity theory for the parabolic variational inequality version of the Allen–Cahn equation.

2.1. EXISTENCE.

Let $\mathcal{K}$ be as in (1.8) and

$$\mathcal{G} := \left\{ \eta \in L^\infty(\Omega) : |\eta(x)| \leq 1 \text{ a.e. } x \in \Omega \right\}.$$

Given $g \in \mathcal{G}$ and $T > 0$ we say that $u \in C[0, T; L^2(\Omega)]$ is a solution to the double obstacle problem provided

$$\left\{ \begin{array}{l} u \in L^2(0, T; H^1(\Omega)), \quad u_t \in L^2(0, T; (H^1(\Omega))'), \\ u(\cdot, t) \in \mathcal{K} \text{ a.e. } t \in (0, T), \quad u(\cdot, 0) = g(\cdot), \\ \int_0^T \left\{ \langle u_t, \eta - u \rangle + (\nabla u, \nabla(\eta - u)) - \frac{1}{\varepsilon^2} (u, \eta - u) \right\} dt \geq 0 \\ \forall \eta \in L^2(0, T; H^1(\Omega)), \quad \eta(t) \in \mathcal{K} \quad \forall t. \end{array} \right. \quad (2.1)$$

We have the following theorem concerning existence and uniqueness.

THEOREM 2.1. (1) For each $g \in \mathcal{G}$ there exists a unique solution to (2.1) and the mapping $g \rightarrow u(t)$ is continuous in $L^2(\Omega)$ for each $t > 0$.

(2) For each $T > 0$ and $\tau > 0$, the following regularity holds:

$$\begin{aligned} \sqrt{t} u_t &\in L^2(0, T; L^2(\Omega)), \\ u &\in C[\tau, T; H^1(\Omega)] \cap L^2(\tau, T; H^2(\Omega)) \cap C^{1+\alpha, (1+\alpha)/2}(\Omega \times (\tau, T)). \end{aligned}$$

(3) There exist $\rho (= \rho(\varepsilon))$ and $t^* (= t^*(\varepsilon))$ such that for each $g \in \mathcal{G}$

$$\begin{aligned} \|u(t)\|_{H^1(\Omega)} &\leq \rho & \forall t \geq t^*, \\ \|u_t\|_{L^2(\Omega)} + \|u(t)\|_{H^2(\Omega)} &\leq \rho & \forall t \geq t^*. \end{aligned}$$

(4) For any $0 < \tau < t$,

$$E(u(t)) + 2 \int_\tau^t \|u_t(s)\| ds = E(u(\tau)).$$

(5) u solves the parabolic problem

$$\left\{ \begin{array}{ll} u_t - \Delta u = \beta \in \frac{1}{\varepsilon^2} \beta(u) & \text{in } \Omega \times (0, \infty), \\ \partial u / \partial n = 0 & \text{on } \partial \Omega \times (0, \infty), \\ u(x, 0) = g(x) & \text{on } \Omega \times \{0\}, \end{array} \right. \quad (2.2)$$

where $\beta(u)$ is a maximal graph given by

$$\beta(u) = \left\{ \begin{array}{ll} \infty & \text{if } u < -1, \\ [-1, 0] & \text{if } u = -1, \\ u & \text{if } u \in (-1, 1), \\ [0, 1] & \text{if } u = 1, \\ -\infty & \text{if } u > 1. \end{array} \right. \quad (2.3)$$

Proof. Existence and regularity are proven using the standard regularization technique for parabolic variational inequalities. For each $\delta > 0$, let $\beta^\delta \in C^\infty(\mathcal{R})$ satisfy

$$\begin{cases} \beta^\delta(r) = r & \forall |r| \leq 1 - \delta, \\ \beta^\delta(\pm 1) = 0, \\ (\beta^\delta(r))' \leq 1 & \forall |r| \leq 1. \end{cases} \quad (2.4)$$

It follows that

$$\begin{aligned} r\beta^\delta(r) &\leq r^2 & \forall |r| \leq 1, \\ \lim_{\delta \rightarrow 0} \beta^\delta(r) &= r & \forall |r| < 1. \end{aligned} \quad (2.5)$$

For convenience we define

$$B_\delta(u) = \int_0^u \beta^\delta(r) dr$$

so that

$$0 \leq B_\delta(u) \leq u^2/2 \quad \forall |u| \leq 1. \quad (2.6)$$

Standard theory for quasi-linear parabolic equations yields the existence of a unique solution to

$$\begin{cases} u^\delta_t - \Delta u^\delta = \frac{1}{\varepsilon^2} \beta^\delta(u^\delta), & x \in \Omega, t > 0, \\ \partial u^\delta / \partial n = 0, & x \in \Omega, t > 0 \\ u^\delta(x, 0) = g(x), & x \in \Omega \end{cases} \quad (2.7)$$

which satisfies, by maximum principle because of (2.4b) and (2.5),

$$|u^\delta(x, t)| \leq 1 \quad \forall (x, t) \in \Omega \times (0, \infty). \quad (2.8)$$

Multiplying (2.7a) by u^δ and integrating yield

$$\frac{1}{2} \frac{d}{dt} \|u^\delta\|^2 + \|\nabla u^\delta\|^2 = \frac{1}{\varepsilon^2} \langle \beta^\delta(u^\delta), u^\delta \rangle \leq \frac{|\Omega|}{\varepsilon^2}$$

so that

$$\int_0^t \|\nabla u^\delta(s)\|^2 ds \leq |\Omega| \left(\frac{1}{2} + \frac{t}{\varepsilon^2} \right). \quad (2.9)$$

Multiplying (2.7a) by u^δ_t and integrating yield

$$\frac{1}{2} \frac{d}{dt} \|\nabla u^\delta\|^2 + \|u^\delta_t\|^2 = \frac{1}{\varepsilon^2} \frac{d}{dt} \int_\Omega B_\delta(u^\delta) dx$$

so that for each $t > \tau > 0$,

$$2 \int_\tau^t \|u^\delta_t(s)\|^2 ds + E^\delta(u^\delta(t)) = E(u^\delta(\tau)) \quad (2.10)$$

where

$$E^\delta(\eta) = \int_{\Omega} \left\{ |\nabla \eta|^2 + \frac{1}{\varepsilon^2} - \frac{2}{\varepsilon^2} B_\delta(\eta) \right\} dx. \quad (2.11)$$

It follows from (2.10), (2.6) that

$$2 \int_{\tau}^t \|u^\delta_t(s)\|^2 ds + \|\nabla u^\delta(t)\|^2 \leq \|\nabla u^\delta(\tau)\|^2 + \frac{|\Omega|}{\varepsilon^2}. \quad (2.12)$$

Similarly multiplying (2.7a) by tu^δ_t and integrating yield

$$\begin{aligned} \int_0^t s \|u^\delta_t(s)\|^2 ds + \frac{1}{2} t \|\nabla u^\delta(t)\|^2 &= \frac{1}{2} \int_0^t \|\nabla u^\delta(s)\|^2 ds \\ &\quad + \frac{1}{\varepsilon^2} t \int_{\Omega} B_\delta(u^\delta(t)) - \frac{1}{\varepsilon^2} \int_0^t \int_{\Omega} B_\delta(u^\delta) \\ &\leq \frac{|\Omega|}{2} \left(\frac{1}{2} + \frac{t}{\varepsilon^2} \right) + \frac{|\Omega|t}{2\varepsilon^2}. \end{aligned}$$

by (2.6) and (2.9). It follows that

$$\|\nabla u^\delta(t)\|^2 \leq |\Omega| \left(\frac{1}{2t} + \frac{2}{\varepsilon^2} \right).$$

Therefore there exists t^* such that for $t > t^*/2$,

$$\|u^\delta(t)\|_{H^1(\Omega)} \leq \frac{3|\Omega|}{\varepsilon^2}. \quad (2.13)$$

Differentiating (2.7a) with respect to t , multiplying by $(t - t^*/2)u^\delta_t$, and using (2.4c) yield

$$\frac{d}{dt} \left((t - t^*/2) \|u^\delta_t\|^2 \right) \leq \left(1 + \frac{2(t - t^*/2)}{\varepsilon^2} \right) \|u^\delta_t\|^2 \quad \forall t > t^*/2,$$

and integrating in time and using (2.12), (2.13) we obtain for $t > t^*/2$,

$$(t - t^*/2) \|u^\delta_t(t)\|^2 \leq \left(1 + \frac{2(t - t^*/2)}{\varepsilon^2} \right) \frac{1}{2} \left(\frac{3|\Omega|}{\varepsilon^2} + \frac{|\Omega|}{\varepsilon^2} \right).$$

Since (2.8) holds we have from (2.7a), (2.5) that

$$\|u^\delta(t)\|_{H^2(\Omega)} \leq C(\|u^\delta_t\| + 1/\varepsilon^2).$$

For different initial data g_1 and g_2 it is an easy calculation to show

$$\|u^\delta_1(t) - u^\delta_2(t)\| \leq e^{c_\varepsilon t} \|g_1 - g_2\| \quad (2.14)$$

for the two solutions $\{u^\delta_i\}_{i=1}^2$ and C_ε a positive constant independent of δ and $\{g_i\}$.

The statement of the theorem is a standard consequence of a compactness argument with

$$u(t) = \lim_{\delta \rightarrow 0} u^\delta(t),$$

$$\beta(u) \ni \beta = w^* - \lim_{\delta \rightarrow 0} \beta_\delta(u^\delta).$$

2.2. ASYMPTOTIC BEHAVIOR AS $t \rightarrow \infty$.

Let $\{\mathcal{S}(t)\}_{t \geq 0}$ be the family of solution operators defined by $\mathcal{S}(t)g := u(t)$ where $u(\cdot)$ solves (2.1) with g as initial data. It follows from Theorem 2.1 that $\{\mathcal{S}(t)\}_{t \geq 0}$ forms a continuous semigroup on $\mathcal{G} \subset L^2(\Omega)$. We denote by $\mathcal{E}$ the set of equilibria defined by

$$\mathcal{E} \equiv \left\{ v \in \mathcal{K} \mid \varepsilon^2(\nabla v, \nabla \eta - \nabla v) \geq (v, \eta - v), \quad \forall \eta \in \mathcal{K} \right\}. \quad (2.15)$$

For each $g \in \mathcal{G}$ we define the ω -limit set $\omega(g)$ by

$$\omega(g) = \left\{ v \in \mathcal{G} \mid \exists \{t_n\} \rightarrow \infty \text{ such that } \|\mathcal{S}(t_n)g - v\| \rightarrow 0 \text{ as } n \rightarrow \infty \right\}. \quad (2.16)$$

THEOREM 2.2. *The semigroup $\{\mathcal{S}(t)\}_{t \geq 0}$ possesses a global attractor $\mathcal{A}$ which is bounded in $\mathcal{K}$. Furthermore for any $g \in \mathcal{G}$ it holds that $\omega(g) \subset \mathcal{E}$ and there exists a constant e such that $e = \mathcal{E}(v)$, $\forall v \in \omega(g)$.*

Proof. We follow the exposition given in Temam (1988). It follows from Theorem 2.1 that $\{\mathcal{S}(t)\}_{t \geq t^*}$ is uniformly compact and that there exists a ρ such that $\mathcal{B} = \{\eta \in \mathcal{K} \mid \|\eta\|_{H^1(\Omega)} \leq \rho\}$ is an absorbing set. Hence the existence of a global attractor is an immediate consequence of Temam [1988; Theorem 1.1, p23].

Since $E(\cdot) \geq 0$, it follows from the results (2), (3), and (4) of Theorem 2.1 that $E(\mathcal{S}(t)g)$ is a bounded decreasing continuous function on (t^*, ∞) and so has a unique limit $e = \lim_{t \rightarrow \infty} E(\mathcal{S}(t)g)$. Since $\cup_{t \geq t^*} \mathcal{S}(t)g$ is relatively compact, there exists a cluster point $v \in \mathcal{K}$ such that $v = \lim_{k \rightarrow \infty} \mathcal{S}(t_k)g \in \omega(g)$ and because $E(\cdot)$ is continuous on $H^1(\Omega)$ and $\mathcal{S}(t)g \in C[\tau, T; H^1(\Omega)]$ for all $T > 0$, it follows that $E(v) = \lim_{k \rightarrow \infty} E(\mathcal{S}(t_k)g) = e$. Hence, $E(v) = e$ for all $v \in \omega(g)$ and since $\omega(g)$ is invariant, $\mathcal{S}(t)v \in \omega(g)$ for all $t \geq 0$. Therefore $E(\mathcal{S}(t)v) = e = E(v)$ for all $t > 0$ and so result (4) implies that $\frac{d}{dt} \mathcal{S}(t)v = 0$ or $v \in \mathcal{E}$. $\square$

2.3. COMPARISON LEMMA.

LEMMA 2.3. *Let $w \in L^2(0, T; H^1(\Omega))$ and $w_t \in L^2(0, T; (H^1(\Omega))')$ satisfy*

$$w(x, t) \leq 1, \quad (x, t) \in \Omega \times (0, T),$$

$$w(x, 0) \leq g(x), \quad x \in \Omega$$

and

$$\int_0^T \left\{ \langle w_t, \eta \rangle + (\nabla w, \nabla \eta) - \frac{1}{\varepsilon^2} (f, \eta) \right\} \leq 0$$

$$\forall \eta \in L^2(0, T; H^1(\Omega)), \quad \eta \geq 0 \text{ a.e. } \Omega \times (0, T) \quad (2.17)$$

where $f(w - u)_+ \leq w(w - u)_+$ and u solves (2.1) with initial data $g \in \mathcal{G}$. Then

$$w \leq u \quad \text{a.e. } \Omega \times (0, T).$$

Proof. Taking $\eta = (w - u)_+$ in (2.17) and $\eta = u + (w - u)_+$ in (2.1c) gives a pair of inequalities which when added yield

$$\int_0^T \left\{ \frac{1}{2} \frac{d}{dt} \|(w - u)_+\|^2 + \|\nabla(w - u)_+\|^2 \right\} dt \leq \int_0^T \frac{1}{\varepsilon^2} \|(w - u)_+\|^2 dt.$$

The assertion follows from Gronwall's inequality. $\square$

In the sequel, we shall also use the following comparison lemma.

LEMMA 2.4. Let u be the solution of (2.1) and Ω' be an open subset of Ω with Lipschitz boundary. Assume that a function $v \in W_{\infty}^{2,1}(\Omega' \times (\tau, T))$ satisfies

$$\begin{cases} v_t - \Delta v \leq \frac{1}{\varepsilon^2} \beta(v) & \text{in } \Omega' \times (\tau, T), \\ v(x, t) \leq 1 & \text{on } \overline{\Omega'} \times [\tau, T], \\ v(x, t) \leq u(x, t) & \text{on } \partial\Omega' \times [\tau, t] \cup \Omega' \times \{0\}. \end{cases}$$

Then

$$v(x, t) \leq u(x, t) \quad \forall (x, t) \in \Omega' \times (\tau, t).$$

Here $f \leq \beta(v)$ means that $f \leq v$ if $v \in (-1, 1)$, $f \leq 1$ if $v = 1$ and $f \leq 0$ if $v = -1$.

Since u solves the parabolic problem (2.2), the proof follows from standard routine techniques, namely, subtracting the parabolic equations from each other, multiplying the resulting equation by $(v - u)_+$, integrating and using Gronwall's inequality.

The comparison lemmas for supersolutions can be stated in a corresponding way and are omitted.

3 Asymptotic behavior of $\varepsilon \rightarrow 0$.

In this section we consider the asymptotic behavior of the solution, denoted by u^ε , to the double obstacle problem (2.1). It is convenient to begin by describing the evolution of an interface according to its mean curvature.

3.1. MOTION BY MEAN CURVATURE.

The family $\{\Gamma(t)\}_{t \in [0, T]}$ of sufficiently smooth hypersurfaces in $\mathcal{R}^N$ is said to form a motion by mean curvature flow starting from Γ_0 if the following holds:

- (i) $\Gamma(t)$ is the boundary of an open set $A(t) \in \mathcal{R}^N$ for $t \in [0, T]$;
- (ii) $\Gamma(0) = \Gamma_0$;

(iii) The speed V of the hypersurface $\Gamma(t)$ in the outward normal direction to $A(t)$ is $-K^m$, the sum of the principle curvatures (mean curvature) of $\Gamma(t)$ where K^m is positive if $A(t)$ is convex.

It is convenient to describe such a flow by introducing the signed distance function $d(x, t) : \mathcal{R}^N \times [0, T] \rightarrow \mathcal{R}^1$ from x to $\Gamma(t)$ so that $d(x, t)$ is negative in $A(t)$ and positive in $\mathcal{R}^N \setminus \overline{A(T)}$. It follows that

$$\begin{aligned} \Gamma(t) &= \{x \in \mathcal{R}^N \mid d(x, t) = 0\}, & |\nabla d(x, t)| &= 1 \text{ a.e. } x \in \mathcal{R}^N, \\ d_t(x, t) &= -V(x, t), & \Delta d(x, t) &= K^m(x, t) \quad \forall x \in \Gamma(t). \end{aligned}$$

The following facts are known.

(1) If $\Gamma_0 \in C^{m+\alpha}$ ($m \geq 2$) is the boundary of a bounded open set then there exists a unique solution of the motion by mean curvature flow starting from Γ_0 in a time interval $(0, T]$ for some $T > 0$. Furthermore the signed distance function $d \in C^{m+\alpha, (m+\alpha)/2}$ in a neighborhood of $\cup_{t \in [0, T]} (\Gamma(t) \times \{t\})$.

(2) If $N = 2$ or Γ_0 is convex then the solution can be extended up to $T = T^*$ where T^* is the time at which $\Gamma(\cdot)$ shrinks to a single point.

For more details, we refer to Hamilton [1982], Gage–Hamilton [1986], Grayson [1987], Huisken [1984], Chen–Giga–Goto [1991], and Evans–Spruck [1991].

3.2. APPROXIMATION OF MEAN CURVATURE FLOW BY THE DOUBLE OBSTACLE PROBLEM.

Let $\{\Gamma(t)\}_{t \in [0, T]}$ be a mean curvature flow such that for each $t \in [0, T]$, $\Gamma(t)$ lies in the interior of a bounded open set Ω and $\text{dist}(\Gamma(t), \partial\Omega) \geq \delta > 0$ for some fixed δ . We shall assume that the associated signed distance function d satisfies

$$\sup_{t \in [0, T]} \sup_{|d| \leq c_0} |\nabla(d_t - \Delta d)| \leq D_0. \quad (3.1)$$

THEOREM 3.1. *Let $\gamma(\varepsilon)$ satisfy*

$$\lim_{\varepsilon \rightarrow 0} \gamma(\varepsilon) = 0$$

and assume that there exists ε_0 such that for $\varepsilon \in (0, \varepsilon_0)$,

$$\frac{\pi}{2}\varepsilon \leq \gamma(\varepsilon) \quad \text{and} \quad \frac{\pi}{2}\varepsilon + 2\gamma(\varepsilon)e^{2D_0T} \leq c_0. \quad (3.2)$$

Let $u^\varepsilon(x, t)$ be the unique solution of (2.1) with initial data $g^\varepsilon \in \mathcal{G} \cap C(\overline{\Omega})$ satisfying

$$\{x : d(x, 0) > \gamma(\varepsilon)\} \subset \{x : g^\varepsilon(x) = 1\}, \quad (3.3)$$

$$\{x : d(x, 0) < -\gamma(\varepsilon)\} \subset \{x : g^\varepsilon(x) = -1\}. \quad (3.4)$$

Then for all $t \in [0, T]$, it holds that

$$\{x : d(x, t) \geq \gamma(\varepsilon)(1 + 2e^{2D_0t})\} \subset \{x : u^\varepsilon(x, t) = 1\}, \quad (3.5)$$

$$\{x : d(x, t) \leq -\gamma(\varepsilon)(1 + 2e^{2D_0t})\} \subset \{x : u^\varepsilon(x, t) = -1\}. \quad (3.6)$$

Proof. The fact that $\{\Gamma(t)\}$ is a mean curvature flow implies

$$d_t - \Delta d = 0 \quad \text{on } \Gamma(t), \quad t \in (0, T]$$

and the assumption (3.1) yields

$$|d_t - \Delta d| \leq D_0 |d|$$

whenever $|d| \leq c_0$.

Set

$$z^\varepsilon(x, t) := d(x, t) - 2\gamma(\varepsilon)e^{2D_0 t}$$

and

$$\Omega^\varepsilon(t) := \{x \in \Omega : |z^\varepsilon(x, t)| < \varepsilon\pi/2\}.$$

It follows from (3.2) that

$$|d(x, t)| \leq c_0 \quad \forall x \in \Omega^\varepsilon(t), \quad t \in (0, T].$$

Hence on $\Omega^\varepsilon(t)$,

$$\begin{aligned} z_t^\varepsilon - \Delta z^\varepsilon &= d_t - \Delta d - 4D_0\gamma(\varepsilon)e^{2D_0 t} \\ &\leq D_0(|d| - 4\gamma(\varepsilon)e^{2D_0 t}) \\ &\leq D_0(\varepsilon\pi/2 - 2\gamma(\varepsilon)) \leq 0 \end{aligned}$$

by (3.2).

We now introduce $U \in C^{1,1}(\mathcal{R})$ given by

$$U(s) = \begin{cases} 1 & s \geq \pi/2, \\ \sin s & s \in (-\pi/2, \pi/2), \\ -1 & s \leq -\pi/2. \end{cases}$$

Set

$$v^\varepsilon(x, t) := U(z^\varepsilon(x, t)/\varepsilon).$$

Observe that $v^\varepsilon(x, 0) > -1$ if and only if $d(x, 0) - 2\gamma(\varepsilon) > -\varepsilon\pi/2$. Thus $v^\varepsilon(x, 0) > -1$ implies that $d(x, 0) > \gamma(\varepsilon)$ and so by (3.3) $g^\varepsilon(x) = 1$. Hence it holds that

$$-1 \leq v^\varepsilon(x, 0) \leq g^\varepsilon(x) = u^\varepsilon(x, 0).$$

Let

$$f(x, t) = \begin{cases} v^\varepsilon(x, t) & v^\varepsilon(x, t) > -1, \\ 0 & v^\varepsilon(x, t) = -1. \end{cases} \quad (3.7)$$

For any $\varphi \in C^1(\Omega \times (0, T))$ and $\varphi \geq 0$ we have

$$\begin{aligned} & \int_0^T \int_{\Omega} \left\{ v_t^\varepsilon \varphi + \nabla v^\varepsilon \nabla \varphi - \frac{1}{\varepsilon^2} f \varphi \right\} dx dt \\ &= \int_0^T \int_{\Omega^\varepsilon(t)} \left\{ v_t^\varepsilon \varphi + \nabla v^\varepsilon \nabla \varphi - \frac{1}{\varepsilon^2} f \varphi \right\} dx dt - \int \int_{\{v^\varepsilon = -1\} \cup \{v^\varepsilon = 1\}} \frac{1}{\varepsilon^2} f \varphi dx dt \\ &\leq \int_0^T \int_{\Omega^\varepsilon(t)} \left\{ v_t^\varepsilon - \Delta v^\varepsilon - \frac{1}{\varepsilon^2} v^\varepsilon \right\} \varphi dx dt, \end{aligned}$$

where we have used the fact that on $\partial\Omega^\varepsilon(t)$, $|z^\varepsilon| = \varepsilon\pi/2$ and $\nabla v^\varepsilon = \frac{1}{\varepsilon} \nabla z^\varepsilon U'(z^\varepsilon/\varepsilon) = 0$.

Calculation yields

$$v_t^\varepsilon - \Delta v^\varepsilon - \frac{1}{\varepsilon^2} v^\varepsilon = U' \frac{z_t^\varepsilon}{\varepsilon} - U' \frac{\Delta z^\varepsilon}{\varepsilon} - U'' \frac{|\nabla z^\varepsilon|^2}{\varepsilon^2} - \frac{1}{\varepsilon^2} U$$

and since on $\Omega^\varepsilon(t)$, $U'' = -U$, $U' > 0$ and $|\nabla z^\varepsilon| = 1$, we find

$$v_t^\varepsilon - \Delta v^\varepsilon - \frac{1}{\varepsilon^2} v^\varepsilon = \frac{1}{\varepsilon} U'(z_t^\varepsilon - \Delta z^\varepsilon) \leq 0 \quad \forall x \in \Omega^\varepsilon(t).$$

Therefore

$$\int_0^T \int_{\Omega} \left\{ v_t^\varepsilon \varphi + \nabla v^\varepsilon \nabla \varphi - \frac{1}{\varepsilon^2} f \varphi \right\} dx dt \leq 0.$$

By (3.7), $f(v^\varepsilon - u^\varepsilon)_+ \leq v^\varepsilon(v^\varepsilon - u^\varepsilon)_+$. It then follows by the comparison Lemma 2.3 that

$$u^\varepsilon(x, t) \geq v^\varepsilon(x, t) \quad \forall (x, t) \in \Omega \times [0, T].$$

Since $d(x, t) \geq \gamma(\varepsilon) + 2\gamma(\varepsilon)e^{2D_0 t}$ implies that $z^\varepsilon(x, t) \geq \varepsilon\pi/2$ which again implies that $v^\varepsilon(x, t) = 1$, it follows that (3.5) holds. The proof of (3.6) is identical. $\square$

REMARK 3.2. It follows from Theorem 3.1 that the set

$$\Gamma^\varepsilon(t) = \{x \in \Omega : u^\varepsilon(x, t) = 0\}$$

satisfies the estimate

$$\text{dist}(\Gamma(t), \Gamma^\varepsilon(t)) \leq c\gamma(\varepsilon) \quad \forall t \in [0, T]$$

for some c independent of ε and t . It follows that choosing

$$\gamma(\varepsilon) \leq \frac{\pi}{2}\varepsilon + c\varepsilon$$

the estimate

$$\text{dist}(\Gamma(t), \Gamma^\varepsilon(t)) \leq \bar{c}\varepsilon$$

holds.

3.3. GENERATION AND PROPAGATION OF AN INTERFACE.

In this subsection, we shall show that, for more general initial data than considered in §3.2, after a short time an interface will be formed near the set $\Gamma_0 = \{x \in \Omega : g(x) = 0\}$ and will propagate according to motion by mean curvature.

LEMMA 3.3. *Let $g \in \mathcal{G} \cap W^{1,\infty}(\Omega)$. Then there exist positive constants $\{c_1, \varepsilon_1, \tau_1\}$ such that if $\varepsilon \in (0, \varepsilon_1)$ then*

$$u^\varepsilon(x, t) \geq 1 - \varepsilon \quad \forall x \in \Omega_\varepsilon^+, t \in [\tau_1 \varepsilon^2 |\ln \varepsilon|, (\tau_1 + 1) \varepsilon^2 |\ln \varepsilon|], \quad (3.8)$$

$$u^\varepsilon(x, t) \leq -1 + \varepsilon \quad \forall x \in \Omega_\varepsilon^-, t \in [\tau_1 \varepsilon^2 |\ln \varepsilon|, (\tau_1 + 1) \varepsilon^2 |\ln \varepsilon|] \quad (3.9)$$

where

$$\begin{aligned} \Omega_\varepsilon^+ &:= \{x \in \Omega : g(x) > c_1 \varepsilon |\ln \varepsilon|\}, \\ \Omega_\varepsilon^- &:= \{x \in \Omega : g(x) < -c_1 \varepsilon |\ln \varepsilon|\}. \end{aligned}$$

Proof. Let $\phi \in C^0(\mathcal{R})$ satisfy

$$\begin{aligned} \phi(0) = \phi(1) = \phi(-2) = 0, \quad \phi'(0) > 0, \quad \phi'(1) < 0, \quad \phi'(-2) < 0, \\ \phi(s) \leq s \quad \forall s \in [-1, 1]; \quad \phi(s) > 0 \quad \forall s \in (0, 1); \quad \phi(s) < 0 \quad \forall s \in (-2, 0). \end{aligned}$$

Set $v^\varepsilon(x, t)$ to be the unique solution of

$$\begin{aligned} v_t^\varepsilon - \Delta v^\varepsilon &= \frac{1}{\varepsilon^2} \phi(v^\varepsilon), & \text{in } \Omega \times (0, T), \\ v^\varepsilon(x, 0) &= g^\varepsilon(x) & \text{on } \Omega, \\ \partial v^\varepsilon / \partial n &= 0 & \text{on } \partial\Omega \times (0, T). \end{aligned}$$

where $g^\varepsilon \in \mathcal{G}$ and

$$g^\varepsilon \begin{cases} = g(x) & \text{if } g(x) \leq 1/2, \\ = 3/4 & \text{if } g(x) \geq 7/8, \\ \leq g(x) & \text{if } g(x) \in (1/2, 7/8). \end{cases}$$

It follows that, by Theorems 1 and 2 of Chen [1991], there exist positive constants $\{c_1, \varepsilon_1, \tau_1\}$ such that if $\varepsilon \in (0, \varepsilon_1)$ then

$$v^\varepsilon(x, t) \geq 1 - \varepsilon, \quad \forall t \in (\tau_1 \varepsilon^2 |\ln \varepsilon|, (\tau_1 + 1) \varepsilon^2 |\ln \varepsilon|), \quad x \in \{y \in \Omega : g^\varepsilon \geq c_1 \varepsilon |\ln \varepsilon|\} = \Omega_\varepsilon^+.$$

Furthermore, since $|v^\varepsilon| \leq 1$ and $\phi(v^\varepsilon) \leq \beta(v^\varepsilon)$, it follows by the comparison lemma that

$$u^\varepsilon(x, t) \geq v^\varepsilon(x, t) \quad \forall (x, t) \in \Omega \times [0, T].$$

Thus (3.8) is proved and a similar argument proves (3.9). $\square$

LEMMA 3.4. *For any $c > 0$ let*

$$Q_{ce} := \{(x, t) : |x| \leq c\varepsilon, 0 < t < c^2 \varepsilon^2\},$$

$$\partial_T Q_{c\varepsilon} = \{(x, t) : |x| = c\varepsilon, 0 \leq t \leq c^2\varepsilon^2\} \cup \{(x, 0) : |x| < c\varepsilon\}.$$

Then there exists a positive constant c such that if for some $x_0 \in \Omega$ and $\delta > 0$, the conditions $(x_0, \delta) + Q_{c\varepsilon} \in \Omega \times (0, T)$ and

$$u^\varepsilon(x + x_0, t + \delta) \geq 1/2 \quad \forall (x, t) \in \partial_T Q_{c\varepsilon}$$

are satisfied, then

$$u^\varepsilon(x_0, \delta + c^2\varepsilon^2) = 1.$$

Proof. Let $\xi(\cdot) \in C^\infty(\mathcal{R})$ satisfy

$$\begin{aligned} \xi(s) &= 1/2 \quad \forall s \leq 0; & \xi(s) &= 1 \quad \forall s \geq 1; \\ 0 \leq \xi'(s) &\leq 2, & |\xi''(s)| &\leq 2 \quad \forall s \in \mathcal{R}^1. \end{aligned}$$

Define

$$v^\varepsilon(x, t) = \xi\left(\frac{t}{c^2\varepsilon^2}\right)\xi(y(x))$$

where

$$y(x) = (\sqrt{2}c\varepsilon - \sqrt{|x|^2 + c^2\varepsilon^2})/(c\varepsilon(\sqrt{2} - 1)).$$

We wish to compare $v^\varepsilon(\cdot, \cdot)$ with $u^\varepsilon(x_0 + \cdot, \delta + \cdot)$ in $Q_{c\varepsilon}$. Clearly,

- (i) $1/4 \leq v^\varepsilon(x, t) \leq 1 \quad \forall (x, t) \in \overline{Q_{c\varepsilon}};$
- (ii) $v^\varepsilon(x, t) \leq 1/2 \leq u^\varepsilon(x_0 + x, \delta + t) \quad \forall (x, t) \in \partial_T Q_{c\varepsilon};$
- (iii) $v^\varepsilon(0, c^2\varepsilon^2) = 1.$

Computation yields

$$\begin{aligned} v_t^\varepsilon - \Delta v^\varepsilon - \frac{1}{\varepsilon^2}v^\varepsilon &= \frac{1}{c^2\varepsilon^2}\xi'\left(\frac{t}{c^2\varepsilon^2}\right)\xi(y) + \frac{Nc^2\varepsilon^2 + (N-1)|x|^2}{(\sqrt{2}-1)c\varepsilon(|x|^2 + c^2\varepsilon^2)^{3/2}}\xi\left(\frac{t}{c^2\varepsilon^2}\right)\xi'(y) \\ &\quad - \frac{|x|^2}{(\sqrt{2}-1)^2c^2\varepsilon^2(|x|^2 + c^2\varepsilon^2)}\xi\left(\frac{t}{c^2\varepsilon^2}\right)\xi''(y) - \frac{1}{\varepsilon^2}v^\varepsilon \\ &\leq \frac{1}{c^2\varepsilon^2} + \frac{N}{(\sqrt{2}-1)c^2\varepsilon^2} + \frac{1}{(\sqrt{2}-1)^2c^2\varepsilon^2} - \frac{1}{4\varepsilon^2} \leq 0 \end{aligned}$$

for all $(x, t) \in Q_{c\varepsilon}$ if c is sufficiently large. It follows by the comparison principle that

$$v^\varepsilon(x, t) \leq u^\varepsilon(x_0 + x, \delta + t) \quad \forall (x, t) \in Q_{c\varepsilon}$$

and in particular,

$$u^\varepsilon(x_0, \delta + c^2\varepsilon^2) \geq v^\varepsilon(0, c^2\varepsilon^2) = 1.$$

The assertion of the lemma thus follows. $\square$

First applying Lemma 3.3 and then applying Lemma 3.4 with $\delta = \tau_1\varepsilon^2|\ln \varepsilon|$ yield

$$u^\varepsilon(x, \tau_1\varepsilon^2|\ln \varepsilon| + c^2\varepsilon^2) = 1 \quad \forall x \in \{x \in \Omega_\varepsilon^+ : \text{dist}(x, \partial\Omega_\varepsilon^+) \geq c\varepsilon\},$$

$$u^\varepsilon(x, \tau_1 \varepsilon^2 |\ln \varepsilon| + c^2 \varepsilon^2) = -1 \quad \forall x \in \{x \in \Omega_\varepsilon^- : \text{dist}(x, \partial \Omega_\varepsilon^-) \geq c\varepsilon\}.$$

Finally applying Theorem 2.1 with $\gamma(\varepsilon) = C\varepsilon |\ln \varepsilon|$, we conclude the following.

THEOREM 3.5. *Assume that $g \in \mathcal{G} \cap W^{1,\infty}(\Omega)$ and*

$$|\nabla g| > c \quad \forall x \in \{y \in \Omega : |g(y)| \leq c\} \quad (3.10)$$

for some $c > 0$. Let $\Gamma_0 = \{y \in \Omega : g(y) = 0\}$ and assume that $\Gamma_0 \subset \Omega$ and $\Gamma_0 \in C^{3+\alpha}$. Then there exists positive constants C and ε_0 such that for any $\varepsilon \in (0, \varepsilon_0)$ and $t \in (C\varepsilon^2 |\ln \varepsilon|, T]$,

$$\begin{aligned} u^\varepsilon(x, t) &= 1 \quad \text{if } d(x, t) > C\varepsilon |\ln \varepsilon|, \\ u^\varepsilon(x, t) &= -1 \quad \text{if } d(x, t) < -C\varepsilon |\ln \varepsilon| \end{aligned}$$

where $d(x, t)$ and T are as in §3.1.

REMARK 3.6. In Theorem 3.5, condition (3.10) is only used to ensure that $\Omega \setminus (\Omega_\varepsilon^+ \cup \Omega_\varepsilon^-)$ is contained in a $O(\varepsilon |\ln \varepsilon|)$ neighborhood of Γ_0 , whereas the regularity assumption on Γ_0 is to ensure the existence of a smooth solution to the motion by mean curvature equation.

4 Asymptotic behavior as $t \rightarrow \infty$ in one space dimension.

4.1. STATEMENT OF MAIN RESULT.

In this section we shall restrict our attention to one space dimension and set $\Omega = (0, 1)$. We shall prove the following theorem.

THEOREM 4.1. *Let $g \in \mathcal{G}$ and let $u_g(t) = S(t)g$ be the unique solution of the double obstacle problem (2.1). Then there exists $f \in C^{1,1}[0, 1] \in \mathcal{E}$ such that*

$$\lim_{t \rightarrow \infty} u_g(x, t) = f(x) \quad \forall x \in [0, 1].$$

We have already established in §2 that $\omega(g) \in \mathcal{E}$, so in order to prove the theorem we need only show that $\omega(g)$ consists of a single element. It is convenient to characterize $\mathcal{E}$ as follows (c.f. Blowey–Elliott [1991a]).

Any $f \in \mathcal{E}$ satisfies: $f \in C^{1,1}[0, 1]$ and

$$\begin{cases} \varepsilon^2 f_{xx} + f &= 0 & \text{for } x \in \Omega_I := \{x \in (0, 1) : |f(x)| < 1\}. \\ f'(x) &= 0 & \text{for } x \in \partial \Omega_T \cup \partial \Omega. \end{cases} \quad (4.1)$$

Let K be the largest integer strictly less than $1/(\varepsilon\pi)$. It is obvious that if $K + 1 = 1/\varepsilon\pi$, then

$$f(x) = a \cos x/\varepsilon \quad \forall a \in [-1, 1]$$

is a solution to (4.1) and furthermore for any $\varepsilon > 0$,

$$f(x) \equiv 0, \quad f(x) \equiv -1, \quad f(x) \equiv 1$$

are equilibria. It is convenient to set

$$\mathcal{E}_0 := \{f : f(x) \equiv 1 \text{ or } f(x) \equiv -1\}$$

and

$$\mathcal{E}^c = \begin{cases} \{f : f(x) \equiv 0\} & \text{if } 1 < (K+1)\varepsilon\pi, \\ \{f : f(x) = a \cos x/\varepsilon, |a| \leq 1\} & \text{if } 1 = (K+1)\varepsilon\pi. \end{cases}$$

Any other solution f of (4.1) has at least one layer connecting $f = 1$ to $f = -1$ in which $|f(x)| < 1$. By the definition of K , there can be at most K layers in a solution other than $f(x) = \cos x/\varepsilon \in \mathcal{E}^c$ when $1 = (K+1)\varepsilon\pi$. Therefore for each $k \in \{1, 2, \dots, K\}$, we let $\mathcal{E}_k$ be the set of k -layered solutions characterized by the $k+1$ parameters $\{a; s_1, \dots, s_k\}$ in the following way. $f \equiv f[a; s_1, \dots, s_k] \in \mathcal{E}_k$ if and only if

$$\left\{ \begin{array}{l} |a| = 1, \quad 0 \leq s_1, \quad s_k \leq 1 - \varepsilon\pi, \quad s_j \leq s_{j+1} - \varepsilon\pi \quad \forall j = 1, \dots, k-1; \\ A_1^k = a, \quad A_{j+1}^k = -A_j^k, \quad j = 1, 2, \dots, k; \\ f(x) = A_j^k \cos(x - s_j^k)/\varepsilon \quad \text{if } s_j^k \leq x \leq s_j^k + \varepsilon\pi; \\ f(x) = a \quad \text{if } 0 \leq x \leq s_1, \quad f(x) = A_{k+1}^k \quad \text{if } s_k + \varepsilon\pi \leq x \leq 1; \\ f(x) = A_{j+1}^k \quad \text{if } s_j + \varepsilon\pi \leq x \leq s_{j+1}, \quad j = 1, 2, \dots, k-1. \end{array} \right. \quad (4.2)$$

This leads to the following conclusion.

LEMMA 4.2.

$$\mathcal{E} = \cup_{k=0}^K \mathcal{E}_k \cup \mathcal{E}^c.$$

4.2. STABILITY OF THE EQUILIBRIA.

In this subsection we study the stability of the equilibria in $\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_K$ and $\mathcal{E}^c$ respectively. A direct calculation yields the following lemma.

LEMMA 4.3. (i) $\inf_{\eta \in H^1(\Omega) \cap \mathcal{G}} E(\eta) = 0$;

(ii) $E(f) = \frac{k\pi}{\varepsilon} =: e_k \quad \forall f \in \mathcal{E}_k, 0 \leq k \leq K$;

(iii) $E(f) = \frac{1}{\varepsilon^2} =: e^c \quad \forall f \in \mathcal{E}^c$;

(iv) $0 = e_0 < e_1 < \dots < e_K < e^c = 1/\varepsilon^2$.

(v) The sets $\{\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_K, \mathcal{E}^c\}$ are isolated in $L^\infty(0, 1)$ in the sense that there exists $\rho > 0$ such that

$$\text{dist}(\mathcal{E}_i, \mathcal{E}_j) > \rho, \quad \text{dist}(\mathcal{E}_j, \mathcal{E}^c) > \rho, \quad \forall i, j, i \neq j.$$

(vi) The set $\mathcal{E}^c$ is connected and for each $k = 1, \dots, K$, $\mathcal{E}_k$ consists of two isolated connected subsets

$$\{f[1; s_1, \dots, s_k] : 0 \leq s_1, \dots, s_k \leq 1 - \varepsilon\pi, \quad s_j \leq s_{j+1} - \varepsilon\pi \quad \forall j = 1, \dots, k-1\}$$

and

$$\{f[-1; s_1, \dots, s_k] : 0 \leq s_1, \dots, s_k \leq 1 - \varepsilon\pi, \quad s_j \leq s_{j+1} - \varepsilon\pi \quad \forall j = 1, \dots, k-1\}.$$

(vii) The k -layered equilibria have the energy π/ε associated with each layer (interface) connecting $+1$ to -1 ; there is no energy associated with the set $\{x : |f(x)| = 1\}$.

Since from §2 the Lyapunov functional $E(\cdot)$ is constant on $\omega(g)$, the following holds:

COROLLARY 4.4. $\omega(g)$ is contained in one and only one of $\{\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_K, \mathcal{E}^c\}$.

DEFINITION 4.5. (1) $f \in \mathcal{E}$ is called *asymptotically stable* if there exists $\delta > 0$ such that

$$\lim_{t \rightarrow \infty} u_g(\cdot, t) = f, \quad \forall g \in B_\delta(f) := \{h \in \mathcal{G} : \|f - h\|_{L^\infty} \leq \delta\}.$$

(2) $f \in \mathcal{E}$ is called *stable* if for every $\delta > 0$ there exists an $\eta > 0$ such that

$$\overline{\lim}_{t \rightarrow \infty} \max_{x \in [0,1]} |u_g(x, t) - f(x)| \leq \delta \quad \forall g \in B_\eta(f).$$

(3) If $f \in \mathcal{E}$ is not stable then it is said to be *unstable*.

(4) An unstable $f \in \mathcal{E}$ is called *M-unstable* if there exists a sequence $g_n \rightarrow f$ in $L^\infty(0, 1)$ such that

$$\lim_{n \rightarrow \infty} \overline{\lim}_{t \rightarrow \infty} \|u_{g_n}(\cdot, t) - f\|_{L^\infty} > 0$$

and a sequence $f_n \rightarrow f$ in $L^\infty(0, 1)$ such that

$$\lim_{n \rightarrow \infty} \overline{\lim}_{t \rightarrow \infty} \|u_{f_n}(\cdot, t) - f(\cdot)\|_{L^\infty} = 0.$$

LEMMA 4.6. If $f \in \mathcal{E}_0$ then f is asymptotically stable.

Proof. For every $a \in (0, 1) \cup (-1, 0)$, set

$$w(a, t) = \begin{cases} ae^{t/\varepsilon^2} & \text{if } 0 \leq t \leq \varepsilon^2 \ln 1/a, \\ \text{sign } a & \text{if } t > \varepsilon^2 \ln 1/a. \end{cases}$$

It follows that $w(a, t) = \mathcal{S}(t)a$ and we can prove, by the comparison Lemma 2.2, that

$$w\left(\inf_{x \in [0,1]} g(x), t\right) \leq u_g(x, t) \leq w\left(\sup_{x \in [0,1]} g(x), t\right)$$

and therefore

$$\text{if } g > 0 \text{ then } \lim_{t \rightarrow \infty} u_g(\cdot, t) = 1,$$

$$\text{if } g < 0 \text{ then } \lim_{t \rightarrow \infty} u_g(\cdot, t) = -1,$$

from which, the lemma follows. $\square$

Remark 4.7. This Lemma is true for the multi-dimensional problem. The proof is identical.

LEMMA 4.8. *Each element of $\mathcal{E}^c$ is M -unstable.*

Proof. Set

$$K' = \begin{cases} K + 1 & \text{if } \varepsilon\pi(K + 1) = 1, \\ K & \text{if } \varepsilon\pi(K + 1) > 1, \end{cases}$$

$$U = \{\eta : \eta(x) = \sum_{j=0}^{K'} a_j \cos(j\pi x)\},$$

and

$$S = \{\eta : \eta(x) = \sum_{j=K'+1}^{\infty} a_j \cos(j\pi x)\}.$$

Let $f \in \mathcal{E}^c$ and

$$g = f + h \in \mathcal{G}$$

where $h(\not\equiv 0) \in U$. A direct calculation yields

$$E(g) = \frac{1}{\varepsilon^2} - \frac{a_0^2}{\varepsilon^2} + \sum_{j=1}^{K'} \frac{a_j^2}{2} \left(j^2 \pi^2 - \frac{1}{\varepsilon^2} \right) < \frac{1}{\varepsilon^2} = E(f).$$

(If $K'\varepsilon\pi = 1$ and $f = \cos x/\varepsilon$, then take $a_{K'} < 0$ and $a_0 \neq 0$ so that $f + h \in \mathcal{G}$.)

It follows that

$$\lim_{t \rightarrow \infty} E(u_g(\cdot, t)) \leq E(g) < E(f).$$

Therefore U is an unstable manifold.

On the other hand, let $h \in S$ be chosen so that

$$g = f + h \in \mathcal{G} \quad \text{and} \quad |g(x)| < 1 \quad \text{a.e. .}$$

It follows that

$$\widetilde{u}_g := f(x) + \sum_{j=K'+1}^{\infty} a_j e^{(1/\varepsilon^2 - j^2 \pi^2)t} \cos(j\pi x)$$

solves

$$\begin{aligned} u_t - u_{xx} &= u/\varepsilon^2, & x \in (0, 1), t > 0; \\ u(x, 0) &= g(x), & x \in (0, 1); & \quad u_x(0, t) = u_x(1, t) = 0, \quad t > 0. \end{aligned}$$

Since $1/\varepsilon^2 - j^2\pi^2 \leq \pi^2((K' + 1)^2 - j^2) < 0$, it follows that

$$\lim_{t \rightarrow \infty} \widetilde{u}_g(\cdot, t) = f.$$

Suppose that $f \neq \pm \cos x/\varepsilon$ then there exists $\delta > 0$ such that $\|f\|_\infty < 1 - \delta$. Choose $h \in S$ such that

$$\sum_{j=K'+1}^{\infty} |a_j| \leq \delta/2.$$

It follows that

$$\|\widetilde{u}_g\|_\infty < 1 - \delta/2 \quad \forall t > 0$$

which implies that $\widetilde{u}_g(\cdot, t) = u_g(\cdot, t)$ and the assertion of the lemma follows.

When $f = \pm \cos x/\varepsilon$, the sequence $\{\pm(1 - 1/n)\cos x/\varepsilon\}_{n=1}^\infty$ is a sequence of equilibria which converges to f as $n \rightarrow \infty$. This completes the proof of the lemma. $\square$

In order to discuss the stability of $\{\mathcal{E}_k\}_{k=1}^K$, it is convenient to introduce

$$\mathcal{E}_k^s = \{f[a; s_1, \dots, s_k] \in \mathcal{E}_k : 0 < s_1, s_k < 1 - \varepsilon\pi, s_j < s_{j+1} - \varepsilon\pi \forall j = 1, \dots, k-1\},$$

$$\mathcal{E}_k^u = \{f[a; s_1, \dots, s_k] \in \mathcal{E}_k : s_1 = 0 \text{ or } s_k = 1 - \varepsilon\pi \text{ or } s_{j+1}^k = s_j^k + \varepsilon\pi \text{ for some } j\},$$

so that

$$\mathcal{E}_k = \mathcal{E}_k^s \cup \mathcal{E}_k^u, \quad \mathcal{E}_k^s \cap \mathcal{E}_k^u = \emptyset.$$

LEMMA 4.9. For each $k \in \{1, \dots, K\}$, every element of $\mathcal{E}_k^u$ is M -unstable.

Proof. We first consider the case $k = 2$. Without loss of generality let $f = f[1; s, s + \varepsilon\pi] \in \mathcal{E}_2^u$. Then $f_n := f[1; s, s + \varepsilon\pi + 1/n]$ (or $f_n := f[1; s - 1/n, s + \varepsilon\pi]$ when $s = 1 - 2\varepsilon\pi$) is a sequence of equilibria which converges to f as $n \rightarrow \infty$. It remains to show that f is unstable.

Let $g \neq f$ satisfy

$$f \leq g \leq 1.$$

By comparison,

$$u_g(\cdot, t) \geq f(\cdot) \quad \forall t \geq 0.$$

so that

$$u_g(x, t) = 1 \quad \text{when} \quad |x - (s + \varepsilon\pi)| \geq \varepsilon\pi.$$

Let $\alpha \in (\pi/2, \pi)$ and set

$$h(x) := \begin{cases} 0 & x - (s + \varepsilon\pi) \leq -\alpha\varepsilon, \\ \sin[\pi(x - (s + \varepsilon\pi) + \alpha\varepsilon)/(2\alpha\varepsilon)] & |x - (s + \varepsilon\pi)| < \alpha\varepsilon, \\ 0 & x - (s + \varepsilon\pi) \geq \alpha\varepsilon. \end{cases}$$

Clearly,

$$\int_s^{s+2\epsilon\pi} \left\{ f' h' - \frac{1}{\epsilon^2} f h \right\} dx = - \int_s^{s+2\epsilon\pi} \left(f'' + \frac{1}{\epsilon^2} f \right) h dx = 0.$$

Take $\delta \in (0, 1)$ small so that

$$g = f + \delta h \in \mathcal{G}.$$

Direct calculation yields

$$\begin{aligned} E(g) &= E(f) + 2\delta \int_s^{s+2\epsilon\pi} \left\{ f' h' - \frac{1}{\epsilon^2} f h \right\} dx + \delta^2 \int_s^{s+2\epsilon\pi} \left\{ h'^2 - \frac{1}{\epsilon^2} h^2 \right\} dx \\ &= E(f) + \delta^2 \int_{(s+\epsilon\pi)-\epsilon\alpha}^{(s+\epsilon\pi)+\epsilon\alpha} h \left(-h'' - \frac{1}{\epsilon^2} h \right) dx \\ &= E(f) + \delta^2 \int_{(s+\epsilon\pi)-\epsilon\alpha}^{(s+\epsilon\pi)+\epsilon\alpha} \frac{h^2}{\epsilon^2} \left(\frac{\pi^2}{4\alpha^2} - 1 \right) dx < E(f). \end{aligned}$$

Therefore if $\tilde{f} \in \omega(g)$, then $E(\tilde{f}) < E(f)$. This establishes the instability of f . The proof for $k = 1$ and $k > 2$ is identical. $\square$

LEMMA 4.10. *For each $k \in \{1, \dots, K\}$, the set $\mathcal{E}_k^s$ is stable.*

Proof. Let $f \in \mathcal{E}_k$ and $g \in B_\eta(f)$. The Sobolev inequality yields

$$\|f - u_g(\cdot, t)\|_\infty \leq \sqrt{2} \|f - u_g\|^{1/2} \|f - u_g\|_{H^1}^{1/2} \leq \sqrt{2\eta\rho} e^{C_\epsilon t/2} \quad \forall t \geq t^* \quad (4.3)$$

where in the second inequality we have used (2.14) and Theorem 2.1(3).

For simplicity, we shall assume that $k = 2$. Since for $f \in \mathcal{E}_k^s$, the layers are isolated and lie in the interior of $(0, 1)$, the argument we use for $k = 2$ can be readily extended to the general case. So without loss of generality, we assume that $f = f[-1; s_1, s_2] \in \mathcal{E}_2^s$ where $0 < s_1 < s_2 - \epsilon\pi < 1 - 2\epsilon\pi$.

For any μ satisfying $0 < \mu < \frac{1}{5} \min\{\epsilon\pi, s_1, s_2 - s_1 - \epsilon\pi, 1 - s_2 - \epsilon\pi\}$, let $w^\mu(y, t)$ be the solution of

$$\begin{cases} w_t^\mu - w_{yy}^\mu = 1/(2\epsilon^2), & (y, t) \in (-\mu, \mu) \times (0, \infty), \\ w^\mu(-\mu, t) = w^\mu(\mu, t) = 0, & t \in (0, \infty), \\ w^\mu(y, 0) = 0, & y \in (-\mu, \mu). \end{cases}$$

One can easily verify that

$$w^\mu(0, t) > w^\mu(y, \tau) \quad \forall 0 \leq \tau < t, \quad y \in (-\mu, \mu).$$

Set $M(\mu) = w^\mu(0, 1)$. Taking smaller μ if necessary, we can assume, without loss of generality, that $0 < M(\mu) \leq 1/2$. For every $x_0 \in (s_1 + \epsilon\pi + \mu, s_2 - \mu)$, consider the function W^μ defined by

$$W^\mu(x, t) = 1 - M(\mu) + w^\mu(x - x_0, t).$$

Clearly,

$$1/2 \leq W^\mu < 1 \quad \forall t \in [0, 1), x \in I := (x_0 - \mu, x_0 + \mu).$$

It follows that

$$W_t^\mu - W_{xx}^\mu - \frac{1}{\varepsilon^2} W^\mu \leq w_t^\mu - w_{yy}^\mu - \frac{1}{2\varepsilon^2} = 0, \quad (x, t) \in I \times (0, 1).$$

Taking η small such that $\sqrt{2\eta\rho}e^{C_\varepsilon(t^*+1)/2} \leq M(\mu)$, one obtains that for all $(x, t) \in \partial I \times (0, 1) \cup I \times \{0\}$,

$$W^\mu(x, t) = 1 - M(\mu) \leq 1 - \eta\rho e^{C_\varepsilon(t^*+t)} \leq u_g(x, t)$$

by (4.3). The comparison Lemma then implies that

$$W^\mu(x, t) \leq u_g(x, t^* + t) \quad \forall (x, t) \in I \times [0, 1].$$

Consequently,

$$u_g(x_0, t^* + 1) = 1 \quad \forall x_0 \in (s_1 + \varepsilon\pi + \mu, s_2 - \mu).$$

Define

$$\eta(\mu) = \frac{1}{2\rho e^{C_\varepsilon(t^*+1)}} \min \left\{ M^2(\mu), 4 \sin^4 \frac{\mu}{2\varepsilon} \right\}.$$

We claim that if $\eta \in (0, \eta(\mu))$ then the inequality

$$u_g(x, t^* + 1) \geq f[-1; s_1 + 2\mu, s_2 - 2\mu](x) \quad \forall x \in (0, 1) \quad (4.4)$$

holds. In fact, since (4.4) holds in the interval $I_1 := [0, s_1 + 2\mu] \cup [s_2 - 2\mu + \varepsilon\pi, 1]$ (where $f[-1; s_1 + 2\mu, s_2 - 2\mu] = -1$) and in the interval $I_2 := [s_1 + \varepsilon\pi + \mu, s_2 - \mu]$ (where $u_g(x, t^* + 1) = 1$), we need only consider the case when $x \in [0, 1] \setminus (I_1 \cup I_2) = (s_1 + 2\mu, s_1 + \varepsilon\pi + \mu) \cup (s_2 - \mu, s_2 + \varepsilon\pi - 2\mu) \equiv I_3 \cup I_4$. When $x \in I_3$, we can use the definition of f in (4.2) to calculate

$$\begin{aligned} & f[-1; s_1, s_2](x) - f[-1; s_1 + 2\mu, s_2 - 2\mu](x) \\ & \geq \min \left\{ \min_{y \in [s_1 + 2\mu, s_1 + \varepsilon\pi]} \left(-\cos \frac{y - s_1}{\varepsilon} + \cos \frac{y - s_1 - 2\mu}{\varepsilon} \right), \right. \\ & \quad \left. \min_{y \in [s_1 + \varepsilon\pi, s_1 + \varepsilon\pi + \mu]} \left(1 + \cos \frac{y - s_1 - 2\mu}{\varepsilon} \right) \right\} \\ & = 2 \sin^2 \frac{\mu}{2\varepsilon}. \end{aligned}$$

It follows that

$$\begin{aligned} & u_g(x, t^* + 1) - f[-1; s_1 + 2\mu, s_2 - 2\mu](x) \\ & \geq f[-1; s_1, s_2](x) - f[-1; s_1 + 2\mu, s_2 - 2\mu](x) - \|u_g(\cdot, t^* + 1) - f[-1; s_1, s_2]\|_{L^\infty} \\ & \geq 2 \sin^2 \frac{\mu}{2\varepsilon} - \sqrt{2\eta\rho}e^{C_\varepsilon(t^*+1)/2} \geq 0 \end{aligned}$$

by (4.3) the the definition of $\eta(\mu)$. The case when $x \in I_4$ can be treated similarly. This establishes (4.4).

It follows from (4.4) that, for all $(x, t) \in (0, 1) \times [t^* + 1, \infty)$,

$$u_g(\cdot, t) \geq \mathcal{S}(t - t^* - 1)f[-1; s_1 + 2\mu, s_2 - 2\mu] = f[-1; s_2 + 2\mu, s_2 - 2\mu].$$

Similarly, we can show that

$$u_g(x, t) \leq f[-1; s_1 - 2\mu, s_2 + 2\mu](x) \quad \forall (x, t) \in (0, 1) \times [t^* + 1, \infty)$$

provided that $\eta \in (0, \eta(\mu))$. Therefore,

$$\overline{\lim}_{t \rightarrow \infty} \|u_g - f\|_\infty \leq \|f[-1; s_1 - 2\mu, s_2 + 2\mu] - f[-1; s_1 + 2\mu, s_2 - 2\mu]\|_\infty.$$

Clearly, as $\eta \rightarrow 0$, we can let $\mu \rightarrow 0$ so that the right hand side goes to zero. Hence,

$$\lim_{\eta \rightarrow 0} \overline{\lim}_{t \rightarrow \infty} \sup_{g \in B_\eta(f)} \|u_g(\cdot, t) - f\|_\infty = 0.$$

The assertion of the lemma follows. $\square$

Remark 4.11. From (4.3), we see that the L^∞ stability is equivalent to L^2 stability.

4.3. COMPLETION OF THE PROOF OF THEOREM 4.1.

It is sufficient to show that ω consists of a singleton. Suppose that f_1 and f_2 belong to $\omega(g)$. By Corollary 4.4, both f_1 and f_2 belong to the same set being one of $\{\mathcal{E}_0, \mathcal{E}_1, \dots, \mathcal{E}_K, \mathcal{E}^c\}$.

In the case $f_1 \in \left(\bigcup_{i=1}^K \mathcal{E}_i^s\right) \cup \mathcal{E}_0$, it follows, by lemma 4.1 and 4.10, that $f_1 = f_2$.

Now we consider the case $f_1 \in \mathcal{E}_k^u$, $k \in \{1, \dots, K\}$. Write $f_1 = f[a; s_1, \dots, s_k]$ and $f_2 = f[\bar{a}; \bar{s}_1, \dots, \bar{s}_k]$. We want to show that $a = \bar{a}$, $s_1 = \bar{s}_1$, $\dots$, and $s_k = \bar{s}_k$.

Since $\omega(g)$ is connected, f_1 and f_2 belong to the same connected component of $\mathcal{E}_k$, so that $a = \bar{a}$. Without loss of generality we assume that $a = \bar{a} = -1$.

We proceed to show that $s_1 = \bar{s}_1$. Assume that this is not true, say, $\bar{s}_1 > s_1 \geq 0$. Set $\mu = (\bar{s}_1 - s_1)/5$. We can use the same argument as in the proof of Lemma 4.10 to deduce that

$$u_g(\cdot, t) \leq f[-1; \bar{s}_1 - 2\mu] \quad \forall t > t_n + t^* + 1$$

where t_n is the time such that $\|u_g(\cdot, t_n) - f_2\|_\infty \leq \eta(\mu)$. It follows that

$$f \leq f[-1, \bar{s}_1 - 2\mu] \quad \forall f \in \omega(g),$$

contradicting to the fact that $f_1 \in \omega(g)$. Therefore $s_1 = \bar{s}_1$. Successively repeating the same argument for $i = 2, \dots, k$, it follows that $s_i = \bar{s}_i$ for all $j = 1, \dots, k$, and therefore $f_1 = f_2$.

It remains to consider the case $f_1, f_2 \in \mathcal{E}^c$. We need only to consider the case when $1 = (K + 1)\varepsilon\pi$ since otherwise $\mathcal{E}^c$ consists of a single element $f \equiv 0$. The following lemma completes the proof of the theorem.

LEMMA 4.12. *Suppose that $1/\varepsilon\pi$ is an integer and $\omega(g) \in \mathcal{E}^c$. Then there exists a constant $a \in [-1, 1]$ such that*

$$\lim_{t \rightarrow \infty} u_g(x, t) = a \cos \frac{x}{\varepsilon} \quad x \in [-1, 1].$$

Proof. Since $\omega(g)$ is connected, we need only consider the case when there exists $a \in (-1, 1)$ such that $a \cos x/\varepsilon \in \omega(g)$ because otherwise $\omega(g) = \{\cos x/\varepsilon\}$ or $\omega(g) = \{-\cos x/\varepsilon\}$ and therefore the lemma holds automatically.

Noting that $u_g(\cdot, t)$ is in $H^2((0, 1))$, we can write u_g as the uniformly convergent Fourier series

$$u_g(x, t) = \frac{1}{2}a_0(t) + \sum_{k=1}^{\infty} a_k(t) \cos(k\pi x)$$

where

$$a_k(t) = 2 \int_0^1 u_g(x, t) \cos(k\pi x) dx, \quad k = 0, 1, 2, \dots,$$

and, by using Theorem 1 (5),

$$\begin{aligned} \frac{d}{dt} a_k(t) &= 2 \int_0^1 \frac{\partial}{\partial t} u_g(x, t) \cos(k\pi x) dx \\ &= 2 \int_0^1 \left(u_{xx} + \frac{1}{\varepsilon^2} u \right) \cos(k\pi x) dx + \frac{1}{\varepsilon^2} \int_0^1 (\beta(u) - u) \cos(k\pi x) dx \\ &= \left(\frac{1}{\varepsilon^2} - k^2 \pi^2 \right) a_k(t) + \frac{1}{\varepsilon^2} e_k(t) \end{aligned} \tag{4.5}$$

where

$$e_k(t) = \int_0^1 (\beta(u) - u) \cos(k\pi x) dx.$$

The function e_k satisfies

$$|e_k(t)| \leq m(t) := \sup_{\tau \geq t} \text{measure} \{x : |u_g(x, \tau)| = 1\}.$$

We claim that

$$\lim_{t \rightarrow \infty} m(t) = 0. \tag{4.6}$$

In fact, otherwise $\text{measure} \{x : |u_g(x, \tau_n)| = 1\} > \delta$ for some positive δ and for a sequence τ_n converging to ∞ . It then follows that there exists an $f \in \omega(g)$ such that $|f(\alpha)| = 1$ for some $\alpha \notin \{0, \varepsilon\pi, 2\varepsilon\pi, \dots, K\varepsilon\pi, 1\}$. Hence $f \notin \mathcal{E}^c$, which is a contradiction. Therefore (4.6) must hold.

Solving the ordinary differential equation (4.5) yields

$$a_k(t) = a_k(t_0)e^{(1/\varepsilon^2 - k^2\pi^2)(t-t_0)} + \frac{1}{\varepsilon^2} \int_{t_0}^t e^{(1/\varepsilon^2 - k^2\pi^2)(t-\tau)} e_k(\tau) d\tau \quad \forall t \geq t_0 > 0.$$

It follows that for $k \geq K+2$,

$$\begin{aligned} |a_k(t)| &\leq a_k(t_0)e^{(1/\varepsilon^2 - k^2\pi^2)(t-t_0)} + \frac{1}{(k\varepsilon\pi)^2 - 1} m(t_0) \\ &\leq 2e^{\pi^2[(K+1)^2 - k^2](t-t_0)} + \frac{m(t_0)}{k^2[(K+1)^{-2} - k^{-2}]} \\ &\leq C(e^{-kt_0} + m(t_0)/k^2) \quad \forall t \geq 2t_0, \end{aligned} \quad (4.7)$$

where in the second inequality we have used the fact that $|a_k| \leq 2$ which follows from the definition of a_k .

Direct calculation yields

$$E(u_g(\cdot, t)) = \frac{1}{\varepsilon^2} - \frac{a_0^2(t)}{4\varepsilon^2} + \sum_{k=1}^K \frac{a_k^2(t)}{2} \left(k^2\pi^2 - \frac{1}{\varepsilon^2} \right) + \sum_{k=K+2}^{\infty} \frac{a_k^2(t)}{2} \left(k^2\pi^2 - \frac{1}{\varepsilon^2} \right)$$

Since $E(u_g(\cdot, t)) \searrow e^c = 1/\varepsilon^2$ and

$$0 \leq \sum_{k=K+2}^{\infty} a_k^2(2t) \left(k^2\pi^2 - \frac{1}{\varepsilon^2} \right) \leq C \left[\frac{e^{-t}}{1 - e^{-t}} + m(t) \right] \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

it follows that

$$\lim_{t \rightarrow \infty} \left[-\frac{1}{4\varepsilon^2} a_0^2(t) + \sum_{k=1}^K \frac{a_k^2(t)}{2} \left(k^2\pi^2 - \frac{1}{\varepsilon^2} \right) \right] = 0;$$

i.e.,

$$\lim_{t \rightarrow \infty} a_k(t) = 0 \quad \forall k = 0, 1, \dots, K. \quad (4.8)$$

Write u_g as

$$u_g = u^+ + u^0 + u^-$$

where

$$\begin{aligned} u^+(x, t) &= \frac{a_0(t)}{2} + \sum_{k=1}^K a_k(t) \cos(k\pi x), \\ u^0(x, t) &= a_{K+1}(t) \cos((K+1)\pi x) = a_{K+1}(t) \cos \frac{x}{\varepsilon}, \\ u^-(x, t) &= \sum_{k=K+2}^{\infty} a_k(t) \cos(k\pi x). \end{aligned}$$

It follows from (4.8) that

$$\lim_{t \rightarrow \infty} \|u^+(\cdot, t)\|_{\infty} = 0 \quad (4.9)$$

and from (4.7) that

$$\|u^-(\cdot, 2t)\|_\infty \leq \sum_{k=K+2}^{\infty} |a_k(2t)| \leq C \left[\frac{e^{-t}}{1 - e^{-t}} + m(t) \right] \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (4.10)$$

Let N be large enough such that

$$\begin{aligned} \|u_g(x, t_N) - a \cos(x/\varepsilon)\|_\infty &\leq \frac{1}{8}(1 - |a|), \\ \|u^\pm(\cdot, t)\|_\infty &\leq \frac{1}{8}(1 - |a|) \quad \forall t \geq t_N. \end{aligned}$$

It follows that $|u_g(x, t_N)| \leq 1 - \frac{7}{8}(1 - |a|)$. Let d be the largest constant such that

$$\|u_g(\cdot, t)\|_\infty < 1 - \frac{1}{8}(1 - |a|) \quad \forall t \in [t_N, t_N + d].$$

Clearly, $d > 0$ and $e_k(t) = 0$ for all $t \in [t_N, t_N + d]$. It follows that $a_{K+1}(t) = a_{K+1}(t_N)$ for all $t \in [t_N, t_N + d]$. Therefore, for all $t \in [t_N, t_N + d]$,

$$\begin{aligned} u_g(x, t) &= u^+(x, t) + a_{K+1}(t_N) \cos(x/\varepsilon) + u^-(x, t). \\ &= u_g(x, t_N) + (u^+(x, t_N + d) - u^+(x, t_N)) + (u^-(x, t_N + d) - u^-(x, t_N)). \end{aligned}$$

This implies that $d = \infty$ since otherwise

$$\begin{aligned} 1 - \frac{1}{8}(1 - |a|) &= \|u_g(\cdot, t_N + d)\|_\infty \\ &\leq \|u(\cdot, t_N)\|_\infty + \|u^+(\cdot, t_N)\|_\infty + \|u^+(\cdot, t_N + d)\|_\infty \\ &\quad + \|u^-(\cdot, t_N)\|_\infty + \|u^-(\cdot, t_N + d)\|_\infty \\ &\leq 1 - \frac{7}{8}(1 - |a|) + 4 \cdot \frac{1}{8}(1 - |a|) = 1 - \frac{3}{7}(1 - |a|), \end{aligned}$$

a contradiction. Therefore,

$$u_g(x, t) = u^+(x, t) + a_{K+1}(t_N) \cos \frac{x}{\varepsilon} + u^-(x, t) \quad \forall t \in (t_N, \infty).$$

Consequently, by (4.9) and (4.10),

$$\lim_{t \rightarrow \infty} u_g(x, t) = a_{K+1}(t_N) \cos \frac{x}{\varepsilon} \quad \forall x \in [0, 1].$$

The assertion of the lemma follows. This also completes the proof of Theorem 4.1. $\square$

Remark 4.12. The fact that $\|u_g\| < 1$ in $[t_N, \infty)$ implies $a_k(t) = a_k(t_N)e^{1/\varepsilon^2 - k^2\pi^2}$ for all $t \geq t_N$, so that $a_k(t) = 0$ for all $k = 0, 1, \dots, K$ and $t \geq t_N$; that is, $u^+ \equiv 0$ for $t \geq t_N$.

References

- [1] S. Allen & J. Cahn, 1979, *A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening*, Acta. Metall., **27**, 1084–1095.
- [2] J. F. Blowey & C. M. Elliott, 1991
 - (a) *The Cahn-Hilliard gradient theory for phase separation with non-smooth free energy, Part I: Mathematical Analysis*, European. J. Appl. Math. **2**, 233–279
 - (b) *The Cahn-Hilliard gradient theory for phase separation with non-smooth free energy, Part II: Numerical Analysis*, to appear in European. J. Appl. Math. .
 - (c) *Curvature dependent phase boundary motion and parabolic obstacle problems*, to appear in Proceedings of IMA Workshop “Degenerate Diffusion”.
- [3] L. Bronsard & R. V. Kohn, 1991, *Motion by mean curvature as the singular limit of Ginzburg–Landau dynamics*, J. Diff. Eqns. **90**, 211–237.
- [4] J. Carr & R. Pego,
 - (a) 1989, *Very slow phase separation in one dimension*, Lecture Notes in Physics (M. Rascle ed.) **344**, Springer–Verlag, 216–226
 - (b) 1990, *Invariant manifolds for metastable pattern in $u_t = \varepsilon^2 u_{xx} - f(u)$* , Proc. Roy. Soc. Edinburgh **116**, 133–160.
- [5] Xinfu Chen, 1990, *Generation and Propagation of interface in reaction–diffusion equations*, to appear in J. Diff. Eqns. .
- [6] Y.G. Chen, Y. Giga, & S. Goto, 1991, *Uniqueness and existence of viscosity solution of generalized mean curvature flow equations*, J. Diff. Geom., **33**, 749–786.
- [7] P. DeMottoni & M. Schatzman, 1989, *Evolution géométrique d’interfaces*, C. R. Acad. Sci. Sér. I Math. **309** , 453–458.
- [8] L. C. Evans, H. M. Soner, & P. E. Souganidis, 1990, *The Allen–Cahn equation and the generalized motion by mean curvature*, Preprint.
- [9] L. C. Evans & J. Spruck, 1991, *Motion by mean curvature I*, J. Diff. Geom., **33**, 635–681.
- [10] P. Fife & L. Hsiao, 1988, *The generation and propagation of internal layers*, Nonlinear Anal. **70**, 31–46.

- [11] G. Fusco, 1990, *A geometric approach to the dynamics of $u_t = \varepsilon^2 u_{xx} + f(u)$ for small ε* , Lecture Notes in Physics (Kirchgassner, ed) **359**, Springer-Verlag, 53–73.
- [12] G. Fusco & J. K. Hale, 1989, *Slow-motion manifolds, dormant instability, and singular perturbations*, J. Dynamics Differential Equations **1**, No. 1, 75–94.
- [13] M. Gage & R. Hamilton, 1986, *The shrinking of convex plane curves by heat equation*, J. Diff. Geom. **23**, 69–96.
- [14] M. Grayson, 1987, *The heat equation shrinks embedded plane curves to points*, J. Diff. Geom. **26**, 285–314.
- [15] R. S. Hamilton, 1982, *Three manifolds with positive Ricci curvature*, J. Diff. Geom. **17**, 255–306.
- [16] G. Huisken, 1984, *Flow by mean curvature of convex surface into point*. J. Diff. Geom. **20**, 237–266. ,
- [17] J. Rubinstein, P. Sternberg, & J. B. Keller, 1989, *Fast reaction, slow diffusion and curve shorting*, SIAM J. Appl. Math. **49**, 116–133.
- [18] R. Temam, 1988, INFINITE-DIMENSIONAL DYNAMICAL SYSTEMS IN MECHANICS AND PHYSICS, Appl. Math. Sci., Springer-Verlag, New York.

#	Author/s	Title
839	Oscar P. Bruno and Fernando Reitich,	Numerical solution of diffraction problems: a method of variation of boundaries
840	Oscar P. Bruno and Fernando Reitich,	Solution of a boundary value problem for Helmholtz equation via variation of the boundary into the complex domain
841	Victor A. Galaktionov and Juan L. Vazquez,	Asymptotic behaviour for an equation of superslow diffusion. The Cauchy problem
842	Josephus Hulshof and Juan Luis Vazquez,	The Dipole solution for the porous medium equation in several space dimensions
843	Shoshana Kamin and Juan Luis Vazquez,	The propagation of turbulent bursts
844	Miguel Escobedo, Juan Luis Vazquez and Enrike Zuazua,	Source-type solutions and asymptotic behaviour for a diffusion-convection equation
845	Marco Biroli and Umberto Mosco,	Discontinuous media and Dirichlet forms of diffusion type
846	Stathis Filippas and Jong-Shenq Guo,	Quenching profiles for one-dimensional semilinear heat equations
847	H. Scott Dumas,	A Nekhoroshev-like theory of classical particle channeling in perfect crystals
848	R. Natalini and A. Tesei,	On a class of perturbed conservation laws
849	Paul K. Newton and Shinya Watanabe,	The geometry of nonlinear Schrödinger standing waves
850	S.S. Sritharan,	On the nonsmooth verification technique for the dynamic programming of viscous flow
851	Mario Taboada and Yuncheng You,	Global attractor, inertial manifolds and stabilization of nonlinear damped beam equations
852	Shigeru Sakaguchi,	Critical points of solutions to the obstacle problem in the plane
853	F. Abergel, D. Hilhorst and F. Issard-Roch,	On a dissolution-growth problem with surface tension in the neighborhood of a stationary solution
854	Erasmus Langer,	Numerical simulation of MOS transistors
855	Haïm Brezis and Shoshana Kamin,	Sublinear elliptic equations in $\mathbf{R}^n$
856	Johannes C.C. Nitsche,	Boundary value problems for variational integrals involving surface curvatures
857	Chao-Nien Chen,	Multiple solutions for a semilinear elliptic equation on $\mathbf{R}^N$ with nonlinear dependence on the gradient
858	D. Brochet, X. Chen and D. Hilhorst,	Finite dimensional exponential attractor for the phase field model
859	Joseph D. Fehribach,	Mullins-Sekerka stability analysis for melting-freezing waves in helium-4
860	Walter Schempp,	Quantum holography and neurocomputer architectures
861	D.V. Anosov,	An introduction to Hilbert's 21st problem
862	Herbert E Huppert and M Grae Worster,	Vigorous motions in magma chambers and lava lakes
863	Robert L. Pego and Michael I. Weinstein,	A class of eigenvalue problems, with applications to instability of solitary waves
864	Mahmoud Affouf,	Numerical study of a singular system of conservation laws arising in enhanced oil reservoirs
865	Darin Beigie, Anthony Leonard and Stephen Wiggins,	The dynamics associated with the chaotic tangles of two dimensional quasiperiodic vector fields: theory and applications
866	Gui-Qiang Chen and Tai-Ping Liu,	Zero relaxation and dissipation limits for hyperbolic conservation laws
867	Gui-Qiang Chen and Jian-Guo Liu,	Convergence of second-order schemes for isentropic gas dynamics
868	Aleksander M. Simon and Zbigniew J. Grzywna,	On the Larché-Cahn theory for stress-induced diffusion
869	Jerzy Luczka, Adam Gadowski and Zbigniew J. Grzywna,	Growth driven by diffusion
870	Mitchell Luskin and Tsorng-Whay Pan,	Nonplanar shear flows for nonaligning nematic liquid crystals
871	Mahmoud Affouf,	Unique global solutions of initial-boundary value problems for thermodynamic phase transitions
872	Richard A. Brualdi and Keith L. Chavey,	Rectangular L -matrices
873	Xinfu Chen, Avner Friedman and Bei Hu,	The thermistor problem with zero-one conductivity II
874	Raoul LePage,	Controlling a diffusion toward a large goal and the Kelly principle
875	Raoul LePage,	Controlling for optimum growth with time dependent returns
876	Marc Hallin and Madan L. Puri,	Rank tests for time series analysis a survey
877	V.A. Solonnikov,	Solvability of an evolution problem of thermocapillary convection in an infinite time interval
878	Horia I. Ene and Bogdan Vernescu,	Viscosity dependent behaviour of viscoelastic porous media
879	Kaushik Bhattacharya,	Self-accommodation in martensite
880	D. Lewis, T. Ratiu, J.C. Simo and J.E. Marsden,	The heavy top: a geometric treatment
881	Leonid V. Kalachev,	Some applications of asymptotic methods in semiconductor device modeling
882	David C. Dobson,	Phase reconstruction via nonlinear least-squares
883	Patricio Aviles and Yoshikazu Giga,	Minimal currents, geodesics and relaxation of variational integrals on mappings of bounded variation
884	Patricio Aviles and Yoshikazu Giga,	Partial regularity of least gradient mappings

- 885 **Charles R. Johnson and Michael Lundquist**, Operator matrices with chordal inverse patterns
886 **B.J. Bayly**, Infinitely conducting dynamos and other horrible eigenproblems
887 **Charles M. Elliott and Stefan Luckhaus**, 'A generalised diffusion equation for phase separation of a multi-compon
mixture with interfacial free energy'
888 **Christian Schmeiser and Andreas Unterreiter**, The derivation of analytic device models by asymptotic methods
889 **LeRoy B. Beasley and Norman J. Pullman**, Linear operators that strongly preserve the index of imprimitivity
890 **Jerry Donato**, The Boltzmann equation with lie and cartan
891 **Thomas R. Hoffend Jr., Peter Smereka and Roger J. Anderson**, A method for resolving the laser induced
local heating of moving magneto-optical recording media
892 **E.G. Kalnins, Willard Miller, Jr. and Sanchita Mukherjee**, Models of q -algebra representations:
the group of plane motions
893 **T.R. Hoffend Jr. and R.K. Kaul**, Relativistic theory of superpotentials for a nonhomogeneous, spatially
isotropic medium
894 **Reinhold von Schwerin**, Two metal deposition on a microdisk electrode
895 **Vladimir I. Oliker and Nina N. Uraltseva**, Evolution of nonparametric surfaces with speed depending on
curvature, III. Some remarks on mean curvature and anisotropic flows
896 **Wayne Barrett, Charles R. Johnson, Raphael Loewy and Tamir Shalom**, Rank incrementation via
diagonal perturbations
898 **Mingxiang Chen, Xu-Yan Chen and Jack K. Hale**, Structural stability for time-periodic one-dimensional
parabolic equations
899 **Hong-Ming Yin**, Global solutions of Maxwell's equations in an electromagnetic field with the temperature-
dependent electrical conductivity
900 **Robert Grone, Russell Merris and William Watkins**, Laplacian unimodular equivalence of graphs
901 **Miroslav Fiedler**, Structure-ranks of matrices
902 **Miroslav Fiedler**, An estimate for the nonstochastic eigenvalues of doubly stochastic matrices
903 **Miroslav Fiedler**, Remarks on eigenvalues of Hankel matrices
904 **Charles R. Johnson, D.D. Olesky, Michael Tsatsomeros and P. van den Driessche**, Spectra with positive
elementary symmetric functions
905 **Pierre-Alain Gremaud**, Thermal contraction as a free boundary problem
906 **K.L. Cooke, Janos Turi and Gregg Turner**, Stabilization of hybrid systems in the presence of feedback delays
907 **Robert P. Gilbert and Yongzhi Xu**, A numerical transmutation approach for underwater sound propagation
908 **LeRoy B. Beasley, Richard A. Brualdi and Bryan L. Shader**, Combinatorial orthogonality
909 **Richard A. Brualdi and Bryan L. Shader**, Strong hall matrices
910 **Håkan Wennerström and David M. Anderson**, Difference versus Gaussian curvature energies; monolayer versus
bilayer curvature energies applications to vesicle stability
911 **Shmuel Friedland**, Eigenvalues of almost skew symmetric matrices and tournament matrices
912 **Avner Friedman, Bei Hu and J.L. Velazquez**, A Free Boundary Problem Modeling Loop Dislocations in
Crystals
913 **Ezio Venturino**, The Influence of Diseases on Lotka-Volterra Systems
914 **Steve Kirkland and Bryan L. Shader**, On Multipartite Tournament Matrices with Constant Team Size
915 **Richard A. Brualdi and Jennifer J.Q. Massey**, More on Structure-Ranks of Matrices
916 **Douglas B. Meade**, Qualitative Analysis of an Epidemic Model with Directed Dispersion
917 **Kazuo Murota**, Mixed Matrices Irreducibility and Decomposition
918 **Richard A. Brualdi and Jennifer J.Q. Massey**, Some Applications of Elementary Linear Algebra in
Combinations
919 **Carl D. Meyer**, Sensitivity of Markov Chains
920 **Hong-Ming Yin**, Weak and Classical Solutions of Some Nonlinear Volterra Integro-differential Equations
921 **B. Leinkuhler and A. Ruehli**, Exploiting Symmetry and Regularity in Waveform Relaxation Convergence
Estimation
922 **Xinfu Chen and Charles M. Elliott**, Asymptotics for a Parabolic Double Obstacle Problem
923 **Yongzhi Xu and Yi Yan**, An Approximate Boundary Integral Method for Acoustic Scattering in Shallow Oceans
924 **Yongzhi Xu and Yi Yan**, Source Localization Processing in Perturbed Waveguides
925 **Kenneth L. Cooke and Janos Turi**, Stability, Instability in Delay Equations Modeling Human Respiration
926 **F. Bethuel, H. Brezis, B.D. Coleman and F. Hélein**, Bifurcation Analysis of Minimizing Harmonic
Maps Describing the Equilibrium of Nematic Phases Between Cylinders
927 **Frank W. Elliott, Jr.**, Signed Random Measures: Stochastic Order and Kolmogorov Consistency Conditions
928 **D.A. Gregory, S.J. Kirkland and B.L. Shader**, Pick's Inequality and Tournaments
929 **J.W. Demmel, N.J. Higham and R.S. Schreiber**, Block LU Factorization
930 **Victor A. Galaktionov and Juan L. Vazquez**, Regional Blow-Up in a Semilinear Heat Equation with
Convergence to a Hamilton-Jacobi Equation
931 **Bryan L. Shader**, Convertible, Nearly Decomposable and Nearly Reducible Matrices
932 **Dianne P. O'Leary**, Iterative Methods for Finding the Stationary Vector for Markov Chains