**NONSTATIONARY FILTRATION IN PARTIALLY
SATURATED POROUS MEDIA**

By

Xinfu Chen
Avner Friedman
and
Tsuyoshi Kimura

IMA Preprint Series # 1099

January 1993

NONSTATIONARY FILTRATION IN PARTIALLY SATURATED POROUS MEDIA

XINFU CHEN*, AVNER FRIEDMAN** AND TSUYOSHI KIMURA**

Abstract. A nonstationary 2-dimensional filtration in a porous medium is considered, whereby part of the medium is saturated, another part is unsaturated but wet, and the remaining part is dry. The saturated/unsaturated and unsaturated/dry interfaces are free boundaries. It is shown that there exists a unique solution and that the saturation function is continuous in the wet portion of the medium; this implies that the two interfaces are separated. Under some (monotonicity type) conditions on the initial and boundary data it is shown that the free boundaries are continuous.

§1. The model. We consider a non-stationary flow in porous medium. Denote by u the hydrostatic potential due to capillary suction, i.e., the pressure, and denote by $c(u)$ the relative volumetric moisture content, i.e., the saturation. If u exceeds a critical value, which we take to be 0, then the porous medium becomes saturated and $c(u)$ is equal to a constant, which is taken to be 1. In the dry portion of the porous medium the pressure is also a constant, which is taken to be -1 . The function $c(u)$ is monotone increasing for $-1 < u < 0$ and, for simplicity, we take it to be $1 + u$. Thus,

$$c(u) = \begin{cases} 1 + u & \text{if } -1 < u < 0 \\ 1 & \text{if } u \geq 0 \end{cases}$$

The pressure satisfies the equation [3; p. 488]

$$(1.1) \quad \frac{\partial}{\partial t} c(u) = \Delta u$$

in the flow region. To account for the free boundary condition at the wet/dry interface (where $u = -1$) we extend the definition of $c(u)$ to $u = -1$ by taking

$$(1.2) \quad c(u) = \min\{u, 0\} + H(u + 1) \quad \text{for } u \geq -1$$

where H is the Heaviside function. As shown in [7], equation (1.1) across the wet/dry interface represents the (physical) equation of continuity, provided gravity is ignored. To

---

*University of Pittsburgh, Department of Mathematics and Statistics, Pittsburgh, Pennsylvania 15260; e-mail address: xinfu@unix.cis.pitt.edu. Partially supported by the National Science Foundation Grant DMS-9200459

**University of Minnesota, Institute for Mathematics and its Applications, Minneapolis, Minnesota 55455. This work is partially supported by the National Science Foundation Grant DMS-87-22187.

**Kao Corporation, Institute for Knowledge and Intelligence Science, Bunka 2-1-3, Sumida-ku, Tokyo 131, Japan

include gravity in the downward direction $-y$, the term $\frac{\partial}{\partial y} H(u + 1)$ needs to be added to the right hand side of (1.1); see [7].

We note that in the unsaturated region equation (1.1) is the same equation satisfied by the temperature in the one-phase Stefan problem.

In the unsaturated/saturated region equation (1.1) becomes

$$(1.3) \quad \frac{\partial \alpha(u)}{\partial t} = \Delta u \quad \alpha(u) = 1 + \min\{u, 0\}$$

and the condition one imposes at the interface $u = 0$ is that $\alpha(u)$ be continuous.

The one-phase Stefan problem has been studied extensively in the literature; see [9] and the references therein; under some initial and boundary conditions the free boundary is C^∞ . Equation (1.3) has been studied in recent years by several authors: [6] [11] [14] [18] [19] for one dimension and [1] [2] [8] [10] [12] [13] [15] for higher dimensions; in the case of 1-dimension the free boundary is known to be in C^∞ .

In this paper we consider the filtration problem in the 2-dimensional half-space

$$\mathbf{R}_2^+ = \{(x, y) ; x \in \mathbf{R}^1, y > 0\}$$

and assume that

$$(1.4) \quad \frac{\partial u(x, 0, t)}{\partial y} = -g(x, t)$$

where

$$(1.5) \quad \begin{aligned} g(x, t) &= g(-x, t) \geq 0, \\ g(x, t) &= 0 \quad \text{if } |x| > x_0 \quad (0 < x_0 < \infty). \end{aligned}$$

This means that the fluid is being injected across the boundary $y = 0$. We also prescribe the initial moisture content

$$(1.6) \quad c(u(x, y, 0)) = \gamma_0(x, y), \quad \gamma_0(x, y) = \gamma_0(-x, y).$$

Finally, we assume that the medium is dry at ∞ :

$$(1.7) \quad \lim_{x^2 + y^2 \rightarrow \infty} u(x, y, t) = -1, \quad t > 0.$$

Since the data are symmetric with respect to x , we also seek a solution which is symmetric with respect to x (Actually by the uniqueness proof given in §2, no other

solutions exist), that is, $u(x, y, t) = u(-x, y, t)$. We can then restrict the problem to $x > 0$, by imposing the boundary condition

$$(1.8) \quad \frac{\partial u(0, y, t)}{\partial x} = 0 .$$

We refer to (1.1), (1.2), (1.4)–(1.8) as *problem (P)*. The model may represent, for example, a filtration of water into soil and a spill over a table cloth, or over other porous media.

We shall need the following notation:

$$\begin{aligned} \Omega_R &= \{(x, y); x > 0, y > 0, x^2 + y^2 < R^2\} , \\ \Gamma_{1,R} &= \{(x, 0), 0 < x < R\}, \quad \Gamma_{2,R} = \{(0, y); 0 < y < R\} , \\ \Gamma_R &= \{(x, y); x > 0, y > 0, x^2 + y^2 = R^2\} , \\ \Omega_{R,T} &= \Omega_R \times \{0 < t < T\}, \quad \Gamma_{i,R,T} = \Gamma_{i,R} \times \{0 < t < T\} , \\ \Gamma_{R,T} &= \Gamma_R \times \{0 < t < T\} . \end{aligned}$$

Definition 1.1. A pair (u, γ) is called a *weak solution* of problem (P) if

$$\begin{aligned} u &\in L^2(0, T; H^1(\Omega_R)) \quad \forall R > 0, T > 0 , \\ \gamma &\in c(u) , \\ u &= -1, \quad \gamma = -1 \quad \text{if} \quad x^2 + y^2 > \sigma(t) \end{aligned}$$

where $\sigma(t)$ is a continuous monotone increasing function of t and, for any $R > \sigma(T)$,

$$(1.9) \quad \iint_{\Omega_{R,T}} (\nabla u \cdot \nabla \zeta - \gamma \zeta_t) = \iint_{\Omega_R} \gamma_0 \zeta(\cdot, 0) + \iint_{\Gamma_{1,R,T}} g \zeta \quad \forall \zeta \in V_{R,T}$$

where

$$V_{R,T} = \{\zeta \in H^1(\Omega_{R,T}), \zeta(\cdot, T) = 0, \zeta|_{\Gamma_{R,T}} = 0\} .$$

Note that $\gamma \in c(u)$ means that $\gamma(x, y, t) = c(u(x, y, t))$ if $u(x, y, t) > -1$,

$$-1 \leq \gamma(x, y, t) \leq 0 \quad \text{if} \quad u(x, y, t) = -1 ,$$

and $u \geq -1$ a.e. (i.e., the graph $c(u)$ is empty if $u < -1$).

In §2 we define a truncated problem (P_R) and prove that it has a unique solution. In §3 we let $R \rightarrow \infty$ and establish the existence and uniqueness of a weak solution to problem

(P). Next, in §4 we prove that $\max\{c(u), 0\} = 1 + \min\{u, 0\}$ is a continuous function. This implies that the two interfaces $S_1 = \partial\{u = 0\} \cap \partial\{u < 0\}$ and $S_2 = \partial\{u > -1\} \cap \partial\{u = -1\}$ are separated by positive distance. In §5 we make the assumptions

$$\frac{\partial \gamma_0}{\partial x} \leq 0, \quad \frac{\partial \gamma_0}{\partial y} \leq 0, \quad \frac{\partial g(x, t)}{\partial x} \leq 0, \quad g(0, t) > 0$$

and prove that S_1 and S_2 have continuous representations $r = \zeta_1(\theta, t)$ and $r = \zeta_2(\theta, t)$ (in polar coordinates (r, θ)); ζ_2 is actually C^∞ , whereas for ζ_1 we only get a logarithmic-type modulus of continuity.

§2. The truncated problem. In this section we assume that

$$(2.1) \quad g \in L^2(\Gamma_{1,R,T}), \quad g \geq 0,$$

$$(2.2) \quad -1 \leq \gamma_0 \leq 1.$$

Definition 2.1. A pair (u, γ) is called a *weak solution* to problem (P_R) if for any $T > 0$

$$(2.3) \quad u \in L^2(0, T; H^1(\Omega_R)), \quad u = -1 \quad \text{on} \quad \Gamma_{R,T},$$

$$(2.4) \quad \gamma \in c(u)$$

and (1.9) is satisfied for any $\zeta \in V_{R,T}$.

THEOREM 2.1. *There exists at most one weak solution to problem (P_R) .*

Proof. Suppose there exist two solutions (u_1, γ_1) and (u_2, γ_2) . For any $t_0 \in (0, T)$ set

$$\zeta_{t_0} = \begin{cases} \int_t^{t_0} [u_1(\cdot, \tau) - u_2(\cdot, \tau)] d\tau & \text{if } t < t_0 \\ 0 & \text{if } t > t_0. \end{cases}$$

Then $\zeta_{t_0} \in V_{R,T}$. Taking $\zeta = \zeta_{t_0}$ in (1.9) for both $u = u_1$ and $u = u_2$ and subtracting one equation from the other, we get

$$\begin{aligned} 0 &= \int_0^{t_0} \int_{\Omega_R} \left[\nabla(u_1 - u_2) \cdot \nabla \int_t^{t_0} (u_1(\cdot, \tau) - u_2(\cdot, \tau)) + (\gamma_1 - \gamma_2)(u_1 - u_2) \right] \\ &= \frac{1}{2} \int_{\Omega_R} \left| \nabla \int_0^{t_0} (u_1(\cdot, \tau) - u_2(\cdot, \tau)) \right|^2 + \iint_{\Omega_{R,t_0}} (\gamma_1 - \gamma_2)(u_1 - u_2). \end{aligned}$$

Since $\gamma_i \in c(u_i)$, $(\gamma_1 - \gamma_2)(u_1 - u_2) \geq 0$. Hence

$$\nabla \int_0^{t_0} [u_1(\cdot, \tau) - u_2(\cdot, \tau)] = 0 \quad \text{a.e.}$$

which implies that $u_1 - u_2 = 0$ a.e. and then, by (1.9), $\gamma_1 - \gamma_2 = 0$ a.e. .

THEOREM 2.2. *There exists a unique weak solution to problem (P_R) .*

Proof. Let c_ε ($\varepsilon > 0$) be smooth approximations to c , as $\varepsilon \rightarrow 0$, so that $c_\varepsilon(0) = 1$, and

$$(2.5) \quad c'_\varepsilon(u) = \begin{cases} \frac{\varepsilon}{(1+u)^3} & \text{if } \varepsilon < u < \infty \\ 1 & \text{if } -1 \leq u \leq 0 \\ \frac{1}{\varepsilon} & \text{if } -\infty < u < -1 - \varepsilon, \end{cases}$$

$$(2.6) \quad c''_\varepsilon(u) \leq 0.$$

Let δ_ε be such that

$$c_\varepsilon(-1 - \delta_\varepsilon) = -1.$$

Note that $\delta_\varepsilon \leq 2\varepsilon$, and $c_\varepsilon \leq 1 + 2\varepsilon$.

Let $\gamma_{0\varepsilon}$ be C^∞ functions that

$$-1 \leq \gamma_{0\varepsilon} \leq 1, \quad \gamma_{0\varepsilon} = -1 \quad \text{in a neighborhood of } \Gamma_R,$$

$$\frac{\partial \gamma_{0\varepsilon}}{\partial x} = 0 \quad \text{on } \Gamma_{2,R},$$

$$\gamma_{0\varepsilon} \rightarrow \gamma_0 \quad \text{in } L^2(\Omega_R) \quad \text{and a.e.}$$

as $\varepsilon \rightarrow 0$. Set $u_{0\varepsilon} = c_\varepsilon^{-1}(\gamma_{0\varepsilon})$. Then

$$-1 - \delta_\varepsilon \leq u_{0\varepsilon} \leq 0,$$

$$u_{0\varepsilon} = -1 - \delta_\varepsilon \quad \text{in a neighborhood of } \Gamma_R,$$

$$\frac{\partial u_{0\varepsilon}}{\partial x} = 0 \quad \text{on } \Gamma_{2,R}.$$

Let g_ε be smooth functions such that

$$g_\varepsilon \geq 0, \quad -g_\varepsilon = \frac{\partial u_{0\varepsilon}}{\partial y} \quad \text{on} \quad \Gamma_{1,R} \cap \{t = 0\},$$

$$g_\varepsilon \rightarrow g \quad \text{in} \quad L^2(\Gamma_{1,R,T}).$$

Consider the parabolic problem

$$(2.7) \quad \frac{\partial}{\partial t} c_\varepsilon(u) = \Delta u \quad \text{in} \quad \Omega_{R,T},$$

$$(2.8) \quad \frac{\partial u}{\partial x} = 0 \quad \text{on} \quad \Gamma_{2,R,T},$$

$$(2.9) \quad -\frac{\partial u}{\partial y} = g_\varepsilon \quad \text{on} \quad \Gamma_{1,R,T},$$

$$(2.10) \quad u = -1 - \delta_\varepsilon \quad \text{on} \quad \Gamma_{R,T},$$

$$(2.11) \quad u = u_{0\varepsilon} \quad \text{at} \quad t = 0.$$

Since the initial and boundary conditions are smooth and satisfy the compatibility conditions for a classical solution, there exists a unique solution u_ε such that

$$u_\varepsilon \in C^{2+\alpha, 1+\alpha/2}(\overline{\Omega}_{R,T}) \cap C^\infty(\Omega_{R,T}).$$

By the maximum principle

$$(2.12) \quad u_\varepsilon \geq -1 - \delta_\varepsilon.$$

Multiplying (2.7) by $u_\varepsilon + 1 + \delta_\varepsilon$ and integrating over $\Omega_{R,T}$ we get

$$(2.13) \quad \int_{\Omega_R} W_\varepsilon(u_\varepsilon(\cdot, T)) + \iint_{\Omega_{R,T}} |\nabla u_\varepsilon|^2 = \int_{\Omega_R} W_\varepsilon(u_\varepsilon(\cdot, 0)) + \iint_{\Gamma_{1,R,T}} g_\varepsilon(u_\varepsilon + 1 + \delta_\varepsilon)$$

where

$$W_\varepsilon(u) = \int_{-1-\delta_\varepsilon}^u c'_\varepsilon(s)(s + 1 + \delta_\varepsilon) \, ds.$$

Notice that $W_\varepsilon(u) \geq 0$ and

$$W_\varepsilon(u_\varepsilon(\cdot, 0)) \leq 1 + C\varepsilon \quad (C \text{ constant})$$

Since by Sobolev's inequalities

$$\|f\|_{2,\Omega_R}^2 + \|f\|_{2,\Gamma_{1,R}}^2 \leq C_R(\|\nabla f\|_{2,\Omega_R}^2 + \|f\|_{2,\Gamma_R}^2) ,$$

we deduce from (2.13) that

$$(2.14) \quad \int_0^T \|u_\varepsilon\|_{H^1(\Omega_R)}^2 \leq C_R \left[1 + \int_0^T \|g_\varepsilon\|_{2,\Gamma_{1,R}}^2 \right] .$$

The function $\gamma_\varepsilon = c_\varepsilon(u_\varepsilon)$ satisfies

$$(2.15) \quad -1 - \delta_\varepsilon \leq \gamma_\varepsilon \leq 1 + 2\varepsilon .$$

We can now choose a sequence $\varepsilon \rightarrow 0$ such that

$$(2.16) \quad \begin{aligned} u_\varepsilon &\rightharpoonup u && \text{weakly in } L^2(\Omega_{R,T}) , \\ \nabla u_\varepsilon &\rightharpoonup \nabla u && \text{weakly in } L^2(\Omega_{R,T}) , \\ \gamma_\varepsilon &\rightharpoonup \gamma && \text{weakly in } L^2(\Omega_{R,T}) , \end{aligned}$$

for all $T > 0$ where u, γ are functions satisfying

$$\begin{aligned} u &\in L^2[0, T; H^1(\Omega_R)] , \quad u \geq -1 , \\ \gamma &\in L^\infty(\Omega_{R,T}) , \quad -1 \leq \gamma \leq 1 \end{aligned}$$

for all $T > 0$. Multiplying (2.7) by $\zeta, \zeta \in V_{R,T}$, integrating over $\Omega_{R,T}$, and passing to the limit with $\varepsilon \rightarrow 0$, we obtain the relation (1.9). In order to complete the proof that (u, γ) is a weak solution it remains to show that (2.4) holds.

Set

$$(2.17) \quad w_\varepsilon = \max\{c_\varepsilon(u_\varepsilon), 0\} = 1 + \min\{u, 0\} .$$

Then $|\nabla w_\varepsilon| \leq |\nabla u_\varepsilon|$ and therefore, by (2.14), for any $0 \leq t_1 < t_2 \leq T$,

$$(2.18) \quad \int_{t_1}^{t_2} \|w_\varepsilon(\cdot, \tau)\|_{H^1(\Omega_R)} \leq C_{R,T} < \infty .$$

Next, for any small $h > 0$,

$$\begin{aligned}
& \int_0^{T-h} \int_{\Omega_R} |w_\varepsilon(\cdot, t+h) - w_\varepsilon(\cdot, t)|^2 \\
& \leq \int_0^{T-h} \int_{\Omega_R} [c_\varepsilon(u_\varepsilon(\cdot, t+h)) - c_\varepsilon(u_\varepsilon(\cdot, t))] [u_\varepsilon(\cdot, t+h) - u_\varepsilon(\cdot, t)] \\
& = \int_0^{T-h} \int_{\Omega_R} \int_t^{t+h} \frac{\partial}{\partial \tau} c_\varepsilon(u_\varepsilon(\cdot, \tau)) \cdot [u_\varepsilon(\cdot, t+h) - u_\varepsilon(\cdot, t)] \\
& = \int_0^{T-h} \int_{\Omega_R} \int_t^{t+h} \Delta u_\varepsilon(\cdot, \tau) [u_\varepsilon(\cdot, t+h) - u_\varepsilon(\cdot, t)] \\
& = \int_0^{T-h} \int_t^{t+h} \left\{ \int_{\Gamma_{1,R}} g(\cdot, \tau) [u_\varepsilon(\cdot, t+h) - u_\varepsilon(\cdot, t)] - \int_{\Omega_R} \nabla u_\varepsilon(\cdot, \tau) \cdot [\nabla u_\varepsilon(\cdot, t+h) - \nabla u_\varepsilon(\cdot, t)] \right\}
\end{aligned}$$

and after applying the Schwarz inequality on the right-hand side,

$$(2.19) \quad \int_0^{T-h} \int_{\Omega_R} |w_\varepsilon(\cdot, t+h) - w_\varepsilon(\cdot, t)|^2 \leq Ch$$

where C is a constant independent of ε . The estimate (2.18) implies that

$$\left\{ \int_{t_1}^{t_2} w_\varepsilon(\cdot, \tau) d\tau \right\} \quad \text{is precompact in } L^2(\Omega_R)$$

for any $0 \leq t_1 < t_2 \leq T$. This together with (2.19) can be used to easily show (see [17]) that $\{w_\varepsilon\}$ is precompact in $L^2(\Omega_{R,T})$. It follows that, for a subsequence,

$$w_\varepsilon \rightarrow w \quad \text{in } L^2(\Omega_{R,T}) \quad \text{and a.e.}$$

for some function $w \in L^2(\Omega_{R,T})$, for any $T > 0$. Consider the set in $\Omega_{R,T}$,

$$C_\mu \equiv \{\mu < w < 1 - \mu\}, \quad \mu > 0.$$

For any small $\lambda > 0$ there is a set A_λ of measure $< \lambda$ such that (for a subsequence) $w_\varepsilon \rightarrow w$ uniformly in $C_\mu \setminus A_\lambda$. Therefore

$$\frac{\mu}{2} < w_\varepsilon < 1 - \frac{\mu}{2} \quad \text{in } C_\mu \setminus A_\lambda$$

if ε is small and, consequently,

$$\frac{\mu}{2} < c_\varepsilon(u_\varepsilon) < 1 - \frac{\mu}{2}.$$

From the definition of c_ε it then follows that u_ε is uniformly convergent on $C_\mu \setminus A_\lambda$. The limiting function is of course u (since $u_\varepsilon \rightarrow u$). Since λ and μ are arbitrary,

$$(2.20) \quad \gamma_\varepsilon = c_\varepsilon(u_\varepsilon) \rightarrow c(u) \quad \text{a.e. in } 0 < w < 1.$$

The same argument shows that, on $\{w = 1\} \setminus \tilde{A}_\lambda$, $c_\varepsilon(u_\varepsilon) > 1 - \mu/2$ and $u_\varepsilon > -\frac{\mu}{2}$ if ε is small enough where $\text{meas}(\tilde{A}_\lambda) < \lambda$. Thus $\liminf_{\varepsilon \rightarrow 0} c_\varepsilon(u_\varepsilon) \geq 1$ and $\liminf_{\varepsilon \rightarrow 0} u_\varepsilon \geq 0$ on $w = 1$. Since $c_\varepsilon(u) \leq 1 + 2\varepsilon$ for all u , we conclude that

$$(2.21) \quad \gamma = 1 \quad \text{on } \{w = 1\} = \{c(u) = 1\}.$$

Next, for any $\delta > 0$,

$$(2.22) \quad \begin{aligned} 0 &= \lim_{\varepsilon \rightarrow 0} |\{w_\varepsilon > \delta, w = 0\}| = \lim_{\varepsilon \rightarrow 0} |\{c_\varepsilon(u_\varepsilon) > \delta, w = 0\}| \\ &= \lim_{\varepsilon \rightarrow 0} |\{u_\varepsilon \geq -1 + \delta, w = 0\}|. \end{aligned}$$

Since $u_\varepsilon \rightarrow u$,

$$\begin{aligned} \int_{\{w=0\}} (u + 1 - \delta)^+ &= \int_{\{w=0\}} (u + 1 - \delta) \text{sgn}(u + 1 - \delta)^+ = \lim_{\varepsilon \rightarrow 0} \int_{\{w=0\}} (u_\varepsilon + 1 - \delta) \text{sgn}(u + 1 - \delta) \\ &\leq \lim_{\varepsilon \rightarrow 0} |\{u_\varepsilon + 1 - \delta > 0, w = 0\}|^{1/2} \left\{ \int_{\{w=0\}} (u_\varepsilon + 1 - \delta)^2 \right\}^{1/2} \rightarrow 0 \end{aligned}$$

by (2.22) and the fact that $\|u_\varepsilon\|_{L^2} \leq C$. It follows that $u \leq -1 + \delta$ on $\{w = 0\}$ for any $\delta > 0$, so that

$$u = -1 \quad \text{on } w = 0;$$

in other words, $c(u) = c(-1) = [-1, 0]$ on $\{w = 0\}$. On the other hand on $\{w = 0\}$ we have $w_\varepsilon \rightarrow 0$ a.e. so that $\overline{\lim} c_\varepsilon(u_\varepsilon) \leq 0$, and thus $\gamma \leq 0$. Since also $\gamma \geq -1$ we conclude that $\gamma \in c(u)$ on $\{w = 0\}$. Recalling (2.20), (2.22), the assertion (2.4) follows.

§3. The complete problem. In this section we extend Theorem 2.2 to problem (P). We shall need the following comparison result.

LEMMA 3.1. *Let $(u^1, \gamma^1), (u^2, \gamma^2)$ satisfy*

$$\begin{aligned} u^i &\in L^2(0, T; H^1(\Omega_R)), \quad u_i = -1 \quad \text{on} \quad \Gamma_{R,T}, \\ \gamma^i &\in c(u^i), \end{aligned}$$

and

$$(3.1) \quad \iint_{\Omega_{R,T}} (\nabla u^1 \cdot \nabla \zeta - \gamma^1 \zeta_t) \geq \int_{\Omega_R} \gamma_0^1 \zeta(\cdot, 0) + \iint_{\Gamma_{1,R,T}} g^1 \zeta,$$

$$(3.2) \quad \iint_{\Omega_{R,T}} (\nabla u^2 \cdot \nabla \zeta - \gamma^2 \zeta_t) \leq \int_{\Omega_R} \gamma_0^2 \zeta(\cdot, 0) + \iint_{\Gamma_{1,R,T}} g^2 \zeta$$

for any $\zeta \in V_{R,T}$, $\zeta \geq 0$, where $\gamma_0^2 \leq \gamma_0^1$, $g^2 \leq g^1$. Then either $u^2 \leq u^1$ or $\gamma^2 \leq \gamma^1$.

Proof. The proof is an extension of the proof of stability for the Stefan problem [9; Chap. 5]. Set

$$u = u^2 - u^1, \quad \gamma = \gamma^2 - \gamma^1, \quad \gamma_0 = \gamma_0^2 - \gamma_0^1, \quad g = g^2 - g^1.$$

Taking ζ in $V_{R,T}$, $\zeta \geq 0$, $[\partial \zeta / \partial n]_{\Gamma_{i,R,T}} = 0$ ($i = 1, 2$) and substituting it in (3.1), (3.2), we get

$$(3.3) \quad \iint_{\Omega_{R,T}} (u \Delta \zeta + \gamma \zeta_t) + \int_{\Omega_R} \gamma_0 \zeta(\cdot, 0) + \iint_{\Gamma_{1,R,T}} g \zeta \geq 0.$$

We shall apply (3.3) to functions $\zeta = \zeta_m$ constructed as follows: Let a_m, b_m be smooth strictly positive functions and let f_m be smooth nonnegative functions. Then the ζ_m are the solutions of

$$(3.4) \quad a_m \frac{\partial \zeta_m}{\partial t} + b_m \Delta \zeta_m = -f_m \quad \text{in} \quad \Omega_{R,T},$$

$$(3.5) \quad \zeta_m(\cdot, T) = 0,$$

$$(3.6) \quad \zeta_m|_{\Gamma_{R,T}} = 0, \quad \frac{\partial \zeta_m}{\partial n}|_{\Gamma_{i,R,T}} = 0 \quad (i = 1, 2).$$

Clearly $\zeta_m \geq 0$. Multiplying (3.4) by $\frac{1}{a_m} \Delta \zeta_m$ and integrating, we easily obtain after using (3.5), (3.6) and the Schwarz inequality,

$$(3.7) \quad \sup_{0 \leq t \leq T} \int_{\Omega_R} |\nabla \zeta_m(\cdot, t)|^2 + \iint_{\Omega_{R,T}} \frac{b_m}{a_m} |\Delta \zeta_m|^2 \leq C \iint_{\Omega_{R,T}} \frac{f_m^2}{a_m b_m}.$$

Similarly, if we multiply (3.4) by $\frac{1}{b_m} \frac{\partial \zeta_m}{\partial t}$ and integrate, we get

$$(3.8) \quad \sup_{0 \leq t \leq T} \int_{\Omega_R} |\nabla \zeta_m(\cdot, t)|^2 + \iint_{\Omega_{R,T}} \frac{a_m}{b_m} \left| \frac{\partial \zeta_m}{\partial t} \right|^2 \leq C \iint_{\Omega_{R,T}} \frac{f_m^2}{a_m b_m}.$$

Substituting $\zeta = \zeta_m$ into (3.3) we obtain, for any function $\alpha \in L^\infty(\Omega_{R,T})$, $|\alpha| \leq 1$,

$$0 \leq \int_{\Omega_R} \gamma_0 \zeta_m(\cdot, 0) + \iint_{\Gamma_{1,R,T}} g \zeta_m + \iint_{\Omega_{R,T}} (-\alpha) f_m + \iint_{\Omega_{R,T}} \left[(u - \alpha b_m) \Delta \zeta_m + (\gamma - \alpha a_m) \frac{\partial \zeta_m}{\partial t} \right].$$

Hence

$$(3.9) \quad \iint_{\Omega_{R,T}} \alpha f_m \leq \int_{\Omega_R} \gamma_0^+ \zeta_m(\cdot, 0) + \iint_{\Gamma_{1,R,T}} g^+ \zeta_m + \iint_{\Omega_{R,T}} \left[|u - \alpha b_m| |\Delta \zeta_m| + |\gamma - \alpha a_m| \left| \frac{\partial \zeta_m}{\partial t} \right| \right].$$

Using (3.7), (3.8) we can estimate the last integral by

$$\left\{ \iint_{\Omega_{R,T}} \frac{f_m^2}{a_m b_m} \right\}^{1/2} \left\{ \left(\iint_{\Omega_{R,T}} \frac{a_m}{b_m} |u - \alpha b_m|^2 \right)^{1/2} + \left(\iint_{\Omega_{R,T}} \frac{b_m}{a_m} |\gamma - \alpha a_m|^2 \right)^{1/2} \right\}.$$

Similarly,

$$\int_{\Omega_R} \gamma_0^+ \zeta_m(\cdot, 0) \leq \|\gamma_0^+\|_{p, \Omega_R} \|\zeta_m\|_{p', \Omega_R} \leq C \|\gamma_0^+\|_{p, \Omega_R} \|\nabla \zeta_m(\cdot, 0)\|_{2, \Omega_R}$$

$$\leq C \|\gamma_0^+\|_{p, \Omega_R} \left\{ \iint_{\Omega_{R,T}} \frac{f_m^2}{a_m b_m} \right\}^{1/2},$$

$$\int_{\Gamma_{1,R,T}} g^+ \zeta_m \leq \int_0^T \|g^+(\cdot, t)\|_{p, \Gamma_{1,R}} \sup_{0 \leq t \leq T} \left\{ \int_{\Gamma_{1,R}} \zeta_n^{p'}(\cdot, t) \right\}^{1/p'}$$

$$\leq C \int_0^T \|g^+(\cdot, t)\|_{p, \Gamma_{1,R}} \left\{ \iint_{\Omega_{R,T}} \frac{f_m^2}{a_m b_m} \right\}^{1/2}.$$

Take

$$\alpha = \begin{cases} 1 & \text{if } u > 0 \text{ or if } u = 0, \gamma > 0 \\ -1 & \text{if } u < 0 \text{ or if } u = 0, \gamma < 0 \\ 0 & \text{if } u = 0, \gamma = 0 \end{cases}$$

and $\widehat{f}_m = a_m b_m \chi_{\{\alpha > 0\}}$. Then

$$\iint_{\Omega_{R,T}} \alpha \widehat{f}_m = \iint_{\Omega_{R,T}} a_m b_m \chi_{\{\alpha > 0\}} = \iint_{\Omega_{R,T}} \frac{\widehat{f}_m^2}{a_m b_m}.$$

We can approximate $\widehat{f}_m$ by smooth functions f_m to which (3.9) and the subsequent estimates of the right-hand side of (3.9) hold and then, by taking the limit, obtain

$$(3.10) \quad \iint_{\Omega_{R,T}} a_m b_m \chi_{\{\alpha > 0\}} \leq C \left\{ \|\gamma_0^+\|_{p, \Omega_R} + \int_0^T \|g^+(\cdot, t)\|_{p, \Gamma_{1,R}} + J_m \right\}$$

where

$$\begin{aligned} J_m &= \left(\iint_{\Omega_{R,T}} \frac{a_m}{b_m} (u - \alpha b_m)^2 \right)^{1/2} + \left(\iint_{\Omega_{R,T}} \frac{b_m}{a_m} (\gamma - \alpha a_m)^2 \right)^{1/2} \\ &\leq \|a_m\|_\infty^{1/2} \left(\iint_{\Omega_{R,T}} \frac{(|u| - b_m)^2}{b_m} \right)^{1/2} + \|b_m\|_2^{1/2} \left(\iint_{\Omega_{R,T}} \frac{(|\gamma| - a_m)^4}{a_m^2} \right)^{1/2} \end{aligned}$$

by the definition of α .

To further estimate J_m we need the following lemma.

LEMMA 3.2. *If $a \in L^\infty(\Omega_{R,T})$, $a \geq 0$ and $b \in L^2(\Omega_{R,T})$, $b \geq 0$, then there exist sequences of C^∞ functions a_m, b_m in $\overline{\Omega}_{R,T}$ satisfying:*

- (i) $a_m > \frac{1}{m}, b_m > \frac{1}{m},$
- (ii) $\|a_m\|_\infty \leq \|a\|_\infty + 1, \|b_m\|_2 \leq \|b\|_2 + 1,$
- (iii) $a_m \rightarrow a, b_m \rightarrow b \text{ in } L^2(\Omega_{R,T}),$

and

$$(iv) \quad \iint_{\Omega_{R,T}} \frac{(a - a_m)^4}{a_m^2} \rightarrow 0, \quad \iint_{\Omega_{R,T}} \frac{(b - b_m)^2}{b_m} \rightarrow 0,$$

as $m \rightarrow \infty$.

Proof. Let

$$\widehat{a}_m = \max \left\{ a, \frac{1}{m} \right\}, \quad \widehat{b}_m = \max \left\{ b, \frac{1}{m} \right\}$$

and let

$$a_{m\varepsilon} = \rho_\varepsilon * \widehat{a}_m, \quad b_{m\varepsilon} = \rho_\varepsilon * \widehat{b}_m \quad (\varepsilon > 0)$$

be mollifiers of $\widehat{a}_m, \widehat{b}_m$. Then

$$\frac{1}{m} \leq a_{m\varepsilon} \leq \|a\|_\infty + 1, \quad b_{m\varepsilon} \geq \frac{1}{m}$$

and

$$\begin{aligned} a_{m\varepsilon} &\rightarrow \widehat{a}_m \quad \text{in } L^4 \quad (\text{in fact in any } L^p), \\ b_{m\varepsilon} &\rightarrow \widehat{b}_m \quad \text{in } L^2. \end{aligned}$$

We take $\varepsilon = \varepsilon_m$ so that

$$\|a_{m\varepsilon} - \widehat{a}_m\|_4 < \frac{1}{m}, \quad \|b_{m\varepsilon} - \widehat{b}_m\|_2 < \frac{1}{m},$$

and set $a_m = a_{m\varepsilon_m}$, $b_m = b_{m\varepsilon_m}$. Then (i)–(iii) clearly hold, whereas (iv) is proved as follows:

$$\begin{aligned} \iint \frac{(b - b_m)^2}{b_m} &= 2 \iint \frac{(\widehat{b}_m - b_m)^2}{b_m} + 2 \iint \frac{(b - \widehat{b}_m)^2}{b_m} \\ &\leq 2m \|b_{m\varepsilon_m} - \widehat{b}_m\|_2^2 + 2m \iint_{\{b < \frac{1}{m}\}} \left(b - \frac{1}{m}\right)^2 \leq \frac{C}{m} \rightarrow 0, \\ \iint \frac{(a - a_m)^4}{a_m^2} &\leq 16 \iint \frac{(\widehat{a}_m - a_m)^4}{a_m^2} + 16 \iint \frac{(a - \widehat{a}_m)^4}{a_m^2} \\ &\leq 16m^2 \|\widehat{a}_m - a_{m\varepsilon_m}\|_4^4 + 16m^2 \iint_{\{a < \frac{1}{m}\}} \left(a - \frac{1}{m}\right)^4 \leq \frac{C}{m^2} \rightarrow 0. \end{aligned}$$

We apply Lemma 3.2 to $a = |\gamma|$, $b = |u|$ and conclude that $J_m \rightarrow 0$ if $m \rightarrow \infty$. Then (3.10) yields

$$\iint_{\Omega_{R,T}} u^+ \gamma^+ \leq C(\|\gamma_0^+\|_{p,\Omega_R} + \int_0^T \|g^+(\cdot, t)\|_{p,\Gamma_{1,R}})^2.$$

Since, by assumption, the right-hand side is equal to zero, it follows that

$$u^+ \gamma^+ = 0$$

This implies that either $u^2 \leq u^1$ or $\gamma^2 \leq \gamma_1$.

In the sequel we assume that

$$(3.11) \quad \gamma_0(x, y) = -1 \quad \text{if} \quad (x^2 + y^2)^{1/2} \geq x_0, \quad -1 \leq \gamma_0 \leq 1,$$

$$(3.12) \quad 0 \leq g(x, t) \leq M, \quad g(x, t) = 0 \quad \text{if} \quad x \geq x_0,$$

where M and x_0 are positive constants.

THEOREM 3.3. *There exists a continuous monotone increasing function $t \rightarrow \sigma(t)$ such that the solutions (u_R, γ_R) of (P_R) satisfy*

$$u_R = -1, \quad \gamma_R = -1 \quad \text{if} \quad x^2 + y^2 > \sigma(t).$$

Proof. For any $r > x_0 + 1$ denote by w_r the solution to

$$\begin{aligned} \Delta w &= 0 \quad \text{in} \quad \Omega_r, \\ \frac{\partial w}{\partial x} &= 0 \quad \text{on} \quad \Gamma_{2,r}, \\ -\frac{\partial w}{\partial y} &= \begin{cases} N & \text{if} \quad 0 \leq x \leq x_0 \\ 0 & \text{if} \quad x_0 < x < r, \end{cases} \\ w &= -1 \quad \text{if} \quad x^2 + y^2 = r^2. \end{aligned}$$

If $r_1 > r_2$ then $w_{r_1} > w_{r_2} > 0$ in Ω_{r_2} , so that $\partial w_r / \partial r \geq 0$ if $x^2 + y^2 \leq r^2$. Note that, by comparison,

$$w_r(x, y) \leq N \int_{-x_0}^{x_0} \Gamma(x, y; \xi, 0) d\xi$$

where Γ is the Neumann function for Δ in $\{y > 0\}$. Hence

$$\lim_{r \rightarrow \infty} w_r(x, y) \equiv W(x, y) \quad \text{exists and is finite.}$$

We choose $N > M$ and so large that

$$W(x, y) > \gamma_0(x, y) \quad \text{in} \quad \Omega_\infty ,$$

and then also

$$(3.13) \quad c(w_{\tilde{r}}) \geq \gamma_0 \quad \text{for some} \quad \tilde{r} \in (x_0, \infty) .$$

The function $w_r(r \cos \theta, r \sin \theta)$ (for $r > \tilde{r}$) is smooth in (r, θ) (also at $\theta = 0$) and therefore

$$K(r) = \sup_{0 \leq \theta \leq \frac{\pi}{2}} |\nabla w_r(r \cos \theta, r \sin \theta)|$$

is Lipschitz continuous in r . Let $s(t)$ be the solution to

$$\frac{ds}{dt} = K(s) , \quad s(0) = \tilde{r}$$

and define

$$\bar{u}(x, y, t) = w_{s(t)}(x, y),$$

$$\bar{\gamma}(x, y, t) = c(\bar{u})$$

if $x^2 + y^2 \leq s^2(t)$. Extend $\bar{u}$ by -1 and $\bar{\gamma}$ by -1 to $x^2 + y^2 > s^2(t)$. Then, by (3.13), $\bar{\gamma} \geq \gamma_0$ everywhere.

If $R > s(T)$ then for any $\zeta \in V_{R,T}$, $\zeta \geq 0$,

$$\begin{aligned} & \iint_{\Omega_{R,T}} (\nabla \bar{u} \cdot \nabla \zeta - \bar{\gamma} \zeta_t) - \iint_{\Omega_R} \gamma_0 \zeta(\cdot, 0) - \iint_{\Gamma_{1,R,T}} g \zeta \\ &= \int_0^T \int_{\Omega_{s(t)}} (c(\bar{u})_t - \Delta \bar{u}) + \int_0^T \int_{\Gamma_{s(t)}} \left(\frac{\partial \bar{u}}{\partial n} + \dot{s} \right) \zeta \\ &+ \int_{\Omega_{s(0)}} [c(\bar{u}(\cdot, 0)) - \gamma_0] \zeta(\cdot, 0) + \int_{\Gamma_{1,x_0,T}} (N - g) \zeta \geq 0 \end{aligned}$$

since

$$c(\bar{u})_t - \Delta \bar{u} = c'(\bar{u}) \frac{\partial w}{\partial r} \frac{ds(t)}{dt} \geq 0 \quad \left(w = w_r \Big|_{r=s(t)} \right) ,$$

$$- \frac{\partial \bar{u}}{\partial n} \leq K(s(t)) = \dot{s} \quad \text{on} \quad \Gamma_{s(t)}$$

and $\bar{\gamma} \geq \gamma_0$. Applying Lemma 3.1 we deduce that $u_R \leq \bar{u} = -1$ if $x^2 + y^2 \geq s^2(t)$. Using (1.9) with ζ supported in $x^2 + y^2 > s^2(t)$ we find that also $\gamma_R = -1$ if $x^2 + y^2 > s^2(t)$.

Suppose (u_R, γ_R) and $(u_{R'}, \gamma_{R'})$ are weak solutions to (P_R) with $R' > R > s^2(T)$. Then by the proof of Theorem 2.1 we find that $u_R = u_{R'}, \gamma_R = \gamma_{R'}$, if $t < T$. Hence

$$(3.14) \quad u = \lim_{R \rightarrow \infty} u_R, \quad \gamma = \lim_{R \rightarrow \infty} \gamma_R$$

form a weak solution to problem (P) . The uniqueness of the solution follows by the proof of Theorem 2.1. We summarize:

THEOREM 3.3. *If (3.11), (3.12) hold then there exists a unique weak solution (u, γ) to problem (P) ; it is given by (3.14). Furthermore, u is uniformly bounded.*

The last statement needs to be proved. Let w be the solution to

$$\Delta w = 0 \quad \text{in } \Omega_R,$$

with boundary conditions

$$\begin{aligned} w_x(0, y) &= 0 \quad \text{if } 0 < y < R, \\ -w_y(x, 0) &= \begin{cases} M_0 & \text{if } 0 < x < x_0 \\ 0 & \text{if } x_0 < x < R, \end{cases} \\ w &= 0 \quad \text{if } x^2 + y^2 = R^2. \end{aligned}$$

If M_0 is large enough then $c(w) > \gamma_0$ and then, by comparison $u_\varepsilon \leq w$. It follows that

$$(3.15) \quad u_\varepsilon \leq C \quad \text{in } \Omega_{R,T}$$

where C a constant is independent of ε, R, T , and then also $u \leq C$.

§4. Continuity of $\max\{c(u), 0\}$. In this section we assume, in addition to (3.11), (3.12), that

$$(4.1) \quad \gamma_0 \in c(u_0) \quad \text{where} \quad \int |\nabla u_0|^2 < \infty,$$

$$(4.2) \quad g \in BV(0, T; L^1(\Omega_R)) \quad \forall R > 0, T > 0.$$

Multiplying (2.7) (for $u = u_\varepsilon$) by $\partial u_\varepsilon / \partial t$ and integrating yields

$$\begin{aligned} 0 &= \iint_{\Omega_{R,t}} c'_\varepsilon(u_\varepsilon) u_{\varepsilon,t}^2 + \frac{1}{2} \int_{\Omega_R} |\nabla u_\varepsilon(\cdot, \tau)|^2 \Big|_{\tau=0}^{\tau=t} - \iint_{\Gamma_{1,R,t}} u_{\varepsilon,t} g_\varepsilon \\ (4.3) \quad &\geq \iint_{\Omega_{R,t}} c'_\varepsilon(u_\varepsilon) u_{\varepsilon,t}^2 + \frac{1}{2} \int_{\Omega_R} |\nabla u_\varepsilon(\cdot, t)|^2 - \frac{1}{2} \int_{\Omega_R} |\nabla u_\varepsilon(\cdot, 0)|^2 \\ &\quad - \|u_\varepsilon\|_{L^\infty} (2\|g_\varepsilon\|_{L^\infty} + \|g_\varepsilon\|_{BV[0,T;L^1(\Omega_R)]}) . \end{aligned}$$

By (4.1), (4.2) we can define $\gamma_{0\varepsilon}$ and g_ε such that the last two terms are bounded independently of ε (here we use also (3.15)). Defining

$$(4.4) \quad w_\varepsilon = \max\{c_\varepsilon(u_\varepsilon), 0\}$$

and noting that

$$|\nabla w_\varepsilon| \leq |\nabla u_\varepsilon|, \quad |w_{\varepsilon,t}| \leq |u_{\varepsilon,t}|$$

we then obtain from (4.3),

$$(4.5) \quad \iint_{\Omega_{R,T}} |w_{\varepsilon,t}|^2 + \sup_{0 \leq t \leq T} \int_{\Omega_R} |\nabla w_\varepsilon(\cdot, t)|^2 \leq C.$$

This estimate will be used to prove the following theorem.

THEOREM 4.1. *The function $w = \max\{c(u), 0\}$ is continuous.*

Proof. Introduce the notation: $X = (x, y)$,

$$\begin{aligned} B_\rho(X) &= \text{ball of radius } \rho \text{ and center } X, \\ Q_\rho(X, t) &= B_\rho(X) \times \left\{ t - \frac{7}{8}\rho^2, t + \frac{1}{8}\rho^2 \right\}, \\ \int_{B_\rho} &= \frac{1}{|B_\rho|} \int_{B_\rho}, \quad \iint_{Q_\rho} = \frac{1}{|Q_\rho|} \iint_{Q_\rho}, \\ w_{B_\rho} &= \int_{B_\rho} w, \quad w_{Q_\rho} = \iint_{Q_\rho} w \end{aligned}$$

where $B_\rho = B_\rho(X)$, $Q_\rho = Q_\rho(X, t)$. From (4.5) we have

$$(4.6) \quad \iint_{\Omega_{R,T}} |w_t|^2 + \sup_{0 \leq t \leq T} \int_{\Omega_R} |\nabla w|^2 \leq C.$$

LEMMA 4.2. *If a function w satisfies (4.6) then for any $(X, t) \in \Omega_{R,T}$*

$$(4.7) \quad \lim_{\rho \rightarrow 0} \iint_{Q_\rho(X, t)} |w - w_{Q_\rho(X, t)}|^2 = 0.$$

Proof. Since $H^1(0, T; L^2(\Omega_R))$ is continuously imbedded in $C^{1/2}(0, T; L^2(\Omega_R))$, there is a version $t \rightarrow w^*(\cdot, t)$ such that for any $0 < t < T$

$$(4.8) \quad \frac{1}{t} \int_t^{t+h} w(\cdot, \tau) - w^*(\cdot, t) \rightarrow 0 \quad \text{in } L^2(\Omega_R)$$

and therefore also a.e. in (X, t) . Since $w \in L^\infty(0, T; H^1(\Omega_R))$, (4.8) implies that $w^*(\cdot, t) \in H^1(\Omega_R)$ for all $t \in (0, T)$ and

$$(4.9) \quad \|w^*(\cdot, t)\|_{H^1(\Omega_R)} \leq C .$$

By Sobolev's imbedding

$$\begin{aligned} \int_{B_\rho(X)} |w^*(\cdot, t) - w_{B_\rho(X)}^*(t)|^2 &\leq C\rho^2 \int_{B_\rho(X)} |\nabla w^*(\cdot, t)|^2 , \\ \sup_{t - \frac{7}{8}\rho^2 < \tau < t + \frac{1}{8}\rho^2} |w(y, \tau) - w(y, t)|^2 &\leq C\rho^2 \int_{t - \frac{7}{8}\rho^2}^{t + \frac{1}{8}\rho^2} w_t^2(y, s) ds . \end{aligned}$$

Therefore, with $Q_\rho = Q_\rho(X, t)$, $B_\rho = B_\rho(X)$,

$$\begin{aligned} \iint_{Q_\rho} |w - w_{Q_\rho}|^2 &= \inf_{c \in \mathbb{R}^1} \iint_{Q_\rho} |w^* - c|^2 \leq \iint_{Q_\rho} |w^* - w_{B_\rho}^*(t)|^2 \\ &\leq \iint_{Q_\rho} \left\{ 2|w^*(y, \tau) - w^*(y, t)|^2 + 2|w^*(y, t) - w_{B_\rho}^*(t)|^2 \right\} \\ &\leq C\rho^4 \left\{ \iint_{Q_\rho} |w_t^*|^2 + \int_{B_\rho} |\nabla w^*(\cdot, t)|^2 \right\} . \end{aligned}$$

Hence

$$(4.10) \quad \lim_{\rho \rightarrow 0} \iint_{Q_\rho} |w - w_{Q_\rho}|^2 \leq C \overline{\lim}_{\rho \rightarrow 0} \left\{ \iint_{Q_\rho} |w_t^*|^2 + \int_{B_\rho(X)} |\nabla w^*(\cdot, t)|^2 \right\} .$$

From the estimate (4.9) and the estimate on w_t in (4.6), which must hold also for w_t^* , it follows that the right-hand side of (4.10) is equal to zero.

LEMMA 4.3. *There exists a small positive constant μ (independent of ρ) such that*
(i) *if*

$$(4.11) \quad \iint_{Q_\rho(X,t)} \left| \left(w - \frac{1}{2} \right)^+ \right|^2 < \mu$$

then $w \leq \frac{3}{4}$ in $Q_{\rho/2}^*(X,t) \equiv B_{\rho/2}(X) \times \left\{ t - \frac{\rho^2}{8}, t + \frac{\rho^2}{8} \right\}$, and

(ii) if

$$(4.12) \quad \iint_{Q_\rho(X,t)} \left| \left(w - \frac{1}{2} \right)^- \right|^2 < \mu$$

then $w \geq \frac{1}{4}$ in $Q_{\rho/2}^*$.

Proof. For the purpose of this proof only we modify the definition of the $c_\varepsilon(u)$ (defined in (2.5)) for $u \geq 0$, taking

$$(4.13) \quad \begin{aligned} c_\varepsilon(u) &= 1 + u & \text{if } -1 \leq u \leq 0 & \text{(as before),} \\ c'_\varepsilon(u) &= \frac{\varepsilon}{(1+u)^3} & \text{if } u \geq 0 & \text{(instead of } u \geq \varepsilon \text{).} \end{aligned}$$

The new functions c_ε are not differentiable at $u = 0$, so that also u_ε are not as smooth as before. This however does not affect any of the previous results.

Let ζ be a smooth function satisfying:

$$\zeta = 1 \quad \text{in} \quad B_s \times \left[t + \frac{\rho^2}{8} - s^2, t + \frac{\rho^2}{8} \right] \equiv \tilde{Q}_s,$$

$$\zeta = 0 \quad \text{on the parabolic boundary of } Q_\rho,$$

$$\zeta \geq 0;$$

we can choose ζ of the form $\zeta_0(t)\zeta_1(r)$, $r = \sqrt{x^2 + y^2}$, with $\zeta'_1 \leq 0, \zeta''_1 \leq 0$, so that $\Delta\zeta \leq 0$. Here $s \in (0, \rho)$ is still to be determined. Let k be a real number, $k \in \left[\frac{1}{4}, \frac{3}{4} \right]$. Multiplying (2.7) (with $u = u_\varepsilon$) by $(w_\varepsilon - k)^+\zeta^2$ and integrating over $Q_{\rho,\tau} \equiv \{(X', t') ; (X', t') \in Q_\rho, t' < \tau\}$ we get

$$(4.14) \quad \begin{aligned} & \iint_{Q_{\rho,\tau}} \frac{\partial}{\partial t} c_\varepsilon(u_\varepsilon) \cdot (w_\varepsilon - k)^+\zeta^2 + \iint_{Q_{\rho,\tau}} [\nabla u_\varepsilon \cdot \nabla (w_\varepsilon - k)^+]\zeta^2 \\ &= - \iint_{Q_{\rho,\tau}} \nabla u_\varepsilon \cdot (w_\varepsilon - k)^+ \nabla \zeta^2. \end{aligned}$$

We claim that

$$(4.15) \quad \left| \iint_{Q_{\rho,\tau}} [\nabla u_\varepsilon \cdot \nabla (w_\varepsilon - k)^+] \zeta^2 - \iint_{Q_{\rho,\tau}} |\nabla (w_\varepsilon - k)^+|^2 \zeta^2 \right| \leq C_\varepsilon \iint_{Q_{\rho,\tau}} |\nabla u_\varepsilon|^2 .$$

Indeed, if $w_\varepsilon > k$, $u_\varepsilon < 0$ then $w_\varepsilon = c(u_\varepsilon) = u_\varepsilon + 1$ (by (4.13)) so that

$$\nabla u_\varepsilon \cdot \nabla (w_\varepsilon - k)^+ = \nabla (w_\varepsilon - k) \cdot \nabla (w_\varepsilon - k)^+ = |\nabla (w_\varepsilon - k)^+|^2 .$$

These equalities hold trivially also if $w_\varepsilon \leq k$. Finally, if $w_\varepsilon > k$ and $u_\varepsilon > 0$ then $|\nabla w_\varepsilon| \leq \varepsilon |\nabla u_\varepsilon|$, by (4.13), and this gives the error term on the right-hand side of (4.15).

Next we have, since $c_\varepsilon(u_\varepsilon) = w_\varepsilon$ when $w_\varepsilon > k$,

$$(4.16) \quad \begin{aligned} \iint_{Q_{\rho,\tau}} \frac{\partial}{\partial t} c_\varepsilon(u_\varepsilon) \cdot (w_\varepsilon - k)^+ \zeta^2 &= \iint_{Q_{\rho,\tau}} \frac{\partial}{\partial t} (w_\varepsilon - k) \cdot (w_\varepsilon - k)^+ \zeta^2 \\ &= \frac{1}{2} \int_{B_\rho} [(w_\varepsilon - k)^+]^2 \zeta^2(\cdot, \tau) - \iint_{Q_{\rho,\tau}} |(w_\varepsilon - k)^+|^2 \zeta \zeta_t . \end{aligned}$$

Consider finally the right-hand side of (4.14). We may assume that $w_\varepsilon > k$ in the integrand; then $u_\varepsilon + 1 > 0$, $w_\varepsilon = c_\varepsilon((u_\varepsilon + 1)^+ - 1)$ and

$$\nabla u_\varepsilon (w_\varepsilon - k)^+ = \nabla (u_\varepsilon + 1)^+ (w_\varepsilon - k)^+ = \nabla G_\varepsilon(u_\varepsilon) \quad \text{where}$$

$$G_\varepsilon(u_\varepsilon) = \int_{\bar{u}}^{u_\varepsilon+1} (c_\varepsilon(s-1) - k)^+ ds \leq C[(c_\varepsilon(u_\varepsilon) - k)^+]^2 \leq C[(w_\varepsilon - k)^+]^2 ;$$

here $\bar{u}$ is determined by $c_\varepsilon(\bar{u} - 1) = k$. We then have (cf. [8; p. 7])

$$\begin{aligned} - \iint_{Q_{\rho,\tau}} \nabla u_\varepsilon \cdot (w_\varepsilon - k)^+ \nabla \zeta^2 &= - \iint_{Q_{\rho,\tau}} \nabla G_\varepsilon(u_\varepsilon) \cdot \nabla \zeta^2 \\ &= -2 \iint_{Q_{\rho,\tau}} \nabla [\zeta G_\varepsilon(u_\varepsilon)] \cdot \nabla \zeta + 2 \iint_{Q_{\rho,\tau}} G_\varepsilon(u_\varepsilon) |\nabla \zeta|^2 \leq 2 \iint_{Q_{\rho,\tau}} G_\varepsilon(u_\varepsilon) |\nabla \zeta|^2 , \end{aligned}$$

since $\Delta \zeta \leq 0$. Combining this with (4.16), (4.15), we obtain from (4.14), after taking $\varepsilon \rightarrow 0$,

$$(4.17) \quad \begin{aligned} \iint_{Q_\rho} |\nabla((w - k)^+ \zeta)|^2 + \sup_{t - \frac{7}{8}\rho^2 \leq \tau \leq t + \frac{1}{8}\rho^2} \int_{B_\rho} |(w - k)^+|^2 \zeta^2(\tau) \\ \leq C \iint_{Q_\rho} |(w - k)^+|^2 [|\zeta_t| + |\nabla \zeta|^2] . \end{aligned}$$

Notice that ζ can be chosen so that

$$|\zeta_t| + |\nabla \zeta|^2 \leq \frac{C}{(\rho - s)^2} .$$

We can now proceed as in [16; Chap. 2, §6]. Indeed we apply (4.17) recursively with ρ replaced by $\rho_n, s = \rho_{n+1}, k = k_n$ where

$$\rho_n = \left(\frac{1}{2} + \frac{1}{2^n} \right) \rho , \quad k_n = \frac{3}{4} - \frac{1}{2^n} , \quad n \geq 2$$

and obtain (cf. [16; Lemma 6.1, p. 106])

$$\iint_{Q_{\rho_{n+1}}} |(w - k_{n+1})^+|^2 \leq C_0 C^n \left\{ \iint_{Q_{\rho_n}} |(w - k_n)^+|^2 \right\}^{3/2} \quad (n \geq 2)$$

where C_0, C are constants independent of μ . Therefore, by [16; Lemma 5.7, p. 96], if (4.11) holds with μ small enough then

$$\iint_{Q_{\rho_\infty}} |(w - k_\infty)^+|^2 = 0 , \quad \text{i.e.,} \quad w \leq \frac{3}{4} \quad \text{in} \quad Q_{\rho/2} .$$

The proof of (ii) is similar.

We return to the proof of Theorem 4.1. For any $(X, t) \in \Omega_{R,T}$ and for any small $\mu > 0$ we have, by Lemma 4.2,

$$\iint_{Q_\rho(X,t)} |w - w_{Q_\rho(X,t)}|^2 < \mu$$

if ρ is small enough. Consider separately the cases:

$$(i) \quad w_{Q_\rho}(X, t) < \frac{1}{2} ,$$

$$(ii) \quad w_{Q_\rho}(X, t) \geq \frac{1}{2} .$$

In case (i),

$$\iint_{Q_\rho(X,t)} \left| \left(w - \frac{1}{2} \right)^+ \right|^2 \leq \iint_{Q_\rho(X,t)} \left| (w - w_{Q_\rho(X,t)})^+ \right|^2 < \mu$$

and Lemma 4.3 gives

$$w \leq \frac{3}{4} \quad \text{in} \quad Q_{\rho/2}^* = Q_{\rho/2}^*(X, t) ,$$

so that $u \leq -\frac{1}{4}$ in $Q_{\rho/2}^*$. Similarly, in case (ii), $w \geq \frac{1}{4}$ and $u \geq -\frac{3}{4}$ in $Q_{\rho/2}^*$.

In case (i) u satisfies

$$\frac{\partial}{\partial t} (u + H(u + 1)) = \Delta u \quad \text{in } Q_{\rho/2}^* .$$

By a result of Caffarelli and Friedman [5] u is a continuous function with modulus of continuity

$$\omega(r) = C 2^{-|\log r|^\lambda} \quad \text{for any } 0 < \lambda < \frac{1}{2} \quad (C > 0) .$$

This implies the continuity of $\max\{c(u), 0\}$ with the same modulus of continuity. In case (ii) u satisfies (1.1) in $Q_{\rho/2}^*$ where $c(u)$ is continuous, $c(u) = 1$ if $u \geq 1$, and $c'(u) > 0$ if $u < 1$. By a result of DiBenedetto and Gariepy [8], $c(u)$ is continuous, and this concludes the proof of Theorem 4.1.

§5. Properties of the free boundary. In this section we assume that

$$(5.1) \quad \frac{\partial \gamma_0}{\partial x} \leq 0, \quad \frac{\partial \gamma_0}{\partial y} \leq 0, \quad \frac{\partial g}{\partial x} \leq 0 .$$

THEOREM 5.1. *Under the assumptions (5.1),*

$$(5.2) \quad \frac{\partial u}{\partial y} \leq 0, \quad \frac{\partial \gamma}{\partial y} \leq 0$$

and

$$(5.3) \quad \frac{\partial u}{\partial x} \leq 0, \quad \frac{\partial \gamma}{\partial x} \leq 0$$

where the derivatives are taken in the distribution sense.

Proof. Differentiating (2.7) with respect to y we get for $v = \partial u_\epsilon / \partial y$ the equation

$$c'_\epsilon(u_\epsilon) v_t - \Delta v + c''_\epsilon(u_\epsilon) u_{\epsilon,t} v = 0 .$$

By the maximum principle we then conclude that $v \leq 0$. Similarly one shows that $\partial u_\epsilon / \partial x \leq 0$. Taking $\epsilon \rightarrow 0$, the assertions of the theorem follow.

Remark 5.1. If $\frac{\partial}{\partial \theta} \gamma_0(r \cos \theta, r \sin \theta) \leq 0$ then also $\frac{\partial}{\partial \theta} u \leq 0$.

By Theorem 5.1 there is a version of γ which is everywhere monotone decreasing in x and in y ; hence also in $r = \sqrt{x^2 + y^2}$. Set

$$(5.4) \quad \zeta_1(\theta, t) = \sup\{r; \gamma(r \cos \theta, r \sin \theta, t) = 1\} ,$$

$$(5.5) \quad \zeta_2(\theta, t) = \inf\{r; \gamma(r \cos \theta, r \sin \theta, t) \leq 0\} ,$$

and introduce the free boundaries

$$S_1 \quad : \quad r = \zeta_1(\theta, t) ,$$

$$S_2 \quad : \quad r = \zeta_2(\theta, t) .$$

By Theorem 4.1, the sets S_1, S_2 are separated by a positive distance, in any compact set in $\Omega_\infty \cap \{0 < t < \infty\}$.

By Theorem 5.1, for each $t > 0$, $\zeta_1(\theta, t)$ and $\zeta_2(\theta, t)$ are Lipschitz continuous in θ . From a result of Caffarelli [4] it also follows that

$$(5.6) \quad \zeta_2(\theta, t) \in C^\infty\left(0, \frac{\pi}{2}\right] \times (0, \infty) .$$

It is also well known (see, for example, [9]) that

$$(5.7) \quad \frac{\partial \zeta_2}{\partial t} \geq 0 .$$

We proceed to study the free boundary S_1 .

THEOREM 5.2. *If $g(x, t) \equiv 0$ for $t_0 \leq t \leq t_0 + h$ then $\zeta_1(\theta, t) = 0$ for $0 < \theta < \frac{\pi}{2}$, $t_0 < t < t_0 + h$.*

Proof. Let $\bar{u}$ be the solution to the Stefan problem

$$\frac{\partial}{\partial t}(\bar{u} + H(\bar{u} + 1)) = \Delta \bar{u} \quad \text{in} \quad \Omega_R \times (t_0, t_0 + h) , \quad R \gg 1 ,$$

$$c(\bar{u}(\cdot, t_0)) = c(\min[u(\cdot, t_0), 0]) ,$$

$$\bar{u}_x(0, y) = 0 , \quad \bar{u}_y(x, 0) = 0 ,$$

$$\bar{u} = -1 \quad \text{on} \quad \Gamma_R \times (t_0, t_0 + h) .$$

By the maximum principle $-1 \leq \bar{u} < 0$ in $\Omega_{R,T}$, so that $\bar{u}$ is actually a solution of

$$\frac{\partial}{\partial t} c(\bar{u}) = \Delta \bar{u} .$$

Since $c(u(X, t_0)) = c(\bar{u}(X, t_0))$ if $u(X, t_0) \leq 0$ and $c(u(X, t_0)) = 1 = c(0) = c(\bar{u}(X, t_0))$ if $u(X, t_0) > 0$, we have $c(u) = c(\bar{u})$ at $t = t_0$ and by uniqueness (Theorem 2.1) $u \equiv \bar{u}$. It follows that $u < 0$ in $\Omega_R \times (t_0, t_0 + h)$ and therefore $\zeta_1(\theta, t) = 0$ if $0 < \theta < \frac{\pi}{2}$, $t_0 < t < t_0 + h$.

THEOREM 5.3. If $g(x, t) \geq b > 0$ for $0 \leq x \leq x_1$, $t_1 < t < t_2$ ($x_1 > 0$, $t_2 > t_1 \geq 0$) then $\zeta_1(\theta, t)$ is continuous in (θ, t) for $t_1 < t < t_2$, $0 \leq \theta \leq \frac{\pi}{2}$.

The proof depends on two lemmas.

LEMMA 5.4. Let $r_0 = \zeta_1(\theta_0, t_0) > 0$ where $0 < \theta_0 < \frac{\pi}{2}$, $t_1 < t_0 < t_2$. Then there is a constant $C > 0$ such that for all τ positive and sufficiently small

$$(5.8) \quad \zeta_1(\theta_0, t_0 + \tau) \geq \zeta_1(\theta_0, t_0) - C\tau^{1/6}.$$

Proof. Set $x^0 = r_0 \cos \theta_0$, $y^0 = r_0 \sin \theta_0$. For any small $\rho \geq 0$ define

$$D_\rho = \{(x, y); 0 < x < x^0 - \rho, 0 < y < y^0 - \rho\},$$

$$\ell_\rho = \partial D_\rho \cap \{x > 0, y > 0\}.$$

Take any small $\delta > 0$ and let w_1 be the harmonic function in D_0 such that $w_1 = -\delta$ on ℓ_0 , $\partial w_1 / \partial x = 0$ on $\partial D_0 \cap \{x = 0\}$, and $\partial w_1 / \partial y = -g_0$ on $\partial D_0 \cap \{y = 0\}$ where

$$0 \leq g_0 \leq b, \quad g_0 = b \quad \text{in} \quad \left[\frac{1}{4} \bar{x}, \frac{3}{4} \bar{x} \right],$$

$$g = 0 \quad \text{if} \quad x < \frac{1}{8} \bar{x} \quad \text{or} \quad x > \frac{7}{8} \bar{x} \quad (\bar{x} = \min\{x^0, x_1\}),$$

and $g \in C^3[0, \infty)$. Then $w_1 \in C^2(\overline{D})$ and, by comparison,

$$(5.9) \quad -\delta + \frac{1}{C_0}(x - x_0)(y - y_0) \leq w_1(x, y) \leq -\delta + C_0(x - x_0)(y - y_0) \quad \text{in} \quad D_0$$

if C_0 is a sufficiently large positive constant. It follows that

$$(5.10) \quad w_1 < 0 \quad \text{in} \quad D_0 \setminus D_{A\delta}, \quad A = [2C_0 \max\{x^0, y^0\}]^{-1}.$$

Let $\zeta \in C^2(\overline{\Omega_R})$ ($R \gg 1$) be a function satisfying:

$$(5.11) \quad \begin{aligned} \zeta &= 0 \quad \text{in} \quad D_{A\delta}, \quad \zeta = 1 \quad \text{in} \quad \Omega_R \setminus D_0, \quad 0 \leq \zeta \leq 1, \\ \zeta_y(x, 0) &= 0, \quad \zeta_x(0, y) = 0, \\ |D\zeta| &\leq \frac{C_1}{\delta}, \quad |D^2\zeta| \leq \frac{C_1}{\delta^2} \quad \text{in} \quad \Omega_R \end{aligned}$$

where C_1 is a positive constant independent of δ . Extend w_1 into $\Omega_R \setminus D_0$ by $w_1 = -\delta$ and define

$$(5.12) \quad w = (1 - \zeta)w_1 - \zeta \quad \text{in} \quad \Omega_R.$$

Since w_1 is in $C^2(\overline{D}_0)$,

$$(5.13) \quad \|w\|_{C^2(\Omega_R)} \leq \frac{C_2}{\delta^2}.$$

We now define

$$\begin{aligned} \bar{u} &= \max \left\{ -1, w - \frac{C_2}{\delta^2}(t - t_0) \right\} \quad \text{in } \Omega_R \times [t_0, t_2], \\ \underline{\gamma} &= \begin{cases} c(\bar{u}) & \text{if } \bar{u} > -1 \\ -1 & \text{if } \bar{u} = -1. \end{cases} \end{aligned}$$

We wish to apply Lemma 3.1 to show that $u \geq \bar{u}$ in $\{\bar{u} < 0\}$.

If $\bar{u}(x, y, t) \geq 0$ then $w(x, y) \geq 0$ and, by (5.10)–(5.12), $(x, y) \in D_{A\delta}$, so that

$$\underline{\gamma}_t - \Delta \bar{u} \leq -\Delta \bar{u} = -\Delta w = 0.$$

If $-1 < \bar{u}(x, y, t) < 0$ then $\bar{u} = w - \frac{C_2}{\delta^2}(t - t_0)$ and hence

$$\underline{\gamma}_t - \Delta \bar{u} = u_t - \Delta \bar{u} = -\frac{C_2}{\delta^2} - \Delta w \leq 0$$

by (5.13). Observe next that $\partial \bar{u} / \partial n \leq 0$ on $\partial\{\bar{u} > -1\} \cap \{\bar{u} = -1\}$ where n is the normal pointing into $\{\bar{u} = -1\}$ and, therefore $\Delta \bar{u}$ is a positive measure on $\partial\{\bar{u} \geq -1\} \cap \{\bar{u} = -1\}$. It then easily follows that

$$\underline{\gamma}_t - \Delta \bar{u} \leq 0$$

in the distribution sense, in a neighborhood of $\partial\{\bar{u} \geq -1\} \cap \{\bar{u} = -1\}$. In conclusion, $\underline{\gamma}_t - \Delta \bar{u} \leq 0$ in $\Omega_R \times [t_0, t_2]$ in the distribution sense.

Since $u(\cdot, t_0) \geq 0$ in D_0 , $\gamma(\cdot, t_0) = 1$ and therefore $\gamma(\cdot, t_0) \geq \underline{\gamma}(\cdot, t_0)$ in D_0 . Also $\gamma(\cdot, t_0) \geq -1 = \underline{\gamma}(\cdot, t_0)$ in $\Omega_R \setminus D_0$ (since $\bar{u}(\cdot, t_0) = -1$ in $\Omega_R \setminus D_0$). Finally $\bar{u}_x = 0$ on $\{x = 0\}$ and $-u_y \leq g$ on $\{y = 0\}$. We can therefore apply Lemma 3.1 with $u^2 = \bar{u}$ and conclude that either $u \geq \bar{u}$ or $\gamma \geq \underline{\gamma}$; in particular, $u \geq 0$ if $\bar{u} \geq 0$.

From (5.9) we know that for any $\lambda > 0$ and $0 < \tau < t_2 - t_0$,

$$\begin{aligned} \bar{u}((r_0 - \lambda) \cos \theta_0, (r_0 - \lambda) \sin \theta_0, t_0 + \tau) &\geq w(x_0 - \lambda \cos \theta_0, y_0 - \lambda \sin \theta_0) - \frac{C_2}{\delta^2} \tau \\ &\geq -\delta + \frac{1}{C_0} \lambda^2 \sin \theta_0 \cos \theta_0 - \frac{C_2}{\delta^2} \tau > 0 \end{aligned}$$

if we take $\delta = \tau^{1/3}$, $\lambda = C_3 \tau^{1/6} = C_3 \delta^{1/2}$ where

$$C_3 = \left[\frac{1 + C_2}{\sin \theta_0 \cos \theta_0} C_0 \right]^{1/2}$$

(notice that $(x_0 - \lambda \cos \theta_0, y_0 - \lambda \sin \theta_0) \in D_{A\delta}$ so that $\zeta = 0$ at this point). We conclude that

$$u((r_0 - C_3 \tau^{1/6}) \cos \theta_0, (r_0 - C_3 \tau^{1/6}) \sin \theta_0, \tau) \geq 0$$

and the assertion (5.8) thus follows.

Remark 5.2. The constant C in (5.8) is uniform in (x^0, y^0, t_0) provided (x^0, y^0, t_0) remains in a compact subset of $\{x > 0, y > 0, t_1 < t < t_2\}$.

LEMMA 5.5. *Let $r_0 = \zeta_1(\theta_0, t_0) > 0$, $0 < \theta_0 < \frac{\pi}{2}$, $t_1 < t_0 < t_2$. Then there is a positive constant C such that*

$$(5.14) \quad \zeta_1(\theta_0, t_0 + \tau) \leq \zeta_1(\theta_0, t_0) + C \left(\log \frac{1}{\tau} \right)^{-1/6}.$$

Proof. By Lemma 5.4, if $t_0 - h \leq t \leq t_0$ (h small) then

$$u(x^0 + \delta, y^0 + \delta, t) < 0 \quad (\delta = Ch^{1/6}).$$

Hence

$$u(x, y^0 + \delta, t) \leq 0, \quad u(x^0 + \delta, y, t) \leq 0$$

for $x^0 + \delta \leq x < \infty$, $y^0 + \delta \leq y < \infty$, $t_0 - h \leq t \leq t_0$.

Denote by $\tilde{R}$ the interior of the subset in $t_0 - h \leq t \leq t_0$ where $x \geq x^0 + \delta$ or $y \geq y^0 + \delta$, and $u > -1$. Denote by $\partial_0 \tilde{R}$ the lateral boundary of $\tilde{R}$ on which $u = -1$ and by $\partial_1 \tilde{R}$ the lateral boundary of $\tilde{R}$ where either $x = x_0 + \delta$ or $y = y_0 + \delta$. Let U be the solution of

$$\begin{aligned} U_t &= \Delta U \quad \text{in } \tilde{R}, \\ U &= -1 \quad \text{on } \partial_0 \tilde{R}, \quad U = 0 \quad \text{on } \partial_1 \tilde{R}, \\ U &= 0 \quad \text{at } t = t_0 - h. \end{aligned}$$

Then $u \leq U$. Hence, by parabolic estimates on U ,

$$(5.15) \quad u(x^0 + 2\delta, y^0 + 2\delta, t_0) \leq U(x^0 + 2\delta, y^0 + 2\delta, t_0) \leq -Ce^{-\frac{c}{h}} \equiv -\tilde{\delta}$$

where C, c are positive constants.

Let R be a large constant such that $u = -1$ if $x^2 + y^2 > R^2$, $t \in [t_1, t_2]$. For each small $\rho \geq 0$ define

$$D_\rho = (0, R + \rho) \times (0, R + \rho) \setminus \{x \geq x^0 + \rho, y \geq y^0 + \rho\}.$$

Let w_1 be harmonic function in $D_{3\delta}$ satisfying:

$$w_{1,x} = 0 \quad \text{on} \quad \partial D_{3\delta} \cap \{x = 0\}, \quad -w_{1,y} = G \geq \sup_{t_1 \leq t \leq t_2} g(\cdot, t) \quad \text{on} \quad \partial D_{3\delta} \cap \{y = 0\},$$

and $w_1 = -\tilde{\delta}$ on the remaining part of $\partial D_{3\delta}$.

Using the conformal mapping $z \rightarrow z^{2/3}$ near the corner of $D_{3\delta}$ at $(x_0 + 3\delta, y_0 + 3\delta)$ we can deduce that

$$(5.16) \quad -\tilde{\delta} + \frac{d^{2/3}}{C_0} \leq w_1(x, y) \leq -\tilde{\delta} + C_0 d^{2/3} \quad \text{in} \quad D_{3\rho} \cap B_\eta(x_0 + 3\delta, y_0 + 3\delta)$$

where $B_\eta(x_1, y_1)$ is a disc with center (x_1, y_1) and radius η , d is the distance from (x, y) to $\partial D_{3\delta}$, and C_0, η are some positive constants.

We can use reflection to estimate w_1 near the other corners of $D_{3\delta}$ and thus deduce that there is a constant $A > 0$ such that

$$(5.17) \quad w_1 < -\frac{\tilde{\delta}}{2} \quad \text{in} \quad D_{3\delta} \setminus D_{3\delta - A\tilde{\delta}^{3/2}}.$$

and (since $\tilde{\delta} < \delta^3$ if $\delta \ll 1$) that

$$(5.18) \quad w_1 > 0 \quad \text{in} \quad D_{2\delta}.$$

From interior estimates for harmonic functions,

$$(5.19) \quad \|w_1\|_{C^k(D_{3\delta - \frac{1}{2}A\tilde{\delta}^{3/2}})} \leq \frac{C_1}{\tilde{\delta}^{\frac{3k}{2}}} \quad (k = 1, 2).$$

Let $\zeta \in C^3(\Omega_{R+1})$ be a function satisfying:

$$(5.20) \quad \begin{aligned} \zeta &= 0 \quad \text{in} \quad D_{3\delta - A\tilde{\delta}^{3/2}}, \quad \zeta = 1 \quad \text{in} \quad \Omega_{R+1} \setminus D_{3\delta - \frac{1}{2}A\tilde{\delta}^{3/2}}, \\ \zeta_x &= 0 \quad \text{on} \quad x = 0 \quad \text{and} \quad \zeta_y = 0 \quad \text{on} \quad y = 0, \\ 0 &\leq \zeta \leq 1, \quad |D\zeta| \leq \frac{C_2}{\tilde{\delta}^{3/2}}, \quad |D^2\zeta| \leq \frac{C}{\tilde{\delta}^3}. \end{aligned}$$

Extend w_1 into $\Omega_{R+1} \setminus D_{3\delta}$ by $w_1 = -\tilde{\delta}$ and define

$$(5.21) \quad w = (1 - \zeta)w_1 - \tilde{\delta}\zeta.$$

Since $w_1 \geq -\tilde{\delta}$, also $w \geq -\tilde{\delta}$. Further, by (5.19), (5.20)

$$(5.22) \quad |D^2 w|_{C^0(\Omega_{R+1})} \leq \frac{C_3}{\tilde{\delta}^3} .$$

We now choose

$$\bar{u} = w + \frac{C_3}{\tilde{\delta}^3}(t - t_0) \quad \text{in} \quad \Omega_R \times \left[t_0, t_0 + \frac{\tilde{\delta}^4}{2C_3} \right) ,$$

$$\bar{\gamma} = c(\bar{u}) .$$

Since $w > -\tilde{\delta}$ in Ω_R and $w \geq 0$ in $D_{2\delta}$ we have, by (5.15), that $\gamma(\cdot, t_0) \leq \bar{\gamma}(\cdot, t_0)$ in Ω_R (indeed $\bar{\gamma}(\cdot, t_0) = 1$ in $D_{2\delta}$ and $\bar{\gamma}(\cdot, t_0) > -1 = \gamma(\cdot, t_0)$ in $\Omega_R \setminus D_{2\delta}$). Next $\bar{u}_x = 0$ on $\{x = 0\}$ and $-\bar{u}_y = G > -u_y$ on $y = 0$. We finally show that $\bar{u}$ is a supersolution.

If $\bar{u}(x, y, t) > 0$, $t \in [t_0, t_0 + \tilde{\delta}^4/(2C_3))$ then $w \geq -\frac{\tilde{\delta}}{2}$ and, by (5.17), $(x, y) \in D_{3\delta - A\tilde{\delta}^{3/2}}$. Consequently

$$\bar{\gamma}_t - \Delta \bar{u} \geq -\Delta \bar{u} = 0 .$$

If $\bar{u}(x, y, t) \leq 0$ then, since $\bar{u} \geq w > -\tilde{\delta}$,

$$\bar{\gamma}_t - \Delta \bar{u} = \bar{u}_t - \Delta \bar{u} = \frac{C_3}{\tilde{\delta}^3} - \Delta w > 0$$

by (5.22). This completes the proof that $\bar{u}$ is a supersolution.

We can then apply Lemma 3.1 to conclude that $u \leq \bar{u}$ in $\{\bar{u} \leq 0\}$. Since for $x > x^0 + 3\delta$, $y > y^0 + 3\delta$ and $\tau = \tilde{\delta}^4/(2C_3)$,

$$\bar{u}(x, y, t_0 + \tau) = w(x, y) + \frac{\tilde{\delta}}{2} = -\tilde{\delta} + \frac{\tilde{\delta}}{2} < 0 ,$$

it follows that $u(x, y, t_0 + \tau) < 0$. This implies that

$$\zeta_1(\theta_0, t_0 + \tau) \leq \zeta_1(\theta_0) + \frac{3\delta}{C_4}$$

where $C_4 = \min\{\cos \theta_0, \sin \theta_0\}$. Choosing $\tilde{\delta}$ as in (5.15) and recalling that $\delta = Ch^{1/6}$, we easily deduce the relation

$$\delta = C_5 \left(\log \frac{C_6}{\tau} \right)^{-1/6} ,$$

and (5.14) follows.

From Lemmas 5.4, 5.5 we get

$$(5.23) \quad |\zeta_1(\theta_0, t_0 + \tau) - \zeta_1(\theta_0, t_0)| \leq C \left(\log \frac{1}{\tau} \right)^{-1/6} ;$$

the constant C is independent of θ_0, t_0 so long as θ_0, t_0 vary in a compact subset of $\left\{ 0 < \theta_0 < \frac{\pi}{2}, t_1 < t_0 < t_2 \right\}$. The constant C degenerates as $\theta_0 \rightarrow 0$ or as $\theta_0 \rightarrow \frac{\pi}{2}$. However the proofs of the lemmas can be modified to establish the continuity of $\zeta_1(\theta, t)$ in t also at $\theta = 0, \pi$.

Since the function $\theta \rightarrow \zeta_1(\theta, t)$ is Lipschitz continuous in θ , uniformly in t , the proof of Theorem 5.3 is complete.

Remark 5.3. If $g(x, t) \geq b > 0$ for $0 \leq t < t_2$ then the continuity of $\zeta_1(\theta, t)$ at $t = 0$ can be established by slight modifications in the previous proofs.

Remark 5.4. The 1-d analog of the problem studied in this paper is:

$$(5.24) \quad \begin{aligned} u_{xx} &= 0, \quad u > 0 \quad \text{if} \quad 0 < x < \zeta_1(t), \quad t > 0, \\ u_t &= u_{xx}, \quad -1 < u < 0 \quad \text{if} \quad \zeta_1(t) < x < \zeta_2(t), \quad t > 0, \\ u &= -1 \quad \text{if} \quad x > \zeta_2(t), \quad t > 0, \\ u(\zeta_1(t), t) &= 0, \quad u(\zeta_2(t), t) = -1 \quad \text{if} \quad t > 0, \\ -u_x(0, t) &= g(t) \quad \text{if} \quad t > 0 \quad (g \geq 0), \\ u(x, 0) &= u_0(x) \quad \text{if} \quad 0 < x < \infty \quad (\partial u_0 / \partial x \leq 0). \end{aligned}$$

All the results established in this paper are clearly valid for the problem (5.24). We shall now show that if $g(t) > 0$ for $0 \leq t < T$, $g(T) = 0$ then, $\zeta_1(t)$ may become discontinuous at $t = T$ even when $g(t)$ is continuous at $t = T$. (Thus the assumption $g \geq b > 0$ made in Theorem 5.3 cannot be dropped, in general.) Let

$$W(x, t, \delta) = \frac{1}{2\delta\sqrt{t}} e^{-x^2/(4t)} - \frac{1}{\delta} - 1 \quad (\delta > 0, \delta \ll 1).$$

Then

$$W = 0 \quad \text{on} \quad x = s_1(t) = 2t \left[\log \frac{1}{4t(1+\delta)^2} \right]^{1/2}, \quad 0 < t < \frac{1}{4(1+\delta)^2},$$

$$W = -1 \quad \text{on} \quad x = s_2(t) = \left[2t \log \frac{1}{4t} \right]^{1/2}, \quad 0 < t < \frac{1}{4}.$$

Set

$$\widehat{G}(\delta, t) = -\frac{\partial W}{\partial x} \Big|_{x=s_1(t)} = \frac{1+\delta}{\delta\sqrt{2t}} \left[\log \frac{1}{4t(1+\delta)^2} \right]^{1/2}, \quad 0 < t < \frac{1}{4(1+\delta)^2}.$$

Define, for $0 < t < 1/[4(1+\delta)^2]$,

$$\widehat{u}(x, t) = \begin{cases} -\widehat{G}(\delta, t)x & \text{if } x < s_1(t) \\ W(x, t, \delta) & \text{if } s_1(t) < x < s_2(t) \\ -1 & \text{if } x > s_2(t), \end{cases}$$

and set

$$\begin{aligned} \underline{u}(x, \tau) &= \widehat{u} \left(x + x_0, \frac{1}{4(1+\delta)^2} + \tau \right), \\ G(\delta, \tau) &= \widehat{G} \left(\delta, \frac{1}{4(1+\delta)^2} + \tau \right) \end{aligned}$$

where

$$x_0 = -\zeta_1(T - \delta) + s_2 \left(\frac{1}{4(1+\delta)^2} - \delta \right).$$

Notice that $G(\delta, \tau) > 0$ if $-\delta < \tau < 0$, $G(\delta, 0) = 0$. The function $\underline{u}$ is a subsolution for the differential equations in (5.24) (we use here the fact that $\dot{s}_2(t) < 0$). Hence if

$$(5.25) \quad g(T + \tau) \geq G(\delta, \tau), \quad -\delta < \tau < 0$$

and

$$(5.26) \quad u(x, T - \delta) > \underline{u}(x, -\delta)$$

then, by comparison,

$$(5.27) \quad u(x, T + \tau) \geq \underline{u}(x, \tau) \quad (-\delta < \tau < 0).$$

To prove (5.26) we note that $\underline{u}(x, -\delta) = -1$ if $x = \zeta_1(T - \delta)$ so that (5.26) is valid for $x \geq \zeta_1(T - \delta)$. Next, if

$$(5.28) \quad \zeta_1(T - \delta) > 2C\delta,$$

where C is large enough constant, then $\underline{u}(0, -\delta) < 0$ and therefore (5.26) holds also for $0 \leq x < \zeta_1(T - \delta)$. We conclude that if (5.25) and (5.28) hold then (5.27) is valid and, in particular,

$$\underline{u}(\zeta_1(T), 0) < 0 \quad \text{if} \quad \zeta_1(T) > 0,$$

that is,

$$\zeta_1(T) + x_0 > s_1 \left(\frac{1}{4(1+\delta)^2} \right) = 0 \quad \text{if} \quad \zeta_1(T) > 0 .$$

Recalling the definition of x_0 we conclude that

$$\zeta_1(T) \geq \zeta_1(T - \delta) - C_1 \delta \quad \text{if} \quad \zeta_1(T) > 0 .$$

where C_1 is a positive constant independent of δ and C . Choosing C so large that $C > C_1$ and recalling (5.28), it follows that

$$(5.29) \quad \zeta_1(T) > C\delta \quad \text{if} \quad \zeta_1(T) > 0 .$$

Consider now a solution of (5.24) with $g(t) > 0$ for $0 < t < T - \delta$. We can choose δ so small that (5.28) is satisfied and then extend $g(t)$ to $[T - \delta, T]$ so that it satisfies (5.25) and $g(T) = 0$. We can then apply (5.29) for each T' smaller than T ; since $\zeta_1(T') > 0$ we then have $\zeta_1(T') \geq C\delta$. Since however $\zeta_1(T) = 0$, $\zeta_1(t)$ is discontinuous at $t = T$. Note that $g(t)$ decreases linearly to zero as $t \uparrow T$. Perhaps a faster decrease will not produce a discontinuity for $\zeta_1(t)$ at $t = T$.

REFERENCES

- [1] H.W. ALT AND S. LUCKHAUS, *Quasilinear elliptic-parabolic differential equations*, Math. Zeit., 183 (1983), 311–341.
- [2] H.W. ALT, S. LUCKHAUS AND A. VISINTIN, *On nonstationary flow through porous media*, Ann. Mat. Pure Appl., 136 (1984), 303–316.
- [3] J. BEAR, *Dynamics of Fluids in Porous Media*, American Elsevier, New York, 1972.
- [4] L.A. CAFFARELLI, *The regularity of free boundaries in higher dimension*, Acta Math., 139 (1977), 155–184.
- [5] L.A. CAFFARELLI AND A. FRIEDMAN, *Continuity of the temperature in the Stefan problem*, Indiana Univ. Math. J., 28 (1979), 53–70.
- [6] J.R. CANNON AND F.A. MOHAMED, *A multifree boundary problem arising in the theory of liquid flow in a porous medium*, Boll. U.M.I. (7) 3–B (1989), 69–93.
- [7] E. DiBENEDETTO AND A. FRIEDMAN, *Periodic behavior for the evolutionary dam problem and related free boundary problems*, Comm. P.D.E., 11 (1986), 1297–1377.
- [8] E. DiBENEDETTO AND R. GARIEPY, *Local behavior of solutions of an elliptic-parabolic equation*, Arch. Rat. Mech. Anal., 97 (1987), 1–18.
- [9] A. FRIEDMAN, *Variational Principles and Free-Boundary Problems*, John Wiley-Interscience, New York, 1982.
- [10] U. HORNUNG, *A parabolic-elliptic variational inequality*, Manuscripta Math., 39 (1982), 155–172.
- [11] J. HULSHOF, *An elliptic-parabolic free boundary problem: continuity of the interface*, Proc. Royal Soc. Edinburgh, 106A (1987), 327–339.
- [12] J. HULSHOF, *Bounded weak solutions of an elliptic-parabolic Neumann problem*, Trans. Amer. Math. Soc., 303 (1987), 211–227.

- [13] J. HULSHOF, *Spherically symmetric solutions of an elliptic-parabolic Neumann problem*, Rocky Mountain J. of Math., 21 (1991), 671–681.
- [14] J. HULSHOF AND L.A. PELETIER, *An elliptic-parabolic free boundary problem*, Nonlinear Anal., 10 (1986), 1327–1346.
- [15] J. HULSHOF AND N. WOLANSKI, *Monotone flows in N -dimensional partially saturated porous media; Lipschitz-continuity of the interface*, 102 (1988), 287–305,.
- [16] O.A. LADYZENSKAJA, V.A. SOLONNIKOV AND N.N. URAL'CEVA, *Linear and Quasilinear Equations of Parabolic Type*, Transl. Math. Monographs, Amer. Math. Soc., Providence, R.I., 1968.
- [17] J. SIMON, *Compact sets in the space $L^p(0, T; B)$* , Ann. Mat. Pure Appl., 146 (1987), 65–96..
- [18] C.J. VAN DUYN, *Nonstationary filtration in partially saturated porous media: Continuity of the free boundary*, Arch. Rat. Mech. Anal., 78 (1982), 261–265.
- [19] C.J. VAN DUYN AND L.A. PELETIER, *Nonstationary filtration in partially saturated porous media*, Arch. Rat. Mech. Anal., 78 (1982), 173–198.

#	Author/s	Title
1021	IMA Summer Program for Graduate Students,	Mathematical Modeling
1022	Wayne Barrett, Charles R. Johnson, and Pablo Tarazaga,	The real positive definite completion problem for a simple cycle
1023	Charles A. McCarthy,	Fourth order accuracy for a cubic spline collocation method
1024	Martin Hanke, James Nagy, and Robert Plemmons,	Preconditioned iterative regularization for ill-posed problems
1025	John R. Gilbert, Esmond G. Ng, and Barry W. Peyton,	An efficient algorithm to compute row and column counts for sparse Cholesky factorization
1026	Xinfu Chen,	Existence and regularity of solutions of a nonlinear nonuniformly elliptic system arising from a thermistor problem
1027	Xinfu Chen and Weiqing Xie,	Discontinuous solutions of steady state, viscous compressible Navier-Stokes equations
1028	E.G. Kalnins, Willard Miller, Jr., and Sanchita Mukherjee,	Models of q -algebra representations: Matrix elements of the q -oscillator algebra
1029	W. Miller, Jr. and Lee A. Rubel,	Functional separation of variables for Laplace equations in two dimensions
1030	I. Gohberg and I. Koltracht,	Structured condition numbers for linear matrix structures
1031	Xinfu Chen,	Hele-Shaw problem and area preserved curve shortening motion
1032	Zhangxin Chen and Jim Douglas, Jr.	Modelling of compositional flow in naturally fractured reservoirs
1033	Harald K. Wimmer,	On the existence of a least and negative-semidefinite solution of the discrete-time algebraic Riccati equation
1034	Harald K. Wimmer,	Monotonicity and parametrization results for continuous-time algebraic Riccati equations and Riccati inequalities
1035	Bart De Moor, Peter Van Overschee, and Geert Schelfhout,	H_2 model reduction for SISO systems
1036	Bart De Moor,	Structured total least squares and L_2 approximation problems
1037	Chjan Lim,	Nonexistence of Lyapunov functions and the instability of the Von Karman vortex streets
1038	David C. Dobson and Fadil Santosa,	Resolution and stability analysis of an inverse problem in electrical impedance tomography – dependence on the input current patterns
1039	C.N. Dawson, C.J. van Duijn, and M.F. Wheeler,	Characteristic-Galerkin methods for contaminant transport with non-equilibrium adsorption kinetics
1040	Bing-Yu Zhang,	Analyticity of solutions of the generalized Korteweg-de Vries equation with respect to their initial values
1041	Neerchal K. Nagaraj and Wayne A. Fuller,	Least squares estimation of the linear model with autoregressive errors
1042	H.J. Sussmann & W. Liu,	A characterization of continuous dependence of trajectories with respect to the input for control-affine systems
1043	Karen Rudie & W. Murray Wonham,	Protocol verification using discrete-event systems
1044	Rohan Abeyaratne & James K. Knowles,	Nucleation, kinetics and admissibility criteria for propagating phase boundaries
1045	Gang Bao & William W. Symes,	Computation of pseudo-differential operators
1046	Srdjan Stojanovic,	Nonsmooth analysis and shape optimization in flow problem
1047	Miroslav Tuma,	Row ordering in sparse QR decomposition
1048	Onur Tokur & Hitay Özbay,	On the computation of suboptimal H^∞ controllers for unstable infinite dimensional systems
1049	Hitay Özbay,	H^∞ optimal controller design for a class of distributed parameter systems
1050	J.E. Dunn & Roger Fosdick,	The Weierstrass condition for a special class of elastic materials
1051	Bei Hu & Jianhua Zhang,	A free boundary problem arising in the modeling of internal oxidation of binary alloys
1052	Eduard Feireisl & Enrique Zuazua,	Global attractors for semilinear wave equations with locally distributed nonlinear damping and critical exponent
1053	I-Heng McComb & Chjan C. Lim,	Stability of equilibria for a class of time-reversible, $D_n \times O(2)$ -symmetric homogeneous vector fields
1054	Ruben D. Spies,	A state-space approach to a one-dimensional mathematical model for the dynamics of phase transitions in pseudoelastic materials
1055	H.S. Dumas, F. Golse, and P. Lochak,	Multiphase averaging for generalized flows on manifolds
1056	Bei Hu & Hong-Ming Yin,	Global solutions and quenching to a class of quasilinear parabolic equations
1057	Zhangxin Chen,	Projection finite element methods for semiconductor device equations
1058	Peter Guttorp,	Statistical analysis of biological monitoring data
1059	Wensheng Liu & Héctor J. Sussmann,	Abnormal sub-Riemannian minimizers

- 1060 **Chjan C. Lim**, A combinatorial perturbation method and Arnold's whiskered Tori in vortex dynamics
- 1061 **Yong Liu**, Axially symmetric jet flows arising from high speed fiber coating
- 1062 **Li Qiu & Tongwen Chen**, $\mathcal{H}_2$ and $\mathcal{H}_\infty$ designs of multirate sampled-data systems
- 1063 **Eduardo Casas & Jiongmin Yong**, Maximum principle for state-constrained optimal control problems covered by quasilinear elliptic equations
- 1064 **Suzanne M. Lenhart & Jiongmin Yong**, Optimal control for degenerate parabolic equations with logistic growth
- 1065 **Suzanne Lenhart**, Optimal control of a convective-diffusive fluid problem
- 1066 **Enrique Zuazua**, Weakly nonlinear large time behavior in scalar convection-diffusion equations
- 1067 **Caroline Fabre, Jean-Pierre Puel & Enrike Zuazua**, Approximate controllability of the semilinear heat equation
- 1068 **M. Escobedo, J.L. Vazquez & Enrike Zuazua**, Entropy solutions for diffusion-convection equations with partial diffusivity
- 1069 **M. Escobedo, J.L. Vazquez & Enrike Zuazua**, A diffusion-convection equation in several space dimensions
- 1070 **F. Fagnani & J.C. Willems**, Symmetries of differential systems
- 1071 **Zhangxin Chen, Bernardo Cockburn, Joseph W. Jerome & Chi-Wang Shu**, Mixed-RKDG finite element methods for the 2-D hydrodynamic model for semiconductor device simulation
- 1072 **M.E. Bradley & Suzanne Lenhart**, Bilinear optimal control of a Kirchhoff plate
- 1073 **Héctor J. Sussmann**, A cornucopia of abnormal subriemannian minimizers. Part I: The four-dimensional case
- 1074 **Marek Rakowski**, Transfer function approach to disturbance decoupling problem
- 1075 **Yuncheng You**, Optimal control of Ginzburg-Landau equation for superconductivity
- 1076 **Yuncheng You**, Global dynamics of dissipative modified Korteweg-de Vries equations
- 1077 **Mario Taboada & Yuncheng You**, Nonuniformly attracting inertial manifolds and stabilization of beam equations with structural and Balakrishnan-Taylor damping
- 1078 **Michael Böhm & Mario Taboada**, Global existence and regularity of solutions of the nonlinear string equation
- 1079 **Zhangxin Chen**, BDM mixed methods for a nonlinear elliptic problem
- 1080 **J.J.L. Velázquez**, On the dynamics of a closed thermosyphon
- 1081 **Frédéric Bonnans & Eduardo Casas**, Some stability concepts and their applications in optimal control problems
- 1082 **Hong-Ming Yin**, $L^{2,\mu}(Q)$ -estimates for parabolic equations and applications
- 1083 **David L. Russell & Bing-Yu Zhang**, Smoothing and decay properties of solutions of the Korteweg-de Vries equation on a periodic domain with point dissipation
- 1084 **J.E. Dunn & K.R. Rajagopal**, Fluids of differential type: Critical review and thermodynamic analysis
- 1085 **Mary Elizabeth Bradley & Mary Ann Horn**, Global stabilization of the von Kármán plate with boundary feedback acting via bending moments only
- 1086 **Mary Ann Horn & Irena Lasiecka**, Global stabilization of a dynamic von Kármán plate with nonlinear boundary feedback
- 1087 **Vilmos Komornik**, Decay estimates for a petrovski system with a nonlinear distributed feedback
- 1088 **Jesse L. Barlow**, Perturbation results for nearly uncoupled Markov chains with applications to iterative methods
- 1089 **Jong-Shenq Guo**, Large time behavior of solutions of a fast diffusion equation with source
- 1090 **Tongwen Chen & Li Qiu**, $\mathcal{H}_\infty$ design of general multirate sampled-data control systems
- 1091 **Satyanad Kichenassamy & Walter Littman**, Blow-up surfaces for nonlinear wave equations, I
- 1092 **Nahum Shimkin**, Asymptotically efficient adaptive strategies in repeated games, Part I: certainty equivalence strategies
- 1093 **Caroline Fabre, Jean-Pierre Puel & Enrique Zuazua**, On the density of the range of the semigroup for semilinear heat equations
- 1094 **Robert F. Stengel, Laura R. Ray & Christopher I. Marrison**, Probabilistic evaluation of control system robustness
- 1095 **H.O. Fattorini & S.S. Sritharan**, Optimal chattering controls for viscous flow
- 1096 **Kathryn E. Lenz**, Properties of certain optimal weighted sensitivity and weighted mixed sensitivity designs
- 1097 **Gang Bao & David C. Dobson**, Second harmonic generation in nonlinear optical films
- 1098 **Avner Friedman & Chaocheng Huang**, Diffusion in network
- 1099 **Xinfu Chen, Avner Friedman & Tsuyoshi Kimura**, Nonstationary filtration in partially saturated porous media
- 1100 **Walter Littman & Baisheng Yan**, Rellich type decay theorems for equations $P(D)u = f$ with f having support in a cylinder
- 1101 **Satyanad Kichenassamy & Walter Littman**, Blow-up surfaces for nonlinear wave equations, II
- 1102 **Nahum Shimkin**, Extremal large deviations in controlled I.I.D. processes with applications to hypothesis testing
- 1103 **A. Narain**, Interfacial shear modeling and flow predictions for internal flows of pure vapor experiencing film condensation
- 1104 **Andrew Teal & Laurent Praly**, Global stabilizability and observability imply semi-global stabilizability by output feedback