**STABILITY OF THE FRACTIONAL STEP Θ -SCHEME
FOR THE NONSTATIONARY NAVIER-STOKES EQUATIONS**

By

Petr Klouček

and

Franz S. Rys

IMA Preprint Series # 1162

July 1993

STABILITY OF THE FRACTIONAL STEP Θ -SCHEME FOR THE NONSTATIONARY NAVIER-STOKES EQUATIONS

PETR KLOUČEK AND FRANZ S. RYS

University of Minnesota and ETH Zürich

ABSTRACT. We present stability and convergence analysis of the fractional step Θ -scheme for the Navier–Stokes equations coupled with the of the finite element approximations. The solutions of the underlying Stokes and Burgers problem are described in detail. The asymptotic solution obtained by the numerical stabilization is compared with the analytical solution of the Lid-driven cavity problem.

INTRODUCTION

We present stability analysis of the nonstationary Navier-Stokes equations fully discretized by the finite element method and fractional step Θ -scheme.

The main advantage of this approach, developed by R.Glowinski in [G II], consists in the separation of the two underlying problems: solution of the nonlinear parabolic equation and the imposed incompressibility constrain. The method consists in the subsequent solution of the Generalized Stokes problem (linear parabolic equation with the incompressibility condition), of the solution of the Regularized Burgers problem (nonlinear parabolic equation), and, of the correction of the incompressibility condition by solving again the Generalized Stokes problem.

For the sake of completeness we have included a detailed description of the numerical solution of these two underlying problems. The central idea behind the use of the conjugate gradient (CG) solution of the Stokes and Burgers problems is to reduce the update procedure to the solution of the Poisson equation. The implementation can be highly optimized since we deal with just one type of the linear system. On the other hand the use of the CG method to solve the Regularized Burgers problem limits the range of the characteristic scales of the numerical flow.

The paper contains also example comparing the numerical asymptotic solution with the analytical solution of the simple Lid-driven cavity flow. The cavity computations suggest that the method works up to $Re = 3000$ (with the Taylor-Hood velocity/pressure approximation). We observed production of a considerable amount

1991 *Mathematics Subject Classification.* 65N10,65N30,76H05,35K55.

Key words and phrases. Fractional step Θ -scheme, Nonstationary incompressible Navier-Stokes equations, Stokes problem, Non-linear Dirichlet problem.

Financial support by the Swiss National Science Foundation (SNF Contract No. 70TP-029733) and by the “Kommission zur Förderung wissenschaftlicher Forschung” (KWF Contracts No. 1930.1 & 1931.1) is gratefully acknowledged.

of oscillations in the kinetic energy prohibiting the convergence of the CG method for the Regularized Burgers problem above this range.

The paper is organized as follows:

- In section 1 we recall the known theoretical results about the Navier-Stokes equations.
- In Section 2 we describe the Fractional step Θ -time stepping scheme. We also describe the solution of the corresponding Generalized Stokes and Regularized Burgers problems.
- In Section 3 we recall the results about the conforming and nonconforming finite element approximations of the Navier–Stokes equations .
- Section 4 contains the proof of the stability of the fractional step Θ -scheme and the convergence of the finite element approximations.
- In Section 5 we recall the analytically solvable Lid-driven cavity problem to discuss some of the properties of the method.

The derivation of the stability conditions leading to the proof of the Theorem is rather long. The reason is that we cannot use all of the properties of the nonlinear term because of the form of the scheme.

1. NAVIER–STOKES EQUATIONS

The flow of a viscous incompressible fluid is frequently described by the following initial–boundary value problem:

$$\begin{aligned}
 \vec{u}_t - \nu \Delta \vec{u} + (\vec{u} \cdot \nabla) \vec{u} + \nabla p &= \vec{f}, & \text{in } \Omega \times (0, T), \\
 \nabla \cdot \vec{u} &= 0, & \text{in } \Omega \times (0, T), \\
 \vec{u} &= \vec{g}, & \text{on } \partial\Omega, \\
 u(0) &= \vec{u}_0, & \text{in } \Omega,
 \end{aligned}
 \tag{NS}$$

where $\vec{u} \equiv \vec{u}(x, t)$, $p \equiv p(x, t)$ are the unknown velocity and pressure, respectively, and Ω is a bounded region in $\mathbb{R}^n$, $n = 2, 3$, with a sufficiently regular boundary $\partial\Omega$. Given data of the initial–boundary value problem are the viscosity, ν , the volume force, $\vec{f} \equiv \vec{f}(x, t)$, the boundary velocity, $\vec{g} = \vec{g}(x, t)$, and the initial velocity, $\vec{u}_0 \equiv \vec{u}_0(x)$. The density of the fluid is normalized. We assume that $\int_{\Omega} \vec{g} \cdot \vec{n} \, dx = 0$, $\vec{n}$ being the unit outward normal on $\partial\Omega$.

Functional setting. The following subsets of the space $[H^1(\Omega)]^n$ will be frequently used

$$\begin{aligned}
 H_g(\Omega) &= \{\vec{v} \in [H^1(\Omega)]^n \mid \vec{v}|_{\partial\Omega} = \vec{g}\}, \\
 V_g(\Omega) &= \{\vec{v} \in [H^1(\Omega)]^n \mid \nabla \cdot \vec{v} = 0, \vec{v}|_{\partial\Omega} = \vec{g}\}.
 \end{aligned}$$

We also use the spaces $H_0(\Omega)$ and $V_0(\Omega)$, which are defined as above, provided the functions have zero trace on the boundary $\partial\Omega$. Sometimes we refer to these spaces as either H or V . Also, $[H_0^1(\Omega)]^n$ is endowed with the product structure and consequently the spaces H, V are equipped with a scalar product and the Hilbert norm given by $((\vec{u}, \vec{v})) \equiv \sum_{i=1}^n \int_{\Omega} \nabla u_i \cdot \nabla v_i \, dx$ and by $\|\vec{u}\| = \sqrt{((\vec{u}, \vec{u}))}$, respectively. All

the considered spaces are Hilbert spaces for this norm. Consequently, the $[L^2(\Omega)]^n$ scalar product and the corresponding norm are given by $(\vec{u}, \vec{u}) = \sum_{i=1}^n \int_{\Omega} u_i v_i dx$ and by $|\vec{u}| = \sqrt{(\vec{u}, \vec{u})}$. For the usual Bochner spaces of the time dependent functions with values in some Banach space X we use the notation:

$$L^p(0, T; X) = \{u \mid u : (0, T) \rightarrow X, \text{ measurable; } \int_0^T \|u(\tau)\|_X^p d\tau < +\infty\},$$

with the standard modification for $p = \infty$. Moreover, we denote

$$b(\vec{u}, \vec{v}, \vec{w}) \equiv ((\vec{u} \cdot \nabla) \vec{v}, \vec{w}).$$

The form $b(., ., .)$ is defined and trilinear continuous on $[H^1(\Omega)]^n \times [H^1(\Omega)]^n \times [H^1(\Omega)]^n$.

In this notation we define the weak solution of the problem (NS) as follows:

Definition. Given $\vec{u}_0 \in V_0$ and $\vec{f} \in L^2(0, +\infty; [L^2(\Omega)]^n)$ the weak solution of the initial boundary value the problem (NS), defined on some interval $[0, T]$, $T > 0$, is a function $\vec{u}$ such that

$$\begin{aligned} \vec{u} &\in L^\infty(0, T; V_g), \\ (\text{WNS}) \quad &(\vec{u}_t, \vec{\varphi}) + \nu((\vec{u}, \vec{\varphi})) + b(\vec{u}, \vec{u}, \vec{\varphi}) = (\vec{f}, \vec{\varphi}), \quad \forall \vec{\varphi} \in V_0, \\ &\vec{u}(0) = \vec{u}_0. \end{aligned}$$

We note that there exists unique weak solution to the problem (WNS) on some time interval $[0, T]$. Moreover, it is known that there exists unique strong solution $u \in L^\infty(0, T; V_g) \cap L^2(0, T; V_0 \cap [H^2(\Omega)]^n)$, $u_t \in L^2(0, T; [L^2(\Omega)]^n)$, for all $T > 0$, in two space dimensions, which is uniformly bounded in the H^1 norm, i.e. $\|\vec{u}(t)\| \leq C < +\infty$, where $C \equiv C(\nu, \vec{g}, \vec{f}, \Omega)$. In addition, the corresponding pressure p satisfying (NS) is in $L^2(0, T; H^1(\Omega))$. Under somewhat stronger assumptions $\vec{u}_0 \in [H^2(\Omega)]^n$ and $\vec{f}_t \in L^2(0, +\infty; [L^2(\Omega)]^n)$ we have $\vec{u} \in L^\infty(0, T; [H^1(\Omega)]^n)$, $u_t \in L^\infty(0, T; [L^2(\Omega)]^n) \cap L^2(0, T; [H^1(\Omega)]^n)$ and $p \in L^\infty(0, T; H^1(\Omega))$.

In the case of the three space dimensions the strong solution is known to exist and to be unique on some interval $[0, T_1]$, where $T_1 = C/(1 + \|\vec{u}_0\|^2)^2$. The weak solution, which exists for all $T > 0$, coincides with the strong solution at least on the interval $[0, T_1]$ but its uniqueness is not proved. These results can be found e.g. in Heywood[H], Ladyzhenskaya[L], Temam[T I].

To avoid the difficulties of the discretization of the divergence-free space V it is convenient to work with the following formulation (in the form of the saddle point problem) which is equivalent to (WNS) for sufficiently regular functions:

Find $\{\vec{u}, p\} \in L^\infty(0, T; H_g) \times L^2(\Omega)$ such that

$$\begin{aligned} (\text{SNS}) \quad &(\vec{u}_t, \vec{\varphi}) + \nu((\vec{u}, \vec{\varphi})) + b(\vec{u}, \vec{u}, \vec{\varphi}) - (p, \nabla \cdot \vec{\varphi}) = (\vec{f}, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0, \\ &(\nabla \cdot \vec{u}, q) = 0, \quad \forall q \in L^2(\Omega), \\ &\vec{u}(0) = \vec{u}_0. \end{aligned}$$

The pressure p is unique in $L^2(\Omega)/\mathbb{R}$ in both definitions (WNS) and (SNS).

2. FRACTIONAL STEP METHOD AND NAVIER-STOKES EQUATIONS

Fractional step (or splitting up) methods can be applied to the Navier–Stokes equations in many different ways. We will use the version advocated in R. Rannacher [R], where some more details and comparisons with other time stepping schemes (Euler, Crank–Nicolson, etc.) can be found.

In order to treat the main difficulties of the problem, the incompressibility and the nonlinearity, separately, we define Generalized Stokes problem and Regularized Burgers problem by splitting up the sum $\nu((\vec{u}, \vec{\varphi})) + b(\vec{u}, \vec{u}, \vec{\varphi})$ into two parts, $\alpha\nu((\vec{u}, \vec{\varphi}))$ and $(1-\alpha)\nu((\vec{u}, \vec{\varphi})) + b(\vec{u}, \vec{u}, \vec{\varphi})$, considered on the space V and the space H , respectively.

Denoting $\vec{u}^n \equiv \vec{u}(x, t_n)$, $\vec{f}^n \equiv \vec{f}(x, t_n)$, simple version of the fractional step method applied to the Navier–Stokes equations reads:

Generalized Stokes problem. Find $\{\vec{u}^{n+\frac{1}{2}}, p^{n+\frac{1}{2}}\} \in \{H_g \times L^2(\Omega)\}$ such that

$$\frac{1}{\Delta t}(\vec{u}^{n+\frac{1}{2}}, \vec{\varphi}) + \alpha\nu((\vec{u}^{n+\frac{1}{2}}, \vec{\varphi})) - (p^{n+\frac{1}{2}}, \nabla \cdot \vec{\varphi}) =$$

$$(FRNS1) \quad \frac{1}{\Delta t}(\vec{u}^n, \vec{\varphi}) - (1-\alpha)\nu((\vec{u}^n, \vec{\varphi})) - b(\vec{u}^n, \vec{u}^n, \vec{\varphi}) + (\vec{f}^n, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0,$$

$$(\nabla \cdot \vec{u}^{n+\frac{1}{2}}, q) = 0, \quad \forall q \in L^2(\Omega).$$

Regularized Burgers problem. Find $\vec{u}^{n+1} \in H_g$ such that

$$(FRNS2) \quad \frac{1}{\Delta t}(\vec{u}^{n+1}, \vec{\varphi}) + (1-\alpha)\nu((\vec{u}^{n+1}, \vec{\varphi})) + b(\vec{u}^{n+1}, \vec{u}^{n+1}, \vec{\varphi}) =$$

$$\frac{1}{\Delta t}(\vec{u}^{n+\frac{1}{2}}, \vec{\varphi}) - \alpha\nu((\vec{u}^{n+\frac{1}{2}}, \vec{\varphi})) + (p^{n+\frac{1}{2}}, \nabla \cdot \vec{\varphi}) + (\vec{f}^{n+\frac{1}{2}}, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0.$$

2.1 Fractional step Θ -scheme for the Navier–Stokes equations.

Further refinement of the splitting up approach consists in introduction of further correction of the solution with respect to the incompressibility condition.

We divide the time interval $[t_n, t_{n+1}]$ of the length Δt into the three subintervals $[t_n, t_{n+\Theta})$, $[t_{n+\Theta}, t_{n+(1-\Theta)})$ and $[t_{n+(1-\Theta)}, t_{n+1}]$ of the lengths $\Theta\Delta t$, $(1-2\Theta)\Delta t$ and $\Theta\Delta t$, respectively. Using this partition, we obtain $\vec{u}^{n+1}$ from $\vec{u}^n$ by first solving (FRNS1) for $\vec{u}^{n+\Theta}$ then by solving (FRNS2) for $\vec{u}^{n+(1-\Theta)}$ with $\vec{u}^{n+\frac{1}{2}}$ replaced by $\vec{u}^{n+\Theta}$ and finally by solving (FRNS1) for $\vec{u}^{n+1}$ with $\vec{u}^n$ replaced by $\vec{u}^{n+(1-\Theta)}$.

Different implementation of this method can be found in Glowinski [G II].

Let $\Theta_b \equiv 1 - 2\Theta$ and $\beta \equiv 1 - \alpha$. Assuming $\Theta \in (0, \frac{1}{2})$ and $\alpha \in (0, 1)$, the scheme in the saddle point formulation reads:

First step. Find $\{\vec{u}^{n+\Theta}, p^{n+\Theta}\} \in H_g \times L^2(\Omega)$ such that

$$\frac{1}{\Theta\Delta t}(\vec{u}^{n+\Theta}, \vec{\varphi}) + \alpha\nu((\vec{u}^{n+\Theta}, \vec{\varphi})) - (p^{n+\Theta}, \nabla \cdot \vec{\varphi}) =$$

$$\frac{1}{\Theta\Delta t}(\vec{u}^n, \vec{\varphi}) - \beta\nu((\vec{u}^n, \vec{\varphi})) - b(\vec{u}^n, \vec{u}^n, \vec{\varphi}) + (\vec{f}^n, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0,$$

$$(\nabla \cdot \vec{u}^{n+\Theta}, q) = 0, \quad \forall q \in L^2(\Omega).$$

Second step. Find $\bar{u}^{n+(1-\Theta)} \in H_g$ such that

$$\begin{aligned} \frac{1}{\Theta_b \Delta t}(\bar{u}^{n+(1-\Theta)}, \vec{\varphi}) + \beta \nu((\bar{u}^{n+(1-\Theta)}, \vec{\varphi})) + b(\bar{u}^{n+(1-\Theta)}, \bar{u}^{n+(1-\Theta)}, \vec{\varphi}) = \\ \frac{1}{\Theta_b \Delta t}(\bar{u}^{n+\Theta}, \vec{\varphi}) - \alpha \nu((\bar{u}^{n+\Theta}, \vec{\varphi})) + (p^{n+\Theta}, \nabla \cdot \vec{\varphi}) + (\bar{f}^n, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0. \end{aligned}$$

Third step. Find $\{\bar{u}^{n+1}, p^{n+1}\} \in H_g \times L^2(\Omega)$ such that

$$\begin{aligned} \frac{1}{\Theta \Delta t}(\bar{u}^{n+1}, \vec{\varphi}) + \alpha \nu((\bar{u}^{n+1}, \vec{\varphi})) - (p^{n+1}, \nabla \cdot \vec{\varphi}) = \frac{1}{\Theta \Delta t}(\bar{u}^{n+(1-\Theta)}, \vec{\varphi}) \\ - \beta \nu((\bar{u}^{n+(1-\Theta)}, \vec{\varphi})) - b(\bar{u}^{n+(1-\Theta)}, \bar{u}^{n+(1-\Theta)}, \vec{\varphi}) + (\bar{f}^{n+1}, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0, \\ (\nabla \cdot \bar{u}^{n+1}, q) = 0, \quad \forall q \in L^2(\Omega). \end{aligned}$$

On each time level we have to solve two saddle point problems and one nonlinear Dirichlet problem.

2.2 Solution of the Stokes problem.

The first and the third step can be solved by the modified version of the conjugate gradient algorithm (c.f. R.Glowinski [G I], Chp. VII, §5.8.7.3.4). This method appears to be quite robust and efficient.

Consider $W \subset L^2(\Omega)$ defined by $W = \{q \in L^2(\Omega) \mid \int_{\Omega} q(x) dx = 0\}$. The solution method is based on the following Proposition, proved in Crouzeix [C].

Proposition. Let operator $\mathcal{A} : L^2(\Omega) \rightarrow L^2(\Omega)$ be defined by

$$\mathcal{A}q = \nabla \cdot \vec{v},$$

where, for $q \in L^2(\Omega)$, the function $\vec{v} \in H_0$ is the solution of the equation

$$(2.2.1) \quad \beta \vec{v} - \nu \Delta \vec{v} = -\nabla q, \quad \text{in } \Omega.$$

For $\beta > 0$ and $\nu > 0$ the operator $\mathcal{A}$ is W -elliptic, self-adjoint and automorphic from W to W . $\square$

We apply the Proposition in the following way. Let $\{\vec{u}, p\} \in H_g \times L^2(\Omega)$ be the weak solution of the following Stokes problem

$$(2.2.2) \quad \begin{aligned} \alpha \vec{u} - \nu \Delta \vec{u} + \nabla p &= \vec{f}, \quad \text{in } \Omega, \\ \nabla \cdot \vec{u} &= 0, \quad \text{in } \Omega, \\ \vec{u} &= \vec{g}, \quad \text{on } \partial\Omega, \end{aligned}$$

and let $\vec{u}_0 \in H_g$ be the weak solution of the Dirichlet problem

$$(2.2.3) \quad \begin{aligned} \beta \vec{u}_0 - \nu \Delta \vec{u}_0 &= \vec{f}, \quad \text{in } \Omega, \\ \vec{u}_0 &= \vec{g}, \quad \text{on } \partial\Omega. \end{aligned}$$

Existence of the strong solutions (i.e. solutions in $H^2(\Omega) \times H^1(\Omega)$ and $H^2(\Omega)$, respectively) in both cases justifies the subtraction of (2.2.2) and (2.2.3) and we obtain

$$\alpha(\vec{u} - \vec{u}_0) - \nu \Delta (\vec{u} - \vec{u}_0) = -\nabla p, \quad \text{in } \Omega,$$

$$\vec{u} - \vec{u}_0 \in H_0.$$

Applying the above Proposition and the incompressibility condition, it follows that the pressure is given by the unique solution in $L^2(\Omega)/\mathbb{R}$ of the equation

$$(\text{SP}) \quad \mathcal{A}p = -\nabla \cdot \vec{u}_0.$$

2.3 Conjugate gradient solution of the problem (SP). For arbitrary $p^0 \in L^2(\Omega)$ and $\vec{u}^0$ being the solution of (2.2.3) we set $z^0 \equiv \nabla \cdot \vec{u}^0$ and $w^0 \equiv z^0$. Then, for $n \geq 0$, we compute $p^{n+1}, z^{n+1}, w^{n+1}$ from p^n, z^n, w^n as follows.

First step (Descent).

$$\rho_n = \frac{(z^n, w^n)}{(\mathcal{A}w^n, w^n)}, \quad p^{n+1} = p^n - \rho_n w^n.$$

Second step (New descent direction).

$$z^{n+1} = z^n - \rho_n \mathbb{A}w^n,$$

$$\gamma_n = \frac{|z^{n+1}|^2}{|z^n|^2},$$

$$w^{n+1} = z^{n+1} + \gamma_n w^n.$$

Since the operator $\mathcal{A}$ is defined through the solution of (2.2.1), we use the following implementation of the algorithm (2.3).

Initialization. Let $p^0 \in H^1(\Omega)$ be arbitrary. Find $\vec{u}^0 \in H_g$ such that

$$\alpha(\vec{u}^0, \vec{\varphi}) + \nu((\vec{u}^0, \vec{\varphi})) = (\vec{f}, \vec{\varphi}) - (\nabla p^0, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0.$$

Find $z^0 \in H^1(\Omega)$ such that

$$(z^0, \phi) = (\nabla \cdot \vec{u}^0, \phi), \quad \forall \phi \in H^1(\Omega),$$

$$w^0 \equiv z^0.$$

Then, for $n \geq 0$, we compute $p^{n+1}, z^{n+1}, w^{n+1}$ from p^n, z^n, w^n as follows.

First step (Descent). Find $\vec{\chi} \in H_0$ such that

$$\alpha(\vec{\chi}^n, \vec{\varphi}) + \nu((\vec{\chi}^n, \vec{\varphi})) = -(\nabla w^n, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0.$$

Find $\vartheta \in H^1(\Omega)$ such that

$$(\vartheta^n, \phi) = (\nabla \cdot \vec{\chi}^n, \phi), \quad \forall \phi \in H^1(\Omega),$$

$$\rho_n = \frac{(z^n, w^n)}{(\vartheta^n, w^n)}, \quad \text{and}$$

$$p^{n+1} = p^n - \rho_n w^n.$$

Second step (New descent direction). Find $z^{n+1} \in H^1(\Omega)$ such that

$$\begin{aligned} (z^{n+1}, \phi) &= (p^{n+1}, \phi), \quad \forall \phi \in H^1(\Omega), \\ \gamma_n &= \frac{|z^{n+1}|^2}{|z^n|^2}, \\ w^{n+1} &= z^{n+1} + \gamma_n w^n. \end{aligned}$$

Stop condition: $\rho_n \|w^n\| \leq \varepsilon \ll 1$.

Remark 1. The above version differs through the introduction of $\vartheta \in H^1(\Omega)$ from the version described in Glowinski [G I]. $\square$

Remark 2. Corresponding velocity is computed by Initialization step with p^0 replaced by p^{n+1} . The Initialization step for the first and the third step of the Θ -scheme reads:

Find $\bar{u}^0 \in H_g$ such that

$$\begin{aligned} \frac{1}{\Theta \Delta t} (\bar{u}^0, \vec{\varphi}) + \alpha \nu ((\bar{u}^0, \vec{\varphi})) &= \frac{1}{\Theta \Delta t} (\bar{u}^m, \vec{\varphi}) - \beta \nu ((\bar{u}^m, \vec{\varphi})) \\ -b(\bar{u}^m, \bar{u}^m, \vec{\varphi}) + (\vec{f}^m, \vec{\varphi}) + (p^m, \nabla \cdot \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0, \quad m = n, \text{ or } n + \Theta. \end{aligned}$$

For $m = n + \Theta$, $\vec{f}^m = \vec{f}^{n+1}$. We use the pressure from the preceding time level as the starting value instead of an arbitrary choice of p^0 . $\square$

2.4 Solution of the Regularized Burgers problem. We use the conjugate gradient method to minimize the cost functional J arising from the least square formulation.

The second step of the Θ -method consists, in general, in solving the following problem. Find $\vec{u} \in H_g$ such that

$$\alpha_1 \vec{u} - \alpha_2 \Delta \vec{u} + (\vec{u} \cdot \nabla) \vec{u} = \vec{f}, \quad \text{in } \Omega, \quad \alpha_1, \alpha_2 > 0.$$

We reformulate this problem to find the minimum of the quadratic functional J .

Let $\vec{\xi} \equiv \vec{\xi}(\vec{w}) \in H_0$ be the solution of the state equation with $\vec{w} \in H_g$:

$$\begin{aligned} \alpha_1 (\vec{\xi}, \vec{\varphi}) + \alpha_2 ((\vec{\xi}, \vec{\varphi})) &= \\ \alpha_1 (\vec{w}, \vec{\varphi}) + \alpha_2 ((\vec{w}, \vec{\varphi})) + b(\vec{w}, \vec{w}, \vec{\varphi}) - (\vec{f}, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0. \end{aligned}$$

We define the functional $J : H_g \rightarrow \mathbb{R}$ by

$$J(\vec{w}) = \frac{\alpha_1}{2} |\vec{\xi}|^2 + \frac{\alpha_2}{2} \| \vec{\xi} \|^2.$$

Now, the problem in the least square formulation reads: Find $\vec{u} \in H_g$ such that

$$(BP) \quad J(\vec{u}) \leq J(\vec{w}), \quad \forall \vec{w} \in H_g.$$

2.5 Conjugate gradient minimization.

We solve the problem (BP) using the well-known Polack-Ribier strategy in implementing the conjugate gradient method to minimize the cost functional.

Initialization. Let $\vec{u}^0 \in H_g$ is arbitrary. Find $\vec{g}^0 \in H_0$ such that

$$\alpha_1(\vec{g}^0, \vec{\varphi}) + \alpha_2((\vec{g}^0, \vec{\varphi})) = \langle J'(\vec{u}^0), \vec{\varphi} \rangle, \quad \forall \vec{\varphi} \in H_0, \\ \vec{z}^0 \equiv \vec{g}^0.$$

Here, $\langle \cdot, \cdot \rangle$ is the duality pairing between H_0 and H_0^* . We have for the right-hand side

$$\langle J'(\vec{u}^0), \vec{\varphi} \rangle = \alpha_1(\vec{\xi}, \vec{\varphi}) + \alpha_2((\vec{\xi}, \vec{\varphi})) + ((\vec{\varphi} \cdot \nabla) \vec{u}, \vec{\xi}) + ((\vec{u} \cdot \nabla) \vec{\varphi}, \vec{\xi}).$$

Then, for $n \geq 0$, we compute $\vec{u}^{n+1}, \vec{g}^{n+1}, \vec{z}^{n+1}$ from $\vec{u}^n, \vec{g}^n, \vec{z}^n$ as follows.

First step (Line minimization). Find $\lambda^n \in \mathbb{R}$ such that

$$J(\vec{u}^n - \lambda^n \vec{z}^n) \leq J(\vec{u}^n - \lambda \vec{z}^n), \quad \forall \lambda \in \mathbb{R}, \\ \vec{u}^{n+1} = \vec{u}^n - \lambda^n \vec{z}^n.$$

Second step (New descent direction). Find $\vec{g}^{n+1} \in H_0$ such that

$$\alpha_1(\vec{g}^{n+1}, \vec{\varphi}) + \alpha_2((\vec{g}^{n+1}, \vec{\varphi})) = \langle J'(\vec{u}^{n+1}), \vec{\varphi} \rangle, \quad \forall \vec{\varphi} \in H_0.$$

Set (Polack–Ribier strategy)

$$\gamma_{n+1} = \frac{\alpha_1(\vec{g}^{n+1}, \vec{g}^{n+1} - \vec{g}^n) + \alpha_2((\vec{g}^{n+1}, \vec{g}^{n+1} - \vec{g}^n))}{\alpha_1 |\vec{g}^n|^2 + \alpha_2 \|\vec{g}^n\|^2}, \quad \text{and} \\ \vec{z}^{n+1} = \vec{g}^{n+1} + \gamma_{n+1} \vec{z}^n.$$

Stop condition: $J(\vec{u}^{n+1}) \leq \varepsilon \ll 1$.

In order to implement this method for the solution of the Regularized Burgers step in the Θ -scheme we have to modify the state equation. Now, $\vec{\xi} \in H_0$ is the solution of the following problem:

$$\frac{1}{\Theta_b \Delta t} (\vec{\xi}, \vec{\varphi}) + \beta \nu ((\vec{\xi}, \vec{\varphi})) = \frac{1}{\Theta_b \Delta t} (\vec{w}, \vec{\varphi}) + \beta \nu ((\vec{w}, \vec{\varphi})) + b(\vec{w}, \vec{w}, \vec{\varphi}) \\ - \frac{1}{\Theta_b \Delta t} (\vec{u}^{n+\Theta}, \vec{\varphi}) + \beta \nu ((\vec{u}^{n+\Theta}, \vec{\varphi})) - (p^{n+\Theta}, \nabla \cdot \vec{\varphi}) - (\vec{f}, \vec{\varphi}), \quad \forall \vec{\varphi} \in H_0.$$

We set $\vec{u}^{n+(1-\Theta)} \equiv \vec{u}^n$ if $J(\vec{u}^n) \leq \varepsilon \ll 1$.

3. SPATIAL APPROXIMATION OF THE SOLUTION

The squeezing property of the solution of (WNS) indicates that the flow is essentially characterized by a finite number of parameters. Taking into account that also the number of determining modes is finite (c.f. P.Constantin *et al.* [CFT]), there is some theoretical justification to the use of the Galerkin approximation of the Navier–Stokes equations in the spaces H_h and V_h , with $\dim H_h = \dim V_h = nN(h)$.

When the solution of (WNS) converge to a permanent regime this approach does not always leads to acceptable results. Permanent regime is usually characterized by the presence of the small scale structures which are difficult to capture by the Galerkin approximation.

The existence of some kind of attracting sets for the solution of the Navier–Stokes equations may lead to the construction of a nonlinear system for the functions representing different modes (scales) of the solution. Capturing the interactions of the different modes can help to deal with difficult flows.

Yet, the Galerkin approximation remains the most reliable approximation method so far, as it is neither clear which modes determine the qualitative properties of the solution (e.g. the time of the transition to turbulence) nor it is established in general how to determine them.

3.1 Mixed finite element approximation.

The choice of finite element spaces is basically driven by the approximation of the Generalized Stokes problem. Let $H_h(\Omega)$ and $L_h^2(\Omega)$, with $\dim H_h(\Omega) = nN(h)$, $\dim L_h^2(\Omega) = M(h)$, are the finite element approximations of the spaces $[H^1(\Omega)]^n, L^2(\Omega)$, where h is parameter of the mesh size. We define $H_{gh}, V_{gh}, H_{0h}, V_{0h}$ by

$$\begin{aligned} H_{gh}(\Omega) &= \{\vec{v}_h \in H_h^1(\Omega) \mid \vec{v}_h|_{\partial\Omega} = \vec{g}\}, \\ V_{gh}(\Omega) &= \{\vec{v}_h \in H_h^1(\Omega) \mid (\nabla \cdot \vec{v}_h, q) = 0, \forall q_h \in L_h^2(\Omega), \vec{v}_h|_{\partial\Omega} = \vec{g}\}, \\ V_{0h}(\Omega) &= \{\vec{v}_h \in H_h^1(\Omega) \mid (\nabla \cdot \vec{v}_h, q) = 0, \forall q_h \in L_h^2(\Omega), \vec{v}_h|_{\partial\Omega} = 0\}, \end{aligned}$$

and $H_{0h} = H_h(\Omega) \cap [H_0^1(\Omega)]^n$.

Replacing the spaces V, H, L^2 by their finite dimensional equivalents in (WNS) and (SNS) we obtain the Galerkin approximations $\vec{u}_h \in L^\infty(0, T; V_{gh})$ and $\{\vec{u}_h, p_h\} \in L^\infty(0, T; H_{gh} \times L_h(\Omega))$ of the weak solution of (WNS) and (SNS).

3.2 Properties of the approximate spaces. Let $\{Q_h\}_h$ be a family of quadrilaterals so that $\bar{\Omega} = \bigcup_{Q \in Q_h} Q$. We set $h(Q) = \text{length of the longest diagonal of } Q$, and $h = \max_{Q \in Q_h} h(Q)$. We suppose

$$(A) \quad \frac{h}{\min_{Q \in Q_h} h(Q)} \leq \beta, \quad \forall Q_h.$$

The decompositions Q_h are assumed to be “face to face”.

Finite element spaces approximating the “velocity space” $[H^1(\Omega)]^n$ and the “pressure space” $L^2(\Omega)$ have, in general, the forms

$$\begin{aligned} H_h(\Omega) &= \{\vec{v}_h \in [L^2(\Omega)]^n \mid \vec{v}_h|_Q \in [P(Q)]^n, \forall Q \in Q_h\}, \\ L_h(\Omega) &= \{\vec{q}_h \in [L^2(\Omega)]^n \mid \vec{q}_h|_Q \in [R(Q)]^n, \forall Q \in Q_h\}, \end{aligned}$$

where $P(Q)$ and $R(Q)$ are spaces of polynomials, the maximum degree of which is bounded uniformly with respects to $Q \in Q_h$ and h . We refer to the approximation H_h as conforming (and we denote this space H_h^1) if the polynomial space $P(Q)$ is such that $H_h \subset [H^1(\Omega)]^n$ and nonconforming otherwise. Moreover, we assume:

(B) Each $\vec{v}_h \in H_h(\Omega)$ satisfies $\int_\Gamma (\vec{v}_h|_Q - \vec{v}_h|_{Q'}) dS = 0$, $\Gamma = Q \cap Q'$, $Q, Q' \in Q_h$ and $\int_\Gamma \vec{v}_h|_Q dS = 0$, $\Gamma = \partial\Omega \cap Q$, $Q \in Q_h$.

(C) For each $\vec{v} \in [H_0^1(\Omega)]^n \cap [H^2(\Omega)]^n$ and $q \in H^1(\Omega)$ there exist approximations $\Pi_h \vec{v} \in H_h(\Omega)$ and $\pi_h q \in L_h(\Omega)$ so that $\|\nabla_h(\vec{v} - \Pi_h \vec{v})\| \leq Ch \|\vec{v}\|_2$ and $\|q - \pi_h q\|_{L^2(\Omega)/\mathbb{R}} \leq Ch \|q\|_{L^2(\Omega)/\mathbb{R}}$.

The “discrete gradient” ∇_h is understood in the piecewise sense with respect to the decomposition of Q_h .

To ensure that the discrete pressure $p_h(t) \in L_h(\Omega)$ depends continuously on the discrete velocity $\vec{u}_h \in V_{gh}(\Omega)$ uniformly in h , we assume classical “inf-sup” condition for the mixed finite element methods (Babuška–Brezzi–Ladyzhenskaya condition):

For every $q_h \in L_h(\Omega)$, there exists a nontrivial function $\vec{\phi}_h \in V_{gh}(\Omega)$ such that

$$(D) \quad \inf_{q_h \in L_h(\Omega)} \sup_{\vec{\phi}_h \in V_{gh}(\Omega)} \frac{(\nabla_h \vec{\phi}_h, q_h)}{\|\vec{\phi}_h\| \cdot |q_h|} \geq \gamma > 0.$$

4. STABILITY AND CONVERGENCE OF THE FRACTIONAL STEP Θ -SCHEME

The proof of the stability and convergence of the scheme follow the framework developed in [T I,II]. The difficulty in obtaining the a priori estimates lies in the presence of the trilinear form on the right-hand side of the Stokes steps.

Technical preliminaries. We assume that the gradient operator has an extension ∇_h on the algebraic sum $[H_0^1]^n + H_h$ so that $\nabla = \nabla_h$ on $[H_0^1]^n$ and $((\cdot, \cdot))_h \equiv (\nabla_h \cdot, \nabla_h \cdot)_{L^2}$ gives the norm on H_h . This extension ∇_h of ∇ is understood in the nonconforming case in the piecewise sense with respect to the decomposition Q_h of Ω . Thus for $\phi \in [H_0^1]^n + H_h$ the term $\nabla_h \phi$ is the L^2 -function over Ω which coincides with $\nabla \phi|_Q$, $\forall Q \in Q_h$.

Since the spaces H_h, V_h are finite dimensional they can be equipped with equivalent norms induced by $[L^2(\Omega)]^n$ and $[H_0^1(\Omega)]^n$. Moreover the condition (A) ensures that this equivalence can be written in the form:

$$(4.1) \quad D_1 |u_h| \leq \|u_h\|_h \leq D_2 h^{-1} |u_h|.$$

The constant D_2 depends on the degree of polynomial approximation. In the following we will assume that $D_1 = D_2 = 1$.

For each $\vec{u}_h, \vec{v}_h, \vec{w}_h \in H_h$ we define

$$(4.2) \quad b_h(\vec{u}_h, \vec{v}_h, \vec{w}_h) \equiv \frac{1}{2} \left(((\vec{u}_h \cdot \nabla_h) \vec{v}_h, \vec{w}_h) - ((\vec{u}_h \cdot \nabla_h) \vec{w}_h, \vec{v}_h) \right).$$

It holds

$$(4.3) \quad b_h(\vec{u}_h, \vec{v}_h, \vec{v}_h) = 0, \quad \forall \vec{u}_h, \vec{v}_h \in H_h.$$

There exists a function $S(h)$ so that

$$(4.4) \quad |b_h(\vec{u}_h, \vec{v}_h, \vec{w}_h)| \leq S(h) |\vec{u}_h| \|\vec{v}_h\|_h |\vec{w}_h|,$$

and $S(h) = D_3 h^{-1}$, in the conforming case, and, $S(h) = D_3 h^{-(1+\epsilon)}$, in the non-conforming case, in 2D, and $S(h) = D_4 h^{-\frac{3}{2}}$ in 3D, where $\epsilon > 0$, $D_3 > 0$ and $D_4 > 0$.

We assume homogeneous boundary conditions in what follows. To approximate the initial condition we project $\vec{u}_0$ to V_{0h} by finding $u_{0h} \in V_{0h}$ such that

$$(4.5) \quad ((\vec{u}_{0h}, \vec{\varphi}_h))_h = ((\vec{u}_0, \vec{\varphi}_h))_h, \quad \forall \vec{\varphi}_h \in V_{0h}.$$

From now on we consider the following fully discrete problems ($m \equiv n + (1 - \Theta)$).

Find $\vec{u}_h^{n+\Theta} \in V_{0h}$ such that

$$(4.6) \quad \begin{aligned} & \frac{1}{\Theta \Delta t} (\vec{u}_h^{n+\Theta}, \vec{\varphi}_h) + \alpha \nu ((\vec{u}_h^{n+\Theta}, \vec{\varphi}_h))_h = \\ & \frac{1}{\Theta \Delta t} (\vec{u}_h^n, \vec{\varphi}_h) - \beta \nu ((\vec{u}_h^n, \vec{\varphi}_h))_h - b_h(\vec{u}_h^n, \vec{u}_h^n, \vec{\varphi}_h) + (\vec{f}^n, \vec{\varphi}_h), \quad \forall \vec{\varphi}_h \in V_{0h}. \end{aligned}$$

Find $\vec{u}_h^m \in H_{0h}$ such that

$$(4.7) \quad \begin{aligned} & \frac{1}{\Theta_b \Delta t} (\vec{u}_h^m, \vec{\varphi}_h) + \beta \nu ((\vec{u}_h^m, \vec{\varphi}_h))_h + b_h(\vec{u}_h^m, \vec{u}_h^m, \vec{\varphi}_h) = \\ & \frac{1}{\Theta_b \Delta t} (\vec{u}_h^{n+\Theta}, \vec{\varphi}_h) - \alpha \nu ((\vec{u}_h^{n+\Theta}, \vec{\varphi}_h))_h + (p_h^{n+\Theta}, \nabla \cdot \vec{\varphi}_h) + (\vec{f}^n, \vec{\varphi}_h), \quad \forall \vec{\varphi}_h \in H_{0h}. \end{aligned}$$

Find $\vec{u}_h^{n+1} \in V_{0h}$ such that

$$(4.8) \quad \begin{aligned} & \frac{1}{\Theta \Delta t} (\vec{u}_h^{n+1}, \vec{\varphi}_h) + \alpha \nu ((\vec{u}_h^{n+1}, \vec{\varphi}_h))_h = \\ & \frac{1}{\Theta \Delta t} (\vec{u}_h^m, \vec{\varphi}_h) - \beta \nu ((\vec{u}_h^m, \vec{\varphi}_h))_h - b_h(\vec{u}_h^m, \vec{u}_h^m, \vec{\varphi}_h) + (\vec{f}^{n+1}, \vec{\varphi}_h), \quad \forall \vec{\varphi}_h \in V_{0h}. \end{aligned}$$

We define $\tilde{f}^n \equiv \frac{1}{\Delta t} \int_{t_{n-1}}^{t_n} f(t) dt$ and the pressure $p_h^{n+\Theta} \in L^2(\Omega)/\mathbb{R}$ we define by

$$(4.9) \quad \begin{aligned} (p_h^{n+\Theta}, \nabla_h \cdot \vec{\varphi}_h) &= \frac{1}{\Theta \Delta t} (\vec{u}_h^{n+\Theta} - \vec{u}_h^n, \vec{\varphi}_h) + \alpha \nu ((\vec{u}_h^{n+\Theta}, \vec{\varphi}_h))_h + \\ &\quad \beta \nu ((\vec{u}_h^n, \vec{\varphi}_h))_h + b_h(\vec{u}_h^n, \vec{u}_h^n, \vec{\varphi}_h) - (\tilde{f}^n, \vec{\varphi}_h), \quad \forall \vec{\varphi}_h \in H_{0h}. \end{aligned}$$

Next, we define the following three families of functions $\vec{u}_{h,\Delta t}^{(i)} : [0, T] \rightarrow V_h$ (resp. H_h for $i = 3$)

$$(4.10) \quad \left. \begin{aligned} \vec{u}_{h,\Delta t}^{(1)}(t) &= \vec{u}_h^n, \\ \vec{u}_{h,\Delta t}^{(2)}(t) &= \vec{u}_h^{n+\Theta}, \\ \vec{u}_{h,\Delta t}^{(3)}(t) &= \vec{u}_h^{n+(1-\Theta)}, \\ \tilde{f}_{\Delta t}(t) &= \tilde{f}^n, \end{aligned} \right\} \quad t \in [n\Delta t, (n+1)\Delta t),$$

for $n = 1, 2, \dots, N$, $N \cdot \Delta t = T$.

Correspondingly, we use the functions $\vec{w}_{h,\Delta t} : [0, T] \rightarrow V_h$:

$$(4.11) \quad \vec{w}_{h,\Delta t}(t) = \vec{u}_h^n + \frac{1}{\Delta t} (t - t_n)(\vec{u}_h^{n+1} - \vec{u}_h^n), \quad t \in [n\Delta t, (n+1)\Delta t).$$

The functions defined above in (4.10,12) satisfy

$$(4.12) \quad \begin{aligned} &\frac{1}{\Theta} \frac{d}{dt} (\vec{w}_{h,\Delta t}, \vec{\varphi}_h) + \alpha \nu ((\vec{u}_{h,\Delta t}^{(1)}(t + \Delta t), \vec{\varphi}_h))_h + \beta \nu ((\vec{u}_{h,\Delta t}^{(1)}(t), \vec{\varphi}_h))_h = \\ &\frac{1}{\Theta \Delta t} (\vec{u}_{h,\Delta t}^{(3)}(t) - \vec{u}_{h,\Delta t}^{(2)}(t), \vec{\varphi}_h) - \beta \nu ((\vec{u}_{h,\Delta t}^{(3)}(t), \vec{\varphi}))_h - \alpha \nu ((\vec{u}_{h,\Delta t}^{(2)}(t), \vec{\varphi}))_h - \\ &b_h(\vec{u}_{h,\Delta t}^{(2)}(t), \vec{u}_{h,\Delta t}^{(2)}(t), \vec{\varphi}_h) - b_h(\vec{u}_{h,\Delta t}^{(3)}(t), \vec{u}_{h,\Delta t}^{(3)}(t), \vec{\varphi}_h) + (\tilde{f}^n + \tilde{f}^{n+1}, \vec{\varphi}_h), \end{aligned}$$

for all $\vec{\varphi}_h \in V_{0h}$, $t \in [n\Delta t, (n+1)\Delta t)$.

The results concerning the stability and convergence of the scheme (4.6–8) are given by the following

Theorem. *Let us assume that there exists unique strong solution of the problem (WNS), with homogeneous Dirichlet boundary condition, on the time interval $(0, T)$. Let us assume that $\Theta \in (0, 1 - \frac{\sqrt{2}}{2})$, $\alpha \in (0, 1)$, $\delta \in (0, \frac{1}{3})$, and, let us assume that the time step Δt and mesh size h are connected with the data by the following stability conditions*

$$(4.13) \quad D_5 \left(\left(\nu \frac{\Delta t}{h^2} \right)^2 + (\Delta t S(h))^2 \Lambda_T \right) \leq (1 - \delta) \Theta \Delta t \alpha \nu, \quad \text{where}$$

$$\Lambda_T \equiv |\vec{u}_0|^2 + D_5 \left(\left(\nu \frac{\Delta t}{h^2} \right)^2 \|\vec{u}_0\|_h^2 + (\Delta t S(h))^2 |\vec{u}_0|^2 \|\vec{u}_0\|_h^2 + \|\tilde{f}\|_{L^2(0,T,L^2(\Omega))}^2 \right),$$

$$D_5 = \left(\frac{\Theta_b}{\delta} \right)^2 \left(1 + \left(\frac{\Theta_b}{\Theta} \right)^2 \right).$$

Then:

- (1) The functions $\bar{u}_{h,\Delta t}^{(i)}$ belong to a bounded set in $L^2(0, T; V_h) \cap L^\infty(0, T; [L^2(\Omega)]^n)$, for $i = 1, 2$ and to a bounded set in $L^2(0, T; H_h) \cap L^\infty(0, T; [L^2(\Omega)]^n)$, for $i = 3$.
- (2) These functions converge strongly to the unique strong solution of (WNS) in $L^2(0, T; [H_0^1(\Omega)]^n) \cap L^q(0, T; [L^2(\Omega)]^n)$, for $q \in [1, \infty)$, as $(h, \Delta t) \rightarrow 0_+$, in the case of conforming approximations and in $L^q(0, T; [L^2(\Omega)]^n)$, for $q \in [2, \infty)$, in the nonconforming case.

The proof of the Theorem is given in the sections 4.1 (stability) and 4.2 (convergence).

4.1 Stability of the scheme. First, we obtain a priori estimates.

Step 1: We take $\bar{\varphi}_h = \bar{u}_h^{n+\Theta}$ in (4.6) to get

$$(4.1.1) \quad \frac{1}{\Theta \Delta t} |\bar{u}_h^{n+\Theta}|^2 + \alpha \nu \|\bar{u}_h^{n+\Theta}\|_h^2 = \frac{1}{\Theta \Delta t} (\bar{u}_h^n, \bar{u}_h^{n+\Theta}) - \beta \nu ((\bar{u}_h^n, \bar{u}_h^{n+\Theta}))_h \\ - b_h(\bar{u}_h^n, \bar{u}_h^n, \bar{u}_h^{n+\Theta}) + (\bar{f}^n, \bar{u}_h^{n+\Theta})$$

Since the right-hand side of (4.1.1) is equal to

$$(4.1.2) \quad -\frac{1}{2\Theta \Delta t} |\bar{u}_h^{n+\Theta} - \bar{u}_h^n|^2 + \frac{1}{2\Theta \Delta t} |\bar{u}_h^{n+\Theta}|^2 + \frac{1}{2\Theta \Delta t} |\bar{u}_h^n|^2 + \frac{\beta \nu}{2} \|\bar{u}_h^{n+\Theta} - \bar{u}_h^n\|_h^2 \\ - \frac{\beta \nu}{2} \|\bar{u}_h^n\|_h^2 - \frac{\beta \nu}{2} \|\bar{u}_h^{n+\Theta}\|_h^2 - b_h(\bar{u}_h^n, \bar{u}_h^n, \bar{u}_h^{n+\Theta}) + (\bar{f}^n, \bar{u}_h^{n+\Theta}),$$

and, since by (4.1) we have $\|\bar{u}_h^{n+\Theta} - \bar{u}_h^n\|_h^2 \leq \frac{1}{h^2} |\bar{u}_h^{n+\Theta} - \bar{u}_h^n|^2$, we obtain

$$(4.1.3) \quad |\bar{u}_h^{n+\Theta}|^2 + (1 + \alpha) \Theta \Delta t \nu \|\bar{u}_h^{n+\Theta}\|_h^2 + \Theta \Delta t \beta \nu \|\bar{u}_h^n\|_h^2 + \\ \left(1 - \frac{\Theta \Delta t \beta \nu}{h^2}\right) |\bar{u}_h^{n+\Theta} - \bar{u}_h^n|^2 \leq |\bar{u}_h^n|^2 - 2\Theta \Delta t b_h(\bar{u}_h^n, \bar{u}_h^n, \bar{u}_h^{n+\Theta}) + 2\Theta \Delta t (\bar{f}^n, \bar{u}_h^{n+\Theta}).$$

We estimate the right-hand side of (4.1.3) by the Young's inequality and (4.3.4).

Because $0 < \Theta < 1 - \frac{\sqrt{2}}{2}$ we have $2\Theta^2 < \Theta_b^2$ and we obtain

$$(4.1.4) \quad \frac{1}{2} |\bar{u}_h^{n+\Theta} - \bar{u}_h^n|^2 + (\Theta_b \Delta t S(h))^2 |\bar{u}_h^n|^2 \|\bar{u}_h^n\|_h^2 + \Theta \Delta t \nu \|\bar{u}_h^{n+\Theta}\|_h^2 + \frac{\Theta \Delta t}{\nu} |\bar{f}^n|^2.$$

The last two inequalities yield

$$(4.1.5) \quad |\bar{u}_h^{n+\Theta}|^2 + \Theta \Delta t \alpha \nu \|\bar{u}_h^{n+\Theta}\|_h^2 + \Theta \Delta t \beta \nu \|\bar{u}_h^n\|_h^2 + \left(\frac{1}{2} - \frac{\Theta \Delta t \beta \nu}{h^2}\right) |\bar{u}_h^{n+\Theta} - \bar{u}_h^n|^2 \leq \\ |\bar{u}_h^n|^2 + (\Theta_b \Delta t S(h))^2 |\bar{u}_h^n|^2 \|\bar{u}_h^n\|_h^2 + \frac{\Theta \Delta t}{\nu} |\bar{f}^n|^2.$$

$$\begin{aligned} & \cdot (\frac{y_n}{u} \cdot \frac{y}{u} f) + (\frac{y_n}{u} \cdot \frac{y}{u} \Delta'_{\Theta + \frac{y}{u}} \frac{y}{u} d) + \frac{y}{u} ((\frac{y_n}{u} \cdot \frac{y}{u} \Theta + \frac{y}{u} n)) \mathcal{A} \mathcal{D} - (\frac{y_n}{u} \cdot \frac{y}{u} \Theta + \frac{y}{u} n) \frac{\mathcal{I} \nabla^q \Theta}{\Gamma} \\ & = \frac{y}{z} \|\frac{y_n}{u}\| \mathcal{A} \mathcal{D} + \frac{y}{z} \|\frac{y_n}{u}\| \frac{\mathcal{I} \nabla^q \Theta}{\Gamma} \end{aligned} \quad (4.16)$$
$$\begin{aligned} & \cdot ({}_{\Theta+u}^y n - \frac{y}{u} n, {}_u f) - \\ & ({}_{\Theta+u}^y n - \frac{y}{u} n, \frac{y}{u} n, {}_u n) {}^y q + {}^y (({}_{\Theta+u}^y n - \frac{y}{u} n, \frac{y}{u} n)) \mathcal{A} \mathcal{G} + {}^y (({}_{\Theta+u}^y n - \frac{y}{u} n, {}_{\Theta+u}^y n)) \mathcal{A} \mathcal{V} + \\ & ({}_{\Theta+u}^y n - \frac{y}{u} n, \frac{y}{u} n - {}_{\Theta+u}^y n) \frac{\mathcal{I} \nabla \Theta}{1} = (({}_{\Theta+u}^y n - \frac{y}{u} n) \cdot {}^y \Delta, {}_{\Theta+u}^y d) = (\frac{y}{u} n \cdot {}^y \Delta, {}_{\Theta+u}^y d) \end{aligned} \tag{4.17}$$
[illegible]

$$\begin{aligned}
& \cdot \left| u^f \right|_z \frac{\lambda \mathcal{O}}{t \nabla^q \Theta} + \frac{y}{z} \| u_n^y \|_z | u_n^y |_z ((y) S t \nabla^q \Theta) \frac{\mathcal{GZ}}{1} \\
& + \frac{y}{z} \| u_n^y \|_z \left(\frac{z^y}{\lambda \mathcal{G} t \nabla^q \Theta} \right) \frac{\mathcal{GZ}}{1} + \frac{y}{z} | u_n^y - \Theta u_n^y |_z \left(\frac{\Theta}{t^y} \right) \frac{\mathcal{GZ}}{1} + \frac{y}{z} | \Theta u_n^y |_z \quad (6.1.9) \\
& \geq \frac{y}{z} | \Theta u_n^y - u_n^y | (\mathcal{GZ} - 1) + \frac{y}{z} \| \Theta u_n^y \| \lambda \mathcal{O} t \nabla^q \Theta + \frac{y}{z} \| u_n^y \| \lambda \mathcal{G} t \nabla^q \Theta + \frac{y}{z} | u_n^y |
\end{aligned}$$

$$\begin{aligned} & ((\underline{u}_n - \ominus + \underline{u}_n, \underline{u}_n^c) - (\underline{u}_n - \ominus + \underline{u}_n, \underline{u}_n^c) \underline{u} q + \iota((\underline{u}_n - \ominus + \underline{u}_n, \underline{u}_n^c)) \mathcal{A} \mathcal{G} + \\ & \iota((\underline{u}_n - \ominus + \underline{u}_n, \underline{u}_n^c)) \mathcal{A} \mathcal{G} + \iota \left| \underline{u}_n - \ominus + \underline{u}_n \right| \frac{\iota \nabla \Theta}{1} = ((\underline{u}_n - \ominus + \underline{u}_n) \cdot \iota \Delta, \ominus + \underline{u}_n^c) = 0 \end{aligned} \quad (1.10)$$

$$\begin{aligned} & \left(\frac{y_n}{\theta + y_n} - \frac{y_n}{\theta + y_n} \right) - \left(\frac{y_n}{\theta + y_n} - \frac{y_n}{\theta + y_n} \right) y_n + y_n \left(\frac{y_n}{\theta + y_n} - \frac{y_n}{\theta + y_n} \right) \lambda \theta + \\ & y_n \left(\frac{y_n}{\theta + y_n} - \frac{y_n}{\theta + y_n} \right) \lambda \theta + \left| \frac{y_n}{\theta + y_n} - \frac{y_n}{\theta + y_n} \right| \frac{\lambda \nabla \Theta}{1} = \left(\frac{y_n}{\theta + y_n} - \frac{y_n}{\theta + y_n} \right) \cdot y_n \Delta \left(\frac{y_n}{\theta + y_n} - \frac{y_n}{\theta + y_n} \right) = 0 \end{aligned}$$

and gives the inequality

$$\begin{aligned} & \geq \frac{y}{z} \left| \frac{y}{u} - \frac{y}{\Theta + u} \right| \frac{\lambda \Theta}{\Theta} + \frac{y}{z} \left| \frac{y}{u} - \frac{y}{\Theta + u} \right| \frac{\lambda \Theta}{\Theta} + \frac{y}{z} \left| \frac{y}{u} - \frac{y}{\Theta + u} \right| \frac{\lambda \Theta}{\Theta} + \frac{y}{z} \left| \frac{y}{u} - \frac{y}{\Theta + u} \right| \frac{\lambda \Theta}{\Theta} \\ & \quad + \frac{y}{z} \left| \frac{y}{u} - \frac{y}{\Theta + u} \right| \frac{\lambda \Theta}{\Theta} + \frac{y}{z} \left| \frac{y}{u} - \frac{y}{\Theta + u} \right| \frac{\lambda \Theta}{\Theta} + \frac{y}{z} \left| \frac{y}{u} - \frac{y}{\Theta + u} \right| \frac{\lambda \Theta}{\Theta} + \frac{y}{z} \left| \frac{y}{u} - \frac{y}{\Theta + u} \right| \frac{\lambda \Theta}{\Theta} \end{aligned} \quad (4.1.11)$$

Using the last estimate, the inequality (4.1.9) reads

[illegible]

Step 3: We take $\vec{\varphi}_h = \vec{u}_h^{n+1} \in V_{0h}$ in (4.8) and we get

$$(4.1.13) \quad \frac{1}{\Theta \Delta t} |\vec{u}_h^{n+1}|^2 + \alpha \nu \|\vec{u}_h^{n+1}\|_h^2 = \frac{1}{\Theta \Delta t} (\vec{u}_h^m, \vec{u}_h^{n+1}) - \beta \nu ((\vec{u}_h^m, \vec{u}_h^{n+1}))_h - b_h(\vec{u}_h^m, \vec{u}_h^m, \vec{u}_h^{n+1}) + (\vec{f}^{n+1}, \vec{u}_h^{n+1}).$$

As in the step 1 we obtain

$$(4.1.14) \quad \begin{aligned} & |\vec{u}_h^{n+1}|^2 + \Theta \Delta t (1 + \alpha) \nu \|\vec{u}_h^{n+1}\|_h^2 + \Theta \Delta t \beta \nu \|\vec{u}_h^m\|_h^2 + \\ & \left(1 - \frac{\Theta \Delta t \beta \nu}{h^2}\right) |\vec{u}_h^{n+1} - \vec{u}_h^m|^2 \leq |\vec{u}_h^m|^2 - 2\Theta \Delta t b_h(\vec{u}_h^m, \vec{u}_h^m, \vec{u}_h^{n+1}) + \\ & 2\Theta \Delta t (\vec{f}^{n+1}, \vec{u}_h^{n+1}). \end{aligned}$$

To evaluate $b_h(\vec{u}_h^m, \vec{u}_h^m, \vec{u}_h^{n+1})$ we use (4.7) with $\vec{\varphi}_h = \vec{u}_h^{n+1} - \vec{u}_h^m$

$$(4.1.15) \quad \begin{aligned} & -b_h(\vec{u}_h^m, \vec{u}_h^m, \vec{u}_h^{n+1}) = -b_h(\vec{u}_h^m, \vec{u}_h^m, \vec{u}_h^{n+1} - \vec{u}_h^m) = \\ & \frac{1}{\Theta_b \Delta t} (\vec{u}_h^m - \vec{u}_h^{n+\Theta}, \vec{u}_h^{n+1} - \vec{u}_h^m) + \beta \nu ((\vec{u}_h^m, \vec{u}_h^{n+1} - \vec{u}_h^m))_h + \\ & \alpha \nu ((\vec{u}_h^{n+\Theta}, \vec{u}_h^{n+1} - \vec{u}_h^m))_h - (\vec{f}^n, \vec{u}_h^{n+1} - \vec{u}_h^m) - (p^{n+\Theta}, \nabla_h \cdot (\vec{u}_h^{n+1} - \vec{u}_h^m)). \end{aligned}$$

Since by the equation for the pressure (4.9)

$$(4.1.16) \quad \begin{aligned} & -(p^{n+\Theta}, \nabla_h \cdot (\vec{u}_h^{n+1} - \vec{u}_h^m)) = -\frac{1}{\Theta \Delta t} (\vec{u}_h^{n+\Theta} - \vec{u}_h^n, \vec{u}_h^{n+1} - \vec{u}_h^m) \\ & -\alpha \nu ((\vec{u}_h^{n+\Theta}, \vec{u}_h^{n+1} - \vec{u}_h^m))_h - \beta \nu ((\vec{u}_h^n, \vec{u}_h^{n+1} - \vec{u}_h^m))_h - b_h(\vec{u}_h^n, \vec{u}_h^n, \vec{u}_h^{n+1} - \vec{u}_h^m) \\ & + (\vec{f}^n, \vec{u}_h^{n+1} - \vec{u}_h^m), \end{aligned}$$

we express $-2\Theta \Delta t b_h(\vec{u}_h^m, \vec{u}_h^m, \vec{u}_h^{n+1})$ in the form

$$(4.1.17) \quad \begin{aligned} & 2\frac{\Theta}{\Theta_b} (\vec{u}_h^m - \vec{u}_h^{n+\Theta}, \vec{u}_h^{n+1} - \vec{u}_h^m) - 2(\vec{u}_h^{n+\Theta} - \vec{u}_h^n, \vec{u}_h^{n+1} - \vec{u}_h^m) \\ & + 2\Theta \Delta t \beta \nu ((\vec{u}_h^m - \vec{u}_h^{n+\Theta}, \vec{u}_h^{n+1} - \vec{u}_h^m))_h - 2\Theta \Delta t b_h(\vec{u}_h^n, \vec{u}_h^n, \vec{u}_h^{n+1} - \vec{u}_h^m). \end{aligned}$$

Now, we obtain the following inequality from (4.1.14) using (4.1.17) and Schwarz' and Young's inequalities and (4.1)

$$(4.1.18) \quad \begin{aligned} & |\vec{u}_h^{n+1}|^2 + \Theta \Delta t \alpha \nu \|\vec{u}_h^{n+1}\|_h^2 + \Theta \Delta t \beta \nu \|\vec{u}_h^m\|_h^2 \\ & + \left(\frac{1}{8} - \frac{\Theta \Delta t \beta \nu}{h^2}\right) |\vec{u}_h^{n+1} - \vec{u}_h^m|^2 \leq |\vec{u}_h^m|^2 + 8 \left(\frac{\Theta_b \Delta t \nu}{h^2}\right)^2 \|\vec{u}_h^n\|_h^2 \\ & + 2 \left(\frac{\Theta}{\Theta_b}\right)^2 |\vec{u}_h^m - \vec{u}_h^{n+\Theta}|^2 + 8 |\vec{u}_h^{n+\Theta} - \vec{u}_h^n|^2 \\ & (2\Theta_b \Delta t S(h))^2 |\vec{u}_h^n|^2 \|\vec{u}_h^n\|_h^2 + \frac{\Theta \Delta t}{\nu} |\vec{f}^{n+1}|^2. \end{aligned}$$

For $0 < \Theta < 1 - \frac{\sqrt{2}}{2}$ there exists $\delta > 0$ such that $2 \left(\frac{\Theta}{\Theta_b} \right)^2 < 1 - 3\delta$. The results of the step 1 and step 2, namely (4.1.12,11) and (4.1.5), provide the desired inequality

$$(4.1.19) \quad \begin{aligned} & |\bar{u}_h^{n+1}|^2 + \Theta \Delta t \alpha \nu \|\bar{u}_h^{n+1}\|_h^2 + \Theta \Delta t \beta \nu \|\bar{u}_h^m\|_h^2 \\ & + \left(\frac{1}{8} - \frac{\Theta \Delta t \beta \nu}{h^2} \right) |\bar{u}_h^{n+1} - \bar{u}_h^m|^2 \leq |\bar{u}_h^n|^2 + D_5 \frac{\Delta t}{2\alpha \nu} \left(|\bar{f}^n|^2 + |\bar{f}^{n+1}|^2 \right) + \\ & D_5 \left(\left(\frac{\Delta t \nu}{h^2} \right)^2 \|\bar{u}_h^n\|_h^2 + (\Delta t S(h))^2 |\bar{u}_h^n|^2 \|\bar{u}_h^n\|_h^2 \right). \end{aligned}$$

Summing up the above inequality for $n = 1, 2, \dots, r$ we obtain

$$(4.1.20) \quad \begin{aligned} & |\bar{u}_h^{r+1}|^2 + \Theta \Delta t \alpha \nu \sum_{n=0}^r \|\bar{u}_h^{n+1}\|_h^2 + \Theta \Delta t \beta \nu \sum_{n=0}^r \|\bar{u}_h^m\|_h^2 \\ & + \left(\frac{1}{8} - \frac{\Theta \Delta t \beta \nu}{h^2} \right) \sum_{n=0}^r |\bar{u}_h^{n+1} - \bar{u}_h^m|^2 \leq |\bar{u}_{0h}|^2 + D_5 \frac{\Delta t}{\alpha \nu} \sum_{n=0}^{r+1} \left(|\bar{f}^n|^2 \right) \\ & + D_5 \left(\nu \frac{\Delta t}{h^2} \right)^2 \sum_{n=0}^r \|\bar{u}_h^n\|_h^2 + D_5 (\Delta t S(h))^2 \sum_{n=0}^r |\bar{u}_h^n|^2 \|\bar{u}_h^n\|_h^2 \end{aligned}$$

To obtain the a priori estimates we proceed by induction in r . Since the left-hand side of (4.1.20) is bounded for $r = 0$ let us assume that

$$(4.1.21) \quad \begin{aligned} & |\bar{u}_h^{r+1}|^2 + \delta \Theta \Delta t \alpha \nu \sum_{n=0}^r \|\bar{u}_h^{n+1}\|_h^2 + \Theta \Delta t \beta \nu \sum_{n=0}^r \|\bar{u}_h^m\|_h^2 \\ & + \left(\frac{1}{8} - \frac{\Theta \Delta t \beta \nu}{h^2} \right) \sum_{n=0}^r |\bar{u}_h^{n+1} - \bar{u}_h^m|^2 \leq |\bar{u}_{0h}|^2 + D_5 \left(\nu \frac{\Delta t}{h^2} \right)^2 \|\bar{u}_{0h}\|_h^2 \\ & + D_5 (\Delta t S(h))^2 |\bar{u}_{0h}|^2 \|\bar{u}_{0h}\|_h^2 + D_5 \frac{\Delta t}{\alpha \nu} \sum_{n=0}^{r+1} \left(|\bar{f}^n|^2 \right). \end{aligned}$$

We show that this inequality is also true for $r+1$ and then the a priori estimate for $\bar{u}_h^n$ follows by recursion. We denote the right-hand side of the inequality (4.1.21) by Λ_r and we observe that by the induction assumption we have

$$(4.1.22) \quad |\bar{u}_h^{r+1}|^2 \leq \Lambda_r \leq \Lambda_T, \quad \text{for } n = 1, 2, \dots, r.$$

This implies that the right-hand side of (4.1.20) for $r+1$ can be estimated by

$$\Lambda_{r+1} + D_5 \left(\left(\nu \frac{\Delta t}{h^2} \right)^2 + (\Delta t S(h))^2 \Lambda_{r+1} \right) \sum_{n=1}^{r+1} \|\bar{u}_h^n\|_h^2.$$

The stability assumption (4.13) now implies that (4.1.21) is also true for $r+1$ and thus for arbitrary N we have

$$(4.1.23) \quad \begin{aligned} & |\bar{u}_h^{n+1}|^2 + \delta \Theta \Delta t \alpha \nu \sum_{n=0}^N \|\bar{u}_h^{n+1}\|_h^2 + \Theta \Delta t \beta \nu \sum_{n=0}^N \|\bar{u}_h^m\|_h^2 \\ & + \left(\frac{1}{8} - \frac{\Theta \Delta t \beta \nu}{h^2} \right) \sum_{n=0}^N |\bar{u}_h^{n+1} - \bar{u}_h^{n+(1-\Theta)}|^2 \leq \Lambda_T. \end{aligned}$$

Inequalities (4.1.5), (4.1.23) and the regularity requirement (4.13) in the form

$$(\Theta_b \Delta t S(h))^2 \Lambda_T \leq D_5 (\Delta t S(h))^2 \Lambda_T \leq \Theta \Delta t \alpha \nu$$

yield

$$(4.1.24) \quad \begin{aligned} |\vec{u}_h^{n+\Theta}|^2 &\leq |\vec{u}_h^n|^2 + \Theta \Delta t \alpha \nu \sum_{n=0}^N \|\vec{u}_h^n\|_h^2 + \frac{\Theta \Delta t}{\nu} \sum_{n=0}^N |\vec{f}^n|^2 \leq \\ &\Lambda_T \left(1 + \frac{1}{\delta} + \frac{\Theta}{\nu} \right). \end{aligned}$$

The next estimate comes from (4.1.12), (4.1.24) and (4.13). We have

$$(4.1.25) \quad \begin{aligned} |\vec{u}_h^n|^2 &\leq \\ |\vec{u}_h^{n+\Theta}|^2 + D_5 \left(\left(\nu \frac{\Delta t}{h^2} \right)^2 + (\Delta t S(h))^2 \Lambda_T \right) \sum_{n=0}^N \|\vec{u}_h^{n+1}\|_h^2 + \frac{\Delta t}{\alpha \nu} |\vec{f}^n|^2 &\leq \\ \Lambda_T \left(1 + \frac{2}{\delta} + \frac{(1+\alpha)\Theta}{\alpha \nu} \right). \end{aligned}$$

Summing up (4.1.11) for $n = 0, 1, \dots, N$, we obtain by (4.1.23)

$$(4.1.26) \quad \begin{aligned} \sum_{n=0}^N |\vec{u}_h^{n+\Theta} - \vec{u}_h^n|^2 + \Theta \Delta t \alpha \nu \sum_{n=0}^N \|\vec{u}_h^{n+\Theta} - \vec{u}_h^n\|_h^2 &\leq \\ \Theta \Delta t \alpha \nu \sum_{n=0}^N \|\vec{u}_h^n\|_h^2 + \frac{\Theta \Delta t}{\alpha \nu} \sum_{n=0}^N |\vec{f}^n|^2 &\leq \Lambda_T \left(\frac{1}{\delta} + \frac{\Theta}{\alpha \nu} \right), \end{aligned}$$

and, also, from (4.1.26) and (4.1.5), we see that

$$(4.1.27) \quad \begin{aligned} \Theta \Delta t \alpha \nu \sum_{n=0}^N \|\vec{u}_h^{n+\Theta}\|_h^2 &\leq \\ \Theta \Delta t \alpha \nu \sum_{n=0}^N \|\vec{u}_h^{n+\Theta} - \vec{u}_h^n\|_h^2 + \Theta \Delta t \alpha \nu \sum_{n=0}^N \|\vec{u}_h^n\|_h^2 &\leq \Lambda_T \left(1 + \frac{2}{\delta} + \frac{\Theta}{\alpha \nu} \right). \end{aligned}$$

Expressing the results of (4.1.23-27) in terms of the functions $\vec{u}_{h,\Delta t}^{(i)}, i = 1, 2, 3$, we have

$$(4.1.28) \quad \begin{aligned} \sup_{t \in [0, T]} \left| \vec{u}_{h,\Delta t}^{(i)}(t) \right|^2 &\leq \Lambda_T \left(1 + \frac{2}{\delta} + \frac{2\Theta}{\alpha \nu} \right), \quad \text{for } i = 1, 2, 3, \\ \int_0^T \left\| \vec{u}_{h,\Delta t}^{(i)}(t) \right\|_h^2 dt &= \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \left\| \vec{u}_{h,\Delta t}^{(i)}(t) \right\|_h^2 dt = \Delta t \sum_{n=0}^{N-1} \left\| \vec{u}_h^{(k)}(t) \right\|_h^2 \leq \\ &\frac{\Lambda_T}{\Theta \alpha \beta \nu} \left(\frac{2}{\delta} + \frac{\Theta}{\alpha \nu} \right), \quad \text{for } i = 1, 2, 3, \end{aligned}$$

where $k = n$ for $i = 1$, $k = n + \Theta$ for $i = 2$ and $k = n + (1 - \Theta)$ for $i = 3$.

The a priori bound for $\vec{w}_{h,\Delta t}$ is obtained using (4.1.23) and the stability assumption (4.13), namely

$$(4.1.29) \quad \begin{aligned} & \left| \vec{u}_{h,\Delta t}^{(1)} - \vec{w}_{h,\Delta t} \right|_{L^2(0,T;[L^2(\Omega)]^n)}^2 = \\ & \frac{1}{(\Delta t)^2} \sum_{n=0}^{N-1} \left| \vec{u}_h^{n+1} - \vec{u}_h^n \right|^2 \int_{t_n}^{t_{n+1}} (t - t_n)^2 dt = \frac{1}{3} \Delta t \sum_{n=0}^{N-1} \left| \vec{u}_h^{n+1} - \vec{u}_h^n \right|^2 \leq \\ & 2\Delta t \sum_{n=0}^N \left| \vec{u}_h^{n+1} - \vec{u}_h^{n+(1-\Theta)} \right|^2 \leq 2\Delta t \frac{\Lambda_T}{\delta}. \end{aligned}$$

We infer from (4.1.28) and (4.1.29) that $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)} \subset V_h$ is a bounded set in $L^2(0,T;[L^2(\Omega)]^n)$ as it follows by the inequality

$$(4.1.30) \quad \int_0^T |\vec{w}_{h,\Delta t}|^2 dt \leq \int_0^T \left\| \vec{u}_{h,\Delta t}^{(1)} \right\|^2 dt + \int_0^T \left| \vec{u}_{h,\Delta t}^{(1)} - \vec{w}_{h,\Delta t} \right|^2 dt.$$

4.2 Convergence of the scheme. The results (4.1.28) and (4.1.29) justify the existence of functions $\vec{u}_*^{(i)}(t)$ and $\vec{w}_*$ such that as $(h, \Delta t) \rightarrow 0_+$ we have

$$(4.2.1) \quad \vec{u}_{h,\Delta t}^{(i)} \rightharpoonup \vec{u}_*^{(i)} \quad \begin{cases} \text{weakly in } L^2(0,T;V), \text{ conforming case,} \\ \text{weakly in } L^2(0,T;[L^2(\Omega)]^n), \text{ nonconforming case,} \\ \text{weakly-* in } L^\infty(0,T;[L^2(\Omega)]^n), \quad \text{for } i = 1, 2, \end{cases}$$

$$(4.2.2) \quad \vec{u}_{h,\Delta t}^{(3)} \rightharpoonup \vec{u}_*^{(3)} \quad \begin{cases} \text{weakly in } L^2(0,T;[H_0^1(\Omega)]^n), \text{ conforming case,} \\ \text{weakly in } L^2(0,T;[L^2(\Omega)]^n), \text{ nonconforming case,} \\ \text{weakly-* in } L^\infty(0,T;[L^2(\Omega)]^n). \end{cases}$$

Since $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)}$ is bounded in $L^2(0,T;[L^2(\Omega)]^n)$ by (4.1.30), we have

$$(4.2.3) \quad \vec{w}_{h,\Delta t} \rightharpoonup \vec{w}_*, \quad \text{weakly in } L^2(0,T;[L^2(\Omega)]^n).$$

The inequality (4.1.26) yields

$$(4.2.4) \quad \begin{aligned} & \left| \vec{u}_{h,\Delta t}^{(1)} - \vec{u}_{h,\Delta t}^{(2)} \right|_{L^2(0,T;[L^2(\Omega)]^n)}^2 = \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \left| \vec{u}_{h,\Delta t}^{(1)} - \vec{u}_{h,\Delta t}^{(2)} \right|^2 dt = \\ & \Delta t \sum_{n=0}^{N-1} \left| \vec{u}_h^n - \vec{u}_h^{n+\Theta} \right|^2 \longrightarrow 0, \quad \text{as } (h, \Delta t) \longrightarrow 0_+. \end{aligned}$$

Similarly, from (4.1.23), we have

$$(4.2.5) \quad \left| \vec{u}_{h,\Delta t}^{(1)} - \vec{u}_{h,\Delta t}^{(3)} \right|_{L^2(0,T;[L^2(\Omega)]^n)}^2 \longrightarrow 0, \quad \text{as } (h, \Delta t) \longrightarrow 0_+.$$

Now, from (4.1.29) and (4.2.3-5), we see that for almost all $t \in (0,T)$

$$(4.2.6) \quad \vec{u}_*^{(i)} = \vec{w}_* = \vec{u}_*, \quad \text{a.e. in } \Omega, \quad \text{for } i = 1, 2, 3.$$

We show that $\vec{w}_*$ is the solution of (WNS) by proving that the family $\{\vec{w}_{h,\Delta t}\}$ is relatively compact in, at least, $L^2(0,T;[L^2(\Omega)]^n)$. We treat the conforming and nonconforming approximations separately.

Conforming finite element approximations. We note that in this case the spaces V_h and H_h are subspaces of $[H_0^1(\Omega)]$. Integrating (4.12) over the time interval $[t, t + \gamma]$, for some $0 < \gamma < 1$, we obtain

$$\begin{aligned}
 (4.2.7) \quad & \frac{1}{\Theta} (\vec{w}_{h,\Delta t}(t + \gamma) - \vec{w}_{h,\Delta t}(t), \vec{\varphi}_h) + \alpha\nu \int_t^{t+\gamma} ((\vec{u}_{h,\Delta t}^{(1)}(s + \Delta t), \vec{\varphi}_h)) ds + \\
 & \beta\nu \int_t^{t+\gamma} ((\vec{u}_{h,\Delta t}^{(1)}(s), \vec{\varphi}_h)) ds = \frac{1}{\Theta\Delta t} \int_t^{t+\gamma} (\vec{u}_{h,\Delta t}^{(3)}(s) - \vec{u}_{h,\Delta t}^{(2)}(s), \vec{\varphi}_h) ds - \\
 & \beta\nu \int_t^{t+\gamma} ((\vec{u}_{h,\Delta t}^{(3)}(s), \vec{\varphi})) ds - \alpha\nu \int_t^{t+\gamma} ((\vec{u}_{h,\Delta t}^{(2)}(s), \vec{\varphi})) ds - \\
 & \int_t^{t+\gamma} \left(b(\vec{u}_{h,\Delta t}^{(2)}(s), \vec{u}_{h,\Delta t}^{(2)}(s), \vec{\varphi}_h) + b(\vec{u}_{h,\Delta t}^{(3)}(s), \vec{u}_{h,\Delta t}^{(3)}(s), \vec{\varphi}_h) \right) ds + \\
 & \int_t^{t+\gamma} (\vec{f}_{\Delta t}(s) + \vec{f}_{\Delta t}(s + \Delta t), \vec{\varphi}_h) ds.
 \end{aligned}$$

Now, we estimate the terms in (4.2.7). Recalling the estimates in (4.1.28), we get for $i = 1, 2, 3$

$$\begin{aligned}
 (4.2.8) \quad & \left| \int_t^{t+\gamma} ((\vec{u}_{h,\Delta t}^{(i)}(s), \vec{\varphi}_h)) ds \right| \leq \|\vec{\varphi}_h\| \int_t^{t+\gamma} \|\vec{u}_{h,\Delta t}^{(i)}(s)\| ds \leq \\
 & \sqrt{\gamma} \|\vec{\varphi}_h\| \left(\int_t^{t+\gamma} \|\vec{u}_{h,\Delta t}^{(i)}(s)\|^2 ds \right)^{\frac{1}{2}} \leq \\
 & \sqrt{\gamma} \|\vec{\varphi}_h\| \left(\int_0^T \|\vec{u}_{h,\Delta t}^{(i)}(s)\|^2 ds \right)^{\frac{1}{2}} \leq \sqrt{\frac{\Lambda_T}{\Theta\alpha\beta\nu}} \left(\frac{2}{\delta} + \frac{\Theta}{\alpha\nu} \right) \sqrt{\gamma} \|\vec{\varphi}_h\|.
 \end{aligned}$$

We majorize the trilinear forms by using the estimate (4.1.28), and we get

$$\begin{aligned}
 (4.2.9) \quad & \left| \int_t^{t+\gamma} b(\vec{u}_{h,\Delta t}^{(i)}(s), \vec{u}_{h,\Delta t}^{(i)}(s), \vec{\varphi}_h) ds \right| \leq \\
 & C \|\vec{\varphi}_h\| \int_t^{t+\gamma} \|\vec{u}_{h,\Delta t}^{(i)}(s)\|^{\frac{1}{2}} \|\vec{u}_{h,\Delta t}^{(i)}(s)\|^{\frac{3}{2}} ds \leq C \sqrt{\Lambda_T} \|\vec{\varphi}_h\| \int_t^{t+\gamma} \|\vec{u}_{h,\Delta t}^{(i)}(s)\|^{\frac{3}{2}} ds \leq \\
 & C \sqrt{\Lambda_T} \|\vec{\varphi}_h\| \sqrt[4]{\gamma} \left(\int_0^T \|\vec{u}_{h,\Delta t}^{(i)}(s)\|^2 ds \right)^{\frac{3}{4}} \leq C \Lambda_T^{\frac{5}{4}} \left(\frac{1}{\Theta\alpha\beta\nu} \left(\frac{2}{\delta} + \frac{\Theta}{\alpha\nu} \right) \right)^{\frac{3}{4}} \sqrt{\gamma} \|\vec{\varphi}_h\|.
 \end{aligned}$$

The first term on the right-hand side of (4.2.7) can be controlled by integrating (4.8) over the time interval $[0, T]$ with we $\vec{\varphi}_h \in V_h$. We have

$$\begin{aligned}
 (4.2.10) \quad & \left| \frac{1}{\Theta\Delta t} \int_t^{t+\gamma} (\vec{u}_{h,\Delta t}^{(3)}(s) - \vec{u}_{h,\Delta t}^{(2)}(s), \vec{\varphi}_h) ds \right| \leq \\
 & \beta\nu \left| \int_t^{t+\gamma} ((\vec{u}_{h,\Delta t}^{(3)}(s), \vec{\varphi}_h)) ds \right| + \alpha\nu \left| \int_t^{t+\gamma} ((\vec{u}_{h,\Delta t}^{(2)}(s), \vec{\varphi}_h)) ds \right| + \\
 & \left| \int_t^{t+\gamma} b_h(\vec{u}_{h,\Delta t}^{(3)}(s), \vec{u}_{h,\Delta t}^{(3)}(s), \vec{\varphi}_h) ds \right| + \left| \int_t^{t+\gamma} (\vec{f}_{\Delta t}(s), \vec{\varphi}_h) ds \right|.
 \end{aligned}$$

Applying the estimates (4.2.8-10) and (4.1.28) to (4.2.7) we obtain

$$(4.2.11) \quad |(\vec{w}_{h,\Delta t}(t+\gamma) - \vec{w}_{h,\Delta t}(t), \vec{\varphi}_h)| \leq C\sqrt{\gamma} \|\vec{\varphi}_h\|, \quad \forall \vec{\varphi}_h \in V_h,$$

where the constant $C > 0$ is independent of $\gamma, \Delta t$ and h . Integrating the above inequality for $t \in [0, T - \gamma]$ we obtain with $\vec{\varphi}_h = \vec{w}_{h,\Delta t}(t+\gamma) - \vec{w}_{h,\Delta t}(t) \in V_h$ the inequality

$$(4.2.12) \quad \int_0^{T-\gamma} |\vec{w}_{h,\Delta t}(s+\gamma) - \vec{w}_{h,\Delta t}(s)|^2 ds \leq C' \sqrt{\gamma \int_0^T \|\vec{w}_{h,\Delta t}^{(1)}(s)\|^2 ds} \leq C'' \sqrt{\gamma},$$

where again the constant $C'' > 0$ does not depend on $\gamma, \Delta t$ and h . We have $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)} \subset [H_0^1(\Omega)]^n \subset [L^2(\Omega)]^n$. Application of the compactness criterion [N, Theorem 2.3.1] and the inequality (4.2.12) imply that the set $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)}$ is relatively compact in $L^q(0, T; [L^2(\Omega)]^n)$, for $q \in [1, 2]$. Because the sequence $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)}$ converges weakly to $\vec{w}_*$ in $L^2(0, T; [L^2(\Omega)]^n)$ by (4.2.3), it converges strongly to this function in $L^q(0, T; [L^2(\Omega)]^n)$, for $q \in [1, 2]$, due to its relative compactness in this space. The strong convergence of the sequence $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)}$ in the space $L^2(0, T; [L^2(\Omega)]^n)$, its uniform bound in $L^\infty(0, T; [L^2(\Omega)]^n)$, and, the Lebesgue dominated convergence theorem, yield finally the strong convergence in $L^q(0, T; [L^2(\Omega)]^n)$, for $q \in [1, \infty)$.

To show the strong convergence in $L^2(0, T; [H_0^1(\Omega)]^n)$, we assume, for a moment, that $\vec{w}_*$ is the unique strong solution of (WNS). Integrating (WNS) over the time interval $[0, T]$ and taking $\vec{\varphi} = \vec{w}_* - \vec{w}_{h,\Delta t}$, we obtain

$$(4.2.13) \quad \begin{aligned} \int_0^T \|\vec{w}_* - \vec{w}_{h,\Delta t}\|^2 dt &= \int_0^T ((\vec{w}_{h,\Delta t}, \vec{w}_* - \vec{w}_{h,\Delta t})) dt - \\ &\int_0^T \left(\frac{d\vec{w}_*}{dt}, \vec{w}_* - \vec{w}_{h,\Delta t} \right) dt - \int_0^T b(\vec{w}_*, \vec{w}_*, \vec{w}_* - \vec{w}_{h,\Delta t}) dt + \int_0^T (\vec{f}, \vec{w}_* - \vec{w}_{h,\Delta t}) dt. \end{aligned}$$

Due to the strong convergence $\vec{w}_* - \vec{w}_{h,\Delta t} \rightarrow 0$ in $L^2(0, T; [L^2(\Omega)]^n)$ the last three terms converge to zero. We estimate similarly as in (4.1.28)

$$(4.2.14) \quad \int_0^T \|\vec{w}_{h,\Delta t}\|^2 dt \leq \frac{\Lambda_T}{\Theta \alpha \beta \nu} \left(\frac{2}{\delta} + \frac{\Theta}{\alpha \nu} \right).$$

Because of the compact imbedding of $L^2(0, T; [H_0^1(\Omega)]^n)$ into $L^2(0, T; [L^2(\Omega)]^n)$ and the already known strong convergence in $L^2(0, T; [L^2(\Omega)]^n)$, we conclude that

$$(4.2.15) \quad \vec{w}_{h,\Delta t} \rightharpoonup \vec{w}_*, \quad \text{weakly in } L^2(0, T; [H_0^1(\Omega)]^n).$$

Thus, for the first term on the right-hand side of (4.2.13) we have

$$\limsup_{(h,\Delta t) \rightarrow 0_+} \int_0^T ((\vec{w}_{h,\Delta t}, \vec{w}_* - \vec{w}_{h,\Delta t})) = - \limsup_{(h,\Delta t) \rightarrow 0_+} \int_0^T \|\vec{w}_{h,\Delta t}\|^2 + \int_0^T \|\vec{w}_*\|^2.$$

The previous strong convergence results and (4.1.28) yield

$$\limsup_{(h,\Delta t) \rightarrow 0_+} \int_0^T \|\vec{w}_{h,\Delta t}\|^2 \geq \limsup_{(h,\Delta t) \rightarrow 0_+} \int_0^T \|\vec{u}_{h,\Delta t}^{(1)}\|^2 \geq \int_0^T \|\vec{u}_*\|^2 = \int_0^T \|\vec{w}_*\|^2,$$

which implies together with (4.2.13) that

$$(4.2.16) \quad \vec{w}_{h,\Delta t} \longrightarrow \vec{w}_*, \quad \text{strongly in } L^2(0, T; [H_0^1(\Omega)]^n). \quad \square$$

Nonconforming finite element approximation. The relative compactness of the set $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)}$ is again essential to obtain the strong convergence. Extending the function $\vec{w}_{h,\Delta t}$ by zero outside the interval $[0, T]$, we show first that there exists a constant $C(\gamma) > 0$, independent of h and Δt , and, such that for $\gamma \in (0, \frac{1}{4})$, it holds

$$(4.2.17) \quad \int_{-\infty}^{+\infty} |\tau^{2\gamma}| |\hat{\vec{w}}_{h,\Delta t}(\tau)| d\tau \leq C(\gamma) \frac{\Lambda_T}{\Theta \alpha \beta \nu} \left(\frac{2}{\delta} + \frac{\Theta}{\alpha \nu} \right) < \infty,$$

where $\hat{\vec{w}}_{h,\Delta t}$ is the Fourier transform of $\vec{w}_{h,\Delta t}$. To show that the above inequality holds, we define a function $\vec{g}_{h,\Delta t} : (0, T) \rightarrow V_h$ by

$$(4.2.18) \quad \begin{aligned} ((\vec{g}_{h,\Delta t}, \vec{\varphi}_h))_h &= \frac{1}{\Delta t} (\vec{u}_{h,\Delta t}^{(3)}(t) - \vec{u}_{h,\Delta t}^{(2)}(t), \vec{\varphi}_h) - \\ &\Theta \alpha \nu ((\vec{u}_{h,\Delta t}^{(1)}(t + \Delta t), \vec{\varphi}_h))_h - \Theta \beta \nu ((\vec{u}_{h,\Delta t}^{(1)}(t), \vec{\varphi}_h))_h - \Theta \beta \nu ((\vec{u}_{h,\Delta t}^{(3)}(t), \vec{\varphi}_h))_h - \\ &\Theta \alpha \nu ((\vec{u}_{h,\Delta t}^{(2)}(t), \vec{\varphi}_h))_h - \Theta b_h (\vec{u}_{h,\Delta t}^{(2)}(t), \vec{u}_{h,\Delta t}^{(2)}(t), \vec{\varphi}_h) - \Theta b_h (\vec{u}_{h,\Delta t}^{(3)}(t), \vec{u}_{h,\Delta t}^{(3)}(t), \vec{\varphi}_h) + \\ &\Theta (\vec{f}_{\Delta t}(t) + \vec{f}_{\Delta t}(t + \Delta t), \vec{\varphi}_h), \quad \forall \vec{\varphi}_h \in V_h, \end{aligned}$$

and $t \in [n\Delta t, (n+1)\Delta t)$. Then, (4.12) can be interpreted as

$$\frac{d}{dt} (\vec{w}_{h,\Delta t}(t), \vec{\varphi}_h) = ((\vec{g}_{h,\Delta t}(t), \vec{\varphi}_h))_h, \quad \forall \vec{\varphi}_h \in V_h, \quad t \in (0, T).$$

We estimate

$$(4.2.19) \quad \begin{aligned} \frac{1}{\Theta} \int_0^T \|\vec{g}_{h,\Delta t}(t)\|_h dt &= \frac{1}{\Theta} \int_0^T \sup_{\|\vec{\varphi}_h\|_h=1} ((\vec{g}_{h,\Delta t}(t), \vec{\varphi}_h))_h dt \leq \\ &\frac{1}{\Delta t} \int_0^T |\vec{u}_{h,\Delta t}^{(3)}(t) - \vec{u}_{h,\Delta t}^{(2)}(t)| dt + 2\nu \int_0^T \left(\|\vec{u}_{h,\Delta t}^{(1)}(t)\|_h^2 + \|\vec{u}_{h,\Delta t}^{(3)}(t)\|_h^2 \right) dt + \\ &\beta \nu \int_0^T \|\vec{u}_{h,\Delta t}^{(3)}(t)\|_h^2 dt + \int_0^T \left(\|\vec{u}_{h,\Delta t}^{(2)}(t)\|_h^2 + \|\vec{u}_{h,\Delta t}^{(3)}(t)\|_h^2 \right) dt + 2 \int_0^T |\vec{f}_{\Delta t}(t)| dt. \end{aligned}$$

We conclude from (4.1.28), (4.2.4,5) and (4.2.19) that

$$(4.2.20) \quad \int_0^T \|\vec{g}_{h,\Delta t}(t)\|_h dt \leq C,$$

where the constant $C > 0$ does not depend on h and Δt .

Extending the function $\vec{g}_{h,\Delta t}$ from $[0, T]$ to $\mathbb{R}$ by zero and using (4.12), we get

$$(4.2.21) \quad \begin{aligned} \frac{d}{dt}(\vec{w}_{h,\Delta t}(t), \vec{\varphi}_h) = \\ ((\vec{g}_{h,\Delta t}(t), \vec{\varphi}_h))_h + (\vec{u}_{0h}, \vec{\varphi}_h)\delta_0 - (\vec{u}_h^N, \vec{\varphi}_h)\delta_T, \quad \forall \vec{\varphi}_h \in V_h. \end{aligned}$$

Here, δ_0, δ_T are the Dirac distributions at 0 and T , respectively, describing possibly occurring discontinuities at these points. Taking the Fourier transform of (4.2.21) we obtain

$$(4.2.22) \quad \begin{aligned} -2\pi i\tau(\hat{\vec{w}}_{h,\Delta t}(\tau), \vec{\varphi}_h) = \\ ((\hat{\vec{g}}_{h,\Delta t}(\tau), \vec{\varphi}_h))_h + (\vec{u}_{0h}, \vec{\varphi}_h) - (\vec{u}_h^N, \vec{\varphi}_h)\exp(-2\pi i\tau T). \end{aligned}$$

This equation implies

$$(4.2.23) \quad \begin{aligned} 2\pi|\tau| \left| \hat{\vec{w}}_{h,\Delta t}(\tau) \right|^2 \leq \left\| \hat{\vec{g}}_{h,\Delta t}(\tau) \right\|_h \left\| \hat{\vec{w}}_{h,\Delta t}(\tau) \right\|_h + |\vec{u}_{0h}| \left| \hat{\vec{w}}_{h,\Delta t}(\tau) \right| \leq \\ \int_0^T \left\| \hat{\vec{g}}_{h,\Delta t}(\tau) \right\|_h d\tau \left\| \hat{\vec{w}}_{h,\Delta t}(\tau) \right\|_h + |\vec{u}_{0h}| \left| \hat{\vec{w}}_{h,\Delta t}(\tau) \right|. \end{aligned}$$

This estimate and (4.2.20) imply the existence of a constant $C > 0$, independent of $h, \Delta t$ and γ , such that

$$|\tau| \left| \hat{\vec{w}}_{h,\Delta t} \right|^2 \leq C \left\| \hat{\vec{w}}_{h,\Delta t} \right\|_h.$$

Since $\gamma < \frac{1}{4}$, it holds

$$|\tau^{2\gamma}| \leq \frac{1 + |\tau|}{1 + |\tau|^{1-2\gamma}}, \quad \forall \tau \in \mathbb{R},$$

and thus, by Parseval equality and the two above inequalities, we have

$$(4.2.24) \quad \begin{aligned} \int_{-\infty}^{+\infty} |\tau^{2\gamma}| \left| \hat{\vec{w}}_{h,\Delta t}(t) \right| dt \leq C(\gamma) \int_0^T \left\| \vec{w}_h(\tau) \right\|_h^2 d\tau \leq \\ C(\gamma) \frac{\Lambda_T}{\Theta_{\alpha\beta\nu}} \left(\frac{2}{\delta} + \frac{\Theta}{\alpha\nu} \right). \end{aligned}$$

The last inequality follows from (4.1.28) and (4.1.29).

The relative compactness of the set $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)}$ in the space $L^2(0, T; [L^2(\Omega)]^n)$ is now assured by [T I, Theorem 2.2, Ch. III, §2], because of (4.1.30) and (4.2.24). Moreover, because $\{\vec{w}_{h,\Delta t}\}_{(h,\Delta t)}$ is bounded in $L^\infty(0, T; [L^2(\Omega)]^n)$, what follows by (4.1.28), the convergence is shown in the space $L^q(0, T; [L^2(\Omega)]^n)$, for $q \in [2, \infty)$. $\square$

Applying (4.2.4,5) and (4.1.29) we finally see that

$$(4.2.25) \quad \vec{u}_{h,\Delta t}^{(i)} \longrightarrow w_*, \quad \text{strongly in } L^q(0, T; [L^2(\Omega)]^n), \quad i = 1, 2, 3,$$

for $q \in [1, \infty)$ in the conforming case and for $q \in [2, \infty)$ in the nonconforming case. Moreover, this convergence holds in $L^2(0, T; [H_0^1(\Omega)]^n)$ for the conforming case, because

$$\int_0^T \left\| \vec{u}^{(i)}(t) - \vec{w}_*(t) \right\|_h^2 dt \leq \int_0^T \left(\left\| \vec{u}^{(i)}(t) - \vec{w}_{h,\Delta t}(t) \right\|_h^2 + \left\| \vec{w}_{h,\Delta t}(t) - \vec{w}_*(t) \right\|_h^2 \right) dt,$$

provided $\vec{w}_*$ is the unique solution of (WNS). This will be shown next.

Passage to the limit. We take $\psi \in C^1[0, T]$, with $\psi(T) = 0$, $\vec{\varphi}_h = \Pi_h \vec{v}$, for some $\vec{v} \in V_0 \cap [\mathcal{D}(\Omega)]^n$ and integrate (4.12) over the time interval $[0, T]$. Using the Green's theorem we obtain

$$\begin{aligned}
 (4.2.26) \quad & -\frac{1}{\Theta} \int_0^T (\vec{w}_{h,\Delta t}, \psi'(t) \Pi_h \vec{v}) dt + \alpha \nu \int_0^T ((\vec{u}_{h,\Delta t}^{(1)}(t + \Delta t), \psi(t) \Pi_h \vec{v}))_h dt + \\
 & \beta \nu \int_0^T ((\vec{u}_{h,\Delta t}^{(1)}(t), \psi(t) \Pi_h \vec{v}))_h dt = \frac{1}{\Theta \Delta t} \int_0^T (\vec{u}_{h,\Delta t}^{(3)}(t) - \vec{u}_{h,\Delta t}^{(2)}(t), \psi(t) \Pi_h \vec{v}) dt \\
 & - \beta \nu \int_0^T ((\vec{u}_{h,\Delta t}^{(3)}(t), \psi(t) \Pi_h \vec{v}))_h dt - \alpha \nu \int_0^T ((\vec{u}_{h,\Delta t}^{(2)}(t), \psi(t) \Pi_h \vec{v}))_h dt - \\
 & \int_0^T b_h(\vec{u}_{h,\Delta t}^{(2)}(t), \vec{u}_{h,\Delta t}^{(2)}(t), \psi(t) \Pi_h \vec{v}) dt - \int_0^T b_h(\vec{u}_{h,\Delta t}^{(3)}(t), \vec{u}_{h,\Delta t}^{(3)}(t), \psi(t) \Pi_h \vec{v}) dt + \\
 & \int_0^T (\vec{f}_{\Delta t}(t) + \vec{f}_{\Delta t}(t + \Delta t), \psi(t) \Pi_h \vec{v}) dt - \frac{1}{\Theta} (\vec{u}_{0h}, \Pi_h \vec{v}) \psi(0).
 \end{aligned}$$

Since, by (4.7), we have

$$\begin{aligned}
 & \frac{1}{\Theta \Delta t} \int_0^T (\vec{u}_{h,\Delta t}^{(3)}(t) - \vec{u}_{h,\Delta t}^{(2)}(t), \psi(t) \Pi_h \vec{v}) dt = -\frac{\Theta_b}{\Theta} \beta \nu \int_0^T ((\vec{u}_{h,\Delta t}^{(3)}(t), \psi(t) \Pi_h \vec{v}))_h dt - \\
 & \frac{\Theta_b}{\Theta} \int_0^T b_h(\vec{u}_{h,\Delta t}^{(3)}(t), \vec{u}_{h,\Delta t}^{(3)}(t), \psi(t) \Pi_h \vec{v}) dt - \frac{\Theta_b}{\Theta} \alpha \nu \int_0^T ((\vec{u}_{h,\Delta t}^{(2)}(t), \psi(t) \Pi_h \vec{v}))_h dt - \\
 & \frac{\Theta_b}{\Theta} \int_0^T (\vec{f}_{\Delta t}(t), \psi(t) \Pi_h \vec{v}) dt,
 \end{aligned}$$

we substitute the above into (4.2.26) and pass directly to the limit due to the strong convergence results (4.2.25). Using the condition (C) (section 3.2) and (4.5) for the approximation of the initial velocity, we obtain

$$\begin{aligned}
 & - \int_0^T (\vec{w}_*, \psi'(t) \vec{v}) dt + \int_0^T ((\vec{w}_*, \psi(t) \vec{v})) dt + \int_0^T b(\vec{w}_*, \vec{w}_*, \psi(t) \vec{v}) dt = \\
 & (\vec{u}(0), \vec{v}) \psi(0) - \int_0^T (\vec{f}(t), \psi(t) \vec{v}) dt.
 \end{aligned}$$

Consequently, (WNS) is satisfied in the distributional sense. Using the approximation of the initial velocity field and the Green's theorem it follows that the initial condition is also satisfied.

Convergence of the whole sequence $\{w_{h,\Delta t}\}_{(h,\Delta t)}$ to the solution of (WNS) is guaranteed by the assumed unicity of the strong solution on the given time interval $[0, T]$. This concludes the last step of the proof. $\square$

5. LID-DRIVEN CAVITY FLOW

In this section we use a benchmark problem originally described by O.Burggraf [B] to display some of the computational properties of the method. The test problem

consists in the solution of the Navier–Stokes equations in the square duct with prescribed shear and body forces. The shear force is non-zero on one face of the duct and zero on the other three. Since the velocity is set to be zero at the corners with the given shear it is a type of a regularized cavity problem. The details of this benchmark problem can be found in T.Shih *et al.* [S].

We use four groups of the two dimensional finite elements that satisfy the conditions (B)–(E). We restrict to the polynomials up to the order ≤ 2 and we use the following notation: P_k , k non-negative integer, for the space of polynomials up to order $\leq k$ and Q_k is the space of polynomials up to order $\leq k$ in each variable.

(I) Q_2/Q_1 element: Conforming piecewise biquadratic velocities and continues piecewise bilinear pressure. The element is referred to as Taylor–Hood finite element.

(II) Q_2/P_1 element: Conforming piecewise biquadratic velocities and discontinues piecewise linear pressure. We refer to this element also as Crouzeix–Raviart element.

(III) rQ_2/P_1 element: Conforming piecewise reduced biquadratic (serendipity family) velocities and discontinues piecewise linear pressure.

(IV) Q_1/P_0 element: Nonconforming piecewise bilinear velocities (continuity at midpoints of sides only) and piecewise constant pressure.

The accuracy for Taylor–Hood, Crouzeix–Raviart and rQ_1/P_1 elements is $\mathcal{O}(h^3) \times \mathcal{O}(h^2)$, and the accuracy is $\mathcal{O}(h^2) \times \mathcal{O}(h)$ for Q_1/P_0 element.

5.1 Description of the test problem. We consider $\Omega = [0, 1] \times [0, 1]$. We define the shear $\vec{g} = (g_1, g_2)$ by

$$\begin{aligned} g_1(x, y) &= \begin{cases} 16(x^4 - 2x^3 + x^2), & x \in \partial\Omega, y = 1 \\ 0, & \text{elsewhere} \end{cases} \\ g_2(x, y) &= 0, \quad x, y \in \partial\Omega. \end{aligned}$$

The body force $\vec{f} = (\vec{f}_1, \vec{f}_2)$ is given by

$$\begin{aligned} \vec{f}_1(x, y) &= 0, \quad x, y \in \Omega \\ \vec{f}_2(x, y) &= \nu \cdot 8 \left(24S(x) + 2s'(x)v''(y) + s'''(x)v(y) \right) + \\ &\quad \nu \cdot 64 \left(S_2(x)V(y) - s(y)s'(y)S_1(x) \right), \end{aligned}$$

where

$$\begin{aligned} s(x) &= x^4 - 2x^3 + x^2, \\ v(y) &= y^4 - y^2, \\ S(x) &= \int s(x) dx = \frac{1}{5}x^5 - \frac{1}{2}x^4 + \frac{1}{3}x^3, \end{aligned}$$

$$S_1(x) = s(x)s''(x) - (s'(x))^2 = -4x^6 + 12x^5 - 14x^4 + 8x^3 - 2x^2,$$

$$S_2(x) = \int s(x)s'(x) dx = \frac{1}{2}(s(x))^2,$$

$$V(x) = v(y)v'''(y) - v'(y)v''(y) = -24y^5 + 8y^3 - 4y.$$

In the above expressions the primes denote the differentiation with respect to x and y , respectively. The solution $\{(u_1^\infty, u_2^\infty), p^\infty\}$ to the stationary Navier–Stokes equations with the above given data is given by:

$$u_1^\infty = 8s(x)v'(y) = 8(x^4 - 2x^3 + x^2)(4y^3 - 2y),$$

$$u_2^\infty = -8s'(x)v(y) = -8(4x^3 - 6x^2 + 2x)(y^4 - y^2),$$

$$p^\infty = \nu \cdot 8(S(x)v'''(y) + s'(x)v'(y)) + 64(S_2(x)(v(y)v''(y) - (v'(y))^2)).$$

Fig. 1: Velocity field, $\nu = 0.01[m^{-2}s^{-1}]$

Fig. 2: Maximum error distribution, $\nu = 0.01[m^{-2}s^{-1}]$

Numerical results. We have computed the test problem for $\nu = 0.01[m^{-2}s^{-1}]$ and $\nu = 0.001[m^{-2}s^{-1}]$. The later case is far beyond the validity range of the analytical solution of the test problem because there is presented permanent time-dependent regime. We display the plots of the velocity field, pressure and error distribution for $\nu = 0.01[m^{-2}s^{-1}]$. Moreover, we indicate the behavior of various errors and $|\vec{u}_t(t)|$. For the case with lower viscosity, the computed pressure seems to be in a good agreement with the analytical solution and this is true even for $\nu = 0.0001[m^{-2}s^{-1}]$, where the velocity field is different from the analytical solution. The numerical results suggest the use of this method for the values of up to $Re \sim 3000$, behind this limit the convergence namely of the Burgers step is very slow (rate about .95) because of oscillations produced by a very weak dumping of the method.

In all the computations, we use the finite element approximations of the group (I,II) and only for a lower viscosity $\sim 0.1[m^{-2}s^{-1}]$ the other approximations seems to be

Fig. 3: Pressure isolines, $\nu = 0.01[m^{-2}s^{-1}]$

Fig. 4: Errors behavior, $\nu = 0.01[m^{-2}s^{-1}]$

acceptable. We did the computations on the scale resolution $h = 1/64$ and we used the time step $\Delta t = 0.1s$, fractional parameter $\theta = 0.25$, splitting parameter $\alpha = 0.5$ (for this sort of problems) and the integration was performed by the 4×4 Gaussian quadrature formula.

Acknowledgement. The authors wish to thank Prof.Dr.P.Rys, LTC, ETH-Zürich and the members of his research group for the hospitality and many valuable discussions. Furthermore thanks are due to Prof.Dr.H.Blum and Dr.J.Harik of Prof.Dr.R.Rannacher's numerical group at University Heidelberg for the software support by FEAT.

REFERENCES

- [B]. O.Burrgraf, *Analytical and numerical studies of the structure of steady separated flows*, J.Fluid Mech. 24 (1966), 113–151.
- [C]. M.Crouzeix, *Approximations et Méthodes Itératives de Résolution d'Inéquations Variationnelles et de Problèmes Non Linéaires*, Cah. de l'IRIA, vol. 12,, 1974.
- [CFT]. P.Constantin,C.Foias,R.Temam, *Attractors representing turbulent flows*, Memoirs of the American Mathematical Society 53(314) (1985).
- [G I]. R.Glowinski, *Numerical Methods for Nonlinear Variational Problems*, Springer-Verlag, 1984.
- [G II]. R.Glowinski, *Le Θ schema*, In: M.O.Bristeau, R.Glowinski, J.Perieux, Numerical methods for the Navier–Stokes equations, Comp. Phy. report 6 (1987), 73–187.
- [H]. J.G.Heywood, *Classical solutions of the Navier–Stokes Equations*, In: Lect. Notes in Math. 771.
- [HR I]. J.G.Heywood, R.Rannacher, *Finite element approximation of the nonstationary Navier–Stokes problem. Part I: Regularity of solutions and second-order error estimates for spatial discretization*, SIAM J. Numer. Anal. 19(2) (1982), 275–311.
- [HR II]. J.G.Heywood, R.Rannacher, *Finite element approximation of the nonstationary Navier–Stokes problem. Part II: Stability of solutions and error estimates uniform in time*, SIAM J. Numer. Anal. 23(4) (1986), 750–777.
- [HR III]. J.G.Heywood, R.Rannacher, *Finite element approximation of the nonstationary Navier–Stokes problem. Part III: Smoothing property and higher order error estimates for spatial discretization*, SIAM J. Numer. Anal. 25(3) (1988), 489–512.
- [L]. O.A.Ladyženskaya, *The mathematical theory of viscous incompressible flow*, Gordon and Breach, New York, 1962.
- [N]. J.Nečas, *Introduction to the theory of Nonlinear Elliptic Equations*, Taubner-Texte zur Mathematik, Band 52, Leipzig, 1983.
- [R]. R.Rannacher, *Numerical analysis of nonstationary fluid flow (A survey)*, Preprint 492 (1988), University Heidelberg.
- [S]. T.M.Shih, H.Tan, B.C.Hwang, *Effects of grid staggering on numerical schemes*, Int. J. Numer. Meth. Fluids 9 (1989), 193–212.
- [T I]. R.Temam, *Navier–Stokes Equations: Theory and numerical analysis*, North Holland, Amsterdam, New York, 1977.
- [T II]. R.Temam, *Navier–Stokes Equations and nonlinear functional analysis*, CBMS-NSF Regional conference series in applied mathematics, Soc. for industr. and appl. math., 1983.
- [T III]. R.Temam, *Inertial Manifolds and Multigrid Method*, Siam, J. Math. Anal 21(1) (1990), 154–178.

206 CHURCH ST. SE, SCHOOL OF MATHEMATICS, UNIVERSITY
OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455 USA

Email: kloucek@math.umn.edu

LABORATORIUM FÜR TECHNISCHE CHEMIE, Eidgenössische
TECHNISCHE HOCHSCHULE, CH-8092 ZÜRICH, SWITZERLAND,

Email: U4567@CZHETH5A.BITNET

#	Author/s	Title
1078	Michael Böhm & Mario Taboada,	Global existence and regularity of solutions of the nonlinear string equation
1079	Zhangxin Chen,	BDM mixed methods for a nonlinear elliptic problem
1080	J.J.L. Velázquez,	On the dynamics of a closed thermosyphon
1081	Frédéric Bonnans & Eduardo Casas,	Some stability concepts and their applications in optimal control problems
1082	Hong-Ming Yin,	$L^{2,\mu}(Q)$ -estimates for parabolic equations and applications
1083	David L. Russell & Bing-Yu Zhang,	Smoothing and decay properties of solutions of the Korteweg-de Vries equation on a periodic domain with point dissipation
1084	J.E. Dunn & K.R. Rajagopal,	Fluids of differential type: Critical review and thermodynamic analysis
1085	Mary Elizabeth Bradley & Mary Ann Horn,	Global stabilization of the von Kármán plate with boundary feedback acting via bending moments only
1086	Mary Ann Horn & Irena Lasiecka,	Global stabilization of a dynamic von Kármán plate with nonlinear boundary feedback
1087	Vilmos Komornik,	Decay estimates for a petrovski system with a nonlinear distributed feedback
1088	Jesse L. Barlow,	Perturbation results for nearly uncoupled Markov chains with applications to iterative methods
1089	Jong-Shenq Guo,	Large time behavior of solutions of a fast diffusion equation with source
1090	Tongwen Chen & Li Qiu,	$\mathcal{H}_\infty$ design of general multirate sampled-data control systems
1091	Satyanad Kichenassamy & Walter Littman,	Blow-up surfaces for nonlinear wave equations, I
1092	Nahum Shimkin,	Asymptotically efficient adaptive strategies in repeated games, Part I: certainty equivalence strategies
1093	Caroline Fabre, Jean-Pierre Puel & Enrique Zuazua,	On the density of the range of the semigroup for semilinear heat equations
1094	Robert F. Stengel, Laura R. Ray & Christopher I. Marrison,	Probabilistic evaluation of control system robustness
1095	H.O. Fattorini & S.S. Sritharan,	Optimal chattering controls for viscous flow
1096	Kathryn E. Lenz,	Properties of certain optimal weighted sensitivity and weighted mixed sensitivity designs
1097	Gang Bao & David C. Dobson,	Second harmonic generation in nonlinear optical films
1098	Avner Friedman & Chaocheng Huang,	Diffusion in network
1099	Xinfu Chen, Avner Friedman & Tsuyoshi Kimura,	Nonstationary filtration in partially saturated porous media
1100	Walter Littman & Baisheng Yan,	Rellich type decay theorems for equations $P(D)u = f$ with f having support in a cylinder
1101	Satyanad Kichenassamy & Walter Littman,	Blow-up surfaces for nonlinear wave equations, II
1102	Nahum Shimkin,	Extremal large deviations in controlled I.I.D. processes with applications to hypothesis testing
1103	A. Narain,	Interfacial shear modeling and flow predictions for internal flows of pure vapor experiencing film condensation
1104	Andrew Teel & Laurent Praly,	Global stabilizability and observability imply semi-global stabilizability by output feedback
1105	Karen Rudie & Jan C. Willems,	The computational complexity of decentralized discrete-event control problems
1106	John A. Burns & Ruben D. Spies,	A numerical study of parameter sensitivities in Landau-Ginzburg models of phase transitions in shape memory alloys
1107	Gang Bao & William W. Symes,	Time like trace regularity of the wave equation with a nonsmooth principal part
1108	Lawrence Markus,	A brief history of control
1109	Richard A. Brualdi, Keith L. Chavey & Bryan L. Shader,	Bipartite graphs and inverse sign patterns of strong sign-nonsingular matrices
1110	A. Kersch, W. Morokoff & A. Schuster,	Radiative heat transfer with quasi-monte carlo methods
1111	Jianhua Zhang,	A free boundary problem arising from swelling-controlled release processes
1112	Walter Littman & Stephen Taylor,	Local smoothing and energy decay for a semi-infinite beam pinned at several points and applications to boundary control
1113	Srdjan Stojanovic & Thomas Svobodny,	A free boundary problem for the Stokes equation via nonsmooth analysis
1114	Bronislaw Jakubczyk,	Filtered differential algebras are complete invariants of static feedback
1115	Boris Mordukhovich,	Discrete approximations and refined Euler-Lagrange conditions for nonconvex differential inclusions
1116	Bei Hu & Hong-Ming Yin,	The profile near blowup time for solution of the heat equation with a nonlinear boundary condition
1117	Jin Ma & Jiongmin Yong,	Solvability of forward-backward SDEs and the nodal set of Hamilton-Jacobi-Bellman Equations
1118	Chaocheng Huang & Jiongmin Yong,	Coupled parabolic and hyperbolic equations modeling age-dependent epidemic dynamics with nonlinear diffusion

- 1119 **Jiongmin Yong**, Necessary conditions for minimax control problems of second order elliptic partial differential equations
- 1120 **Eitan Altman & Nahum Shimkin**, Worst-case and Nash routing policies in parallel queues with uncertain service allocations
- 1121 **Nahum Shimkin & Adam Schwartz**, Asymptotically efficient adaptive strategies in repeated games, part II: Asymptotic optimality
- 1122 **M.E. Bradley**, Well-posedness and regularity results for a dynamic Von Kármán plate
- 1123 **Zhangxin Chen**, Finite element analysis of the 1D full drift diffusion semiconductor model
- 1124 **Gang Bao & David C. Dobson**, Diffractive optics in nonlinear media with periodic structure
- 1125 **Steven Cox & Enrique Zuazua**, The rate at which energy decays in a damped string
- 1126 **Anthony W. Leung**, Optimal control for nonlinear systems of partial differential equations related to ecology
- 1127 **H.J. Sussmann**, A continuation method for nonholonomic path-finding problems
- 1128 **Yung-Jen Guo & Walter Littman**, The null boundary controllability for semilinear heat equations
- 1129 **Q. Zhang & G. Yin**, Turnpike sets in stochastic manufacturing systems with finite time horizon
- 1130 **I. Györi, F. Hartung & J. Turi**, Approximation of functional differential equations with time- and state-dependent delays by equations with piecewise constant arguments
- 1131 **I. Györi, F. Hartung & J. Turi**, Stability in delay equations with perturbed time lags
- 1132 **F. Hartung & J. Turi**, On the asymptotic behavior of the solutions of a state-dependent delay equation
- 1133 **Pierre-Alain Gremaud**, Numerical optimization and quasiconvexity
- 1134 **Jie Tai Yu**, Resultants and inversion formula for N polynomials in N variables
- 1135 **Avner Friedman & J.L. Velázquez**, The analysis of coating flows in a strip
- 1136 **Eduardo D. Sontag**, Control of systems without drift via generic loops
- 1137 **Yuan Wang & Eduardo D. Sontag**, Orders of input/output differential equations and state space dimensions
- 1138 **Scott W. Hansen**, Boundary control of a one-dimensional, linear, thermoelastic rod
- 1139 **Robert Lipton & Bogdan Vernescu**, Homogenization of two phase emulsions with surface tension effects
- 1140 **Scott Hansen & Enrique Zuazua**, Exact controllability and stabilization of a vibrating string with an interior point mass
- 1141 **Bei Hu & Jiongmin Yong**, Pontryagin Maximum principle for semilinear and quasilinear parabolic equations with pointwise state constraints
- 1142 **Mark H.A. Davis**, A deterministic approach to optimal stopping with application to a prophet inequality
- 1143 **M.H.A. Davis & M. Zervos**, A problem of singular stochastic control with discretionary stopping
- 1144 **Bernardo Cockburn & Pierre-Alain Gremaud**, An error estimate for finite element methods for scalar conservation laws
- 1145 **David C. Dobson & Fadil Santosa**, An image enhancement technique for electrical impedance tomography
- 1146 **Jin Ma, Philip Protter, & Jiongmin Yong**, Solving forward-backward stochastic differential equations explicitly — a four step scheme
- 1147 **Yong Liu**, The equilibrium plasma subject to skin effect
- 1148 **Ulrich Hornung**, Models for flow and transport through porous media derived by homogenization
- 1149 **Avner Friedman, Chaocheng Huang, & Jiongmin Yong**, Effective permeability of the boundary of a domain
- 1150 **Gang Bao**, A uniqueness theorem for an inverse problem in periodic diffractive optics
- 1151 **Angelo Favini, Mary Ann Horn, & Irena Lasiecka**, Global existence and uniqueness of regular solutions to the dynamic von Kármán system with nonlinear boundary dissipation
- 1152 **E.G. Kalnins & Willard Miller, Jr.**, Models of q -algebra representations: q -integral transforms and “addition theorems”
- 1153 **E.G. Kalnins, V.B. Kuznetsov & Willard Miller, Jr.**, Quadrics on complex Riemannian spaces of constant curvature, separation of variables and the Gaudin magnet
- 1154 **A. Kersch, W. Morokoff & Chr. Werner**, Selfconsistent simulation of sputtering with the DSMC method
- 1155 **Bing-Yu Zhang**, A remark on the Cauchy problem for the Korteweg-de Vries equation on a periodic domain
- 1156 **Gang Bao**, Finite element approximation of time harmonic waves in periodic structures
- 1157 **Tao Lin & Hong Wang**, Recovering the gradients of the solutions of second-order hyperbolic equations by interpolating the finite element solutions
- 1158 **Zhangxin Chen**, L^p -posteriori error analysis of mixed methods for linear and quasilinear elliptic problems
- 1159 **Todd Arbogast & Zhangxin Chen**, Homogenization of compositional flow in fractured porous media
- 1160 **L. Qiu, B. Bernhardsson, A. Rantzer, E.J. Davison, P.M. Young & J.C. Doyle**, A formula for computation of the real stability radius
- 1161 **Maria Inés Treparsky**, Adaptive control of linear discrete time systems with external disturbances under inaccurate modelling: A case study
- 1162 **Petr Klouček & Franz S. Rys**, Stability of the fractional step Θ -scheme for the nonstationary Navier-Stokes equations
- 1163 **Eduardo Casas, Luis A. Fernández & Jiongmin Yong**, Optimal control of quasilinear parabolic equations
- 1164 **Darrell Duffie, Jin Ma & Jiongmin Yong**, Black’s consol rate conjecture
- 1165 **D.G. Aronson & J.L. Vazquez**, Anomalous exponents in nonlinear diffusion