**NEW COARSE GRID OPERATORS FOR HIGHLY
OSCILLATORY COEFFICIENT ELLIPTIC PROBLEMS**

By

Bjorn Engquist

and

Erding Luo

IMA Preprint Series # 1305

March 1995

**INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
UNIVERSITY OF MINNESOTA**

**514 Vincent Hall
206 Church Street S.E.
Minneapolis, Minnesota 55455**

NEW COARSE GRID OPERATORS FOR HIGHLY OSCILLATORY COEFFICIENT ELLIPTIC PROBLEMS

Bjorn Engquist* Erding Luo †

Abstract

New coarse grid operators are developed for elliptic problems with highly oscillatory coefficients. These types of operators are necessary when the character of the differential equation on coarser grids (or longer wavelengths) is different from that on the fine grid. Elliptic problems for composite materials and different classes of hyperbolic problems are practical examples.

The new coarse grid operators can be constructed directly based on the homogenized differential operators or hierarchically computed from the finest grid. Numerical examples are presented showing that the homogenized form of the equations is very useful in the design of coarse grid operators for multigrid methods.

1 INTRODUCTION

Multigrid methods are usually not effective when applied to problems for which the standard coarse grid operators have significantly different properties from those of the fine grid operators [1,3,5,7,8]. In some of these problems the coarse grid operators should be constructed based on other

*Department of Mathematics, University of California at Los Angeles, LA, CA90024. Research supported by ARPA/ONR N00014-92-J-1890 and NSF DMS91-03104.

†IMA, University of Minnesota, Mpls, MN 55415. Research supported by NSF through IMA.

principles than just simple restriction from the finest grid in order to restore the high efficiency of the multigrid method. Elliptic and parabolic equations with strongly variable coefficients and some hyperbolic equations are such problems. One feature in these problems is that the smallest eigenvalues in absolute value do not correspond to very smooth eigenfunctions. It is thus not easy to represent these eigenfunctions on the coarser grids.

We shall investigate elliptic equations with highly oscillatory coefficients,

$$\sum_j \frac{\partial}{\partial x_j} a_j(x, \frac{x}{\epsilon}) \frac{\partial}{\partial x_j} u_\epsilon(x) = f(x), \quad (1)$$

with $a_j(x, \eta)$ strictly positive, continuous and 1-periodic in each component of η . The parameter $\epsilon > 0$ should be regarded as small. This is one class of the problems discussed above for which there exists a fairly complete analytic theory such that a rigorous treatment is possible. This homogenization theory describes the dependence of the large scale features in the solutions on the smaller scales in the coefficients [2]. We shall consider model problems but there are also important practical applications of these equations in the study of elasticity and heat conduction for composite materials.

In this paper we analyze the multigrid methods for equation (1) by introducing new coarse grid operators, based on local or global homogenized operators of the equation. These operators can be numerically calculated from the finest grid operator, by solving a so-called cell problem [2].

The rest of the paper is organized as follows: we introduce our model problem and the theory of homogenization in section 2. In section 3, we briefly outline the multigrid method, then we derive our new coarse grid operators and discuss the convergence theory. In the last section, we present our numerical experiments. In a number of numerical tests we compare the convergence rate for different choices of parameter and coarse grid operators applied to a two dimensional elliptic model problem. We also consider a more realistic problem of heat conduction in a composite structure.

2 Model Problem

2.1 The Partial Differential Equation

Elliptic problems of the form

$$-\nabla \cdot a^\epsilon(x, y) \nabla u_\epsilon = f(x, y), \quad (x, y) \in \Omega = [0, 1] \times [0, 1], \quad (2)$$

will be considered, subject to Dirichlet boundary condition

$$u_\epsilon|_{\partial\Omega} = 0.$$

The function $a^\epsilon(x, y) = a(x/\epsilon, y/\epsilon)$ is strictly positive. And, $f(x, y)$ in the following is always assumed to be smooth. Since it contains a small parameter ϵ , the coefficient highly varies. The small parameter ϵ is the length of oscillation. From homogenization theory [2], it follows,

$$\max_{(x, y) \in \Omega} |u_\epsilon - u| \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0,$$

where u satisfies the following effective equation, which doesn't contain any oscillatory coefficients,

$$-A_{11} \frac{\partial^2 u}{\partial x^2} - (A_{12} + A_{21}) \frac{\partial^2 u}{\partial y \partial x} - A_{22} \frac{\partial^2 u}{\partial y^2} = f(x, y), \quad (x, y) \in \Omega, \quad (3)$$

subject to the same boundary condition. Here, the constant coefficients can be calculated from the following formula,

$$A_{ij} = \int_{\Omega} a(s_1, s_2) \left(\delta_{ij} - \frac{\partial \kappa_j}{\partial s_i} \right) ds_1 ds_2, \quad i, j = 1, 2,$$

and the auxiliary periodic functions κ_j are given by,

$$-\nabla_s \cdot a(s_1, s_2) \nabla_s \kappa_j = \frac{\partial a(s_1, s_2)}{\partial s_j}, \quad j = 1, 2.$$

For simplicity, we consider a model problem. That is, the equation (2) with diagonal oscillatory coefficient,

$$a^\epsilon(x, y) = a\left(\frac{x - y}{\epsilon}\right). \quad (4)$$

From (3), we know that the corresponding homogenized equation is,

$$-\frac{(\mu + \bar{a})}{2} \frac{\partial^2 u}{\partial x^2} + (\mu - \bar{a}) \frac{\partial^2 u}{\partial x \partial y} - \frac{(\mu + \bar{a})}{2} \frac{\partial^2 u}{\partial y^2} = f(x, y), \quad (x, y) \in \Omega, \quad (5)$$

where μ represents the harmonic average of coefficient $a^\epsilon(x, y)$, denoted by $\mu = m(1/a^\epsilon)^{-1}$; and $\bar{a}$ represents the arithmetical average of coefficient $a^\epsilon(x, y)$, denoted by $\bar{a} = m(a^\epsilon)$. Here, the mean value $m(f)$ of a ϵ -periodic function is defined as,

$$m(f) = \frac{1}{\epsilon} \int_0^\epsilon f(x) dx.$$

2.2 Finite Difference Discretization

We discretize the unit square Ω into $N \times N$ equal cells with $(N-1) \times (N-1)$ grid points by taking grid step size h in both the x and y directions, where $h = \frac{1}{N}$. Particularly, we take such grid size h , which is smaller but has the same order as the wavelength ϵ . What's more, we require that the ratio of the wavelength to the grid size to be an irrational number, [4-6,8]. We call such a case the ergodic case, in which the difference operator would sample quite well the original differential operator. This is required in order to achieve the convergence of the different scheme to the solution of the differential equation, [4].

Denote $x_i = ih, y_j = jh$, and

$$a_{ij}^h = a^\epsilon(x_i - \frac{h}{2}, y_j) = a^\epsilon(x_i - \frac{h}{2} - y_j), \quad (6)$$

$$b_{ij}^h = a^\epsilon(x_i, y_j - \frac{h}{2}) = a^\epsilon(x_i - y_j + \frac{h}{2}), \quad (7)$$

$$f_{ij}^h = f(x_i, y_j), \quad i, j = 0, \dots, N. \quad (8)$$

The standard 5-point finite difference stencil of (2) and (4) at the h -grid level is

$$-D_+^i a_{ij}^h D_-^i u_{ij}^h - D_+^j b_{ij}^h D_-^j u_{ij}^h = f_{ij}^h. \quad (9)$$

D_+^i and D_-^i are the standard forward and backward divided differences in x direction, respectively; similarly, D_+^j and D_-^j are for y direction.

Using vector notation, we can rewrite (9) as

$$L_h U_h = F_h, \quad (10)$$

where L_h, A_j^h are tridiagonal matrices, written as

$$L_h = \frac{1}{h^2} \text{Tridiag}[B_{j-1}^h, A_j^h, B_j^h]_{j=1, \dots, N-1}, \quad (11)$$

$$A_j^h = \text{Tridiag}[-a_{i-1,j}^h, a_{i-1,j}^h + a_{ij}^h + b_{ij}^h + b_{ij-1}^h, -a_{ij}^h]_{i=1, \dots, N-1},$$

and B_j^h is a diagonal matrix, $B_j^h = \text{Diag}[-b_{i,j}^h]_{i=1, \dots, N-1}$, $j = 1, \dots, N-1$. Also,

$$U_h = (u_{11}^h, u_{21}^h, \dots, u_{N-1,1}^h, \dots, u_{1N-1}^h, u_{2N-1}^h, \dots, u_{N-1,N-1}^h)^T,$$

$$F_h = (f_{11}^h, f_{21}^h, \dots, f_{N-1,1}^h, \dots, f_{1N-1}^h, f_{2N-1}^h, \dots, f_{N-1,N-1}^h)^T.$$

3 The Multigrid Method

3.1 The Algorithm

The application of the two-level multigrid method to the equation (9) at the n -th iteration usually consists of the following three steps,

- (1) Pre-smoothing : Compute $U_h^{n+\frac{1}{2}}$ by applying γ_1 steps of a given iteration method to (9) that starts with U_h^n at the finest grid. For convenience, we introduce the following notation:

$$U_h^{n+\frac{1}{2}} = S^{\gamma_1}(U_h^n, L_h, f_h);$$

- (2) Coarser grid correction :

- Restrict the residual: $d_H = I_h^H(f_h - L_h U_h^{n+\frac{1}{2}})$;
- Solve the correction e_H : $L_H e_H = d_H$;
- Interpolate: $\tilde{U} = U_h^{n+\frac{1}{2}} + I_H^h e_H$;

- (3) Post-smoothing:

$$U_h^{n+1} = S^{\gamma_2}(\tilde{U}, L_h, f_h).$$

From the above, one can immediately obtain the iteration operator M of the two-grid multigrid method:

$$M = S^{\gamma_2}(I - I_H^h L_H^{-1} I_h^H L_h) S^{\gamma_1}. \quad (12)$$

For the full multigrid method, instead of solving the correction exactly on the coarse H -grid level, we solve it by introducing another coarser grid, and applying the two-grid level multigrid method again. The same procedure can be repeated several times, until the coarsest level. In this paper, we always take the coarse grid step size to be as twice as the previous one. And, we consider the full multigrid cycle, denoted by $V(\gamma_1, \gamma_2)$, with γ_1 pre-smoothing steps and γ_2 post-smoothing steps at each level, till the coarsest grid level with grid step size $\frac{1}{2}$.

For a equation with smoothing coefficients, the standard coarse grid operator approximates quite well the fine grid operator, which, therefore, makes the spectral radius of M small. However, for a partial differential equation with non-smoothing coefficients, this is no longer true. In order to restore the high efficiency of multigrid methods, certain modifications are necessarily needed. For instance, appropriate smoothing iteration methods (such as line Gauss-Seidel method), combined with adapted interpolation operator are often suggested to interface problems. [1, 3]. Meanwhile, to ensure the approximation of the coarse grid operator to the fine grid operator, the variational coarse grid operator is recommended. Nevertheless, the coarse grid operator constructed in this way is awkward and costly. What we do in this paper is to introduce a new principle to define the coarse grid operator, so that a fast multigrid method can be obtained to solve the numerical solution of elliptic equations (1) with highly oscillatory coefficients.

3.2 Construction of Coarse Grid Operators

By considering the homogenized equation and the asymptotic behavior of eigenvalue problem [1,2,6,8,9], we realize that the small eigenvalues of the original oscillatory operator can be approximated by the corresponding homogenized eigenvalues. After a few steps of fine grid smoothing, the error is dominated by the low frequency modes, which can be approximated by the corresponding homogenized ones at the coarser grid level. Based on this idea, we construct the coarse grid operator directly from the homogenized operator. There are two ways to define it.

(1) Global Homogenized Coarse Grid Operator:

If we construct the coarse grid operator directly from the discretization of the corresponding homogenized operator (5), we get the *Global Homogenized Coarse Grid Operator* at H -grid level,

$$L_H = [-\frac{(\mu + \bar{a})}{2}D_+^i D_-^i - \frac{(\mu + \bar{a})}{2}D_+^j D_-^j + \theta(\mu - \bar{a})D_0^i D_0^j]_{i,j=1,\dots,\frac{1}{H}-1}, \quad (13)$$

where D_0 denotes the standard center divided difference. And, θ is a parameter.

However, as we may notice, there are many intermediate frequency modes which could still exist in the error in the process of the multigrid method after fine grid smoothing, unless a large number of smoothing iterations are used. These modes can not be approximated by the homogenized eigenmodes at the coarse grid level. Directly applying the global homogenized coarse grid operator is not very suitable in such a situation, and may delay the convergent rate. Based on this, a revised coarse grid operator is then introduced as follows.

(2) Local Homogenized Coarse Grid Operator:

In order to approximate the fine grid operator better, we construct another grid operator, called here the *Local Homogenized Coarse Grid Operator*, which still maintains the form of the homogenized operator as the global homogenized coarse grid operator does. But, the coefficients here are calculated locally. To show details of the derivation, we first divide the whole domain into many cells. For instance, at point (i, j) on the coarse H -grid level (see Figure 1 and 2), we define 4 cells, denoted by **EH**, **WH**, **SH**, **NH**. We calculate the homogenized operator in each cell, and then use it to define the coarse grid operator. For our model problem, the coarse grid operator at H -grid level has the following form,

$$L_H = [-D_+^i a_{ij}^H D_-^i - D_+^j b_{ij}^H D_-^j + \theta D_0^i c_{ij}^H D_0^j + \theta D_0^j c_{ij}^H D_0^i]_{i,j=1,\dots,\frac{1}{H}-1}, \quad (14)$$

where the coefficients are calculated as follows.

The coefficient a_{ij}^H is the coefficient of $\frac{\partial^2 u}{\partial x^2}$ in the homogenized equation (5) generated in **WH**-cell. That is,

$$a_{ij}^H = \frac{1}{2}(\mu(WH) + \bar{a}(WH)),$$

where $\bar{a}(WH)$ and $\mu(WH)$ denote the arithmetical average and the harmonic average of all a_{ij}^h, b_{ij}^h in **WH**-cell, respectively.

The coefficient b_{ij}^H is the coefficient of $\frac{\partial^2 u}{\partial y^2}$ in the homogenized equation (5) generated in **SH**-cell. That is,

$$b_{ij}^H = \frac{1}{2}(\mu(SH) + \bar{a}(SH)),$$

where $\bar{a}(SH)$ and $\mu(SH)$ denote the arithmetical average and the harmonic average of all a_{ij}^h, b_{ij}^h in **SH**-cell, respectively.

The coefficient c_{ij}^H is the coefficient of $\frac{\partial^2 u}{\partial x \partial y}$ in the homogenized equation (5) generated in cells **EH, NH, SH, WH**. That is,

$$c_{ij}^H = \frac{1}{2}(\mu(EH, NH, SH, WH) - \bar{a}(EH, NH, SH, WH)),$$

where $\bar{a}(EH, NH, SH, WH)$ and $\mu(EH, NH, SH, WH)$ denote the arithmetical average and the harmonic average of all a_{ij}^h, b_{ij}^h in **EH, NH, SH, WH**-cells, respectively.

Figure 1: Coefficients for H -grid level at (i, j) .

Figure 2: Construction of 4 cells on the coarse H -grid level at (i, j) with respect to the finest h -grid level at (iL, jL) .

3.3 Construction of Interpolation

We apply the harmonic interpolation I_H^h , which is based on the continuity of $a^\epsilon(x, y) \frac{\partial u_\epsilon}{\partial x}$, and $a^\epsilon(x, y) \frac{\partial u_\epsilon}{\partial y}$, [1]. That is (See Figure 3), by setting

$$a_{2i-12j}^h D_-^i u_{2i-12j}^h = a_{2i2j}^h D_-^i u_{2i2j}^H,$$

$$b_{2i2j-1}^h D_-^j u_{2i2j-1}^h = b_{2i2j}^h D_-^j u_{2i2j}^H,$$

we solve

$$u_{2i-12j}^h = \frac{a_{2i-12j}^h u_{2i-22j}^H + a_{2i2j}^h u_{2i2j}^H}{a_{2i-12j}^h + a_{2i2j}^h},$$

$$u_{2i2j-1}^h = \frac{b_{2i2j-1}^h u_{2i2j-2}^H + b_{2i2j}^h u_{2i2j}^H}{b_{2i2j-1}^h + b_{2i2j}^h}.$$

At point $(2i-1, 2j-1)$, we use the following weighted interpolation,

$$u_{2i-12j-1}^h = \frac{a_{2i-12j-1}^h u_{2i-22j-1}^h + a_{2i2j-1}^h u_{2i2j-1}^h + b_{2i-12j-1}^h u_{2i-12j-2}^h + b_{2i-12j}^h u_{2i-12j}^h}{a_{2i-12j-1}^h + a_{2i2j-1}^h + b_{2i-12j-1}^h + b_{2i-12j}^h},$$

where $i, j = 1, \dots, N/2 - 1$.

For the restriction operator I_h^H , we take the transpose of the prolongation, i.e.,

$$I_h^H = (I_H^h)^T.$$

Figure 3: Interpolation from H -grid level to h -grid level.

3.4 Discussion of Theoretical Results

Using the Global Homogenized Coarse Grid Operator, the convergence of the two level method applied to elliptic equations with Dirichlet boundary conditions has been analyzed. [6]. In that paper, two classes of two dimensional elliptic equations are especially studied. The first case is the equation with coefficient oscillatory in x direction only; The second case is the equation with coefficient oscillatory diagonally. We prove that when both ϵ and h go to zeros, without requiring the ratio of h to ϵ to be small, as long as the numerical grids sample well the oscillating coefficients so that there are quite a lot cancellations among the high oscillations, one can improve the number of smoothing iteration in the two level method by using the global homogenized coarse grid operator. More precisely, in order that the two level method converges under the global homogenized coarse grid operator, the iteration number γ which is needed is as follows:

1. For the first case,

$$\gamma \geq Ch^{-1-1/3} \ln h;$$

2. For the second case,

$$\gamma \geq Ch^{-1-2/3} \ln h,$$

if the ratio of h to ϵ is a strict irrational number. For the detail of the proof, we refer to [6].

The theoretical results obtained seem to be a little bit pessimistic. However, the process of the theoretical proof indicates the role of homogenized operator in the convergence of multigrid methods. From a number of numerical experiments, [5], we can see that faster convergence rate in practice can be achieved than that required in the theoretical results. The theoretical convergence results are also better than what can be achieved if the coarse grid operator is constructed directly via averaging, [8]. If we replace the homogenized operator with the averaging coarse grid operator, by same analysis, all we can get is that the number of iteration number needed is $O(h^2)$, which is not even asymptotically better than just direct Jacobi method.

4 Numerical Results

4.1 Example 1

We study the mean rate [10] of convergence for the $V(\gamma_1, \gamma_2)$ -cycle multigrid method by using the homogenized coarse grid operators and harmonic interpolation to (9). The mean rate of convergence for $V(\gamma_1, \gamma_2)$ is defined by

$$\rho = \left(\frac{\|L_h u^i - f_h\|_h}{\|L_h u^1 - f_h\|_h} \right)^{\frac{1}{i-1}} \quad (15)$$

where i is the smallest integer satisfying $\|L_h u^i - f_h\|_h \leq 1 \times 10^{-9}$.

In all the numerical experiments, unless noted otherwise, we consider the coefficient $a^\epsilon(x - y) = 2.1 + 2 \sin(2\pi(x - y)/\epsilon)$, where the harmonic average μ and arithmetical average $\bar{a}$ are

$$\mu = 0.64,$$

$$\bar{a} = 2.1.$$

The smoothing iteration operator S is based on the following damped Jacobi iteration,

$$S = I - \omega h^2 L_h. \quad (16)$$

The finest grid points are on a 256×256 mesh, the step at the finest grid level h has an irrational relation with the wavelength ϵ , i.e., $\epsilon = \sqrt{2}h$. The step of the coarsest level equals to $1/2$, and ω in (16) is 0.095, which is tested numerically to be the best. In our numerical experiments, we always

compare two cases by taking two different values for θ , where θ is a parameter introduced in (13) and (14). When $\theta = 1$, it means that the coarse grid operator used is the homogenized operator; When $\theta = 0$, it means that the coarse grid operator used is the operator in (13) and (14) without the cross terms. In the latter case, the coarse grid operator is no longer the homogenized operator.

In Figure 4, we plot spectral radius ρ for $V(\gamma, \gamma)$ as a function of smoothing step γ . Here, we notice that the rate of convergence for the multigrid method is faster for $\theta = 1$ than that of $\theta = 0$. It is clearer when the smoothing step is taken to be large.

Figure 4: Spectral radius ρ as a function of smoothing step γ .

In Figure 5, we plot $\log(\text{residual})$ with respect to the number of $V(3, 3)$ -cycles.

In Figure 4, 5, we applied the local homogenized coarse grid operator for all coarse grid levels. In fact, after a few coarse grid levels, the local homogenized coarse grid operator can be replaced by the global homogenized coarse grid operator. That's, we only need apply the local homogenized coarse grid operator at a few first coarse grid levels, then apply the global homogenized coarse grid operator at the rest of the coarse grid levels. Therefore, the computational number for the construction of the coefficients can be reduced. In Figure 6, we plot the spectral radius as a function of a level variable, after which we switch from the local homogenized coarse grid operator to the global homogenized coarse grid operator. This figure gives us some idea about how the process of eigenmodes of error in the multigrid method is reduced. Namely, after the finest grid level smoothing, there indeed exist

Figure 5: $\log(\text{residual})$ as a function of number of $V(3,3)$.

many intermediate eigenmodes which are not quite close to the homogenized ones.

Figure 6: Spectral radius ρ as a function of a level variable for $V(3,3)$. And, x -axis represents the level number from finest to coarsest.

In Figure 7, we fix pre-smoothing to be 3, and calculate the spectral radius as a function of post-smoothing for $V(3, \gamma)$. From Figure 7, we can see that the more steps we provide for the post-smoothing, the more efficiency of the method with the cross terms. In fact this improvement occurs using either pre- or post-smoothing than that without the cross term's.

In Figure 8, we plot the spectral radius as a function of variable θ for $V(3,3)$. From this picture, we can see that with cross term, the convergence of the multigrid methods can be much improved.

In Figure 9, we plot the spectral radius as a function of $\frac{1}{h}$ for $V(3,3)$, where h is the grid step size of the finest level.

Figure 7: Spectral radius ρ as a function of post-smoothing γ for $V(3, \gamma)$.

Figure 8: Spectral radius ρ as a function of variable θ .

We have showed by numerical experiments in vary different aspects that the homogenized operator is the correct one to define the coarse grid operator. With it, the rate of convergence of multigrid methods can be much improved. So far, in order to isolate the influence of the coarse grid approximation we have kept the smoothing operator fixed. It obviously also affect the performance. If we use SOR iteration method in (16), the convergence rate can be improved. We compare the convergence rate by choosing damped Jacobi iteration and SOR iteration in Figure 10.

4.2 Example 2

Consider a practical problem, described in Figure 11. This can be viewed as a wall with a composite material for insulation in the center. We are interested in the heat conduction in such a composite structure. The governing equation

Figure 9: Spectral radius ρ as a function of $\frac{1}{h}$.

Figure 10: Spectral radius ρ as a function of smoothing steps γ for $V(\gamma, \gamma)$. Lines with circle for SOR iteration; otherwise, for Jacobi.

has the form as (2),

$$-\frac{\partial}{\partial x}C(x, y)\frac{\partial U}{\partial x} - \frac{\partial}{\partial y}C(x, y)\frac{\partial U}{\partial y} = 100,$$

in a rectangular domain $(x, y) \in \Omega = (0, 1) \times (0, 2)$. The boundary conditions and other parameters are given as indicated in Figure 11. Although the coefficient doesn't satisfy the periodic assumption, which is needed in homogenization theory, it still preserves some essential property as before from probabilistic point view. Namely, it is highly oscillatory in the middle part of the domain. Standard discretization of the equation contains almost random distributed magnitude coefficients by taking $\epsilon = \sqrt{2}h$. In such a case, the coarse grid operators can still be generated by the similar idea as we introduce in the previous section. Since the conductivity here is strongly vary in x direction, and has two interfaces in y direction, the structure of the

Figure 11: Model

homogenized coarse grid operator has approximately the following form,

$$-A_{11} \frac{\partial^2 U}{\partial x^2} - A_{22} \frac{\partial^2 U}{\partial y^2},$$

where A_{11} and A_{22} are defined by the following *MGH*. Different way of calculations of the coefficients for A_{11} and A_{22} leads us to different coarse grid operators. The following are specially what we are interested in. We name them in order:

MGH : A_{11} is generated by the local harmonic average of the coefficients, and A_{22} is generated by the local arithmetical average of the coefficients;

MGA : Both A_{11} and A_{22} are generated by the local arithmetical average of the coefficients;

SOR : Without any coarse grid correction, only apply SOR smoothing iteration directly.

In Table 1, we calculate the main rate ρ of the multigrid method $V(2L, 2L)$, with $\omega = 1.7$ in SOR. We compare the main rate ρ of convergence of the

multigrid method, using the above three different methods. The number of grid points at finest level is 2^L . The grid step at coarsest level in the multigrid methods is $\frac{1}{2}$.

L	3	4	5	6	7	8
MGH	0.1514	0.3779	0.4005	0.3922	0.4752	0.4832
MGA	0.1416	0.3084	0.5168	0.5518	0.6358	0.7066
SOR	0.4596	0.7028	0.8843	0.9595		

Table 1: Spectral radius ρ .

All the examples indicate that the homogenized coarse grid operators are very useful to improve the rate of convergence of multigrid methods to elliptic equations with oscillatory coefficients. The effect is more pronounced in the asymptotic region. That is, the number of smoothing iteration is large.

Acknowledgment. We thank Barry Smith for helpful comments and suggestions.

References

- [1] R.E. Alcouffe, A. Brandt, J.E. Dendy, and J.W. Painter, The Multi-Grid Method for the Diffusion Equation with Strongly Discontinuous Coefficients. *SIAM J. Sci. Stat. Comput.* **2**, 4(1981), p. 430-454.
- [2] A. Bensonssan, J.L. Lions, and G. Papanicolaou, *Asymptotic Analysis for Periodic Structure* (North-Holland, Amsterdam, 1987).
- [3] A. Brandt, Multi-level Adaptive Solutions to Boundary-Value Problems. *Mathematics of Comput.* **31**, 138(1981), p. 333-390.
- [4] B. Engquist, Computation of Oscillatory Solutions for Partial Differential Equations. *Lecture Notes 1270* (Springer Verlag, 1989), p. 10-22.
- [5] B. Engquist, and E. Luo, Multigrid Methods For Differential Equations With Highly Oscillatory Coefficients. *Proceedings of the Sixth Copper Mountain Conference on Multigrid Methods*, 1993, p. 175-190.

- [6] B. Engquist, and E. Luo, Convergence of the Two level Method for Elliptic Equation with Coefficients Highly Oscillatory, 1994 (unpublished).
- [7] M. Khalil, and P. Wesseling, Vertex-Centered and Cell-Centered Multigrid for Interface Problems. *J. of Comput. Physics.* **98**, 1(1992), p. 1-10.
- [8] E. Luo, Multigrid Method for Elliptic Equation with Oscillatory Coefficients. *Ph.D. Thesis, Department of Mathematics, UCLA*, 1993.
- [9] F. Santosa, and M. Vogelius, First Order Corrections to the Homogenized Eigenvalues of a Periodic Composite Medium. *SIAM J. Appl. Math.* **53**, 6(1993), p. 1636-1668.
- [10] R.S. Varga, *Matrix Iterative Analysis* (Prentice-Hall. Englewood Cliffs, NJ., 1962).

#	Author/s	Title
1215	Petr Klouček ,	On the existence of the entropic solutions for the transonic flow problem
1216	David A. Schmidt & Chjan C. Lim ,	Full sign-invertibility and symplectic matrices
1217	Piermarco Cannarsa & Maria Elisabetta Tessitore ,	Infinite dimensional Hamilton-Jacobi equations and Dirichlet boundary control problems of parabolic type
1218	Zhangxin Chen ,	Multigrid algorithms for mixed methods for second order elliptic problems
1219	Zhangxin Chen ,	Expanded mixed finite element methods for linear second order elliptic problems I
1220	Gang Bao ,	A note on the uniqueness for an inverse diffraction problem
1221	Moshe Fridman ,	A two state capital asset pricing model
1222	Paolo Baldi ,	Exact asymptotics for the probability of exit from a domain and applications to simulation
1223	Carl Dou & Martin Hildebrand ,	Enumeration and random random walks on finite groups
1224	Jaksa Cvitanic & Ioannis Karatzas ,	On portfolio optimization under “drawdown” constraints
1225	Avner Friedman & Yong Liu ,	A free boundary problem arising in magnetohydrodynamic system
1226	Dominic Welsh ,	Randomised approximation schemes for Tutte-Gröthendieck invariants
1227	Zhangxin Chen, Bernardo Cockburn, Carl L. Gardner, & Joseph W. Jerome ,	Quantum hydrodynamic simulation of hysteresis in the resonant tunneling diode
1228	E.G. Kalnins, G.C. Williams, & Willard Miller, Jr. ,	Intrinsic characterisation of the separation constant for spin one and gravitational perturbations in Kerr geometry
1229	Zhangxin Chen ,	Large-scale averaging analysis of multiphase flow in fractured reservoirs
1230	Bruce Hajek & Babu Narayanan ,	Multigraphs with the most edge covers
1231	K.B. Athreya ,	Entropy maximization
1232	F.I. Karpelevich & Yu.M. Suhov ,	Functional equations in the problem of boundedness of stochastic branching dynamics
1233	E. Dibenedetto & V. Vespri ,	On the singular equation $\beta(u)_t = \Delta u$
1234	M.Ya. Kelbert & Yu.M. Suhov ,	The Markov branching random walk and systems of reaction-diffusion (Kolmogorov-Petrovskii-Piskunov) equations
1235	M. Hildebrand ,	Random walks on random regular simple graphs
1236	W.S. Don & A. Solomonoff ,	Accuracy enhancement for higher derivatives using Chebyshev collocation and a mapping technique
1237	D. Gurarie ,	Symmetries and conservation laws of two-dimensional hydrodynamics
1238	Z. Chen ,	Finite element methods for the black oil model in petroleum reservoirs
1239	G. Bao & A. Friedman ,	Inverse problems for scattering by periodic structure
1240	G. Bao ,	Some inverse problems in partial differential equations
1241	G. Bao ,	Diffraction optics in periodic structures: The TM polarization
1242	C.C. Lim & D.A. Schmidt ,	On noneven digraphs and symplectic pairs
1243	H.M. Soner, S.E. Shreve & J. Cvitanić ,	There is no nontrivial hedging portfolio for option pricing with transaction costs
1244	D.L. Russell & B-Yu Zhang ,	Exact controllability and stabilizability of the Korteweg-de Vries equation
1245	B. Morton, D. Enns & B-Yu Zhang ,	Stability of dynamic inversion control laws applied to nonlinear aircraft pitch-axis models
1246	S. Hansen & G. Weiss ,	New results on the operator Carleson measure criterion
1247	V.A. Malyshev & F.M. Spijksma ,	Intrinsic convergence rate of countable Markov chains
1248	G. Bao, D.C. Dobson & J.A. Cox ,	Mathematical studies in rigorous grating theory
1249	G. Bao & W.W. Symes ,	On the sensitivity of solutions of hyperbolic equations to the coefficients
1250	D.A. Huntley & S.H. Davis ,	Oscillatory and cellular mode coupling in rapid directional solidification
1251	M.J. Donahue, L. Gurvits, C. Darken & E. Sontag ,	Rates of convex approximation in non-Hilbert spaces
1252	A. Friedman & B. Hu ,	A Stefan problem for multi-dimensional reaction diffusion systems
1253	J.L. Bona & B-Y. Zhang ,	The initial-value problem for the forced Korteweg-de Vries equation
1254	A. Friedman & R. Gulliver ,	Organizers , Mathematical modeling for instructors
1255	S. Kichenassamy ,	The prolongation formula for tensor fields
1256	S. Kichenassamy ,	Fuchsian equations in Sobolev spaces and blow-up
1257	H.S. Dumas, L. Dumas, & F. Golse ,	On the mean free path for a periodic array of spherical obstacles
1258	C. Liu ,	Global estimates for solutions of partial differential equations
1259	C. Liu ,	Exponentially growing solutions for inverse problems in PDE
1260	Mary Ann Horn & I. Lasiecka ,	Nonlinear boundary stabilization of parallelly connected Kirchhoff plates
1261	B. Cockburn & H. Gau ,	A posteriori error estimates for general numerical methods for scalar conservation laws
1262	B. Cockburn & P-A. Gremaud ,	A priori error estimates for numerical methods for scalar conservation laws. Part I: The general approach

- 1263 **R. Spigler & M. Vianello**, Convergence analysis of the semi-implicit euler method for abstract evolution equations
- 1264 **R. Spigler & M. Vianello**, WKB-type approximation for second-order differential equations in C^* -algebras
- 1265 **M. Menshikov & R.J. Williams**, Passage-time moments for continuous non-negative stochastic processes and applications
- 1266 **C. Mazza**, On the storage capacity of nonlinear neural networks
- 1267 **Z. Chen, R.E. Ewing & R. Lazarov**, Domain decomposition algorithms for mixed methods for second order elliptic problems
- 1268 **Z. Chen, M. Espedal & R.E. Ewing**, Finite element analysis of multiphase flow in groundwater hydrology
- 1269 **Z. Chen, R.E. Ewing, Y.A. Kuznetsov, R.D. Lazarov & S. Maliassov**, Multilevel preconditioners for mixed methods for second order elliptic problems
- 1270 **S. Kichenassamy & G.K. Srinivasan**, The structure of WTC expansions and applications
- 1271 **A. Zinger**, Positiveness of Wigner quasi-probability density and characterization of Gaussian distribution
- 1272 **V. Malkin & G. Papanicolaou**, On self-focusing of short laser pulses
- 1273 **J.N. Kutz & W.L. Kath**, Stability of pulses in nonlinear optical fibers using phase-sensitive amplifiers
- 1274 **S.K. Patch**, Recursive recovery of a family of Markov transition probabilities from boundary value data
- 1275 **C. Liu**, The completeness of plane waves
- 1276 **Z. Chen & R.E. Ewing**, Stability and convergence of a finite element method for reactive transport in ground water
- 1277 **Z. Chen & Do Y. Kwak**, The analysis of multigrid algorithms for nonconforming and mixed methods for second order elliptic problems
- 1278 **Z. Chen**, Expanded mixed finite element methods for quasilinear second order elliptic problems II
- 1279 **M.A. Horn & W. Littman**, Boundary control of a Schrödinger equation with nonconstant principal part
- 1280 **R.E. Ewing, Yu.A. Kuznetsov, R.D. Lazarov & S. Maliassov**, Substructuring preconditioning for finite element approximations of second order elliptic problems. I. Nonconforming linear elements for the Poisson equation in a parallelepiped
- 1281 **S. Maliassov**, Substructuring preconditioning for finite element approximations of second order elliptic problems. II. Mixed method for an elliptic operator with scalar tensor
- 1282 **V. Jakšić & C.-A. Pillet**, On model for quantum friction II. Fermi's golden rule and dynamics at positive temperatures
- 1283 **V. M. Malkin**, Kolmogorov and nonstationary spectra of optical turbulence
- 1284 **E.G. Kalnins, V.B. Kuznetsov & W. Miller, Jr.**, Separation of variables and the XXZ Gaudin magnet
- 1285 **E.G. Kalnins & W. Miller, Jr.**, A note on tensor products of q -algebra representations and orthogonal polynomials
- 1286 **E.G. Kalnins & W. Miller, Jr.**, q -algebra representations of the Euclidean, pseudo-Euclidean and oscillator algebras, and their tensor products
- 1287 **L.A. Pastur**, Spectral and probabalistic aspects of matrix models
- 1288 **K. Kastella**, Discrimination gain to optimize detection and classification
- 1289 **L.A. Peletier & W.C. Troy**, Spatial patterns described by the Extended Fisher-Kolmogorov (EFK) equation: Periodic solutions
- 1290 **A. Friedman & Y. Liu**, Propagation of cracks in elastic media
- 1291 **A. Friedman & C. Huang**, Averaged motion of charged particles in a curved strip
- 1292 **G. R. Sell**, Global attractors for the 3D Navier-Stokes equations
- 1293 **C. Liu**, A uniqueness result for a general class of inverse problems
- 1294 **H-O. Kreiss**, Numerical solution of problems with different time scales II
- 1295 **B. Cockburn, G. Gripenberg, S-O. Londen**, On convergence to entropy solutions of a single conservation law
- 1296 **S-H. Yu**, On stability of discrete shock profiles for conservative finite difference scheme
- 1297 **H. Behncke & P. Rejto**, A limiting absorption principle for separated Dirac operators with Wigner Von Neumann type potentials
- 1298 **R. Lipton B. Vernescu**, Composites with imperfect interface
- 1299 **E. Casas**, Pontryagin's principle for state-constrained boundary control problems of semilinear parabolic equations
- 1300 **G.R. Sell**, References on dynamical systems
- 1301 **J. Zhang**, Swelling and dissolution of polymer: A free boundary problem
- 1302 **J. Zhang**, A nonlinear nonlocal multi-dimensional conservation law
- 1303 **M.E. Taylor**, Estimates for approximate solutions to acoustic inverse scattering problems
- 1304 **J. Kim & D. Sheen**, A priori estimates for elliptic boundary value problems with nonlinear boundary conditions
- 1305 **B. Engquist & E. Luo**, New coarse grid operators for highly oscillatory coefficient elliptic problems
- 1306 **A. Boutet de Monvel & I. Egorova**, On the almost periodicity of solutions of the nonlinear Schrödinger equation with the cantor type spectrum
- 1307 **A. Boutet de Monvel & V. Georgescu**, Boundary values of the resolvent of a self-adjoint operator: Higher order estimates
- 1308 **S.K. Patch**, Diffuse tomography modulo Graßmann and Laplace
- 1309 **A. Friedman & J.J.L. Velázquez**, Liouville type theorems for fourth order elliptic equations in a half plane