**ASYMPTOTIC BEHAVIOR OF SOLUTIONS OF
COAGULATION-FRAGMENTATION MODELS**

By

Avner Friedman

and

Fernando Reitich

IMA Preprint Series # 1479

June 1997

**INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
UNIVERSITY OF MINNESOTA**

**514 Vincent Hall
206 Church Street S.E.
Minneapolis, Minnesota 55455**

ASYMPTOTIC BEHAVIOR OF SOLUTIONS OF COAGULATION-FRAGMENTATION MODELS

AVNER FRIEDMAN* AND FERNANDO REITICH†

Abstract. In this paper we study the evolution of the number density $n(x, c, t)$ of droplets of volume x with chemical concentration c subject to both coalescence (coagulation) and rupture (fragmentation, breakage). It is proved that as $t \rightarrow \infty$ the concentration tends to a limit value c_∞ and the measure $xn(x, c, t)dxdc$ converges to $\delta(c - c_\infty)dL(x)$ where $\delta(y)$ is the Dirac measure and dL is a measure satisfying the appropriate equilibrium equation.

1. The model. Consider a mixture of a large number of incompressible spherical droplets of varying volume size x in a solution (e.g., water or air). When the mixture is well stirred the distribution of droplet size can be taken to be homogeneous in space. We may then introduce the notion of density number $n(x)$:

$$n(x)dx = \text{number of droplets in unit volume}$$

$$\text{whose volume size lies between } x \text{ and } x + dx;$$

$n(x)$ is independent of the location of the volume element.

Since the droplets are in continuous motion, they may collide and, as a result, coalesce. The motion may also cause rupture. We assume that if two droplets of volumes x and ξ coalesce, they will form a droplet of volume $x + \xi$. The rate of coalescence is given by a function $K(x, \xi)$, which is called the *coalescence* or *coagulation kernel*.

We also introduce the *fragmentation* or *breakage kernel* $B(x, \xi)$ which is the rate that a droplet of volume x breaks into two droplets of volumes ξ and $x - \xi$.

Throughout this paper we assume that

$$\begin{aligned} (1.1) \quad & K(x, \xi) \text{ is continuous and } \geq 0 \text{ for } x, \xi \geq 0, \\ & K(x, \xi) = K(\xi, x), \\ & K(x, \xi) \leq C_0, \\ & B(x, \xi) \text{ is continuous and } \geq 0 \text{ for } x \geq \xi \geq 0, \\ & B(x, \xi) = B(x, x - \xi), \\ & \int_0^x B(x, \xi)d\xi \leq C_1, \\ & \int_0^x \xi B(x, \xi)d\xi \leq C_2x. \end{aligned}$$

Remark 1.1. Note that if volume is preserved under rupture, then

$$\int_0^x B(x, \xi)\xi d\xi = x,$$

* University of Minnesota, Institute for Mathematics and its Applications, Minneapolis, Minnesota 55455.

† Department of Mathematics and Center for Research in Scientific Computation, North Carolina State University, Box 8205, Raleigh, NC 27695-8205.

so that the last condition in (1.1) is very natural.

Due to coagulation and fragmentation the number density will actually vary with time, and we introduce

$$n(x, t) = \text{number density at time } t.$$

The following conservation law holds:

$$\begin{aligned} \frac{d}{dt}n(x, t)dx = & \text{number gained through coalescence} \\ & + \text{number gained through rupture} \\ & - \text{number lost through coalescence} \\ & - \text{number lost through rupture.} \end{aligned}$$

This can be expressed by the evolution equation:

$$\begin{aligned} (1.2) \quad \frac{d}{dt}n(x, t) = & \frac{1}{2} \int_0^x K(x - \xi, \xi)n(x - \xi, t)n(\xi, t)d\xi + \int_x^\infty B(\xi, x)n(\xi, t)d\xi \\ & - n(x, t) \int_0^\infty K(x, \xi)n(\xi, t)d\xi - n(x, t) \int_0^x \frac{\xi}{x} B(x, \xi)d\xi. \end{aligned}$$

Here the underlying assumption is that, although $n(x, t)$ varies with t , the mixture is maintained homogeneous in space at any moment (due to continuous fast stirring).

Together with (1.2) we prescribe an initial condition

$$(1.3) \quad n(x, 0) = n_0(x), \quad n_0(x) \geq 0.$$

There is substantial literature on coagulation and coagulation-fragmentation models. Melzak [10] proved a general existence, uniqueness and positivity of a solution for all $t > 0$. For other results we refer to [2], [4], [9] and the references therein.

We shall actually be interested in a situation where each droplet carries a chemical species which is uniformly distributed within it; the concentration c of the chemical species is a variable quantity. We then introduce

$$\begin{aligned} n(x, c, t)dxdc = & \text{number of droplets at time } t \\ & \text{with volume size in the interval } (x, x + dx) \\ & \text{and concentration (of the chemical} \\ & \text{species) in the interval } (c, c + dc). \end{aligned}$$

Analogously to (1.2), the number density $n(x, c, t)$ evolves as follows:

$$\begin{aligned}
(1.4) \quad \frac{d}{dt}n(x, c, t) &= \frac{1}{2} \int_0^x \int_0^\infty K(x - \xi, \xi) n\left(x - \xi, \frac{xc - \xi\gamma}{x - \xi}, t\right) n(\xi, \gamma, t) \frac{x}{x - \xi} d\gamma d\xi \\
&\quad + \int_x^\infty B(\xi, x) n(\xi, c, t) d\xi \\
&\quad - n(x, c, t) \int_0^\infty \int_0^\infty K(x, \xi) n(\xi, \gamma, t) d\gamma d\xi \\
&\quad - n(x, c, t) \int_0^x \frac{\xi}{x} B(x, \xi) d\xi,
\end{aligned}$$

where we define

$$n(x, c', t) = 0 \quad \text{if } c' < 0.$$

Here we used the fact that rupture does not change the concentration. On the other hand, if particles (ξ, γ) and (η, β) coalesce to produce (x, c) , then

$$x = \xi + \eta, \quad xc = \xi\gamma + \eta\beta$$

so that

$$\eta = x - \xi, \quad \beta = \frac{xc - \xi\gamma}{x - \xi}$$

and $d\gamma d\xi d\eta d\beta = d\gamma d\xi \frac{x}{x - \xi} dx dc$, since the Jacobian determinant

$$\begin{vmatrix} \frac{\partial \eta}{\partial x} & \frac{\partial \eta}{\partial c} \\ \frac{\partial \beta}{\partial x} & \frac{\partial \beta}{\partial c} \end{vmatrix} \quad \text{is equal to } \frac{x}{x - \xi};$$

this explains the appearance of $x/(x - \xi)$ in the first integral on the right hand side of (1.4).

Remark 1.2. The last integral on the right-hand side of (1.2), or (1.4), can also be written in the form

$$\frac{1}{2} \int_0^x B(x, \xi) d\xi$$

since

$$\begin{aligned}
\frac{1}{2} \int_0^x B(x, \xi) d\xi &= \frac{1}{2} \frac{1}{x} \int_0^x [\xi B(x, \xi) + (x - \xi) B(x, \xi)] d\xi \\
&= \frac{1}{2} \frac{1}{x} \left[\int_0^x \xi B(x, \xi) d\xi + \int_0^x (x - \xi) B(x, x - \xi) d\xi \right] \\
&= \frac{1}{x} \int_0^x \xi B(x, \xi) d\xi.
\end{aligned}$$

Together with (1.4) we prescribe an initial condition

$$(1.5) \quad n(x, c, 0) = n_0(x, c), \quad n_0(x, c) \geq 0.$$

The model (1.4) arises in the process of making a photographic film. The film is made of several emulsion layers. One of them consists of oil droplets in aqueous solution. The mixture, prior to being coated on the film substrate, is subjected to coalescence and rupture by means of stirring. A key problem is to determine the rate of coalescence (which depends on parameters, such as the amount of surfactant added at the initial time). One of the methods to determine coalescence is based on adding chemiluminescent species to the mixture:

The optical signal, which is the number of photons emitted per unit time from a droplet, is a function, $f(x, c)$, of the species concentration c in the droplet and the droplet volume x . One can measure the total signal

$$(1.6) \quad S(t) = \int_0^\infty \int_0^\infty f(x, c) n(x, c) dx dc,$$

and this is used to evaluate the rate of coalescence [8, Chap. 6].

Note that if two drops with parameters (x_1, c_1) and (x_2, c_2) coalesce to form a drop with parameters $(\tilde{x}, \tilde{c})$ then

$$\tilde{x} = x_1 + x_2, \quad \tilde{c} = \frac{x_1 c_1 + x_2 c_2}{x_1 + x_2}.$$

Typically $f(x, c) = xc^2$, and then

$$f(\tilde{x}, \tilde{c}) < f(x_1, c_1) + f(x_2, c_2),$$

so that the total signal should decrease.

The model (1.4) was developed by David Ross; see [8, Chap. 6]. He conjectured that as $t \rightarrow \infty$ the concentrations c will all tend to a constant, say c_∞ . With this in mind, one defines the *degree of coalescence*

$$(1.7) \quad D(t) = \int_0^\infty \int_0^\infty (c - c_\infty)^2 x n(x, c, t) dx dc$$

and one would like to prove rigorously that

$$(1.8) \quad D(t) \rightarrow 0 \quad \text{if } t \rightarrow \infty.$$

Consider next the measures $d\mu_t$ defined by

$$(1.9) \quad d\mu_t(x, c) = x n(x, c, t) dx dc$$

If (1.8) is true then we may go one step further and try to prove that as $t \rightarrow \infty$

$$(1.10) \quad d\mu_t(x, c) \rightarrow \delta(c - c_\infty) dL(x)$$

in the sense of weak convergence of measures, where $\delta(y)$ is the Dirac measure and

$$(1.11) \quad dL \text{ is a solution of the equilibrium equation (1.2).}$$

In this paper we shall prove, under mild additional conditions on K and B that (1.8) holds. We shall also establish (1.10), (1.11) under the additional condition that

$$(1.12) \quad \begin{aligned} K(x, \xi) &\leq \eta(x + \xi), \quad \gamma_0 \equiv \inf_{x>0} \frac{1}{x} \int_0^x \xi B(x, \xi) d\xi > 0, \text{ where} \\ 2\eta \int_0^\infty \int_0^\infty x n_0(x, c) dx dc &< \gamma_0. \end{aligned}$$

The structure of the paper is as follows:

In §2 we establish existence, uniqueness and positivity of the solution of (1.4), (1.5) for all $t > 0$. Assuming that

$$(1.13) \quad n_0(x) = \int_0^\infty n_0(x, c) dc$$

we also prove that the function

$$(1.14) \quad n(x, t) = \int_0^\infty n(x, c, t) dc$$

is the unique solution of (1.2), (1.3). In §3 we prove conservation of mass

$$\int_0^\infty x n(x, t) dx = \text{const.}$$

We also show that if $n_0(x, c) = 0$ whenever $c > c_*$ (c_* constant) then $n(x, c, t) = 0$ for $c > c_*$. Using these results we derive in §4 a formula for dS/dt (see (4.2)) when S is defined in (1.6) and $f(x, c)$ is any function satisfying $|f(x, c)| \leq C(1 + x)$. This formula is the key to the proofs of both (1.8) (in §5) and (1.10), (1.11) (in §6).

We finally remark that a coalescence problem for $n(x, c, t)$ was studied in [1], with c being the concentration of surfactant which inhibits coalescence. The limit behavior of the solution and the mathematical method for deriving it are entirely different from the limit behavior and the methods of the present paper.

2. Existence and uniqueness. We shall need the following assumptions:

$$(2.1) \quad \begin{aligned} n_0(x) &\text{ is continuous for } x \geq 0, \\ 0 \leq n_0(x) &\leq A_0, \quad \int_0^\infty n_0(x) dx \leq A_0, \end{aligned}$$

$$(2.2) \quad \begin{aligned} n_0(x, c) &\text{ is continuous for } x \geq 0, \quad c \geq 0, \\ 0 \leq n_0(x, c) &\leq A_1, \quad \int_0^\infty \int_0^\infty n_0(x, c) dx dc \leq A_1 \end{aligned}$$

where A_0, A_1 are positive constants.

THEOREM 2.1. *If (1.1), (2.1) hold then there exists a unique solution of (1.2), (1.3) with $n(x, t)$, $\partial n(x, t)/\partial t$ continuous for $x \geq 0$, $t \geq 0$, such that, for each $T > 0$,*

$$(2.3) \quad \sup_{x \geq 0} |n(x, t)| + \int_0^\infty |n(x, t)| dx \leq C(T) < \infty \text{ if } 0 \leq t \leq T.$$

The solution has the following additional properties:

$$(2.4) \quad n(x, t) \text{ is analytic in } t, \quad t \geq 0,$$

$$(2.5) \quad n(x, t) \geq 0.$$

This theorem is due to Melzak [10]. We note that if K and B have compact support and $n_0(x) > 0$ for all $x > 0$, then the proof of (2.5) is much simpler than the proof given in [10] (or [3]). Indeed, if x is large then from (1.2) we see that $dn(x, t)/dt = 0$ so that $n(x, t) = n_0(x) > 0$. Hence if (2.5) is not true, with strict inequality, then there is a smallest $t = t_0 > 0$ such that $n(x, t) > 0$ if $t < t_0$ for all $x \geq 0$ and $n(x_0, t_0) = 0$ for a finite point x_0 . From (1.2) we then deduce that

$$\frac{dn(x_0, t_0)}{dt} > 0,$$

which is a contradiction. Finally, the solution $n(x, t)$ for general K , B , n_0 , can be obtained as a pointwise limit of solutions with compact K , B and strictly positive $n_0(x)$ (cf. [3] or [4]), so that (2.5) is true in general.

THEOREM 2.2. *If (1.1), (2.2) hold then there exists a unique solution of (1.4), (1.5) with $n(x, c, t)$, $\partial n(x, c, t)/\partial t$ continuous for $x \geq 0$, $c \geq 0$, $t \geq 0$, such that, for each $T > 0$,*

$$(2.6) \quad \sup_{x \geq 0, c \geq 0} |n(x, c, t)| + \int_0^\infty \int_0^\infty |n(x, c, t)| dx dc \leq C(T) < \infty \text{ if } 0 \leq t \leq T.$$

The solution has the following additional properties:

$$(2.7) \quad n(x, c, t) \text{ is analytic in } t, \quad t \geq 0,$$

$$(2.8) \quad n(x, c, t) \geq 0.$$

Proof. The proof is similar to the proof of Theorem 2.1 provided we can treat the first term on the right-hand side of (1.4) analogously to the first term on the right-hand side of (1.2); this means that we have to control the factor $x/(x - \xi)$ in the integrand.

Introduce the operator

$$J(m, n) = \frac{1}{2} \int_0^x \int_0^\infty K(x - \xi, x) m \left(x - \xi, \frac{xc - \xi\gamma}{x - \xi} \right) \frac{x}{x - \xi} n(\xi, \gamma) d\gamma d\xi$$

and break up the integral into

$$\int_0^{x/2} \int_0^\infty + \int_{x/2}^x \int_0^\infty$$

6

In the second integral substitute

$$\gamma \rightarrow \gamma' = \frac{xc - \xi\gamma}{x - \xi}, \quad \text{or} \quad \gamma = \frac{xc - (x - \xi)\gamma'}{\xi}, \quad d\gamma = -\frac{x - \xi}{\xi} d\gamma'.$$

We then obtain

$$\begin{aligned} J(m, n) &= \frac{1}{2} \int_0^{x/2} \int_0^\infty K(x - \xi, \xi) m \left(x - \xi, \frac{xc - \xi\gamma}{x - \xi}, t \right) n(\xi, \gamma, t) \frac{x}{x - \xi} d\gamma d\xi \\ &\quad - \frac{1}{2} \int_{x/2}^x \int_0^\infty K(x - \xi, \xi) n \left(\xi, \frac{xc - (x - \xi)\gamma'}{\xi}, t \right) m(x - \xi, \gamma', t) \frac{x}{\xi} d\gamma' d\xi. \\ &= J_1 - J_2. \end{aligned}$$

Since the factors $\frac{x}{x - \xi}$ and $\frac{x}{\xi}$ in J_1 and J_2 respectively are uniformly bounded, we can easily establish, as in [10], the bound

$$(2.9) \quad |J(m, n)|_{L^\infty} \leq C |n|_{L^\infty} \int \int |m| + C |m|_{L^\infty} \int \int |n|$$

where

$$\begin{aligned} |f|_{L^\infty} &= \sup_{x \geq 0, c \geq 0} |f(x, c, t)|, \\ \int \int |f| &= \int_0^\infty \int_0^\infty |f(x, c, t)| dc dx. \end{aligned}$$

Next we estimate

$$\int \int |J(m, n)| dc dx$$

by substituting

$$c \rightarrow c' = \frac{xc - \xi\gamma}{x - \xi}, \quad dc' = \frac{x}{x - \xi} dc.$$

we obtain

$$(2.10) \quad \int \int |J(m, n)| \leq C \int \int |m| \cdot \int \int |n|$$

With (2.9), (2.10) at hand, we can now proceed as in [10], with just minor changes, to establish Theorem 2.2. The positivity of $n(x, c, t)$ can be proved, more simply, by the argument following Theorem 2.1.

THEOREM 2.3. *If (1.1), (2.1), (2.2) hold and, in addition,*

$$(2.11) \quad n_0(x) = \int_0^\infty n_0(x, c) dc,$$

then

$$(2.12) \quad \int_0^\infty n(x, c, t) dc = n(x, t).$$

where $n(x, c, t)$ and $n(x, t)$ are the solutions of (1.4), (1.5) and (1.2), (1.3), respectively.

Proof. We first proceed formally to integrate (1.4) with respect to c . In the first integral on the right-hand side we substitute

$$(2.13) \quad c \rightarrow c' = \frac{xc - \xi\gamma}{x - \xi}, \quad dc' = \frac{x}{x - \xi} dc$$

to get

$$(2.14) \quad \frac{1}{2} \int_0^\infty K(x - \xi, \xi) n(x - \xi, t) n(\xi, t) d\xi.$$

where $n(x, t)$ is the function defined by the left-hand side of (2.12). All the other integrals yield immediately the corresponding integrals of (1.2). Hence, by uniqueness, the left-hand side of (2.12) is the solution of (1.2), (1.3).

To prove the theorem rigorously we rewrite (1.4) in integrated form:

$$(2.15) \quad n(x, c, t) - n_0(x, c) = \int_0^t dt \left\{ \frac{1}{2} \int_0^x \int_0^\infty K \cdots + \cdots \right\}.$$

Since

$$\int_0^\infty \int_0^\infty n(x, c, t) dc dx \leq C(T) < \infty \text{ if } 0 \leq t \leq T,$$

the integral

$$\int_0^\infty n(x, c, t) dc$$

is finite a.e. in x , for each t , and we may integrate (2.15) with respect to c , $0 \leq c < \infty$. Each of the four multiple integrals thus obtained is well defined and is finite. We now proceed with the substitution (2.13) to obtain the term (2.14, and the proof of the theorem then follows as before.

3. Conservation and invariant laws.

THEOREM 3.1. (*Conservation law*). *If (1.1), (2.1) hold and*

$$(3.1) \quad \int_0^\infty x n_0(x) dx < \infty,$$

then

$$(3.2) \quad \int_0^\infty xn(x, t)dx = \int_0^\infty xn_0(x)dx$$

for all $t > 0$.

Proof. The assertion (3.2) means that

$$(3.3) \quad \frac{d}{dt} \int_0^\infty xn(x, t)dx = 0.$$

This relation follows formally as a special case of Theorem 4.1 with $f(x, c) = x$. However, in order to apply the proof rigorously we need to know a priori that, for any $T > 0$,

$$(3.4) \quad \int_0^\infty xn(x, t)dx \leq C(T) < \infty \quad \text{if } 0 \leq t \leq T.$$

If K , B and n_0 have compact support then (3.4) is true and then so is (3.2). We now approximate general K , B , n_0 by K_j , B_j , $n_{0,j}$ with compact support and denote the corresponding solutions by $n_j(x, t)$. Then, on one hand we have

$$\int_0^\infty xn_j(x, t)dx = \int_0^\infty xn_{0,j}(x)dx \leq C$$

for all j , and on the other hand $n_j(x, t) \rightarrow n(x, t)$ pointwise (cf. [4]). It follows that $n(x, t)$ satisfies (3.4) and consequently (3.3), or (3.2), holds.

THEOREM 3.2. *If (1.1), (2.2) hold and*

$$(3.5) \quad n_0(x, c) = 0 \quad \text{for } x \geq 0, c > c_* \ (c_* > 0),$$

then

$$(3.6) \quad n(x, c, t) = 0 \quad \text{for } x \geq 0, c \geq c_*, t \geq 0.$$

Proof. If $c > c_*$ then in the first integral on the right-hand side of (1.4)

$$\text{either } \gamma > c_* \text{ or else } \frac{xc - \xi\gamma}{x - \xi} > c_*.$$

Assuming that K has compact support and using (2.3), we conclude that the integral is bounded by $CN(t)$, where

$$N(t) = \sup_{x \geq 0, c \geq c_*} n(x, c, t).$$

If B has compact support then the second integral on the right-hand side of (1.4) is bounded by

$$C \sup_{\xi > x} n(\xi, c, t), \quad \text{or by } CN(t).$$

From (1.4) it then follows that

$$\frac{dn(x, c, t)}{dt} \leq CN(t).$$

If $x \geq 0$, $c > c_*$, so that

$$N(t) \leq N(0) + C \int_0^t N(t') dt'.$$

Since $N(0) = 0$, the assertion (3.6) follows.

Having proved (3.6) in the case where K and B have compact support, we can now establish the same result for general K , B by approximation (cf. the proof of Theorem 3.1).

For convenience we collect all the assumptions made so far on $n_0(x, c)$ and $n_0(x)$:

$$(3.7) \quad \left\{ \begin{array}{l} n_0(x) \text{ and } n_0(x, c) \text{ are continuous functions for } x \geq 0, c \geq 0, \\ 0 \leq n_0(x) \leq A_0, \quad \int_0^\infty (1+x)n_0(x)dx < \infty, \\ 0 \leq n_0(x, c) \leq A_1, \quad \int_0^\infty n_0(x, c)dc = n_0(x), \\ n_0(x, c) = 0 \text{ if } c \geq c_* \text{ } (c_* > 0). \end{array} \right.$$

COROLLARY 3.3. *If (1.1) and (3.7) are satisfied then (3.2) and (3.6) hold.*

4. A formula for dS/dt . Let $f(x, c)$ be any function satisfying

$$(4.1) \quad |f(x, c)| \leq C(1+x) \text{ if } x \geq 0, c \leq c_*$$

and arbitrary otherwise. By (2.6) and Corollary 3.3 it follows that the integral

$$(4.2) \quad S(t) = \int_0^\infty \int_0^\infty f(x, c)n(x, c, t)dxdc$$

exists for all $t \geq 0$.

THEOREM 4.1. *If (1.1), (3.7) hold and f satisfies (4.1), then*

$$(4.3) \quad \begin{aligned} \frac{dS(t)}{dt} &= \frac{1}{2} \int_0^\infty dx \int_0^\infty d\xi \int_0^\infty dc \int_0^\infty d\gamma K(x, \xi)n(x, c, t)n(\xi, \gamma, t) \\ &\quad \times \left[f\left(x + \xi, \frac{xc + \xi\gamma}{x + \xi}\right) - f(x, c) - f(\xi, \gamma) \right] \\ &\quad + \int_0^\infty dx \int_0^\infty dc \int_x^\infty B(\xi, x)n(\xi, c, t) \left[f(x, c) - \frac{x}{\xi}f(\xi, c) \right] d\xi. \end{aligned}$$

Note that, by (2.6) and Corollary 3.3, all the integrals on the right-hand side of (4.3) are absolutely convergent.

Proof. Formally

$$(4.4) \quad \frac{dS(t)}{dt} = \int_0^\infty \int_0^\infty f(x, c) \frac{dn(x, c, t)}{dt} dx dc.$$

By substituting dn/dt from (1.4) and using (2.6), (3.2), (4.1), we find that each of the resulting four integrals is absolutely convergent.

A standard argument then shows that $S(t)$ is differentiable and dS/dt is given by (4.4).

We can next write

$$(4.5) \quad \frac{dS}{dt} = I_1 + I_2$$

where

$$\begin{aligned} I_1 &= \int_0^\infty dc \left[\int_0^\infty f(x, c) dx \int_x^\infty B(\xi, x) n(\xi, c, t) d\xi \right. \\ &\quad \left. - \int_0^\infty f(x, c) n(x, c, t) dx \int_0^x \frac{\xi}{x} B(x, \xi) d\xi \right], \\ I_2 &= \frac{1}{2} \int_0^\infty dx \int_0^\infty f(x, c) dc \int_0^x \int_0^\infty K(x - \xi, \xi) n \left(x - \xi, \frac{xc - \xi\gamma}{x - \xi}, t \right) n(\xi, \gamma, t) \frac{x}{x - \xi} d\gamma d\xi \\ &\quad - \int_0^\infty dc \int_0^\infty dx \int_0^\infty n(x, c, t) K(x, \xi) n(\xi, \gamma, t) f(x, c) d\gamma d\xi \\ &= I_{21} - I_{22}. \end{aligned}$$

To evaluate I_1 we change the notation of the variables $x \rightarrow \xi$ and $\xi \rightarrow x$ in the second integral in $[\dots]$ to get

$$\int_0^\infty d\xi \int_0^\xi \frac{x}{\xi} B(\xi, c) f(\xi, c) n(\xi, c, t) dx.$$

This, in turn, is equal, by changing the order of integration, to

$$\int_0^\infty dx \int_x^\infty \frac{x}{\xi} B(\xi, x) f(\xi, c) n(\xi, c, t) d\xi.$$

Hence

$$(4.6) \quad I_1 = \int_0^\infty dc \int_0^\infty dx \int_x^\infty B(\xi, x) \left[f(x, c) - \frac{x}{\xi} f(\xi, c) \right] n(\xi, c, t) d\xi$$

Next we turn to I_{21} and change variables $x \rightarrow x'$ where $x - \xi = x'$ (for each fixed ξ) and $c \rightarrow c'$ by

$$c = \frac{x'c' + \xi\gamma}{x' + \xi}, \quad \text{or} \quad \frac{xc - \xi\gamma}{x - \xi} = c'.$$

We get

$$I_{21} = \frac{1}{2} \int_0^\infty dx' \int_0^\infty d\xi \int_0^\infty dc' \int_0^\infty d\gamma K(x', \xi) n(x', c, t) n(\xi, \gamma, t) f\left(\xi + x', \frac{x'c' + \xi\gamma}{x' + \xi}\right).$$

We shall use the relations

$$(4.7) \quad \begin{aligned} \int_0^\infty dc' \int_0^\infty d\gamma \Phi &= \int_0^\infty dc' \int_0^{c'} d\gamma \Phi + \int_0^\infty dc' \int_{c'}^\infty d\gamma \Phi \\ &= \int_0^\infty dc' \int_0^{c'} d\gamma \Phi + \int_0^\infty d\gamma \int_0^\gamma dc' \Phi, \end{aligned}$$

which is valid with any integrand Φ . Since after integration, in I_{21} , with respect to x' and ξ we get an expression $\Psi(\gamma, c')$ which is symmetric in γ, c' , we can write

$$\int_0^\infty d\gamma \int_0^\gamma dc' \Psi(\gamma, c') = \int_0^\infty dc' \int_0^{c'} d\gamma \Psi(\gamma, c').$$

Hence for the purpose of calculating I_{21} we may replace the right-hand side of (4.7) by

$$2 \int_0^\infty dc' \int_0^{c'} d\gamma \Phi.$$

Consequently

$$(4.8) \quad I_{21} = \int_0^\infty dx' \int_0^\infty d\xi \int_0^\infty dc' \int_0^{c'} d\gamma K(x', \xi) n(x', c', t) n(\xi, \gamma, t) f\left(x' + \xi, \frac{x'c' + \xi\gamma}{x' + \xi}, t\right).$$

Next we can write

$$(4.9) \quad \begin{aligned} I_{22} &= \int_0^\infty dx \int_0^\infty d\xi \int_0^\infty dc \int_0^c d\gamma K(x, \xi) n(x, c, t) n(\xi, \gamma, t) f(x, c) \\ &+ \int_0^\infty dx \int_0^\infty d\xi \int_0^\infty dc \int_0^\infty d\gamma K(x, \xi) n(x, c, t) n(\xi, \gamma, t) f(x, c) \\ &= L_1 + L_2 \end{aligned}$$

In L_2 we write

$$\int_0^\infty dc \int_c^\infty d\gamma = \int_0^\infty d\gamma \int_0^\gamma dc$$

to get

$$L_2 = \int_0^\infty dx \int_0^\infty d\xi \int_0^\infty d\gamma \int_0^\gamma dc K(x, \xi) n(x, c, t) n(\xi, \gamma, t) f(x, c).$$

Changing the notation of variables $\gamma \leftrightarrow c$ and $x \leftrightarrow \xi$ we find, using the symmetry of $K(x, \xi)$, that

$$L_2 = \int_0^\infty d\xi \int_0^\infty dx \int_0^\infty dc \int_0^c d\gamma K(x, \xi) n(\xi, \gamma, t) n(x, c, t) f(\xi, c).$$

We substitute this expression into (4.9) and, upon renaming x', c' in (4.8) by x, c , obtain

$$\begin{aligned} I_2 = I_{21} - I_{22} &= \int_0^\infty dx \int_0^\infty d\xi \int_0^\infty dc \int_0^c d\gamma K(x, \xi) n(x, c, t) n(\xi, \gamma, t) \\ &\times \left[f\left(x + \xi, \frac{xc + \xi\gamma}{x + \xi}\right) - f(x, c) - f(\xi, \gamma) \right]. \end{aligned}$$

Next we write in I_2

$$\cdots \int_0^\infty dc \int_0^c d\gamma \quad \text{as} \quad \cdots \int_0^\infty d\gamma \int_\gamma^\infty dc$$

and change notation $c \leftrightarrow \gamma$, $x \leftrightarrow \xi$. We find that I_2 is equal to a similar integral in which

$$\int_0^\infty dc \int_0^c d\gamma \text{ is replaced by } \int_0^\infty dc \int_c^\infty d\gamma$$

(here we again use the symmetry of $K(x, \xi)$). It follows that I_2 is equal to $\frac{1}{2}$ times the first integral on the right-hand side of (4.3). Recalling also (4.6), (4.5), the assertion (4.3) follows.

Formula (4.3) will play a key role in the subsequent sections. Note that the choice $f = x$ gives the conservation law (3.2) which was already proved. On the other hand the choice $f = xc$ gives a new conservation law:

$$(4.10) \quad \int_0^\infty \int_0^\infty xc n(x, c, t) dx dc = \int_0^\infty \int_0^\infty xc n_0(x, c) dx dc \text{ for all } t \geq 0.$$

COROLLARY 4.2. *If $g(c)$ is a convex function then*

$$\frac{d}{dt} \int_0^\infty \int_0^\infty g(c) x n(x, c, t) dx dc \leq 0.$$

Indeed, the second integral on the right-hand side of (4.3) is zero whereas the first integral is ≤ 0

COROLLARY 4.3. *If $n_0(x, c) = 0$ for $0 \leq c \leq \tilde{c}$, $x \geq 0$ then $n(x, c, t) = 0$ for $0 \leq c \leq \tilde{c}$, $x \geq 0$, $t \geq 0$.*

Indeed, taking $g(c)$ to be the convex function $(\tilde{c} - c)^+$ we conclude that

$$\int_0^\infty \int_0^{\tilde{c}} (\tilde{c} - c)^+ x n(x, c, t) dx dc \leq \int_0^\infty \int_0^{\tilde{c}} (\tilde{c} - c)^+ x n_0(x, c) dx dc = 0$$

and the assertion follows.

The proof of Theorem 4.1 can also be applied (or, rather, specialized) to functions of the form

$$(4.11) \quad \tilde{S}(t) = \int_0^\infty f(x)n(x,t)dx.$$

We then get:

THEOREM 4.4. *If (1.1), (3.7) hold and $f(x)$ satisfies*

$$(4.12) \quad |f(x)| \leq C(1+x),$$

then

$$(4.13) \quad \begin{aligned} \frac{d\tilde{S}(t)}{dt} &= \frac{1}{2} \int_0^\infty \int_0^\infty K(x,\xi)n(x,t)n(\xi,t) [f(x+\xi) - f(x) - f(\xi)] dx d\xi \\ &+ \int_0^\infty dx \int_x^\infty B(\xi,x)n(\xi,t) \left[f(x) - \frac{x}{\xi}f(\xi) \right] d\xi. \end{aligned}$$

5. $D(t) \rightarrow 0$ if $t \rightarrow \infty$. In this section we impose additional mild assumptions on K , B and n_0 :

$$(5.1) \quad \int_0^\infty x^2 n_0(x) dx < \infty$$

and

$$(5.2) \quad \begin{aligned} (a) \quad & K(x,\xi) \leq C(x+\xi); \\ (b) \quad & \partial_x K(x,\xi) \geq -C; \\ (c) \quad & \exists \text{ a positive constant } \gamma_1 \text{ such that, for any } N \gg 1, \\ & K(x,\xi) \geq \frac{\gamma_1}{N}(x+\xi) \text{ if } 0 \leq x, \xi \leq N; \\ (d) \quad & \int_0^\xi B(\xi,x)x(\xi-x)dx \geq \gamma_2 \xi^2 \quad \forall \xi > 0, \text{ for some } \gamma_2 > 0. \end{aligned}$$

Remark 1.1 indicates that condition (5.2)(d) is quite natural. Note also that (5.2)(c) is satisfied if

$$K(x,\xi) \geq \begin{cases} \delta(x+\xi) & \text{for } x+\xi < 1 \\ \delta & \text{for } x+\xi > 1 \end{cases}$$

where δ is a small positive constant.

Set

$$(5.3) \quad M_0 = \int_0^\infty \int_0^\infty x n_0(x,c) dx dc, \quad M_1 = \int_0^\infty \int_0^\infty x c n_0(x,c) dx dc,$$

$$(5.4) \quad c_\infty = \frac{M_1}{M_0}.$$

In this section we prove:

THEOREM 5.1. *If (1.1), (3.7) and (5.1), (5.2) hold, then the function*

$$(5.5) \quad D(t) \equiv \int_0^\infty \int_0^\infty x(c - c_\infty)^2 n(x, c, t) dx dc$$

converges to zero as $t \rightarrow \infty$.

We need several lemmas.

LEMMA 5.2. *There exists a constant C independent of t such that*

$$(5.6) \quad \int_0^\infty x^2 n(x, t) dx \leq C \quad \forall t > 0.$$

Proof. Consider first the case where

$$(5.7) \quad K, B \text{ and } n_0 \text{ have compact support.}$$

Then, for any $T > 0$,

$$\tilde{S}(t) \equiv \int_0^\infty x^2 n(x, t) dx \leq C(T) < \infty \quad \text{if } 0 \leq t \leq T,$$

and we may apply (4.13) with $f(x) = x^2$ (The proof is the same as that of Theorem 4.4.). We get

$$\begin{aligned} \frac{d\tilde{S}}{dt} &\leq C \int_0^\infty \int_0^\infty n(x, t) n(\xi, t) x \xi dx d\xi \\ &\quad + \int_0^\infty n(\xi, t) d\xi \int_0^\xi B(\xi, x) (x^2 - x\xi) dx, \end{aligned}$$

since $K(x, \xi) \leq C$. Using Theorem 3.1 and (5.2) (d) we obtain

$$\frac{d\tilde{S}}{dt} \leq C - \gamma_2 \int_0^\infty \xi^2 n(\xi, t) d\xi = C - \gamma_2 \tilde{S}(t).$$

Consequently

$$(5.8) \quad \tilde{S}(t) \leq \tilde{S}(0) e^{-\gamma_2 t} + \frac{C}{\gamma_2}.$$

and (5.6) follows with a constant which is independent of the assumptions in (5.7). Approximating general K, B, n_0 by functions with compact support and applying (5.8) to the corresponding solutions, we obtain, in the limit, the assertion (5.6).

Introduce the function

$$(5.9) \quad S_2(t) = \int_0^\infty \int_0^\infty x c^2 n(x, c, t) dx dc.$$

Then

$$(5.10) \quad D(t) = S_2(t) - 2c_\infty M_1 + C_\infty^2 M_0.$$

LEMMA 5.3. *The function $S_2(t)$ satisfies:*

$$(5.11) \quad \frac{dS_2(t)}{dt} \leq 0,$$

$$(5.12) \quad \frac{d^2 S_2(t)}{dt^2} \leq C$$

for all $t > 0$, and

$$(5.13) \quad \frac{dS_2(t)}{dt} \rightarrow 0 \quad \text{if } t \rightarrow \infty.$$

Proof. Using (4.3) with $f(x, c) = xc^2$ and noting that

$$(5.14) \quad f\left(x + \xi, \frac{xc + \xi\gamma}{x + \xi}\right) - f(x, c) - f(\xi, \gamma) = -\frac{x\xi}{x + \xi}(c - \gamma)^2,$$

we get

$$(5.15) \quad \frac{dS_2}{dt} = -\frac{1}{2} \int_0^\infty dx \int_0^\infty d\xi \int_0^\infty dc \int_0^\infty d\gamma \frac{K(x, \xi)}{x + \xi} x\xi(c - \gamma)^2 n(x, c, t) n(\xi, \gamma, t),$$

so that $dS_2/dt \leq 0$.

To prove (5.12) we want to apply Theorem 4.1 with

$$(5.16) \quad f(x, c) = \frac{K(x, \xi)}{x + \xi} x(c - \gamma)^2; \quad \xi, \gamma \text{ are parameters.}$$

for simplicity we first take $B = 0$. Then

$$\begin{aligned} \frac{d^2 S_2}{dt^2} &= -\frac{1}{2} \int_0^\infty dx \int_0^\infty d\xi \int_0^\infty dc \int_0^\infty d\gamma \xi n(\xi, \gamma, t) \int_0^\infty d\zeta \int_0^\infty d\sigma K(x, \xi) n(x, c, t) n(\xi, \sigma, t) \\ &\quad \times \left[\frac{K(x + \zeta, \xi)}{x + \zeta + \xi} (x + \zeta) \left(\frac{xc + \zeta\sigma}{x + \zeta} - \gamma \right)^2 - \frac{K(x, \xi)}{x + \xi} x(c - \gamma)^2 \right. \\ &\quad \left. - \frac{K(\zeta, \xi)}{\zeta + \xi} \zeta(\sigma - \gamma)^2 \right]. \end{aligned}$$

Using

$$\frac{xc + \zeta\sigma}{x + \zeta} - \gamma = \frac{x(c - \gamma) + \zeta(\sigma - \gamma)}{x + \zeta}$$

and a calculation similar to (5.14), we find that the expression in brackets is equal to

$$\begin{aligned} &\frac{K(x + \zeta, \xi)}{x + \zeta + \xi} \left(\frac{-x\zeta(c - \sigma)^2}{x + \zeta} \right) + \left(\frac{K(x + \zeta, \xi)}{x + \zeta + \xi} - \frac{K(x, \xi)}{x + \xi} \right) x(c - \gamma)^2 \\ &\quad + \left(\frac{K(x + \zeta, \xi)}{x + \zeta + \xi} - \frac{K(\zeta, \xi)}{\zeta + \xi} \right) \zeta(\sigma - \gamma)^2. \end{aligned}$$

By (5.2)(a), $K(x, \zeta)$ times the first term is bounded by $cx\zeta$, and, by (5.2)(a) and (5.2)(b), $K(x, \zeta)$ times each of the remaining two terms is bounded from below by $-Cx\zeta$.

It follows that $d^2 S_2/dt^2$ is bounded from above by

$$C \left(\int_0^\infty \int_0^\infty xn(x, c, t) dx dt \right)^3 \leq \text{const.}$$

If $B \neq 0$ then we have to estimate the corresponding positive contribution to $d^2 S_2/dt^2$:

$$(5.17) \quad \frac{1}{2} \int_0^\infty dx \int_0^\infty d\xi \int_0^\infty dc \int_0^\infty d\gamma \xi n(\xi, \gamma, t) f(x, c) n(x, c, t) \int_0^x \frac{\zeta}{x} B(x, \zeta) d\zeta$$

where $f(x, c)$ is defined as in (5.16). Since $K(x, \xi) \leq C(x + \xi)$, we have $f(x) \leq Cx$. Recalling also the last condition in (1.1) we find that the expression in (5.17) is bounded by

$$C \left(\int_0^\infty \int_0^\infty xn(x, c, t) dx dc \right)^2 \leq \text{const.}$$

This completes the proof of (5.12).

It remains to prove (5.13). If the assertion is not true then there is a sequence $t_n \rightarrow \infty$ such that

$$0 \geq S'_2(t_n) \rightarrow -\alpha < 0 \text{ if } n \rightarrow \infty.$$

Using (5.12) we get

$$S_2(t_n + \epsilon) - S_2(t_n) = \epsilon S'_s(t_n) + \frac{\epsilon^2}{2} S''_2(\tilde{t}_n) \leq -\frac{\alpha\epsilon}{2}$$

if ϵ is positive and small enough. It follows that we have $f(x, c) \leq Cx$.

$$S_2(t_n + \epsilon) \leq S_2(t_1) - \frac{n\epsilon}{2} < 0$$

if n is sufficiently large, a contradiction.

For any $N \gg 1$ introduces the quantities

$$\begin{aligned} M_{0,N} &= \int_0^N dx \int_0^\infty xn(x, c, t) dc, \\ M_{1,N}(x) &= \int_0^N dx \int_0^\infty cxn(x, c, t) dc \end{aligned}$$

and

$$S_{2,N}(t) = \int_0^N dx \int_0^\infty c^2 xn(x, c, t) dc.$$

By Lemma 5.2,

$$(5.18) \quad \begin{cases} |S_2(t) - S_{2,N}(t)| \leq \frac{C}{N} \text{ for all } t \geq 0, \\ |M_{0,N} - M_0| + |M_{1,N} - M_1| \leq \frac{C}{N}. \end{cases}$$

LEMMA 5.4. *There holds:*

$$(5.19) \quad \int_0^\infty dx \int_0^\infty dc \int_0^\infty d\xi \int_0^\infty d\gamma x \xi (c - \gamma)^2 n(x, c, t) n(\xi, \gamma, t) = 2 (S(t)M_0 - M_1^2)$$

and, in particular,

$$(5.20) \quad S(t)M_0 \geq M_1^2.$$

The proof is obvious.

Proof of Theorem 5.1. by (5.15)

$$\frac{dS_2(t)}{dt} = -\frac{1}{2} \iiint\limits_{x < N, \xi < N} dx dc d\xi d\gamma \frac{K(x, \xi)}{x + \xi} x \xi (c - \gamma)^2 n(x, c, t) n(\xi, \gamma, t) + J_N$$

where, since $K(x, \xi) \leq C$,

$$|J_N| \leq C \iiint\limits_{\xi > N} dx dc d\xi d\gamma x n(x, c, t) n(\xi, \gamma, t) \leq \frac{C}{N^2};$$

the last inequality follows by Lemma 5.2. Using (5.2)(c) we conclude that

$$\frac{dS_2}{dt} \leq -\frac{\gamma_1}{N} \iiint\limits_{x < N, \xi < N} dx dc d\xi d\gamma x \xi (c - \gamma)^2 n(x, c, t) n(\xi, \gamma, t) + \frac{C}{N^2}.$$

Evaluating the last integral as in (5.19), we get

$$\frac{dS_2}{dt} \leq -\frac{2\gamma_1}{N} (S_{2,n}(t)M_{0,N} - M_{1,N}^2) + \frac{C}{N^2}$$

so that

$$S_{2,N}(t)M_{0,N} - M_{1,N}^2 \leq -\frac{N}{2\gamma_1} S_2'(t) + \frac{C}{N}.$$

Using (5.18) we deduce that

$$S_2(t)M_0 - M_1^2 \leq -\frac{N}{2\gamma_1} S_2'(t) + \frac{C}{N}.$$

Taking $t \rightarrow \infty$ and using (5.13) we find that

$$\overline{\lim}_{t \rightarrow \infty} (S(t)M_0 - M_1^2) \leq \frac{C}{N}$$

Taking $N \rightarrow \infty$ and recalling (5.20), we find that

$$S(t)M_0 - M_1^2 \rightarrow 0 \quad \text{if } t \rightarrow \infty.$$

Consequently, as $t \rightarrow \infty$,

$$D(t) \rightarrow \frac{M_1^2}{M_0} - 2C_\infty M_1 + C_\infty^2 M_0 = 0, \quad \text{by (5.10) and (5.4),}$$

and the proof of Theorem 5.1 is complete

Remark 5.5. Dubovskii and Stewart [4] proved existence and uniqueness for (1.2), (1.3) also in case $K(x, \xi)$ is unbounded, provided

$$\begin{aligned} K(x, \xi) &\leq C(x + \xi), \\ n_0(x) &\leq Ce^{-\lambda x} \quad (\lambda > 0); \end{aligned}$$

The solution is exponentially decreasing, and uniqueness is in the class of exponentially decreasing functions. Their result can be extended to (1.4), (1.5). Suppose now that K also satisfies:

$$K(x, \xi) \geq \gamma_3(x + \xi), \quad \gamma_3 > 0.$$

Then the proof of Theorem 5.1 remains valid without actually splitting integrals by $x \gtrless N$ or $\xi \gtrless N$. Thus we get, in this case,

$$\frac{dS_2}{dt} \leq -\gamma_3 (S_2(t)M_0 - M_1^2).$$

The non-negative function

$$T(t) = S_2(t) - \frac{M_1^2}{M_0}$$

then satisfies

$$\frac{dT}{dt} + 2\gamma_3 T \leq 0$$

so that

$$0 \leq T(t) \leq T(0)e^{-2\gamma_3 t}.$$

We conclude that, as $t \rightarrow \infty$,

$$D(t) \rightarrow 0 \text{ exponentially fast.}$$

Note that in this case we do not require the assumption (5.2)(d).

6. Asymptotic behavior of $xn(x, c, t)dxdc$. In this section we add one more assumption on K, B, n_0 :

$$(6.1) \quad K(x, \xi) \leq \eta(x + \xi) \text{ where } 2M_0\eta < \gamma_2, \quad \gamma_2 \text{ as in (5.2)(d).}$$

THEOREM 6.1. *Let (1.1), (3.7), (5.1), (5.2) and (6.1) hold. Then there exists a non-negative measure $dL(x)$ such that*

$$(6.2) \quad \lim_{t \rightarrow \infty} \int_0^\infty \int_0^\infty g(x, c) xn(x, c, t) dx dc = \int_0^\infty g(x, c_\infty) dL(x)$$

for any bounded continuous function $g(x)$.

Introducing the measures

$$d\mu_t(x, c) = xn(x, c, t) dx dc$$

we can restate (6.2) in the form:

$$(6.3) \quad d\mu_t(x, c) \rightarrow \delta(c - c_\infty) dL(x) \text{ as } t \rightarrow \infty, \text{ in the sense of weak convergence,}$$

where $\delta(y)$ is the Dirac measure.

The proof requires several lemmas.

LEMMA 6.2. *Under the assumptions of Theorem 6.1,*

$$(6.4) \quad \lim_{t \rightarrow \infty} \int_0^\infty g(x) xn(x, t) dx \text{ exists}$$

for any bounded continuous function $g(x)$.

Proof. Consider first the function

$$S_0(t) = \int_0^\infty x^2 n(x, t) dx$$

By (4.13)

$$\begin{aligned} \frac{dS_0}{dt} &= \int_0^\infty \int_0^\infty K(x, \xi) n(x, t) n(\xi, t) x \xi dx d\xi \\ &\quad + \int_0^\infty dx \int_x^\infty B(\xi, x) n(\xi, t) (x^2 - x\xi) d\xi \\ &\leq \eta \int_0^\infty \int_0^\infty (x + \xi) n(x, t) n(\xi, t) x \xi dx d\xi \\ &\quad + \int_0^\infty n(\xi, t) d\xi \int_0^\infty B(\xi, x) (x^2 - x\xi) dx \\ &\leq 2\eta S_0(t) M_0 - \gamma_2 S_0(t) \end{aligned}$$

so that

$$(6.5) \quad \frac{dS_0}{dt} \leq -\delta S_0(t), \quad \delta = \gamma_2 - 2\eta M_0 > 0.$$

Suppose now that $g(x)$ is a uniformly Lipschitz continuous function and write

$$(6.6) \quad \begin{aligned} \int_0^\infty g(x)xn(x,t)dx &= \int_0^\infty (g(x)x + Nx^2)n(x,t)dx \\ &\quad - N \int_0^\infty x^2n(x,t)dx \equiv \tilde{S}_1(t) - NS_0(t). \end{aligned}$$

By (6.5), $S_0(t)$ is monotone decreasing, and hence

$$(6.7) \quad \lim_{t \rightarrow \infty} S_0(t) \text{ exists.}$$

Next, by Theorem 4.4 with $f(x) = xg(x)$,

$$\begin{aligned} \frac{d}{dt} \int_0^\infty g(x)xn(x,t)dx &\leq \frac{1}{2} \int_0^\infty dx \int_0^\infty d\xi \eta(x+\xi)n(x,t)n(\xi,t)x\xi \\ &\quad \times \left[\frac{g(x+\xi) - g(x)}{\xi} + \frac{g(x+\xi) - g(\xi)}{x} \right] \\ &\quad + \int_0^\infty dx \int_x^\infty B(\xi,x)n(\xi,t)x[g(x) - g(\xi)]d\xi. \end{aligned}$$

The first integral on the right-hand side is bounded by

$$C \int_0^\infty x^2n(x,t)dx \cdot \int_0^\infty \xi n(\xi,t)d\xi$$

and the second integral is bounded by

$$\int_0^\infty n(\xi,t)d\xi \int_0^\xi B(\xi,x)xC|x-\xi|dx \leq C \int_0^\infty \xi^2n(\xi,t)d\xi.$$

Recalling (6.5), it follows that

$$\frac{d\tilde{S}_1}{dt} \leq -\delta NS_0(t) + C_0S_0(t) < 0 \quad \text{if } N > \frac{C_0}{\delta},$$

and consequently

$$\lim_{t \rightarrow \infty} \tilde{S}_1(t) \text{ exists.}$$

Combining this with (6.7) we deduce from (6.6) that (6.4) holds for any uniformly Lipschitz continuous function $g(x)$.

Consider next the case where $g(x)$ is any bounded continuous function and let g_ϵ be a mollifier of g :

$$|g_\epsilon(x) - g(x)| < \epsilon \text{ and } |g'_\epsilon(x)| \leq \frac{C}{\epsilon} \quad \text{for all } x \geq 0.$$

Write

$$\begin{aligned} \int_0^\infty g(x)xn(x,t)dx &= \int_0^\infty g_\epsilon(x)xn(x,t)dx \\ &+ \int_0^\infty (g(x) - g_\epsilon(x))xn(x,t)dx \equiv T_1(t) + T_2(t). \end{aligned}$$

Clearly

$$|T_2(t)| < \epsilon.$$

By what was already proved above

$$A_\epsilon \equiv \lim_{t \rightarrow \infty} T_1(t) \text{ exists.}$$

Hence

$$\lim_{t \rightarrow \infty} \left| \int_0^\infty g(x)xn(x,t)dx - A_\epsilon \right| \leq \epsilon.$$

Since the A_ϵ are uniformly bounded, we can take a sequence $\epsilon = \epsilon_m \rightarrow 0$ such that $A_{\epsilon_m} \rightarrow A$, and obtain

$$\overline{\lim}_{t \rightarrow \infty} \left| \int_0^\infty g(x)xn(x,t)dx - A \right| = 0.$$

This completes the proof of Lemma 6.2.

LEMMA 6.3. *Under the assumptions of Theorem 6.1,*

$$(6.8) \quad \lim_{t \rightarrow \infty} \int_0^\infty \int_0^\infty g(x,c)xn(x,c,t)dx dc \text{ exists}$$

for any bounded continuous function $g(x,c)$.

The proof is similar to the proof of Lemma 6.2 and will be omitted. Note that here we work (cf.(6.6)) with

$$(g(x,c) + Nx^2) \text{ and } -Nx^2.$$

The limit in (6.8) is a bounded linear functional Φ on the space of bounded continuous functions g equipped with the L^∞ norm. By [6, pp. 261–262] we can write

$$\begin{aligned} (6.9) \quad \lim_{t \rightarrow \infty} \int_0^\infty \int_0^\infty g(x,c)xn(x,c,t)dx dc \\ = \int \int g(x,c)dN(x,c) \equiv \Phi(g) \end{aligned}$$

where dN is a bounded finitely additive regular set function. It will be also useful to view the right-hand side of (6.9) as a distribution, and use the properties

$$(6.10) \quad \begin{aligned} |\Phi(g)| &\leq C|g|_{L^\infty}, \\ \Phi(g) &\geq 0 \text{ if } g \geq 0. \end{aligned}$$

From Theorem 5.1 it follows that

$$\int \int (c - c_\infty)^2 dN(x, c) = 0$$

and therefore also

$$(6.11) \quad \int \int g(x, c)(c - c_\infty)^2 dN(x, c) = 0$$

for any bounded continuous function $g(x, c)$.

If $g(x, c)$ vanishes in a neighborhood of the line $\{c = c_\infty\}$ then we can apply (6.11) to the function

$$\tilde{g}(x, c) = \begin{cases} \frac{g(x, c)}{(c - c_\infty)^2} & \text{on } \text{supp } g \\ 0 & \text{elsewhere} \end{cases}$$

and conclude that

$$\Phi(g) = \int \int \tilde{g}(x, c)(c - c_\infty)^2 dN(x, c) = 0.$$

It follows that the support of the distribution Φ lies in the set $\{c = c_\infty\}$. By a standard result in the theory of distributions (e.g. [7, p. 69, Theorem 31], Φ must then have the form

$$\Phi(g) = \sum_{0 \leq i, j \leq l} \frac{\partial^{i+j}}{\partial c^i \partial x^j} \int g(x, c_\infty) dk_{ij}(x)$$

where the derivatives $\partial^{i+j}/\partial c^i \partial x^j$ are taken in the sense of distributions, and dk_{ij} are measures. In view of (6.10) we must have $l = 0$ and, furthermore, dk_{00} is a non-negative measure. This completes the proof of Theorem 6.1.

The equilibrium equation (1.1) is

$$(6.12) \quad \begin{aligned} &\frac{1}{2} \int_0^\infty K(x - \xi, \xi) n_\infty(x - \xi) n_\infty(\xi) d\xi + \int_x^\infty B(\xi, x) n_\infty(\xi) d\xi \\ &- n_\infty(x) \int_0^\infty K(x, \xi) n_\infty(\xi) d\xi - n_\infty(x) \int_0^x \frac{\xi}{x} B(x, \xi) d\xi = 0. \end{aligned}$$

If $n_\infty(x)$ is a bounded, non-negative, $L^1(0, \infty)$ -solution of (6.12), then the measure $dL(x) = x n_\infty(x) dx$ satisfies:

$$(6.13) \quad \begin{aligned} &\frac{1}{2} \int_0^\infty dL(\xi) \int_0^\infty dL(x) K(x, \xi) \left[\frac{g(x + \xi) - g(x)}{\xi} + \frac{g(x + \xi) - g(\xi)}{x} \right] \\ &+ \int_0^\infty dL(\xi) \int_0^\xi B(\xi, x) \frac{x}{\xi} (g(x) - g(\xi)) dx = 0 \end{aligned}$$

for any Lipschitz function g . Indeed, this follows from Theorem 4.4 applied to $f(x) = xg(x)$.

DEFINITION. A bounded non-negative measure $dL(x)$ is called a *weak solution* of (6.12) if (6.13) holds for any continuously differentiable function $g(x)$ with uniformly bounded derivative.

THEOREM 6.4. *Under the assumptions of Theorem 6.1 the measure $dL(x)$ is a weak solution of (6.12).*

Proof. We first establish (6.13) in case

$$(6.14) \quad |g'(x)| + |g''(x)| \leq C, \quad g(x) \text{ monotone increasing.}$$

The starting point is the assertion that

$$(6.15) \quad \frac{d}{dt} \int_0^\infty xg(x)n(x,t) \rightarrow 0 \text{ if } t \rightarrow \infty.$$

To prove it we set

$$\tilde{S}_2(t) = \int_0^\infty xg(x)n(x,t)dx$$

and proceed as in the proof of Lemma 5.3. By Lemma 6.2

$$(6.16) \quad \lim_{t \rightarrow \infty} \tilde{S}_2(t) \text{ exists.}$$

We next claim that

$$(6.17) \quad \frac{d^2 \tilde{S}_2(t)}{dt^2} \leq C.$$

To prove it we use Theorem 4.4 with $f(x) = xg(x)$ to get:

$$(6.18) \quad \begin{aligned} \frac{d\tilde{S}_2(t)}{dt} &= \frac{1}{2} \int_0^\infty dx \int_0^\infty d\xi K(x, \xi) x\xi n(x, t) n(\xi, t) \\ &\quad \times \left[\frac{g(x + \xi) - g(x)}{\xi} + \frac{g(x + \xi) - g(\xi)}{x} \right] \\ &\quad + \int_0^\infty dx \int_x^\infty B(\xi, x) n(\xi, t) x(g(x) - g(\xi)) d\xi \equiv J_1(t) + J_2(t), \end{aligned}$$

and continue to differentiate once more in t . Consider first the case $B = 0$. Then we can proceed similarly to the proof of (5.13) with $f(x, c)/x$ in (5.16) replaced by

$$f(x) = K(x, \xi) \left[\frac{g(x + \xi) - g(x)}{\xi} + \frac{g(x + \xi) - g(\xi)}{x} \right]$$

where ξ is viewed as a parameter. Using (5.2)(b) and the assumptions on g in (6.14) we easily deduce that

$$\frac{d^2 \tilde{S}_2}{dt^2} \leq C.$$

If $B(x, \xi) \not\equiv 0$ then the additional terms in $d^2 \tilde{S}_2/dt^2$ are also bounded, as can easily be shown.

Having proved (6.16), (6.17), we can now quickly prove (6.15). Indeed, if (6.15) is not true then there is a sequence $t_n \rightarrow \infty$ such that

$$\beta_n \equiv \frac{d\tilde{S}_2(t_n)}{dt} \rightarrow \beta \neq 0 \quad \text{if } n \rightarrow \infty.$$

It follows that

$$\begin{aligned} \tilde{S}_2(t_n - \beta_n \epsilon) - \tilde{S}_2(t_n) &= -\beta_n^2 \epsilon + \frac{1}{2} \beta_n^2 \epsilon^2 \tilde{S}_2''(\tilde{t}_n) \\ &\leq -\beta_n^2 \epsilon + C \epsilon^2 < -\frac{1}{2} \beta^2 \epsilon \end{aligned}$$

if ϵ is positive and small and n is large; this is a contradiction to (6.16).

So far we have proved that the left-hand side of (6.18) converges to zero as $t \rightarrow \infty$. Next we evaluate the right-hand side of (6.18) as $t \rightarrow \infty$. By Theorem 6.1

$$\begin{aligned} (6.19) \quad J_2(t) &= \int_0^\infty \xi n(\xi, t) d\xi \left[\frac{1}{\xi} \int_0^\xi B(\xi, x) x (g(x) - g(\xi)) dx \right] \\ &\rightarrow \int_0^\infty dL(\xi) \int_0^\xi B(\xi, x) \frac{x}{\xi} (g(x) - g(\xi)) dx. \end{aligned}$$

To evaluate $J_1(t)$ introduce the functions

$$\begin{aligned} F(x, t) &= \int_0^\infty d\xi K(x, \xi) \xi n(\xi, t) \frac{g(x + \xi) - g(x)}{\xi}, \\ F_\infty(x) &= \int_0^\infty dL(\xi) K(x, \xi) \frac{g(x + \xi) - g(x)}{\xi}. \end{aligned}$$

The family of functions $\{F(x, t), t > 0\}$ is uniformly bounded and equicontinuous, and, by Theorem 6.1,

$$F(x, t) \rightarrow F_\infty(x) \quad \text{if } t \rightarrow \infty.$$

It follows that the convergence is uniform in any boundary interval $0 \leq x \leq R$. Using also Lemma 5.2, we easily deduce that

$$\int_0^\infty x n(x, t) [F(x, t) - F_\infty(x)] dx \rightarrow 0 \quad \text{if } n \rightarrow \infty.$$

Since, by Theorem 6.1,

$$\int_0^\infty dx n(x, t) F_\infty(x) \rightarrow \int_0^\infty dL(x) F_\infty(x)$$

we conclude that

$$\int_0^\infty xn(x, t)F(x, t)dx \rightarrow \int_0^\infty dL(x)F_\infty(x)$$

if $t \rightarrow \infty$. Combining this result with (6.19), we deduce that the right-hand side of (6.18) converges to the left-hand side of (6.13), and since the left-hand side of (6.18) converges to zero, the equality (6.13) follows.

So far we have assumed that g satisfies (6.14). However by approximation we can establish (6.13) for any monotone increasing g with first bounded and continuous derivative. Finally, if g is not monotone increasing, we can decompose it into a difference of two monotone functions, and this completes the proof of Theorem 6.4.

Remark 6.1. If $n_0(x)$ is a stationary solution then $n(x, t) = n_0(x)$ and so

$$\int_0^\infty n(x, c, t)dc = n_0(x).$$

It follows that

$$\int_0^\infty \int_0^\infty g(x)xn(x, c, t)dx dc = \int_0^\infty g(x)xn_0(x)dx$$

and

$$xn(x, c, t)dx dc \rightarrow \delta(c - c_\infty)(xn_0(x)dx).$$

Remark 6.2. It was observed by Carr [2] that if K and B are related by

$$K(x, y)Q(x)Q(y) = B(x + y, y)Q(x + y)$$

where Q is a positive function, then

$$n_\infty(x) = e^{\lambda x}Q(x)$$

is an equilibrium solution. Dubovskii and Stewart [5] proved that if $K(x, y)$ and $B(x, y)$ are linear functions then $n(x, t)$ converges to an equilibrium solution $n_\infty(x)$; their methods are entirely different from the methods of the present paper.

Acknowledgement. (1) We would like to thank David Ross from Eastman Kodak for suggesting the problems studied in this paper.

(2) The first author is partially supported by the National Science Foundation Grant DMS #94-01251. The second author is partially supported by AFOSR through contract number F49620-95-1-0113 and by NSF through grant number DMS-9622555.

Acknowledgement and Disclaimer. Effort sponsored by the Air Force Office of Scientific Research, Air Force Materials Command, USAF, under grant no. F49620-95-1. The US Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation thereby. The view and conclusions contained herein are those of the authors and should not be interpreted as necessarily representing the official policies or endorsements, either expressed or interpreted, of the Air Force Office of Scientific Research or the US Government.

REFERENCES

- [1] O. Bruno, A. Friedman and F. Reitich, *Asymptotic behavior for a coalescence problem*, Trans. Amer. Math. Soc., 338 (1993), 133–158.
- [2] J. Carr, *Asymptotic behavior of solutions to the coagulation-fragmentation equations I. The strong fragmentation case*, Proc. Royal Soc. Edinburgh, Sect. A 121 (1992), 231–244.
- [3] P.B. Dubovskii, *Mathematical Theory of Coagulation, Lecture Notes Series Number 23*, Research Institute of Mathematics, Global Analysis Research Center, Seoul National University, Seoul, Korea (1994).
- [4] P.B. Dubovskii and I.W. Stewart, *Existence, uniqueness and mass conservation for the coagulation-fragmentation equation*, Math. Methods Appl. Sciences, 19 (1996), 571–591.
- [5] P.B. Dubovskii and I.W. Stewart, *Trend to equilibrium for the coagulation-fragmentation equation*, Math. Methods Appl. Science, 19 (1996), 761–772.
- [6] N. Dunford and J.T. Schwartz, *Linear Operators, Part I*, 4th edition, Interscience, New York (1967).
- [7] A. Friedman, *Generalized Functions*, Prentice-Hall, Englewood Cliffs, N.J. (1963).
- [8] A. Friedman, *Mathematics in Industrial Problems, Part 10*, in IMA Volume Series, Springer-Verlag, to appear.
- [9] A. Friedman and F. Reitich, *A hyperbolic inverse problem arising in the evolution of combustion aerosol*, Archive Rat. Mech. Anal., 110 (1990), 313–350.
- [10] Z.A. Melzak, *A scalar transport equation*, Trans. Amer. Math. Soc., 85 (1957), 547–560.

#	Author/s	Title
=====		
1398	B. Brighi & M. Chipot,	Approximation of infima in the calculus of variations
1399	G. Avalos,	Concerning the well-posedness of a nonlinearly coupled semilinear wave and beam-like equation
1400	R. Lipton,	Variational methods, bounds and size effects for composites with highly conducting interface
1401	B.T. Hayes & P.G. LeFloch,	Non-classical shock waves in scalar conservation laws
1402	K.T. Joseph & P.G. LeFloch,	Boundary layers in weak solutions to hyperbolic conservation laws
1403	Y. Diao, C. Ernst, & E.J.J. Van Rensburg,	Energies of knots
1404	Xiaofeng Ren,	Multi-layer local minimum solutions of the bistable equation in an infinite tube
1405	Vlastimil Pták,	Krylov sequences and orthogonal polynomials
1406	T. Aktosun, M. Klaus, & C. van der Mee,	Factorization of scattering matrices due to aprtitioning of potentials in one-dimensional Schrödinger-type equations
1407	C.-S. Man & R. Paroni,	On the separation of stress-induced and texture-induced birefringence in acoustoelasticity
1408	D.N. Arnold, R.S. Falk, & R. Winther,	Preconditioning discrete approximations of the Reissner-Mindlin plate model
1409	M.A. Kouritzin,	On exact filters for continuous signals with discrete observations
1410	R. Lipton,	The second Stekloff eigenvalue and energy dissipation inequalities for functionals with surface energy
1411	R. Lipton,	The second Stekloff eigenvalue of an inclusion and new size effects for composites with imperfect interface
1412	W. Littman & B. Liu,	The regularity and singularity of solutions of certain elliptic problems on polygonal domains
1413	C.R. Collins,	Spurious oscillations are not fatal in computing microstructures
1414	M.A. Horn,	Sharp trace regularity for the solutions of the equations of dynamic elasticity
1415	A. Friedman, B. Hu & Y. Liu,	A boundary value problem for the Poisson equation with multi-scale oscillating boundary
1416	P. Bauman, D. Phillips & Q. Tang,	Stable nucleation for the Ginzburg-Landau system with an applied magnetic field
1417	J.M. Berg,	A strain profile for robust control of microstructure using dynamic recrystallization
1418	P. Klouček,	Toward the computational modeling of nonequilibrium thermodynamics of the Martensitic transformations
1419	S. Chawla & S.M. Lenhart,	Application of optimal control theory to in Situ bioremediation
1420	B. Li & M. Luskin,	Nonconforming finite element approximation of crystalline microstructure
1421	H. Kang & J.K. Seo,	Inverse conductivity problem with one measurement: Uniqueness of balls in $\mathbb{R}^3$
1422	Avner Friedman & Robert Gulliver,	Organizers, Mathematical modeling for instructors, July 29 – August 16, 1996
1423	G. Friesecke,	Pair correlations and exchange phenomena in the free electron gas
1424	Y.A. Li & P.J. Olver,	Convergence of solitary-wave solutions in a perturbed Bi-Hamiltonian dynamical system I. Compactons and Peakons II. Complex Analytic Behavior III. Convergence to Non-Analytic Solutions
1425	C. Huang,	On boundary regularity of vortex patches for 3D incompressible euler systems
1426	C. Huang,	A free boundary problem with nonlinear jump and kinetics on the free boundaries
1427	X. Chen, C. Huang & J. Zhao,	A nonlinear parabolic equation modeling surfactant diffusion
1428	A. Friedman & B. Hu,	Optimal control of chemical vapor deposition reactor
1429	A. Friedman & B. Hu,	A non-stationary multi-scale oscillating free boundary for the Laplace and heat equations
1430	X. Chen,	Existence, uniqueness, and asymptotic stability of traveling waves in nonlocal evolution equations
1431	J. Yong,	Finding adapted solutions of forward-backward stochastic differential equations – Methods of continuation
1432	J. Yong,	Linear forward-backward stochastic differential equations
1433	D.A. Dawson & M.A. Kouritzin,	Invariance principles for parabolic equations with random coefficients
1434	R. Lipton,	Energy minimizing configurations for mixtures of two imperfectly bonded conductors
1435	D.C. Dobson & F. Santosa,	Nondestructive evaluation of plates using Eddy current methods
1436	W. Littman & B. Liu,	On the spectral properties and stabilization of acoustic flow
1437	S. Sarkar & S. Sundar Sarkar,	Normal distribution as a method for data replication in a parallel data server
1438	S. Sarkar & S. Sundar Sarkar,	Parallel view materialization with dynamic load balancing: A graph theoretic approach
1439	S. Sarkar & S. Sundar Sarkar,	Internet and relational databases in a multi-tier client/server model
1440	J. Liang & S. Subramaniam,	Numerical computing of molecular electrostatics through boundary integral equations
1441	J. Wu,	Inviscid limits and regularity estimates for the solutions of the 2-D dissipative quasi-geostrophic equations
1442	P. Constantin & J. Wu,	Statistical solutions of the Navier-Stokes equations on the phase space of vorticity and the inviscid limits
1443	M.A. Kouritzin,	Stochastic processes and perturbation problems defined by parabolic equations with a small parameter
1444	M.A. Kouritzin,	Approximations for singularly perturbed parabolic equations of arbitrary order
1445	A. Novick-Cohen	Triple junction motion for Allen-Cahn/Cahn-Hilliard systems
1446	P. Klouček,	Approximations of the laminated microstructures

- 1447 S. Sarkar & S.S. Sarkar, A graph theoretic approach for parallel view materialization with dynamic load balancing
- 1448 S. Chawla, A minmax problem for parabolic systems with competitive interactions
- 1449 B. Luong & F. Santosa, Quantitative imaging of corrosion in plates by Eddy current methods
- 1450 R. Jordan & B. Turkington, Ideal magnetofluid turbulence in two dimensions
- 1451 M. Fels & P.J. Olver, Moving coframes. I. A practical algorithm
- 1452 S.Y. Maliassov, On the Schwarz alternating method for eigenvalue problems
- 1453 R. Lipton, Design of particle reinforced heat conducting composites with interfacial thermal barriers
- 1454 J. Berg, A. Yezzi, & A. Tannenbaum, Phase transitions, curve evolution, and the control of semiconductor manufacturing processes
- 1455 G. Avalos & I. Lasiecka, Uniform decay rates of solutions to a structural acoustics model with nonlinear dissipation
- 1456 M. Nitsche, Siemens/IMA technical report
- 1457 L. Wang, J.A. Cox, & A. Friedman, Model analysis of homogeneous optical waveguides by boundary integral method
- 1458 C.P. Fung & S. Lototsky, Nonlinear filtering: Separation of parameters and observations using Galerkin approximation and Wiener chaos decomposition
- 1459 S. Northshield, Several proofs of Ihara's theorem
- 1460 T. Aktosun, M. Klaus & C. van der Mee, Wave scattering in one dimension with absorption
- 1461 F. Santosa, M. Vogelius, & J.-M. Xu, An effective nonlinear boundary condition for a corroding surface. Identification of the damage based on electrostatic data
- 1462 J. Wu, Well-posedness of a semilinear heat equation with weak initial data
- 1463 J. Wu, Quasi-geostrophic type equations with weak initial data
- 1464 J. Ma & J. Yong, Approximate solvability of forward-backward stochastic differential equations
- 1465 T.-P. Tsai, On Leray's self-similar solutions of the Navier-Stokes equations satisfying local energy estimates
- 1466 M.K. Gobbert, T.P. Merchant, L.J. Borucki, & T.S. Cale, A multiscale simulator for low pressure chemical vapor deposition
- 1467 M.C. Tesi, E.J. Janse van Rensburg, E. Orlandini, & S.G. Whittington, Torsion of polygons in $\mathcal{Z}^3$
- 1468 M. Grinfeld & A. Novick-Cohen, The viscous Cahn-Hilliard equation: Morse decomposition and structure of the global attractor
- 1469 M. Mascagni, Polynomial versus matrix methods for leap-ahead in shift-register type pseudorandom number generators
- 1470 M. Mascagni, Parallel linear congruential generators with prime moduli
- 1471 B. Li & M. Luskin, Approximation of a Martensitic laminate with varying volume fractions
- 1472 D. Yang, Stabilized schemes for mixed finite element methods with applications to elasticity and compressible flow problems
- 1473 P.J. Olver & V.V. Sokolov, Integrable evolution equations on associative algebras
- 1474 A. Bondarenko, Singular structure of the fundamental solution of the transport equation
- 1475 S. Sarkar & S.S. Sarkar, Views and data mining in a parallel data server
- 1476 P.E. Bigeleisen & M. Cheney, Models for an anesthesia breathing circuit
- 1477 R. Lipton, Influence of interfacial surface conduction on the DC electrical conductivity of particle reinforced composites
- 1478 B. Cockburn & C.-W. Shu, The local discontinuous Galerkin method for time-dependent convection-diffusion systems
- 1479 A. Friedman & F. Reitich, Asymptotic Behavior of Solutions of Coagulation-Fragmentation Models
- 1480 A. Friedman & C. Huang, Averaged Motion Approach to Brownian Dynamics
- 1481 T. Aktosun, M. Klaus & C. van der Mee, Inverse Scattering in One-Dimensional Nonconservative Media
- 1482 X. Feng, Absorbing Boundary Conditions for Electromagnetic Wave Propagation
- 1483 J.E. Flaherty, R.M. Loy, M.S. Shephard, B.K. Szymanski, J.D. Teresco & L.H. Ziantz, Adaptive Local Refinement with Octree Load-Balancing for the Parallel Solution of Three-Dimensional Conservation Laws
- 1484 D. Hömberg & J. Sokolowski, Optimal Control of Laser Hardening
- 1485 M. Kružík, Numerical Approach to Double Well Problems
- 1486 S. Lototsky, Parameter Estimation for Stochastic Parabolic Equations: Asymptotic Properties of a Two-Dimensional Projection Based Estimate
- 1487 D.E. Norman, A Model for Chemically Reacting Fluid Flows: Weak Solutions and Global Attractors
- 1488 S. Sarkar & S.S. Sarkar, Data Mining on Views in Relational Databases Using Bayesian Networks
- 1489 D. Yang, Dynamic Finite Element Methods for Second Order Parabolic Equations
- 1490 S. Evje & K.H. Karlsen A Note on Viscous Splitting of Degenerate Convection-Diffusion Equations
- 1491 X. Feng, Analysis of A Domain Decomposition Method for The Nearly Elastic Wave Equations Based on Mixed Finite Element Methods
- 1492 B. Cockburn & C.-W. Shu, The Runge-Kutta Discontinuous Galerkin Method for Conservation Laws V: Multidimensional Systems
- 1493 P. Korman & Y. Li, On The Exactness of an S-Shaped Bifurcation Curve
- 1494 M. Kružík, DiPerna—Majda Measures and Uniform Integrability