STOCHASTIC DIFFERENTIAL EQUATIONS IN
DUALS OF NUCLEAR SPACES WITH SOME APPLICATIONS

BY

G. KALLIANPUR

IMA Preprint Series # 244

May 1986

INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
UNIVERSITY OF MINNESOTA
514 Vincent Hall
206 Church Street S.E.
Minneapolis, Minnesota 55455

#	Author(s)	Title
120	D.R.J. Chillingworth,	Three Introductory Lectures on Differential Topology and its Applications
121	Giorgio Vergara Caffarelli,	Green's Formulas for Linearized Problems with Live Loads
122	F. Chiarenza and M. Garofalo,	Unique Continuation for Nonnegative Solutions of Schrödinger Operators
123	J.L. Ericksen,	Constitutive Theory for some Constrained Elastic Crystals
124	Minoru Murata,	Positive solutions of Schrödinger Equations
125	John Maddocks and Gareth P. Barry,	A Model for Twinning
126	M. Kaneko and M. Wooders,	The Core of a Game with a Continuum of Players and Finite Coalitions: Nonemptiness with Bounded Sizes of Coalitions
127	William Zame,	Equilibria in Production Economies with an Infinite Dimensional Commodity Space
128	Myrna Holtz Wooders,	A Tiebout Theorem
129	Abstracts for the Workshop on Theory and Applications of Liquid Crystals	
130	Yoshihiko Giga,	A Remark on A Priori Bounds for Global Solutions of Semilinear Heat Equations
131	M. Chipot and G. Vergara-Caffarelli,	The N-Membranes Problem
132	P.L. Lions and P.E. Souganidis,	Differential Games and Directional Derivatives of Viscosity Solutions of Isaacs' Equations II
133	G. Capriz and P. Giovine,	On Virtual Effects During Diffusion of a Dissipative Medium in a Suspension
134	Fall Quarter Seminar Abstracts	
135	Umberto Mosco,	Wiener Criterion and Potential Estimates for the Obstacle Problem
136	Chi-Sing Ma,	Dynamic Admissible States, Negative Absolute Temperature, and the Entropy Maximum Principle
137	Abstracts for the Workshop on Oscillation Theory, Computation, and Methods of Compensated Compactness	
138	Arie Leizarowitz,	Tracking Nonperiodic Trajectories with the Overtaking Criterion
139	Arie Leizarowitz,	Convex Sets in R^n as Affine Images of some Fixed Set in R^n
140	Arie Leizarowitz,	Stochastic Tracking with the Overtaking Criterion
141	Abstracts from the Workshop on Amorphous Polymers and Non-Newtonian Fluids	
142	Winter Quarter Seminar Abstracts	
143	D.G. Aronson and J.L. Vazquez,	The Porous Medium Equation as a Finite-speed Approximation to a Hamilton-Jacobi Equation
144	E. Sanchez-Palencia and R. Weinberger,	On the Edge Singularities of a Composite Conducting Medium
145	Jon C. Luke,	Soliton Solutions in a Class of Fully Discrete Nonlinear Wave Equations
146	Chi-Sing Ma and R. Cohen,	A Coordinate-Free Approach to the Kinematics of Membranes
147	J.L. Lions,	Asymptotic Problems in Distributed Systems
148	Reiner Lauterbach,	An Example of Symmetry Breaking with Submaximal Isotropy Subgroup
49	Abstracts from the Workshop on Metastability and Incompletely Posed Problems	
50	B. Bozar-Karakiewicz and Jerry Bona,	Wave-dominated Shelves: A Model of Sand-Ridge Formation by Progressive, Infragravity Waves
51	Abstracts from the Workshop on Dynamical Problems in Continuum Physics	
52	V.I. Oikar,	The problem of Embedding S^n into R^n with Prescribed Gauss
53	R. Batra,	The force on a Lattice Defect in an Elastic Body
54	J. Fleckinger and Michael Lapidus,	Eigenvalues of Elliptic Boundary Value Problems with and Infinite Weight Function
55	R. Kohn and M. Vogelius,	Thin Plates with Rapidly Varying Thickness, and Their relation to Structural Optimization
56	M. Gurtin,	Some Results and Conjectures in the Gradient Theory of Phase Transitions
57	A. Novick-Cohen,	Energy Methods for the Cahn-Hilliard Equation
158	M. Siroli and U. Mosco,	Wiener Estimates for Parabolic Obstacle Problems
159	E. Bennett and M. Zeme,	Prices and Bargaining in Cooperative Games
160	M.A. Harris and Y. Sibuya,	The n-th Roots of Solutions of Linear Ordinary Differential Equations
161	William F. Beatty,	Some Dynamical Problems in Continuum Physics
162	P. Sautman and D. Phillips,	Large-Time Behavior of Solutions to a Scalar Conservation Law in Several Space Dimensions
163	A. Novick-Cohen,	Interfacial Instabilities in Directional Solidification of Dilute Binary Alloys: The Kuramoto-Sivashinsky Equations
164	R.F. Weinberger,	On Metastable Patterns in Parabolic Systems
165	D. Arnold and R.S. Falk,	Continuous Dependence on the Elastic Coefficients for a Class of Anisotropic Materials
166	J.J. Sackman,	The Boundary Value Problems for Non-linear Elliptic Equation and the Maximum Principle for Euler-Lagrange Equations
167	Ingo Müller,	Gases and Rubbers
168	Ingo Müller,	Pseudoelasticity in Shape Memory Alloys - an Extreme Case of Thermoelasticity
169	Luis Morgado,	Persistence and Smoothness of Hyperbolic Invariant Manifolds for Functional Differential Equations
170	A. Bancelian and M. Vogelius,	Homogenization Limits of the Equations of Elasticity in Thin Domains
171	H.C. Simpson and S.J. Spector,	On Hadamard Stability in Finite Elasticity
172	J.L. Vazquez and C. Vazquez,	Isolated Singularities of the Solutions of the Schrödinger Equation with a Radial Potential
173	G. Dal Maso and B. Mosco,	Wiener's Criterion and Γ -Convergence
174	John H. Maddocks,	Stability and Folds
175	R. Hardt and D. Kinderlehrer,	Existence and Partial Regularity of Static Liquid Crystal Configurations
176	M. Morokawa,	Construction of Smooth Ergodic Cocycles for Systems with Fast Periodic Approximations
177	J.L. Ericksen,	Stable Equilibrium Configurations of Elastic Crystals
178	Patricio Aviles,	Local Behavior of Solutions of Some Elliptic Equations
179	S.-N. Chow and R. Lauterbach,	A Bifurcation Theorem for Critical Points of Variational Problems
180	R. Pego,	Phase Transitions: Stability and Admissibility in One Dimensional Nonlinear Viscoelasticity
181	Mariano Gagliata,	Quadratic Functions and Partial Regularity
182	J. Bona,	Fully Discrete Galerkin Methods for the Korteweg De Vries Equation
183	J. Maddocks and J. Keller,	Mechanics of Robes
184	F. Bernis,	Qualitative Properties for some nonlinear higher order Nonlinear Higher Order Parabolic Equations with Absorption
185	F. Bernis,	Finite Speed of Propagation and Asymptotic Rates for some Nonlinear Higher Order Parabolic Equations with Minimal Strategy Spaces
186	S. Reichstein and S. Reiter,	Game Forms with Minimal Strategy Spaces
187	T. Ding,	An Answer to Littewood's Problem on Boundedness
188	J. Rubinstein and R. Maierl,	Dispersion and Convection in Periodic Media
189	M.H. Fleming and P.E. Souganidis,	Asymptotic Series and the Method of Vanishing Viscosity
190	H. Beirão da Veiga,	Existence and Asymptotic Behavior for Strong Solutions of Navier-Stokes Equations in the Whole Space
191	L.A. Caffarelli, J.L. Vazquez, and M.L. Woiwinski,	Lipschitz Continuity of Solutions and Interfaces of the N-Dimensional Porous Medium Equation
192	R. Johnson,	m-Functions and Floquet Exponents for Linear Differential Systems
193	F.V. Atkinson and L.A. Peletier,	Ground States and Dirichlet Problems for $-\Delta = F(U)$ in R^n
194	G. Dal Maso, U. Mosco,	The Wiener Modulus of a Radial Measure
195	H. A. Levine and H.F. Weinberger,	Inequalities between Dirichlet and Neuman Eigenvalues
196	J. Rubinstein,	On the Macroscopic Description of Slow Viscous Flow Past a Random Array of Spheres
197	G. Dal Maso and B. Mosco,	Wiener Criteria and Energy Decay for Relaxed Dirichlet Problems

#	Author(s)	Title	#	Author(s)	Title
198	V. Ollker and P. Waltman,	On the Monge-Ampere Equation Arising in the Reflect Mapping Problem	237	W.H. Fleming, S.J. Sheu and H.M. Soner,	On Existence of the Dominant Eigenfunction and its Application to the Large Deviation Properties of an Ergodic Markov Process
199	M. Chipot, D. Kinderlehrer and L. Caffarelli,	Some Smoothness Properties of Linear Laminates	238	R. Jensen and P.E. Sogkanidis,	A Regularity Result for Viscosity Solutions of Hamilton-Jacobi Equations in one Space Dimension
200	Y. Giga and R. Kohn,	Characterizing Blow-up Using Similarity Variables	239	B. Boczar-Karakli, J.L. Bona and D.L. Cohen,	Intersection of Shallow-Water Waves and Bottom Topography
201	P. Cannarsa and H. M. Soner,	On the Singularities of the Viscosity Solutions to Hamilton-Jacobi-Bellman Equations	240	F. Colom'us and M. Klenemann,	Infinite Time Optimal Control and Periodicity
202	Andrew Majda,	Nonlinear Geometric Optics for Hyperbolic Systems of Conservation Laws	241	Harry Kesten,	Scaling Relations for 2D-Percolation
203	G. Buttazzo, G. Dal Maso and U. Mosco,	A Derivation Theorem for Capacities with Respect to a Radon Measure	242	A. Leizarowitz,	Infinite Horizon Optimization for Markov Process with Finite States Spaces
204	S. Corin, M. Mehrotra,	On the identification of Material Symmetry for Anisotropic Elastic Materials	243	Louis H.Y. Chen,	The Rate of Convergence in A Central Limit Theorem for Dependent Random Variables with Arbitrary Index Set
205	R.W.R. Darling,	Constructing Nonhomomorphic Stochastic Flows.			
206	M. Chipot,	On the Reynolds Lubrication Equation			
207	R.V. Kohn and G.W. Milton,	On Bounding the Effective Conductivity of Anisotropic Composites			
208	I.J. Bakelman,	Notes Concerning the Torsion of Hardening Rods and Its N-Dimensional Generalizations			
209	I.J. Bakelman,	The Boundary Value Problems for Non-Linear Elliptic Equations II.			
210	Guangfu Gong & Mingping Qian,	On the Large Deviation Functions of Markov Chains			
211	Arie Leizarowitz,	Control Problems with Random and Progressively Known Targets			
212	R.W.R. Darling,	Ergodicity of a Measure-Valued Markov Chain Induced by Random Transformations			
213	G. Gong, M. Qian & Zhonglin Zhao,	Killed Diffusions and its Conditioning			
214	Arie Leizarowitz,	Controlling Diffusion Processes on Infinite Horizon with the Overtaking Criterion			
215	Millard Seatty,	The Poisson Function of Finite Elasticity			
216	David Ternan,	Traveling Wave Solutions Arising from a Combustion Model			
217	Yuh-Jia Lee,	Sharp Inequalities and Regularity of Heat Semi-Group on Infinite Dimensional Spaces			
218	D. Stroock,	Lecture Notes			
219	Claudio Canuto,	Spectral Methods and Maximum Principle			
220	Thomas O'Brien,	A Two Parameter Family of Pension Contribution Functions and Stochastic Optimization			
221	Takayuki Hida,	Analysis of Brownian Functionals			
222	Leonid Flunkicz,	On Informational Decentralization and Efficiency of Resource Allocation Mechanisms			
223	E.B. Fabes and D.W. Stroock,	A New Proof of Moser's Parabolic Harnack Inequality via the Old Ideas of Nash			
224	Minoru Murata,	Structure of Positive Solution to $(-\Delta + V)u = 0$ in R^n			
225	Paul Dupuis,	Large Deviations Analysis of Reflected Diffusions and Constrained Stochastic Approximation Algorithms in Convex Sets			
226	F. Ferris,	Existence Results for Doubly Nonlinear Higher Order Parabolic Equations on Unbounded Domains.			
227	S. Orey and S. Pelikan,	Large Deviations Principles for Stationary Processes			
228	R. Gulliver and S. Hildebrandt,	Boundary Configurations Spanning Continua of Minimal Surfaces.			
229	J. Baxter, G. Dal Maso & U. Mosco,	Stopping Times and T-Convergence.			
230	Julio Boulliet,	Self-Similar Solutions, Having Jumps and Intervals of Constancy of a Diffusion-heat Conduction Equation			
231	R. Hardt, D. Kinderlehrer & F.-H. Lin,	A Remark About the Stability of Smooth Equilibrium Configurations of Static Liquid Crystals.			
232	M. Chipot and M. Luskin,	The Compressible Reynolds Lubrication Equation			
233	J.H. Maddocks,	A Model for Disclinations in Nematic Liquid Crystal			
234	C. Foias, G.R. Sell and R. Temam,	Inertial Manifolds for Nonlinear Evolutionary Equations			
235	P.L. Chow,	Expectation Functionals Associated with Some Stochastic Evolution Equations			
236	Giuseppe Buttazzo,	Reinforcement by a Thin Layer with Oscillating Thickness			

STOCHASTIC DIFFERENTIAL EQUATIONS IN
DUALS OF NUCLEAR SPACES WITH SOME APPLICATIONS

by

G. Kallianpur

Center for Stochastic Processes
University of North Carolina
Chapel Hill, North Carolina

INTRODUCTION

In these lectures I have aimed at giving an elementary introduction to certain types of stochastic differential equations in infinite dimensional spaces. The material is relatively self contained and should be easily accessible to graduate students. I have freely drawn on some of the existing work in the field, especially the pioneering work of K. Itô as well as some extensions appearing in the papers of I. Mitoma, R. Wolpert and myself, the thesis of S.K. Christensen as well as some recent work which I have been doing jointly with V. Perez-Abreu.

My warm thanks are due to Professor Victor Perez-Abreu for writing up these notes.

G. Kallianpur

March 25, 1986

LECTURE I

COUNTABLY HILBERTIAN NUCLEAR SPACES

In this lecture we introduce Countably Hilbertian Nuclear (CHN) spaces and give some examples to illustrate why these infinite dimensional spaces are convenient for the study of some practical problems, e.g. those occurring in stochastic evolutions.

Let Φ be a (real) linear space whose topology τ is given by an increasing sequence $\|\cdot\|_r$ $r \in \mathbb{N}$ of Hilbertian norms. Let Φ_r be the Hilbert space completion of Φ w.r.t. $\|\cdot\|_r$ and assume that

$$\Phi = \bigcap_{r=1}^{\infty} \Phi_r.$$

Then (Φ, τ) is a Frechet space with metric

$$\rho(\phi, \psi) = \sum_{n=1}^{\infty} \frac{1}{2^n} \frac{\|\phi - \psi\|_n}{1 + \|\phi - \psi\|_n}$$

and (Φ, ρ) is called a Countably Hilbertian Space. Since for $n < m$

$$\|\phi\|_n \leq \|\phi\|_m \quad \phi \in \Phi$$

then

$$\Phi_m \subseteq \Phi_n \quad m > n.$$

A countably Hilbertian space Φ is called nuclear if for each $n \geq 0$ there exists $m > n$ such that the canonical injection $i: \Phi_m \hookrightarrow \Phi_n$ is Hilbert-Schmidt i.e. if $\{\phi_j\}_{j \geq 1}$ is a complete orthonormal system (CONS) in Φ_m then we have

$$\sum_{j=1}^{\infty} \|\phi_j\|_n^2 < \infty.$$

Let Φ'_n be the dual (Hilbert) space of Φ_n and for $f \in \Phi'_n$ let

$$\|f\|_{-n} = \sup_{\|\phi\|_n \leq 1} |f[\phi]|$$

since for $n \leq m$ $\|\phi\|_n \leq \|\phi\|_m$ then

$$\Phi'_n \subseteq \Phi'_m \quad n \leq m.$$

Let Φ' be the topological dual space of Φ with the strong topology, which is given by the complete system of neighborhoods of zero given by sets of the form, $\{f \in \Phi' : \|f\|_B < \epsilon\}$, $\epsilon > 0$, where,

$$\|f\|_B = \sup\{|f[\phi]| : \phi \in B\} \quad B \text{ a bounded set in } \Phi.$$

Then

$$\Phi' = \bigcup_{n=1}^{\infty} \Phi'_n.$$

It is important to note that this topology cannot be given by a countable family of seminorms.

Suppose there is an inner product $\langle \cdot, \cdot \rangle_H$ on Φ which is continuous in the τ -topology of Φ . Let H be the Hilbert space completion of Φ w.r.t. $\langle \cdot, \cdot \rangle_H$. Then the triplet

$$\Phi \hookrightarrow H \hookrightarrow \Phi'$$

is called a rigged Hilbert space or a Gelfand triplet. The Hilbert space H may be one of the Hilbert spaces Φ_r defining the topology of Φ but this is not always the case as we shall illustrate later on.

Example 1.1 Let $\mathcal{S}(R)$ be the space of rapidly decreasing functions on R . The usual definition of $\mathcal{S}(R)$ is the following:

$$\phi \in \mathcal{S}(\mathbb{R}) \text{ iff } \phi \in C^\infty \text{ and } \sup_{x \in \mathbb{R}} |x^\alpha \phi^{(\beta)}(x)| < \infty \quad \alpha, \beta \in \mathbb{N}$$

and the topology of $\mathcal{S}(\mathbb{R})$ (Fréchet) is given by the family of seminorms

$$\{\|\phi\|_{\alpha, \beta} = \sup_{x \in \mathbb{R}} |x^\alpha \phi^{(\beta)}(x)|, \alpha, \beta \in \mathbb{N}\}.$$

The space $\mathcal{S}(\mathbb{R})$ can also be defined using the following sequence of Hilbertian norms: Let

$$H_n(x) = \frac{(-1)^n}{n!} e^{x^2/2} \left(\frac{d}{dx} \right)^n e^{-x^2/2}$$

and

$$g(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}.$$

Then the sequence of Hermite functions $\{\phi_n\}_{n \geq 1}$

$$\phi_{n+1}(x) = \sqrt{g(x)} (n!)^{1/2} H_n(x) \quad n \geq 0 \quad (1.1)$$

is a CONS in $L^2(\mathbb{R})$. Then ([5])

$$\mathcal{S}(\mathbb{R}) = \{\phi \in L^2(\mathbb{R}) : \|\phi\|_p^2 < \infty \quad \forall p \in \mathbb{R}\}.$$

Let $\mathcal{S}_p$ be the completion of $\mathcal{S}(\mathbb{R})$ w.r.t. $\|\cdot\|_p$, then $\mathcal{S}_0 = L^2(\mathbb{R})$

$$\mathcal{S}_p \subset \mathcal{S}_q \quad p > q$$

and

$$\begin{aligned} \mathcal{S}(\mathbb{R}) &= \bigcap_p \mathcal{S}_p \\ \mathcal{S}(\mathbb{R})' &= \bigcup_p \mathcal{S}'_p \end{aligned}$$

The Schwarz topology on $\mathcal{S}$ is the topology determined by $\{\|\cdot\|_p, p \in \mathbb{R}\}$ or equivalent by the countable family $\{\|\cdot\|_p : p \in \mathbb{N}\}$.

Let $f, g \in \mathcal{S}$, then for all $p \in \mathbb{R}$

$$|f[g]|^2 = \left| \sum_n \langle f, \phi_n \rangle_0 \langle g, \phi_n \rangle_0 \right|^2 \leq$$

$$\left(\sum_n \langle f, \phi_n \rangle_0^2 \left(n + \frac{1}{2}\right)^{-2p} \right) \left(\sum_n \langle g, \phi_n \rangle_0^2 \left(n + \frac{1}{2}\right)^{2p} \right) = \|f\|_{-p}^2 \|g\|_p^2$$

which shows that $\mathcal{S}_{-p}$ is the dual of $\mathcal{S}_p$ and therefore

$$\mathcal{S}' = \bigcup_{p \geq 0} \mathcal{S}_{-p}.$$

Finally since for any $p \in \mathbb{R}$ $\{(n + \frac{1}{2})^{-p} \phi_n\}$ is a CONS for $\mathcal{S}_p$ then if $p > q + \frac{1}{2}$

$$\sum_{n=1}^{\infty} \|(n + \frac{1}{2})^{-p} \phi_n\|_q^2 = \sum_{n=1}^{\infty} (n + \frac{1}{2})^{-2(p-q)} < \infty$$

i.e. the canonical injection $\mathcal{S}_p \hookrightarrow \mathcal{S}_q$ is Hilbert-Schmidt for $p > q + \frac{1}{2}$.

Hence $\mathcal{S}(\mathbb{R})$ is a CHNS and if $H = L^2(\mathbb{R})$

$$\mathcal{S}(\mathbb{R}) \hookrightarrow_H \mathcal{S}(\mathbb{R})'$$

is a rigged Hilbert space. $\mathcal{S}(\mathbb{R})'$ is the familiar space of tempered distributions.

The following observation will be useful later (see Remark 1.2): If L is the operator on $H = L^2(\mathbb{R})$ defined by

$$-L = \frac{d^2}{dx^2} - \frac{x^2}{4}$$

then $L\phi_n = \lambda_n \phi_n$ where $\lambda_n = n - \frac{1}{2}$ and for $r > \frac{1}{2}$

$$\sum_{j=1}^{\infty} \|(I+L)^{-r} \phi_j\|_H^2 = \sum_{j=1}^{\infty} (1 + \lambda_j)^{-2r} = \sum_{j=1}^{\infty} (j + \frac{1}{2})^{-2r} < \infty. \quad (1.2)$$

In these lectures we consider the following model for deterministic evolution:

Model. Suppose we have a rigged Hilbert space $\Phi \hookrightarrow H \hookrightarrow \Phi'$ on which is defined a continuous linear operator $A: \Phi \rightarrow \Phi$ and a strongly continuous semigroup $(T_t)_{t \geq 0}$ on H such that the following conditions hold:

- a) $T_t \Phi \subseteq \Phi \quad \forall t \geq 0.$
- b) The restriction $T_t|_{\Phi} : \Phi \rightarrow \Phi$ is τ -continuous $\forall t \geq 0.$
- c) $t \rightarrow T_t \phi$ is continuous $\forall \phi \in \Phi.$
- d) The generator of T_t on H coincide with A on $\Phi.$

If Φ, H, Φ' are already given, (a)-(d) is a restriction on the type of T_t that can be considered. However, it is important to observe that in practical problems, physical considerations give no idea of the rigged Hilbert space $\Phi \hookrightarrow H \hookrightarrow \Phi'$ and only the Hilbert space H and the semigroup T_t are naturally given in the problem, and hence the Schwarz space cannot be chosen in advance.

The following example gives a method of choosing Φ and Φ' when T_t is given and satisfies certain conditions. Then we will introduce some examples where we can set up an infinite dimensional stochastic differential equation by choosing Φ and Φ' suitably so as to satisfy the above conditions.

Given a rigged Hilbert space $\Phi \hookrightarrow H \hookrightarrow \Phi'$, a semigroup $(T_t)_{t \geq 0}$ satisfying the above conditions will be called compatible with (Φ, H, Φ') or equivalently we will refer to (Φ, H, T_t) as a compatible family.

EXAMPLE 1.2 (A class of examples of (Φ, H, Φ', T_t)). Let H be a real separable Hilbert space and $A = -L$ be a closed densely defined self adjoint operator on H s.t. $\langle -L\phi, \phi \rangle_H \leq 0$ for $\phi \in \text{Dom}(L)$. Let T_t be the semigroup on H determined by A . Further assume that some power of the resolvent of L is a

Hilbert-Schmidt operator i.e.

$$\exists r_1 \text{ s.t. } (\lambda I + L)^{-r_1} \text{ is Hilbert-Schmidt} \quad (1.3)$$

This condition helps to find an appropriate CHNS Φ for the model, as we shall now indicate: it implies that there is a CONS $\{\phi_j\}_{j \geq 1}$ in H s.t.

$$L\phi_j = \lambda_j \phi_j \quad \forall j \geq 1$$

and $0 < \lambda_1 < \lambda_2 < \dots$. Take $\lambda = 1$ and define

$$\begin{aligned} \Phi &= \{\phi \in H: \|(I + L)^r \phi\|_H^2 < \infty \quad \forall r \in \mathbb{R}\} \\ &= \{\phi \in H: \sum_{j=1}^{\infty} (1 + \lambda_j)^{2r} \langle \phi, \phi_j \rangle_H^2 < \infty \quad \forall r \in \mathbb{R}\} \end{aligned} \quad (1.4)$$

Define the inner product $\langle \cdot, \cdot \rangle_r$ on Φ by

$$\langle \phi, \psi \rangle_r := \sum_{j=1}^{\infty} (1 + \lambda_j)^{2r} \langle \phi, \phi_j \rangle_H \langle \psi, \phi_j \rangle_H \quad (1.5)$$

and

$$\|\phi\|_r^2 = \langle \phi, \phi \rangle_r.$$

Let Φ_r be the $\|\cdot\|_r$ - completion of Φ . We then have

$$\Phi = \bigcap_r \Phi_r, \quad \Phi' = \bigcup_r \Phi'_r$$

and for $r \leq s$, $\|\phi\|_r \leq \|\phi\|_s$ and so $\Phi_s \subseteq \Phi_r$ with $\Phi_0 = H$. Condition (1.3) implies that the canonical injection $\Phi_p \hookrightarrow \Phi_r$ is Hilbert Schmidt for $p > r + r_1$ and therefore Φ is a CHNS.

For each $r > 0$ Φ_{-r} and Φ_r are in duality under the pairing

$$\eta[\phi] := \sum_{j=1}^{\infty} \langle \eta, \phi_j \rangle_{-r} \langle \phi, \phi_j \rangle_r \quad \eta \in \Phi_{-r}, \quad \phi \in \Phi_r \quad (1.6)$$

and therefore $\phi_{-r} = \phi_r'$. We also have that $\{\phi_j\}_{j \geq 1}$ is a complete orthogonal system (not normal) in ϕ_r for all $r \in \mathbb{R}$.

From now on we will write $\langle \cdot, \cdot \rangle_0 = \langle \cdot, \cdot \rangle_H$ and

$$\theta_1 = \sum_{j=1}^{\infty} (1 + \lambda_j)^{-2r_1} < \infty. \quad (1.7)$$

Now we shall prove that the semigroup T_t satisfies conditions (a)-(d) for our above model for deterministic evolution:

Condition (1.3) implies that $-L$ generates a contraction semigroup T_t on H . For $\phi \in \Phi$ and $t \geq 0$ we have

$$T_t \phi = \sum_{j=1}^{\infty} e^{-t\lambda_j} \langle \phi, \phi_j \rangle_0 \phi_j \in \Phi$$

which implies (a). Next for $t \geq 0$ and $\phi \in \Phi$

$$\|T_t \phi\|_r^2 = \sum_{j=1}^{\infty} e^{-2t\lambda_j} (1 + \lambda_j)^{2r} \langle \phi, \phi_j \rangle_0^2 \leq \sum_{j=1}^{\infty} (1 + \lambda_j)^{2r} \langle \phi, \phi_j \rangle_0^2 = \|\phi\|_r^2.$$

Then since $\psi_n \rightarrow 0$ in Φ if and only if $\|\psi_n\|_r \rightarrow 0 \quad \forall r \in \mathbb{R}$,

$$\|T_t \psi_n\|_r \rightarrow 0 \quad \forall r \in \mathbb{R} \Rightarrow T_t \psi_n \rightarrow 0 \text{ in } \Phi$$

and therefore condition (b) is satisfied.

Now for $s, t \in \mathbb{T}$ and $\phi \in \Phi$

$$\|T_t \phi - T_s \phi\|^2 = \sum_{j=1}^{\infty} (e^{-t\lambda_j} - e^{-s\lambda_j})^2 \langle \phi, \phi_j \rangle_0^2 (1 + \lambda_j)^{2r}$$

and for each $j \geq 1$

$$(e^{-t\lambda_j} - e^{-s\lambda_j})^2 \langle \phi, \phi_j \rangle_0^2 (1 + \lambda_j)^2 \leq 4 \langle \phi, \phi_j \rangle_0^2 (1 + \lambda_j)^{2r}.$$

Since $e^{-t\lambda_j}$ is continuous on t for all $j \geq 1$ then by the Dominated convergence theorem

$$\lim_{t \rightarrow s} \|T_t \phi - T_s \phi\|_r \rightarrow 0 \quad \forall \phi \in \Phi, \quad r \in \mathbb{R}$$

which implies (c).

Now to prove (d) let $\phi \in \Phi$ and define $\psi_n = \sum_{j=1}^n \langle \phi, \phi_j \rangle_0 \phi_j$. Then $\psi_n \rightarrow \phi$ on Φ ,

$$-L\psi_n = \sum_{j=1}^n \langle \phi, \phi_j \rangle_0 L\phi_j = - \sum_{j=1}^n \lambda_j \langle \phi, \phi_j \rangle_0 \phi_j$$

and for $m > n$ and $r \in \mathbb{R}$

$$\left\| \sum_{j=n+1}^m -\lambda_j \langle \phi, \phi_j \rangle_0 \phi_j \right\|_r^2 = \sum_{j=n+1}^{\infty} \lambda^2 (1 + \lambda_j)^{2r} \langle \phi, \phi_j \rangle_0^2 \rightarrow 0 \quad \text{as } n, m \rightarrow \infty.$$

Hence if $\psi = - \sum_{j=1}^{\infty} \lambda_j \langle \phi, \phi_j \rangle_0 \phi_j$ we have $-L\psi_n \rightarrow \psi$ on Φ . But since $-L$ is closed in H and $\|\cdot\|_H$ is ϕ -continuous, then $\phi \in \text{Dom}(L)$ and $\psi = -L\phi$ $\phi \in \Phi$, i.e.

$$-L\phi = - \sum_{j=1}^{\infty} \lambda_j \langle \phi, \phi_j \rangle_0 \phi_j$$

and

$$\|-L\phi\|_r^2 \leq \|\phi\|_{r+1}^2 \quad \forall \phi \in \Phi \quad r \in \mathbb{R}.$$

Hence, $-L\Phi \subseteq \Phi$ and $-L$ is Φ -continuous which implies (d).

Remarks

1.1. A compatible family (Φ, H, T_t) or (Φ, H, L) is called a special compatible family if the generator L satisfies condition (1.3) and Φ is constructed as in Example 1.2, i.e. Φ is given by (1.4).

1.2. The Schwarz space $\mathcal{S}(R)$ of Example 1.1 may be obtained in the framework of the last example by taking

$$-L = \frac{d^2}{dx^2} - \frac{x^2}{4}, \quad H = L^2(R), \quad \lambda_j = j - \frac{1}{2} \quad j \geq 1$$

and $\{\phi_j\}$ the Hermite functionals given by (1.1). Then from (1.2) we have that $-L$ satisfies condition (1.3) for $r_1 > \frac{1}{2}$ and $(\mathcal{L}(R), L^2(R), L)$ is a special compatible family.

The following example occurs in core conductor theory and will be later considered in connection with applications to neuronal behavior.

EXAMPLE 1.3 Let $H = L^2([0, b], dx)$ and consider the deterministic problem

$$\begin{aligned} \frac{\partial V}{\partial t} &= -\alpha V + \beta \Delta V \quad t > 0 \quad 0 < x < b, \\ &(\alpha, \beta > 0 \text{ constants}), \\ V(0, x) &= v_0(x), \\ \frac{\partial V}{\partial x}(t, 0) &= \frac{\partial V}{\partial x}(t, b) = 0. \end{aligned} \tag{I}$$

Let (T_t) be the continuous contraction semigroup on H defined by (I), i.e.

$$(T_t f)(x) = \int_0^b G(x, y; t) f(y) dy \quad f \in H$$

where

$$\begin{aligned} G(x, y; t) &= \sum_{n=0}^{\infty} e^{-\lambda_n t} \phi_n(x) \phi_n(y) \\ \lambda_n &= \alpha + \beta \left(\frac{n\pi}{b} \right)^2 \quad n \geq 1, \end{aligned} \tag{1.8}$$

and

$$\phi_n(x) = \left(\frac{2}{b} \right)^{1/2} \cos \left(\frac{n\pi x}{b} \right) \quad n \geq 1. \tag{1.9}$$

Let $A = -L$ be the generator of T_t . Then (I) has the solution

$$V(t,x) = \int_0^b G(x,y;t) v_0(y) dy, \quad t > 0 \quad (1.10)$$

for $v_0 \in \text{Dom}(L)$.

Now consider the stochastic model

$$\frac{\partial V}{\partial t} = -\alpha V + \beta \Delta V + \dot{W}_{t,x} \quad (II)$$

where $\dot{W}_{t,x}$ is a Gaussian white noise in space time. We shall show how to set up model II as a Φ' -valued stochastic differential equation for a suitable definition of Φ .

For λ_n and ϕ_n as in (1.8) and (1.9) respectively and

$$L = -\alpha I + \beta \Delta \quad (\Delta = \frac{d^2}{dx^2})$$

define Φ by the method explained in Example 1.2 i.e.

$$\Phi = \{ \phi \in H : \sum_{j=1}^{\infty} (1 + \lambda_j)^r \langle \phi, \phi_j \rangle_0^2 < \infty \quad \forall r \in \mathbb{R} \}$$

where

$$\langle \phi, \psi \rangle_0 = \int_0^b \phi(x) \psi(x) dx.$$

If $p > m + \frac{1}{4}$, since $\{(1 + \alpha + \beta(\frac{n\pi}{b})^2)^{-p} \phi_n\}$ is a CONS for Φ_p ,

$$\sum_{n=1}^{\infty} \|(1 + \alpha + \beta(\frac{n\pi}{b})^2)^{-p} \phi_n\|_m^2 = \sum_{n=1}^{\infty} (1 + \alpha + \beta(\frac{n\pi}{b})^2)^{-2(p-m)} < \infty$$

and then the injection map $\Phi_p \hookrightarrow \Phi_m$ is Hilbert Schmidt for $p > m + \frac{1}{4}$.

Hence Φ is a CHNS and is the linear space of all infinite differentiable real functions f such that $f^{(k)}(0) = f^{(k)}(b) = 0$ for all $k = 0, 1, 2, \dots$.

Furthermore

a) $T_t \phi \subseteq \phi$

b) $L\phi \subseteq \phi$

c) $L|_{\phi} = (-\alpha I + \beta \Delta)$

and the operator $(\lambda I + L)^{-r_1}$ is Hilbert-Schmidt if $r_1 > \frac{1}{4}$:

$$\sum_{j=1}^{\infty} \|(\lambda I + L)^{-r_1} \phi_j\|_0^2 = \sum_{j=1}^{\infty} (\lambda + \alpha + \beta(\frac{j\pi}{b}))^{-2r_1} < \infty$$

Now we indicate how to set up model II as a ϕ' -valued stochastic differential equation. We proceed heuristically. From (1.10) we have that (II) can be written as

$$\frac{d}{dt} T_t v_0 = -T_t L v_0 + \dot{W}_{t,x}.$$

Define for $t > 0$

$$\xi_t[\phi] = \int_0^b V(t,x) \phi(x) dx \quad \phi \in \phi,$$

then for $\phi \in \phi$

$$\begin{aligned} \frac{d}{dt} \xi_t[\phi] &= \int_0^b \frac{\partial V}{\partial t} \phi(x) dx \\ &= -\int_0^b LV(t,x) \phi(x) dx + \int_0^b \phi(x) \dot{W}_{t,x} dx \end{aligned}$$

i.e.

$$d\xi_t[\phi] = -\xi_t[L\phi]dt + dW_t[\phi] \quad (1.11)$$

where

$$W_t[\phi] := \int_0^b \phi(x) \dot{W}_{t,x} dx, \quad (1.12)$$

$$\xi_0[\phi] = \int_0^b v_0(x) \phi(x) dx$$

and

$$EW_t[\phi]W_s[\psi] = \min(t,s)\langle\phi,\psi\rangle_H \quad \text{for } \phi,\psi \in \Phi. \quad (1.13)$$

If on Φ' we define T'_t by $T'_t f[\phi] = f[T_t \phi]$ for $f \in \Phi'$ then $L'f[\phi] = f[L\phi]$ and one may write model II as

$$d\xi_t = -L'\xi_t dt + dW_t$$

and

(II)'

ξ_0 given as above.

where W_t is a Φ' -valued Wiener process with covariance $\langle\cdot,\cdot\rangle_H$ as defined in the next section, and a special case of a Φ' -martingale.

Model (II)' is an example of a Φ' -valued linear stochastic differential equation for which we want to solve $\xi_t \in \Phi'$ and

$$\xi \in C(R_+; \Phi')$$

where $C(R_+; \Phi')$ is the space of all Φ' -valued processes on R with continuous paths in the strong topology of Φ' .

LECTURE II

MARTINGALES TAKING VALUES IN DUALS OF NUCLEAR SPACES

Throughout this lecture we assume that $(\Omega, \mathcal{F}, P)$ is a complete probability space with a right continuous filtration $(\mathcal{F}_t)_{t \geq 0}$. Also, unless otherwise explicitly stated, we assume that Φ is a given Countably Hilbertian nuclear space defined by a chain of Hilbert spaces $\Phi_p \subseteq \Phi_q$ $q \leq p$ with norms $\|\cdot\|_p \geq \|\cdot\|_q$ and strong duals $\Phi'_q \subseteq \Phi'_p$. We denote by Φ' the strong dual of Φ and by $\mathcal{B} = \mathcal{B}(\Phi')$ the σ -field of Φ' generated by the sets

$$\{f \in \Phi' : f[\phi] < a\} \quad \phi \in \Phi \quad a \in \mathbb{R}.$$

Definition 2.1 A mapping $X_t(\omega): \mathbb{R}_+ \times \Omega \rightarrow \Phi'$ is a Φ' -process if $X_t(\omega)[\phi]$ is a real valued process $\forall \phi \in \Phi$, i.e.

$$\{\omega: X_t(\omega) \in B\} \in \mathcal{F} \quad \forall B \in \mathcal{B}.$$

In this lecture we introduce two special cases of Φ' -valued processes, namely the Φ' -Wiener process and Φ' -martingales. We give several examples of Wiener processes and illustrate how some infinite dimensional extensions of the real valued Brownian motion (as the cylindrical Brownian motion and a sequence of independent Brownian motions) may be seen as nuclear space valued Wiener processes.

In the case of Φ' -martingales we will make the assumption that $EX_t[\phi]^2 < \infty \quad \forall \phi \in \Phi, t > 0$. This condition is not necessary and will be assumed only for simplicity in order to show the kind of techniques used in the study of Φ' -processes. Examples of these techniques are the following two lemmas. They will be referred in the future as the regularization Theorem and the Baire category argument.

Lemma 2.1 (Regularization) (Itô) Let $Y: \Phi \rightarrow L^2(\Omega, \mathcal{F}, P)$ be a continuous linear map. Then there exists a Φ' -valued random variable $\tilde{Y}$ s.t.

$$\tilde{Y}(\omega)[\phi] = Y(\phi)(\omega) \quad P \text{ a.s. } \forall \phi \in \Phi.$$

Moreover there is a $q > 0$ s.t. $P(\tilde{Y} \in \Phi'_q) = 1$.

Proof. Let $V(\phi) = E(Y(\phi)^2) \quad \forall \phi \in \Phi$. Since V is continuous there exists $r > 0$ and $\delta > 0$ s.t. if $\|\phi\|_r \leq \delta$ then $V(\phi) < 1$. Hence if $\theta = 1/\delta$ we have

$$V(\phi) \leq \theta^2 \|\phi\|_r^2 \quad \forall \phi \in \Phi. \quad (2.1)$$

Let $q > r$ be such that the injection mapping $\Phi_q \hookrightarrow \Phi_r$ is Hilbert-Schmidt and let $\{\phi_j\}_{j \geq 1} \subset \Phi$ be a CONS for Φ_q . Then from (2.1)

$$E\left(\sum_{j=1}^{\infty} Y(\phi_j)^2\right) \leq \theta^2 \sum_{j=1}^{\infty} \|\phi_j\|_r^2 < \infty$$

i.e. if $\Omega_1 = \{\omega: \sum_{j=1}^{\infty} (Y(\phi_j)(\omega))^2 < \infty\}$ then $P(\Omega_1) = 1$. Define

$$Y(\omega) = \begin{cases} \sum_{j=1}^{\infty} Y(\phi_j)(\omega) \phi_j & \text{if } \omega \in \Omega_1 \\ 0 & \text{otherwise} \end{cases}$$

where $\{\phi_j\}_{j \geq 1}$ is the CONS of Φ'_q dual to $\{\phi_j\}_{j \geq 1}$. Then $\tilde{Y}$ is a Φ' -valued random variable s.t. $Y \in \Phi'_q$ a.s. and

$$\tilde{Y}(\omega)[\phi] = \sum_{j=1}^{\infty} Y(\phi_j)(\omega) \langle \phi, \phi_j \rangle_q \quad \phi \in \Phi \text{ a.s.} \quad (2.2)$$

Letting $\psi_n := \sum_{j=1}^n \langle \phi, \phi_j \rangle_q \phi_j$, $\|\psi_n - \phi\|_r \leq \|\psi_n - \phi\|_q \xrightarrow{n \rightarrow \infty} 0$ so that from (2.1)

$$E \left[\sum_{j=1}^n Y(\phi_j) \langle \phi, \phi_j \rangle_q - Y(\phi) \right]^2 \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (2.3)$$

Finally, from (2.2) and (2.3) we have

$$E[\tilde{Y}(\omega)[\phi] - Y(\phi)(\omega)]^2 = 0$$

i.e.

$$\tilde{Y}[\phi] = Y(\phi) \quad \text{a.s.} \quad \forall \phi \in \Phi$$

Q.E.D.

From the proof of the above lemma we obtain the following result.

Corollary 2.1. Let $Y: \Phi \rightarrow L^2(\Omega, \mathcal{F}, P)$ be a continuous linear map. Then there exist $\theta > 0$ and $r > 0$ s.t.

$$E(Y(\phi))^2 \leq \theta^2 \|\phi\|_r^2 \quad \forall \phi \in \Phi.$$

Lemma 2.2 Let $V(\cdot): \Phi \rightarrow [0, \infty)$ satisfy the following conditions:

- (1) V is lower semicontinuous, i.e. $\phi_n \rightarrow \phi \Rightarrow V(\phi) \leq \liminf V(\phi_n)$
- (2) $V(\phi + \psi) \leq V(\phi) + V(\psi) \quad \forall \phi, \psi \in \Phi$
- (3) $V(a\phi) = |a|V(\phi) \quad \forall a \in \mathbb{R}, \phi \in \Phi$
- (4) $V(\phi) < \infty \quad \forall \phi \in \Phi$

Then $V(\phi)$ is a continuous function in Φ and there exist $\theta > 0$ and $r > 0$ s.t.

$$V(\phi) \leq \theta \|\phi\|_r \quad \forall \phi \in \Phi$$

Proof.

Let $D_n = \{\phi \in \Phi: V(\phi) \leq n\}$

Since V is a lower semicontinuous function on Φ , then for each $n \geq 1$ D_n is

a closed set of Φ . Condition 4) implies

$$\Phi = \bigcup_{n=1}^{\infty} D_n \quad (2.4)$$

Then by the Baire category theorem, since Φ is a complete metric space, it is never the union of a countable number of nowhere dense sets. Therefore there exist $\phi_0 \in \Phi$, $\delta_1 > 0$ and a positive integer r such that

$$U_{\phi_0} = \{\phi \in \Phi: \|\phi - \phi_0\|_r < \delta_1\} \subseteq D_{n_0}.$$

Hence, for any $\phi \in \Phi$, $\phi \neq 0$ if $\delta < \delta_1$

$$\delta \frac{\phi}{\|\phi\|_r} + \phi_0 \in U_{\phi_0}$$

and

$$\phi_0 - \delta \frac{\phi}{\|\phi\|_r} \in U_{\phi_0}$$

and therefore they belong to D_{n_0} , i.e.

$$V(\delta \frac{\phi}{\|\phi\|_r} + \phi_0) \leq n_0 \quad (2.5)$$

and

$$V(\phi_0 - \delta \frac{\phi}{\|\phi\|_r}) \leq n_0 \quad (2.6)$$

But using 3) with $a = -1$

$$V(\delta \frac{\phi}{\|\phi\|_r} - \phi_0) = V(\phi_0 - \frac{\delta \phi}{\|\phi\|_r}).$$

Then by 2)

$$V(2\delta \frac{\phi}{\|\phi\|_r}) \leq V(\delta \frac{\phi}{\|\phi\|_r} + \phi_0) + V(\delta \frac{\phi}{\|\phi\|_r} - \phi_0) \leq 2n_0$$

and hence using 3), if $\theta = n_0/\delta$

$$V(\phi) \leq \theta \|\phi\|_r \quad \forall \phi \in \Phi$$

Then the continuity of V follows since using 2)

$$|V(\phi) - V(\psi)| \leq V(\phi - \psi) \leq \theta \|\phi - \psi\|_r \quad \phi, \psi \in \Phi.$$

Q.E.D.

As an example of a typical application of the above lemma (Baire category argument) we have the following result.

Corollary 2.2. Let $(X_t)_{t \geq 0}$ be a Φ' -valued stochastic process s.t.

$$EX_t^2[\phi] < \infty \quad \forall \phi \in \Phi, \quad t \geq 0.$$

Then for each $t \geq 0$ there exist $\theta_t > 0$ and $r_t > 0$ s.t.

$$E(X_t)[\phi])^2 \leq \theta_t^2 \|\phi\|_{r_t}^2 \quad \forall \phi \in \Phi.$$

Proof. For each $t \geq 0$ define

$$V_t(\phi) = \{E(X_t[\phi])^2\}^{1/2} \quad \phi \in \Phi$$

Applying Fatou's lemma we have that $V_t(\phi)$ is a lower semicontinuous function on Φ . Then since V_t satisfies properties (2)-(4) in Lemma 2.2 the corollary follows applying the named lemma.

Q.E.D.

A result of this type, involving a Baire category argument, was first used in the study of Φ' -valued stochastic processes in Mitoma [10].

WIENER PROCESS TAKING VALUES IN THE DUAL OF A NUCLEAR SPACE

Definition 2.2 A strongly sample continuous Φ' -valued stochastic process

$W = (W_t)_{t \geq 0}$ on $(\Omega, \mathcal{F}, P)$ is called a centered Φ' -Wiener process with

covariance $Q(\cdot, \cdot)$ if W_t satisfies the following three conditions:

a) $W_0 = 0$ a.s.

b) W_t has independent increments, i.e. the random variables

$$W_{t_1}[\phi_1], (W_{t_2} - W_{t_1})[\phi_2], \dots, (W_{t_n} - W_{t_{n-1}})[\phi_n]$$

are independent $\forall \phi_1, \dots, \phi_n \in \Phi, 0 \leq t_1 \leq t_2 \leq \dots \leq t_n, n \geq 1$.

c) For each $t \geq 0$ and $\phi \in \Phi$

$$E(e^{iW_t[\phi]}) = e^{-1/2(tQ(\phi, \phi))} \quad (2.7)$$

where Q is a covariance functional, i.e. a positive definite symmetric bilinear form continuous on $\Phi \times \Phi$.

Remarks

2.1. Let $(W_t)_{t \geq 0}$ be a Φ' -Wiener process with covariance Q . Then

$$\{W_t[\phi], \phi \in \Phi, t \geq 0\}$$

is a centered Gaussian system and

$$E(W_t[\phi]W_s[\psi]) = \min(s, t)Q(\phi, \psi) \quad \phi, \psi \in \Phi, s, t \geq 0. \quad (2.8)$$

2.2 A Φ' -valued process $(Z_t)_{t \geq 0}$ is a Non-centered Wiener process if there exists $m \in \Phi'$ s.t. $Z_t - tm$ is a centered Wiener process.

We now prove the existence of a Φ' -valued Wiener process following Perez-Abreau [13].

Theorem 2.1 Let $\{Y(t, \phi), \phi \in \Phi, t \geq 0\}$ be a centered Gaussian system of random variables s.t.

$$E(Y(t, \phi)Y(s, \psi)) = \min(t, s)Q(\phi, \psi) \quad (2.9)$$

where Q is a covariance functional on $\Phi \times \Phi$. Then the following is true:

- 1). There exists a centered Φ' -valued Wiener process $W = (W_t)_{t \geq 0}$ with covariance functional Q s.t.

$$Y(t, \phi) = W_t[\phi] \quad \text{a.s.} \quad \forall \phi \in \Phi, \quad t \geq 0$$

- 2). There exists $q > 0$ depending only on Q and not on t s.t.

$$W. \in C(R_+, \Phi'_q) \quad \text{a.s.}$$

where $C(R_+, \Phi'_q)$ is the space of strongly continuous functions from R_+ to Φ'_q .

Proof. Letting $V^2(\phi) = Q(\phi, \phi)$ $\phi \in \Phi$, V is a function that satisfies conditions (1)-(4) of Lemma 2.2. Then there exist $\theta > 0$ and $r > 0$ s.t.

$$Q(\phi, \phi) \leq \theta \|\phi\|_r^2 \quad \forall \phi \in \Phi. \quad (2.10)$$

By the nuclearity of Φ there exists $q > r$ s.t. the injection $\Phi_q \hookrightarrow \Phi_r$ is a Hilbert-Schmidt map i.e. if $\{\phi_j\}_{j \geq 1} \subseteq \Phi$ is a CONS for Φ_q then

$$\sum_{j=1}^{\infty} \|\phi_j\|_r^2 < \infty. \quad (2.11)$$

Hence by (2.9), (2.10), (2.11) and the monotone convergence theorem we have

$$E\left(\sum_{j=1}^{\infty} Y(t, \phi_j)^2\right) = t \sum_{j=1}^{\infty} Q(\phi_j, \phi_j) \leq t \theta \sum_{j=1}^{\infty} \|\phi_j\|^2 < \infty.$$

Thus if $\Omega_t = \{\omega \in \Omega: \sum_{j=1}^{\infty} Y(t, \phi_j)^2 < \infty\}$ then $P(\Omega_t) = 1$.

Let $\{\hat{\phi}_j\}_{j \geq 1}$ be the CONS of Φ'_q dual to $\{\phi_j\}_{j \geq 1}$ and define

$$\begin{aligned} \sum_{j=1}^{\infty} Y(t, \phi_j)(\omega) \phi_j & \quad \omega \in \Omega_t \\ \tilde{W}_t(\omega) &= \\ 0 & \quad \text{otherwise.} \end{aligned}$$

Then $\tilde{W}_t \in \Phi'_q$ a.s. for all $t \geq 0$,

$$\tilde{W}_t(\omega)[\phi_j] = Y(t, \phi_j)(\omega) \quad \text{for } \omega \in \Omega_t$$

and

$$\tilde{W}_t(\omega)[\phi] = \sum_{j=1}^{\infty} \tilde{W}_t(\omega)[\phi_j] \phi_j[\phi] \quad (2.12)$$

For $\phi \in \Phi_q$, $\phi = \sum_{j=1}^{\infty} \langle \phi, \phi_j \rangle_q \phi_j = \sum_{j=1}^{\infty} \phi_j[\phi] \phi_j$ it follows from (2.10) and

$$\|\phi\|_r \leq \|\phi\|_q \quad r \leq q$$

that

$$\begin{aligned} E(Y(t, \phi) - \sum_{j=1}^n \phi_j[\phi] Y(t, \phi_j))^2 &= tQ(\phi - \sum_{j=1}^n \phi_j[\phi] \phi_j; \phi - \sum_{j=1}^n \phi_j[\phi] \phi_j) \\ &\leq t\theta \|\phi - \sum_{j=1}^n \phi_j[\phi] \phi_j\|_q^2 \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Hence, from this and (2.12) we have

$$\tilde{W}_t[\phi] = Y(t, \phi) \quad \text{a.s. } \forall \phi \in \Phi, \quad t \geq 0.$$

It remains to show that $\tilde{W}$ has a strongly continuous version.

We recall that if G is a real valued Gaussian random variable with zero mean and variance σ^2 , then

$$E(G^4) = 3\sigma^4$$

Then writing $X_t^j = Y(t, \phi_j)$

$$E(\chi_t^j - \chi_s^j)^4 = 3|t - s|^2 \{Q(\phi_j, \phi_j)\}^2. \quad (2.13)$$

Hence, using Holder's inequality, (2.13), (2.10) and (2.11) we have

$$\begin{aligned} E \|\tilde{W}_t - \tilde{W}_s\|_{-q}^4 &= E \left(\sum_{j=1}^{\infty} (\chi_t^j - \chi_s^j)^2 \right)^2 \\ &= E \left(\sum_{j=1}^{\infty} (\chi_t^j - \chi_s^j)^4 + 2 \sum_{j < k} (\chi_t^j - \chi_s^j)^2 (\chi_t^k - \chi_s^k)^2 \right) \\ &\leq \sum_{j=1}^{\infty} E(\chi_t^j - \chi_s^j)^4 + 2 \sum_{j < k} (E(\chi_t^j - \chi_s^j)^4 E(\chi_t^k - \chi_s^k)^4)^{1/2} \\ &= 3|t - s|^2 \left\{ \sum_{j=1}^{\infty} Q(\phi_j, \phi_j)^2 + 2 \sum_{j < k} Q(\phi_j, \phi_j) Q(\phi_k, \phi_k) \right\} \\ &= |t - s|^2 \left\{ \sum_{j=1}^{\infty} Q(\phi_j, \phi_j) \right\}^2 \equiv K^2 |t - s|^2 \end{aligned}$$

i.e.

$$E \|\tilde{W}_t - \tilde{W}_s\|_{-q}^4 \leq K^2 |t - s|^2, \quad K \text{ constant } s, t \in \mathbb{R}_+. \quad (2.14)$$

Then the proof of the theorem is completed using the following variant, given in Itô [6], of Kolmogorov's theorem on the existence of continuous versions for stochastic processes.

Theorem 2.2 (Itô [6]). Let X be a Φ'_q -valued process. If there exist positive constants α, β, K s.t.

$$E(\|X_t - X_s\|_{-q}^\alpha) \leq K |t - s|^{1+\beta}$$

then X has a continuous Φ'_q -version.

In the case of a special compatible family the index q in (2) of Theorem 2.1 can be chosen in the following manner.

Corollary 2.3 Let (Φ, H, T_t) be a special compatible family and assume the hypotheses of Theorem 2.1. Then there exists a Φ' -valued Wiener process $W = (W_t)_{t \geq 0}$ with covariance functional Q such that

$$W. \in C(R_+; \Phi'_q) \text{ a.s.}$$

for any $q \geq r_1 + r_2$ where r_1 is given by (1.3) in Example (1.2) and r_2 is such that

$$Q(\phi, \phi) \leq \theta \|\phi\|_{r_2}^2 \quad \forall \phi \in \Phi$$

for some $\theta > 0$.

Proof

Form the proof of Theorem 2.1, writing r_2 instead of r in (2.10), q was taken such that the injection $\Phi_q \hookrightarrow \Phi_{r_2}$ is a Hilbert-Schmidt map. But in the case of a special compatible family (see Example 1.2) the injection $\Phi_q \hookrightarrow \Phi_{r_2}$ is Hilbert-Schmidt for $q \geq r_1 + r_2$.

Q.E.D.

Some examples of Φ' -valued Wiener processes are now introduced.

EXAMPLE 2.1. Let (Φ, H, L) be a special compatible family (see Example 1.2 and Remark 1.1). Recall that (see (1.7)) there exists $r_1 > 0$ s.t.

$$\theta_1 := \sum_{j=1}^{\infty} (1 + \lambda_j)^{-2r_1} < \infty$$

and the injection $\Phi_q \hookrightarrow \Phi_r$ is a Hilbert-Schmidt map for $q \geq r + r_1$. Let $\langle \cdot, \cdot \rangle_0$ be the inner product in H and define

$$Q_0(\phi, \psi) = \langle \phi, \psi \rangle_0 \quad \phi, \psi \in \Phi.$$

Then from Theorem 2.1 there exists a Φ' -valued Wiener process $(W_t)_{t \geq 0}$ with

covariance functional Q_0 s.t.

$$W \in C(R_+; \Phi'_q) \text{ a.s. if } q > r_1$$

and will be called a Standard Wiener process.

More generally, if $r > 0$ and

$$Q_r(\phi, \psi) := \langle \phi, \psi \rangle_r \quad \phi, \psi \in \Phi$$

then there exists a Φ' -valued process $W \in C(R_+; \Phi'_q)$ for $q > r + r_1$.

As will be shown in later examples, in applications the Q is not always given by one of the inner products on the Hilbert spaces defining Φ .

Nevertheless since Q is continuous on $\Phi \times \Phi$, then, as in the proof of Theorem 2.1, there exist $\theta > 0$ and $r > 0$ s.t.

$$Q(\phi, \phi) \leq \theta \|\phi\|_r^2 \quad \forall \phi \in \Phi$$

and therefore there exists a Φ' -valued Wiener process W s.t.

$$W \in C(R_+; \Phi'_q) \text{ for } q > r + r_1.$$

EXAMPLE 2.2. Let $\mathcal{S}(R)$ be the Schwarz space of Example 1.1 (see also Remark 1.2). Then $(\mathcal{S}, L^2(R), -d^2/dx^2 + x^2/4)$ is a special compatible family where $(\phi_j)_{j \geq 1}$ are the Hermite functionals given by (1.1), $\lambda_j = j - 1/2$, $j \geq 1$, $\langle \cdot, \cdot \rangle_0$ is the inner product on $L^2(R)$ and $r_1 > 1/2$. Taking $\Phi = \mathcal{S}(R)$ and $H = L^2(R)$ in the last example, from (1.2) we have that if $Q_0(\phi, \psi) = \langle \phi, \psi \rangle_0$ then the standard Wiener process $W = (W_t)_{t \geq 0}$ in $\mathcal{S}(R)'$ is such that $W \in C(R_+; \mathcal{S}'_q)$ for $q > 1/2$. Clearly, there is no smallest q such that this happens.

For $\phi \in \Phi$ define

$$W_t^{(1)}[\phi] = W_t[D^2\phi] \text{ where } D = \frac{d}{dx}.$$

Then the covariance functional of the Φ' -valued Wiener process $W^{(1)} = (W_t^{(1)})_{t \geq 0}$

is

$$Q^{(1)}(\phi, \psi) = Q_0(D^2\phi, D^2\psi) = \langle D^2\phi, D^2\psi \rangle_0.$$

We shall show that $W_t^{(1)} \in C(R_+; \mathcal{G}_q')$ for $q > 3/2$. In general we will prove the following: Let $Q_p(\phi, \psi) = \langle \phi, \psi \rangle_p$ $\phi, \psi \in \Phi$ $p > 0$, and let $W = (W_t)_{t \geq 0}$ be the corresponding Φ' -valued Wiener process. Define

$$W_t^{(1)}[\phi] = W_t[D^2\phi] \quad (2.15)$$

then $W^{(1)}$ is a Φ' -valued Wiener process s.t. $W^{(1)} \in C(R_+; \mathcal{G}_q')$ for $q > p + 3/2$:

Clearly

$$Q^{(1)}(\phi, \psi) = \langle D^2\phi, D^2\psi \rangle_p \quad \phi, \psi \in \Phi \quad (2.16)$$

then from Example 1.1 for $\phi \in \Phi$

$$\begin{aligned} Q^{(1)}(\phi, \phi) &= \langle D^2\phi, D^2\phi \rangle_p = \sum_{n=1}^{\infty} (n + 1/2)^{2p} \langle D^2\phi, \phi_n \rangle_0^2 \\ &= \sum_{n=1}^{\infty} (n + 1/2)^{2p} \langle \phi, D^2\phi_n \rangle_0^2 \end{aligned} \quad (2.17)$$

But from (1.1) with the notation of Example (1.1)

$$\begin{aligned} \frac{d}{dx} \phi_{n+1} &= \sqrt{n!} \{ \sqrt{g(x)} H_{n-1}(x) + H_n(x) [-\frac{x}{2} e^{-x^2/4} (2\pi)^{-1/4}] \} \\ &= \sqrt{n!} \{ \sqrt{g(x)} H_{n-1}(x) - \frac{x}{2} \sqrt{g(x)} H_n(x) \} \\ &= \sqrt{n!} \sqrt{g(x)} H_{n-1}(x) + \frac{\sqrt{n!}}{2} \sqrt{g(x)} \{ -(n+1) H_{n+1}(x) - H_{n-1}(x) \} \\ &= \frac{\sqrt{n!}}{2} \frac{\phi_n}{\sqrt{(n-1)!}} - \frac{\sqrt{n!}}{2} (n+1) \frac{\phi_{n+2}}{\sqrt{(n+1)!}} = \frac{\sqrt{n}}{2} \phi_n - \frac{\sqrt{n+1}}{2} \phi_{n+2} \end{aligned}$$

i.e.

$$\frac{d}{dx} \phi_{n+1} = \frac{\sqrt{n}}{2} \phi_n - \frac{\sqrt{n+1}}{2} \phi_{n+2} .$$

Hence

$$\frac{d^2}{dx^2} \phi_{n+1} = \frac{\sqrt{n(n-1)}}{4} \phi_{n-1} - \left(\frac{2n+1}{4} \right) \phi_{n+1} + \frac{\sqrt{(n+1)(n+2)}}{4} \phi_{n+3}$$

and

$$\langle \phi, D^2 \phi_n \rangle = \frac{\sqrt{(n-1)(n-2)}}{4} \langle \phi, \phi_{n-2} \rangle_0 - \left(\frac{2n-1}{4} \right) \langle \phi, \phi_n \rangle_0 + \frac{\sqrt{n(n+1)}}{4} \langle \phi, \phi_{n+2} \rangle_0 \quad (2.19)$$

$$= a_n \langle \phi, \phi_{n-2} \rangle_0 + b_n \langle \phi, \phi_n \rangle_0 + c_n \langle \phi, \phi_{n+2} \rangle_0$$

where $a_n, b_n, c_n = O(n)$.

Then from this and (2.17)

$$\begin{aligned} Q^{(1)}(\phi, \phi) &\leq \alpha_1 \sum_n \left(n + \frac{1}{2}\right)^{2p+2} \langle \phi, \phi_{n-2} \rangle_0^2 + \alpha_2 \sum_n \left(n + \frac{1}{2}\right)^{2p+2} \langle \phi, \phi_n \rangle_0^2 \\ &\quad + \alpha_3 \sum_n \left(n + \frac{1}{2}\right)^{2p+2} \langle \phi, \phi_{n+2} \rangle_0^2 \end{aligned}$$

$$\leq \alpha \|\phi\|_{p+1}^2, \quad (\alpha, \alpha_1, \alpha_2, \alpha_3 \text{ constants}).$$

Since the injection $\mathcal{H}_q \hookrightarrow \mathcal{H}_{p+1}$ is Hilbert-Schmidt for $q > p + 1 + 1/2 = p + 3/2$, we have shown that the ϕ' -valued Wiener process given by (2.15) is such that

$$W^{(1)} \in C(R_+; \mathcal{H}'_q) \text{ a.s. for } q > p + \frac{3}{2}.$$

Cylindrical Brownian motion

Definition 2.3 Let H be any real separable Hilbert space with norm $\|\cdot\|_H$. A family $\{B_t(h): t \in R_+, h \in H\}$ of real valued random variables is called a

Cylindrical Brownian motion (c.B.m.) on H if the following conditions hold:

(1) For each $h \in H$ $h \neq 0$, $\|h\|_H^{-1} B_t(h)$ is a one dimensional standard Wiener process.

(2) For any $\alpha_1, \alpha_2 \in \mathbb{R}$ and $h_1, h_2 \in H$

$$B_t(\alpha_1 h_1 + \alpha_2 h_2) = \alpha_1 B_t(h_1) + \alpha_2 B_t(h_2) \quad \text{a.s.}$$

(3) For each $t \geq 0$ and $h \in H$ $B_t(h)$ is an $\mathcal{F}_t$ -martingale where

$$\mathcal{F}_t = \sigma(B_s(k) : s \leq t, k \in H) .$$

From (1) we have that $\forall t \geq 0$ and $h \in H$

$$E(B_t(h))^2 = t \|h\|_H^2$$

and therefore since by Sazonov's theorem $\exp(-\frac{t}{2} \|h\|^2)$ is not the characteristic functional of a countably additive probability measure on H , there does not exist a process $\bar{B}_t$ in H s.t.

$$B_t(h) = \langle \bar{B}_t, h \rangle_H .$$

From (1), (2) and (3) we have that

$$E(B_t(h) B_s(k)) = \min(s, t) \langle h, k \rangle_H . \quad (2.18)$$

If $\{e_n\}_{n \geq 1}$ is a CONS in H , $B_t^n = B_t(e_n)$ is a sequence of independent one dimensional Brownian motions, and if $h \in H$

$$h = \sum_{n=1}^{\infty} \langle h, e_n \rangle_H e_n$$

for

$$h^{(n)} = \sum_{i=1}^n \langle h, e_i \rangle_H e_i$$

$$B_t(h^{(n)}) = \sum_{j=1}^n \langle h, e_j \rangle_H B_t^j.$$

For $n > m$,

$$E[B_t(h^{(n)}) - B_t(h^{(m)})]^2 = t \sum_{j=m+1}^n \langle h, e_j \rangle_H^2 \rightarrow 0 \quad m \rightarrow \infty$$

and therefore

$$B_t(h) = \sum_{j=1}^{\infty} \langle h, e_j \rangle_H B_t^j \quad \text{a.s.} \quad \forall t > 0 \quad \text{and} \quad h \in H. \quad (2.19)$$

Conversely if $B_t^j (j \geq 1)$ is a sequence of independent Brownian motions then B_t defined by (2.19) is a c.B.m. on H . (Note that the RHS of (2.19) converges a.s. for each $t \geq 0$ and every $h \in H$).

The following result relates Φ' -valued Wiener processes and cylindrical Brownian motions.

Theorem 2.3 Let $W = (W_t)_{t \geq 0}$ be a Φ' -valued Wiener process with covariance functional Q . Then W defines a Rigged Hilbert space

$$\Phi \hookrightarrow H_Q \hookrightarrow \Phi'$$

and a cylindrical Brownian motion $\hat{W}$ on H_Q where H_Q is the Q -completion of Φ .

Proof. Since by assumption Q is a positive definite continuous bilinear symmetric form on $\Phi \times \Phi$, $Q(\cdot, \cdot)$ defines an inner product on $\Phi \times \Phi$. Let H_Q be the Q -completion of Φ . Since Φ is separable then H_Q is separable.

Let $\{\phi_j\}_{j \geq 1} \subseteq \Phi$ be a CONS for H_Q . Then $(W_t^j)_{j \geq 1}$, where $W_t^j = W_t[\phi_j]$, is a sequence of independent standard Wiener processes since

$$E(W_t^j W_s^k) = \min(s, t) Q(\phi_j, \phi_k) = \min(s, t) \delta_{jk}$$

For $h \in H_Q$ $h = \sum_{j=1}^{\infty} \langle h, \phi_j \rangle_Q \phi_j$,

$$\hat{W}_t[h] := \sum_{j=1}^{\infty} \langle h, \phi_j \rangle_Q W_t^j$$

defines a c.B.m. on H_Q as in (2.19).

Φ' -VALUED MARTINGALES

Most of the material in this section is taken from Mitoma [10].

Definition 2.4 A Φ' -valued process $M = (M_t)_{t \geq 0}$ is a Φ' -martingale w.r.t. $(\mathcal{F}_t)$ if for each $\phi \in \Phi$ $M_t[\phi]$ is a martingale w.r.t. $(\mathcal{F}_t)$.

Since it will help in our later work, we shall also assume the additional condition

$$E(M_t[\phi])^2 < \infty \quad \forall \phi \in \Phi \quad t > 0. \quad (2.20)$$

Theorem 2.4 Let M be a Φ' -valued martingale w.r.t. $(\mathcal{F}_t)$. Then there exists a Φ' -valued version $\tilde{M}$ of M s.t. the following conditions hold:

(1) For each $T > 0$ there exists $p = p_T > 0$ s.t.

$$\tilde{M}^T \in D([0, T]; \Phi'_p) \quad \text{a.s.}$$

where $D([0, T]; \Phi'_p)$ is the Skorohod space of right continuous functions with left hand limits (r.c.l.l.) from $[0, T]$ to Φ'_p .

(2) $\tilde{M}$ is r.c.l.l. in the strong Φ' -topology, i.e.

$$\tilde{M} \in D([0, \infty); \Phi') \quad \text{a.s.}$$

Proof. (1) Fix $T > 0$ and define $V_T^2(\phi) = E(M_T[\phi]^2)$. Then by Corollary 2.2 there exist $\theta = \theta_T > 0$ and $r = r_T > 0$ s.t.

$$V_T(\phi) \leq \theta \|\phi\|_r \quad \forall \phi \in \Phi \quad (2.21)$$

Let D be a countable dense subset of $[0, T]$. Then by (1.3) VI in [3], for $\phi \in \Phi$

$$E(\sup_{t \in D} M_t[\phi]^2) \leq 4 \sup_{0 \leq t \leq T} (E M_t[\phi]^2) = 4(E M_T[\phi]^2). \quad (2.22)$$

Let $q \geq r$ be such that the injection map $\phi_q \hookrightarrow \phi_r$ is Hilbert-Schmidt, i.e. if $\{\phi_j\}_{j \geq 1} \subseteq \Phi$ is a CONS for ϕ_q then

$$\sum_{j=1}^{\infty} \|\phi_j\|_r^2 < \infty.$$

Then from (2.21) and (2.22) we have

$$E\left(\sum_{j=1}^{\infty} \sup_{t \in D} (M_t[\phi_j])^2\right) = \sum_{j=1}^{\infty} E(\sup_{t \in D} (M_t[\phi_j])^2) \leq 4\theta^2 \sum_{j=1}^{\infty} \|\phi_j\|_r^2 < \infty.$$

So, if $\Omega_1 = \{\omega \in \Omega: \sum_{j=1}^{\infty} \sup_{t \in D} M_t[\phi_j]^2(\omega) < \infty\}$, $P(\Omega_1) = 1$.

Since each real valued martingale $M_t[\phi_j]$ has a right continuous modification X_t^j , writing

$$\Omega_t^j := \{\omega \in \Omega: X_t^j(\omega) = M_t[\phi_j](\omega)\}, \text{ we have}$$

$P(\Omega_t^j) = 1$ for $t \in D$. Then the set defined by

$$\Omega_2 := \left(\bigcap_{t \in D} \bigcap_{j \geq 1} \Omega_t^j\right) \cap \Omega_1$$

has probability one and if $\omega \in \Omega_2$

$$\sum_{j=1}^{\infty} \sup_{0 \leq t \leq T} (X_t^j(\omega))^2 < \infty.$$

Let $\{\phi_j\}_{j \geq 1}$ be the CONS of Φ'_q dual to $\{\phi_j\}_{j \geq 1}$ and for $0 \leq t \leq T$ define

$$\begin{aligned} \sum_{j=1}^{\infty} \chi_t^j(\omega) \phi_j & \quad \omega \in \Omega_2 \\ \tilde{M}_t(\omega) &= \\ 0 & \quad \text{otherwise} \end{aligned}$$

Then for $0 \leq t \leq T$ $P(\tilde{M}_t \in \Phi'_q) = 1$ and $\tilde{M}_t(\omega)[\phi] = M_t(\omega)[\phi]$ for all $\phi \in \Phi$ $\omega \in \Omega_2$, i.e. $\tilde{M}_t = M_t$ a.s.

Next since for $s, t \in [0, T]$ and $j \geq 1$

$$|\chi_t^j(\omega) - \chi_s^j(\omega)|^2 \leq 4 \sup_{0 \leq t \leq T} (\chi_t^j(\omega))^2, \quad (\omega \in \Omega_2),$$

by the dominated convergence theorem, fixing ω in Ω_2 ,

$$\begin{aligned} \lim_{s \uparrow t} \|\tilde{M}_t(\omega) - \tilde{M}_s(\omega)\|_{-q}^2 &= \lim_{s \uparrow t} \left\| \sum_{j=1}^{\infty} (\chi_t^j(\omega) - \chi_s^j(\omega)) \phi_j \right\|_{-q}^2 \\ &= \lim_{s \uparrow t} \sum_{j=1}^{\infty} |\chi_t^j(\omega) - \chi_s^j(\omega)|^2 = \sum_{j=1}^{\infty} \lim_{s \uparrow t} |\chi_t^j(\omega) - \chi_s^j(\omega)|^2 = 0, \end{aligned}$$

the last assertion following from the right continuity of $\chi_t^j(\omega)$. In a similar fashion the fact that $\tilde{M}_t$ has left hand limits in the $\|\cdot\|_{-q}$ -norm is shown.

Thus we have proved that for each $T > 0$ there exists $q_T > 0$ s.t. M_t has a r.c.l.l. version $\tilde{M}_t$ in the $\|\cdot\|_{q_T}$ -norm, i.e.

$$\tilde{M}^T \in D([0, T]; \Phi'_q).$$

(2) Let $T_n \uparrow \infty$, then by (1) there exists q_n s.t. M_t has a version $\tilde{M}^{T_n}$ with

$$\tilde{M}^{T_n} \in D([0, T_n]; \Phi'_{q_n}) \text{ a.s.}$$

With the notation used in the proof of (1) let $\Omega_3 = \bigcap_{n=1}^{\infty} \Omega_2^n$. If $\omega \in \Omega_3$ define for $0 \leq t < \infty$

$$\tilde{M}_t(\omega) = \tilde{M}_t^{T_n}(\omega) \quad \text{for } T_{n-1} < t \leq T_n, \quad (T_0 = 0).$$

Then $P(\tilde{M} \in \Phi') = 1$ and $\tilde{M}_t(\omega) = M_t(\omega)$ for $\omega \in \Omega_3$.

Hence for $t > 0$ and $\epsilon > 0$ there exists $\delta_t > 0$ s.t. if $t > s + \delta_t$

$$\|\tilde{M}_t(\omega) - \tilde{M}_s(\omega)\|_{-q_n} < \epsilon$$

for $t < T_n$ and therefore for any bounded set $B \subset \Phi$

$$\sup_{\phi \in B} |(\tilde{M}_t(\omega) - \tilde{M}_s(\omega))[\phi]| < \epsilon, \quad t > s + \delta_t$$

i.e. $\tilde{M}_t$ is strongly right continuous. A similar argument shows that it has left hand limits. Q.E.D.

Remarks

2.3) The above theorem can be proved without the assumption (2.20). The proof is very similar to the one given above using (1.1)VI of [3] instead of (2.22).

2.4) If M_t is a Φ' -valued martingale s.t. for each $\phi \in \Phi$

$$\sup_{0 \leq t < \infty} E(M_t[\phi]^2) < \infty, \tag{2.23}$$

there exists $q > 0$ s.t. M_t has a version $\tilde{M}_t \in D([0, \infty), \Phi'_q)$ a.s. This is seen using the fact that if D is a countably dense subset of R_+ then from (2.22)

$$E\left(\sup_{t \in D} (M_t[\phi]^2)\right) \leq 4E(M_\infty[\phi]^2) < \infty.$$

The next theorem is proved in a very similar way to Theorem 2.4.

Theorem 2.5 Let M be a Φ' -valued martingale such that for each $\phi \in \Phi$ $M_t[\phi]$ has a continuous version. Then there exists a Φ' -valued version $\tilde{M}$ of M such that the following conditions hold:

(1) For each $T > 0$ there exists $p = p_T > 0$ s.t.

$$\tilde{M}_\cdot^T \in C([0, T]; \Phi'_p) \text{ a.s.}$$

(2) $\tilde{M}$ is continuous in the strong Φ' -topology i.e.

$$\tilde{M}_\cdot \in C([0, \infty); \Phi') \text{ a.s.}$$

(3) If $\sup_{0 \leq t < \infty} E(M_t[\phi]^2) < \infty$ then there exists $p > 0$ s.t.

$$\tilde{M}_\cdot \in C([0, \infty); \Phi'_p) \text{ a.s.}$$

An example of a Φ' -martingale is the Φ' -valued Wiener process with

$$\mathcal{F}_t = \sigma(W_s[\phi]; 0 \leq s \leq t, \phi \in \Phi)$$

and for which (Theorem 2.1) there exists a continuous version on $C([0, \infty); \Phi'_q)$ for some $q > 0$. This shows that condition (2.23) in Theorem 2.5(3) is not necessary for a Φ' -valued martingale to have a version in $C([0, \infty); \Phi'_p)$ for some $p > 0$.

The following example (due to G. Kallianpur and S. Ramaswamy) gives a Φ' -valued strongly continuous gaussian martingale M_t for which the following is not true: There exists p independent of t s.t.

$$M_t \in \Phi'_p \quad \forall t > 0, \text{ a.s.}$$

EXAMPLE 2.3 Consider the CHNS of Example 1.2 i.e. (Φ, H, L) is a special compatible family where $(I + L)^{-r_1}$ is a Hilbert-Schmidt operator for some $r_1 > 0$

$$L\phi_j = \lambda_j \phi_j$$

$$\sum_{j=1}^{\infty} (1 + \lambda_j)^{-2r_1} < \infty$$

for $\{\phi_j\}_{j \geq 1} \subseteq \Phi$ a CONS for H .

Define for $s \in [0, \infty)$ and $\phi \in \Phi$

$$f(s, \phi) = \sum_{j=1}^{\infty} (1 + \lambda_j)^s \langle \phi_j, \phi \rangle_0$$

Let $(B_s)_{s \geq 0}$ be a real valued standard Brownian motion. Since for each $t > 0$ and $\phi \in \Phi$

$$\int_0^t f(s, \phi)^2 ds < \infty$$

then the Wiener integral

$$X_{t, \phi} = \int_0^t f(s, \phi) dB_s$$

is a Gaussain martingale for each $\phi \in \Phi$. Since $f(s, \phi)$ is linear and continuous in ϕ then the linear random functional

$$X_{t, \phi}: \Phi \rightarrow L_2(\Omega)$$

is Φ -continuous. Hence, by the regularization lemma there exists a Φ' -valued random variable X_t s.t.

$$X_t[\phi] = X_{t, \phi} \quad \text{a.s.} \quad \forall \phi \in \Phi$$

Then for $\mathcal{F}_t = \mathcal{F}_t^B$ $(X_t, \mathcal{F}_t)_{t \geq 0}$ is a Φ' -valued martingale such that $\forall \phi \in \Phi$ $X_t[\phi]$ has a continuous version. Hence by Theorem 2.5 X has a strongly con-

tinuous version also denoted by χ .

Now suppose there exists $p > 0$ s.t. $\chi_t \in \Phi_p'$ a.s. $\forall t \geq 0$. Take $p > r_1$ and consider

$$\phi^{(n)} = \sum_{j=1}^n (1 + \lambda_j)^{-p-r_1} \phi_j.$$

Then $\{\phi^{(n)}\}$ converges in Φ_p to ϕ say, and therefore $\chi_t[\phi^{(n)}] \xrightarrow{n \rightarrow \infty} \chi_t[\phi]$.

But since χ_t is L^2 -continuous

$$E(\chi_t[\phi^{(n)}]^2) \xrightarrow{n \rightarrow \infty} E(\chi_t[\phi])^2 < \infty \quad \forall t \geq 0 \quad (2.24)$$

the finiteness of the limit being a consequence of $\chi_t[\phi]$ being a Gaussian random variable. On the other hand, if $t > p + r_1$,

$$\begin{aligned} E(\chi_t[\phi^{(n)}]^2) &= E\left(\sum_{j=1}^n (1 + \lambda_j)^{-p-r_1} \chi_{t, \phi_j}\right) \\ &= \int_0^t (f(s, \sum_{j=1}^n (1 + \lambda_j)^{-p-r_1} \phi_j))^2 ds = \\ &= \int_0^t \left(\sum_{j=1}^n (1 + \lambda_j)^{-p-r_1+s}\right)^2 ds \\ &> \int_{p+r_1}^t \left(\sum_{j=1}^n (1 + \lambda_j)^{-p-r_1+s}\right)^2 ds \end{aligned}$$

Then by Fatou's lemma

$$\liminf E(\chi_t[\phi^{(n)}]^2) = \infty$$

which, in view of (2.24) implies that $E(\chi_t[\phi])^2 = \infty$, a contradiction.

STOCHASTIC INTEGRALS TAKING VALUES IN A DUAL OF A NUCLEAR SPACE

Consider a Φ' -valued right continuous martingale $(M_t)_{t \geq 0}$ w.r.t. a filtration $(\mathcal{F}_t)_{t \geq 0}$ such that

$$E(M_t[\phi])^2 < \infty \quad \forall \phi \in \Phi \quad \text{and} \quad t \geq 0.$$

Then for each $\phi \in \Phi$ there exists a predictable right continuous, non negative increasing process A_t^ϕ such that $X_t[\phi]^2 - A_t^\phi$ is a martingale, which implies that

$$E(X_t[\phi])^2 = EA_t^\phi \quad (2.25)$$

Let $\mathcal{L}^2(M)$ be the class of real valued predictable processes $f: \mathbb{R}_+ \times \Omega \rightarrow \mathbb{R}$ such that

$$E \int_0^t f_s^2(\omega) dA_s^\phi(\omega) < \infty \quad \forall \phi \in \Phi.$$

Before introducing the definition of the stochastic integral of a real valued process w.r.t. the Φ' -valued martingale M_t we prove the following lemma.

Lemma 2.3 For $f \in \mathcal{L}^2(M)$ and $t \geq 0$ define

$$V_t^f(\phi) = E \int_0^t f_s^2(\omega) dA_s^\phi(\omega) \quad (2.26)$$

Then V_t^f is a continuous function of ϕ .

Proof. We first assume that f is a bounded function, i.e. $|f_t(\omega)| < K \quad \forall t \geq 0 \quad \text{and} \quad \omega \in \Omega$. Then by (2.25) and Corollary 2.2

$$V_t^f(\phi) \leq K EA_t^\phi = KE(X_t[\phi])^2 \leq K \theta_t \|\phi\|_{r_t} \quad \forall \phi \in \Phi$$

which implies that $V_t^f(\phi)$ is ϕ -continuous.

Next if $f \in \mathcal{L}^2(M)$ define

$$f_K(t, \omega) = f(t, \omega) 1_{[-K, K]}(f(t, \omega)) \quad t \geq 0, \quad \omega \in \Omega.$$

Then $|f_K(t, \omega)| \leq K$ and $f_K^2(t, \omega) \uparrow f^2(t, \omega) \quad \forall \quad t \geq 0, \quad \omega \in \Omega.$

Denote by ν_ϕ the measure on the σ -field of predictable sets defined by the relation

$$\int g d\nu_\phi = E \int_0^t g_s dA_s^\phi$$

where g is a non-negative predictable function. We have, using the monotone convergence theorem that

$$E \int_0^t f^2 dA^\phi = \int_{[0, t] \times \Omega} f^2 d\nu_\phi = \lim_{k \rightarrow \infty} \int_{[0, t] \times \Omega} f_k^2 d\nu_\phi = \lim_{k \rightarrow \infty} E \int_0^t f_k^2 dA^\phi.$$

Hence $V_t^f(\phi)$ is the limit of the increasing sequence of continuous functions $V_t^{f_k}(\phi)$, and therefore $V_t^f(\phi)$ is lower semicontinuous.

Then the Baire category argument (Lemma 2.1) implies that $V_t^f(\phi)$ is ϕ -continuous. Q.E.D.

We now define a ϕ' -valued stochastic integral for $f \in \mathcal{L}^2(M)$.

Definition 2.5 Let $f \in \mathcal{L}^2(M)$. For $\phi \in \Phi$ define

$$Y_t(\phi) := \int_0^t f_s dM_s[\phi]$$

where the RHS is the real valued stochastic integral w.r.t. the martingale $M_t[\phi]$. Then $Y_t(\phi)$ is a real valued martingale with a right continuous version and by Lemma 2.3

$$E(Y_t(\phi))^2 = E \int_0^t f_s^2 dA_s^\phi$$

is a continuous function of ϕ . Hence the linear random functional $Y_t(\cdot): \Phi \rightarrow L^2(\Omega, \mathcal{F}, P)$ is continuous and by the regularization Lemma there exists $I_t(f) \in \Phi'$ a.s. such that

$$I_t(f)[\phi] = Y_t(\phi) \text{ a.s. } \forall \phi \in \Phi, t \geq 0.$$

By Theorem 2.4 there exists a version, also denoted by $I_t(f)$, which is r.c.l.l. in the strong Φ' -topology. The right continuous Φ' -valued martingale $I_t(f)$

$$I_t(f) := \int_0^t f_s dM_s$$

is defined to be the stochastic integral of f w.r.t. the Φ' -valued martingale M_t .

We now introduce a Φ' -valued stochastic integral of a Φ' -valued process w.r.t. a real valued martingale.

Definition 2.6 Let $m = (m_t)$ be a real valued right continuous martingale such that $E(m_t^2) < \infty \forall t \geq 0$. Let $A_s = \langle m \rangle_s$ the integrable increasing process of M_t . Let $(F_t)_{t \geq 0}$ be a predictable Φ' -valued process such that

$$E \int_0^t F_s[\phi]^2 dA_s < \infty \quad \forall \phi \in \Phi \text{ and } t \geq 0.$$

Write

$$Y_t(\phi) = \int_0^t F_s[\phi] dM_s$$

then

$$E(Y_t(\phi))^2 = E \int_0^t F_s[\phi]^2 dA_s$$

and by the Baire category argument, the regularization lemma and Theorem 2.4 there exists a Φ' -valued right continuous martingale $J_t(f)$ s.t.

$$J_t(f)[\phi] = Y_t(\phi) \quad \text{a.s.} \quad \forall \phi \in \Phi, \quad t \geq 0.$$

We define $J_t(F) := \int_0^t F_s dM_s$ as the Φ' -valued stochastic integral.

LECTURE III

ORNSTEIN-UHLENBECK STOCHASTIC DIFFERENTIAL EQUATIONS ON DUALS OF NUCLEAR SPACES

We now introduce a special class of linear stochastic differential equations with values in duals of nuclear spaces, namely Ornstein-Uhlenbeck type processes with a nuclear valued martingale as a driving term.

Let $(\Omega, \mathcal{F}, P)$ be a complete probability space with a right continuous filtration $(\mathcal{F}_t)_{t \geq 0}$. Let $\Phi \hookrightarrow H \hookrightarrow \Phi'$ be a rigged Hilbert space, $A: \Phi \rightarrow \Phi$ a continuous linear operator and $(T_t)_{t \geq 0}$ a strongly continuous semigroup such that (Φ, H, T_t, A) is a compatible family (see Lecture I). Let $(M_t)_{t \geq 0}$ be a Φ' -valued martingale (Definition 2.4) which is right continuous with left hand limits (r.c.l.l.). Consider the stochastic differential equation

$$\begin{aligned} d\xi_t &= A' \xi_t dt + dM_t \quad t > 0 \\ \xi_0 &= \eta \end{aligned} \tag{3.1}$$

where η is a $\mathcal{F}_0$ -measurable Φ' -valued random variable and $A': \Phi' \rightarrow \Phi'$ is defined by the relation

$$A'f[\phi] = f[A\phi] \quad \forall f \in \Phi', \phi \in \Phi.$$

A special case of the SDE (3.1) is

$$\begin{aligned} d\xi_t &= A' \xi_t dt + dW_t \quad t > 0 \\ \xi_0 &= \eta \end{aligned} \tag{3.2}$$

where W_t is a Φ' -valued centered Wiener process (Definition 2.2) with covariance Q . From Theorem 2.1 $W. \in C(R_+; \Phi'_q)$ for some $q > 0$ where q depends only on Q and not on t .

In this lecture we solve the SDE (3.2) (Theorem 3.1). The general martingale case (Theorem 3.2) was considered in Christensen [1].

Definition 3.1 We say that the SDE (3.2) has a Φ' -valued solution $\xi = (\xi_t)_{t \geq 0}$ if the following four conditions hold:

- i) (ξ_t) is $\mathcal{F}_t$ -adapted and Φ' -valued.
- ii) $\xi \in C(R_+; \Phi')$ a.s.
- iii) $\xi_t[\phi] = \gamma[\phi] + \int_0^t \xi_s[A_s \phi] ds + W_t[\phi] \quad \forall \phi \in \Phi \text{ a.s. } \forall t \geq 0.$
- iv) For each $T > 0$

$$E\left(\sup_{0 \leq t \leq T} |\xi_t[\phi]|^2\right) < \infty \quad \forall \phi \in \Phi$$

Remark 1. Condition (iv) above is implied by the following condition: for each $T > 0$

$$E \int_0^T (\xi_s[A_s \phi])^2 ds < \infty \quad \forall \phi \in \Phi$$

Proposition 3.1. If $(\xi_t)_{t \geq 0}$ is a solution of the SDE (3.2) then for each $T > 0$ there exist $q_T > 0$ and a version of ξ (denoted also by ξ) such that

$$\xi_{\cdot}^T \in C([0, T]; \Phi'_{q_T}) \quad \text{a.s.}$$

and

$$\xi_t[\phi] = \gamma[\phi] + \int_0^t \xi_s[A_s \phi] ds + W_t[\phi] \quad \forall \phi \in \Phi, \quad 0 \leq t \leq T \quad \text{a.s.}$$

Proof. Given $T > 0$ define

$$G_T^2(\phi) := E\left(\sup_{0 \leq t \leq T} |\xi_t[\phi]|^2\right).$$

Then by condition (iv) in Definition 3.1 $G_T(\phi) < \infty \quad \forall \phi \in \Phi$ and clearly

$G_T(\phi_1 + \phi_2) \leq G_T(\phi_1) + G_T(\phi_2)$, $G_T(a\phi_1) = |a|G_T(\phi_1)$, $\phi_1, \phi_2 \in \Phi$, $a \in \mathbb{R}$. Next $\sup_{0 \leq t \leq T} |\xi_t[\phi]|$ is a lower semicontinuous function of ϕ . Hence by Fatou's Lemma $G_T(\phi)$ is also a lower semicontinuous function of ϕ . Then by a Baire category argument there exist $\theta_T > 0$ and $r_T > 0$ s.t.

$$E\left(\sup_{0 \leq t \leq T} |\xi_t[\phi]|^2\right) \leq \theta_T \|\phi\|_{r_T}^2 \quad \forall \phi \in \Phi$$

Let $p_T > r_T$ such that the injection map $\Phi_{p_T} \hookrightarrow \Phi_{r_T}$ is Hilbert-Schmidt and let $\{\phi_j\}_{j \geq 1} \subseteq \Phi$ be a CONS for Φ_{p_T} with dual basis $\{\phi_j'\}_{j \geq 1}$ a CONS for Φ_{p_T}' . Then

$$E\left(\sum_{j=1}^{\infty} \sup_{0 \leq t \leq T} |\xi_t[\phi_j]|^2\right) \leq \theta_T \sum_{j=1}^{\infty} \|\phi_j\|_{r_T}^2 < \infty$$

Define

$$\Omega_T = \{\omega: \sum_{j=1}^{\infty} \sup_{0 \leq t \leq T} |\xi_t(\omega)[\phi_j]|^2 < \infty\}$$

Then $P(\Omega_T) = 1$. Next define

$$\begin{aligned} & \sum_{j=1}^{\infty} \xi_t(\omega)[\phi_j] \phi_j \quad \omega \in \Omega_T \\ \hat{\xi}_t(\omega) = & \\ & 0 \quad \omega \notin \Omega_T \end{aligned}$$

Hence, $\hat{\xi}_t \in \Phi_{p_T}'$ $0 \leq t \leq T$ a.s. and $\hat{\xi}_t(\omega)[\phi] = \xi_t(\omega)[\phi] \quad \forall \phi \in \Phi$ and $\omega \in \Omega_T$.

Moreover by the dominated convergence theorem if $t, t_0 \in [0, T]$

$$\lim_{t \rightarrow t_0} \|\hat{\xi}_t(\omega) - \hat{\xi}_{t_0}(\omega)\|_{-p_T}^2 = \lim_{t \rightarrow t_0} \sum_{j=1}^{\infty} (\xi_t(\omega)[\phi_j] - \xi_{t_0}(\omega)[\phi_j])^2 = 0$$

i.e. $\hat{\xi}_\bullet^T \in C([0, T]; \Phi_{p_T}')$ a.s. and therefore

$$P(\omega: M_T(\omega) := \sup_{0 \leq t \leq T} \|\hat{\xi}_t\|_{-p_T}^2 < \infty) = 1.$$

From now on we will write ξ_t instead of $\hat{\xi}_t$.

Next for $\omega \in \Omega_T$ and $0 \leq t \leq T$ define

$$Y_t(\omega)[\phi] = \int_0^t \xi_s(\omega)[A_s \phi] ds$$

We will show that $Y^T(\omega) \in C([0, T]; \Phi'_{\ell_T})$ for some $\ell_T > 0$:

$$|Y_t(\omega)[\phi]| \leq M_T(\omega) \int \|A_s \phi\|_{p_T} ds. \quad (3.3)$$

Then using the continuity of the map $s \rightarrow A_s \phi \forall \phi \in \Phi$, by a Baire category argument there exist $\theta'_T > 0$ and $n_T > p_T$ s.t.

$$\sup_{0 \leq t \leq T} |Y_t(\omega)[\phi]|^2 \leq M_T^2(\omega) \theta'^2 \|\phi\|_{n_T}^2 \quad \forall \phi \in \Phi.$$

Then $Y_t(\omega) \in \Phi'_{n_T} \forall 0 \leq t \leq T \omega \in \Omega_T$. Let $\ell_T > n_T$ be such that $\Phi_{\ell_T} \hookrightarrow \Phi_{n_T}$ is Hilbert-Schmidt and let $\{e_j\}_{j \geq 1} \subseteq \Phi$ be a CONS for Φ_{ℓ_T} with dual basis $\{\hat{e}_j\}_{j \geq 1}$ a CONS for Φ'_{ℓ_T} . Then

$$\sum_{j=1}^{\infty} \sup_{0 \leq t \leq T} |Y_t(\omega)[e_j]|^2 \leq M_T^2(\omega) \theta_T^2 \sum_{j=1}^{\infty} \|e_j\|_{n_T}^2 < \infty.$$

Hence since from (3.3) $Y_t(\omega)[\phi]$ is a continuous function of t on $0 \leq t \leq T$ for each $\phi \in \Phi$, by the dominated convergence theorem we have

$$\lim_{t \rightarrow t_0} \|Y_t(\omega) - Y_{t_0}(\omega)\|_{\ell_T}^2 = \lim_{t \rightarrow t_0} \sum_{j=1}^{\infty} (Y_t[e_j] - Y_{t_0}[e_j])^2 = 0 \quad t, t_0 \in [0, T]$$

i.e. $Y^T(\omega) \in C([0, T]; \Phi'_{\ell_T}) \quad \omega \in \Omega_T$.

Then we have shown that $\int_0^t A'_s \xi_s ds \in C([0, T]; \Phi'_{\ell_T})$ a.s. for some $\ell_T > 0$. Hence taking $q_T = \max(r_0, q, p_T, \ell_T)$ we have that

$$Z_t = \gamma + \int_0^t A'_s \xi_s ds + W_t \in C([0, T]; \Phi'_{q_T}) \quad \text{a.s.}$$

Then by conditions (ii) and (iii) in Definition 3.1, for each $T > 0$

$$P(Z_t = \xi_t \quad 0 \leq t \leq T) = 1 \quad \text{Q.E.D.}$$

Theorem 3.1 Let (Φ, H, T_t) be a compatible family and assume that there exists $r_0 > 0$ s.t.

$$E \|\eta\|_{-r_0}^2 < \infty.$$

Then (3.2) has a unique solution given by

$$\xi_t := T_t' \eta + \int_0^t A' T_{t-s} W_s ds + W_t \quad \text{a.s.}$$

i.e.

$$\xi_t[\phi] = \eta[T_t \phi] + \int_0^t W_s[T_{t-s} A \phi] ds + W_t[\phi] \quad \forall \phi \in \Phi \quad (3.4)$$

satisfying the following properties.

a) For each $T > 0$ there exists $p = p_T$ s.t.

$$\xi_{\cdot}^T \in C([0, T], \Phi_{p_T}') \quad \text{a.s.} \quad \text{and} \quad E \left(\sup_{0 \leq t \leq T} \|\xi_t\|_{-p_T}^2 \right) < \infty.$$

b) If $\{e_j\}_{j \geq 1} \subseteq \Phi$ is a CONS in Φ_p and $\xi_t^j := \xi_t[e_j]$ then

$$\sum_{j=1}^{\infty} \xi_t^j e_j \text{ converges uniformly in } [0, T] \text{ in } \Phi_p' \text{ a.s.}$$

c) If in addition (Φ, H, T) is a special compatible family (see Example 1.2 and Remark 1.1) from Corollary 2.3 we have that $q \geq r_1 + r_2$. Then the solution $\xi = (\xi_t)$ of the SDE (3.2) is such that $\xi \in C(R_+; \Phi_p')$ a.s. where $p \geq \max(r_1 + r_2, r_0)$ is independent of t .

We will give the proof of (a) and (b). The proof of (c) is given in Kallianpur and Wolpert [7].

The following Lemma will be useful in the proof of the theorem.

Lemma 3.1 (Christensen [1]) For each $F \in \Phi'$ and $s < t$

$$F[T_{t-s}\phi] - F[\phi] = \int_s^t F[T_{u-s}A\phi]du = \int_s^t F[T_{t-u}A\phi]du \quad \forall \phi \in \Phi. \quad (3.5)$$

Proof. Since we have a compatible family (Φ, H, T_t) $T_t|_{\Phi}$ is a strongly continuous semigroup on the Φ -topology and if

$$\Psi = \{\phi \in \Phi: \lim_{\varepsilon \downarrow 0} \frac{T_{\varepsilon}\phi - \phi}{\varepsilon} = A\phi \text{ in the } \Phi\text{-topology}\}$$

then Ψ is dense in Φ .

Then for all $\phi \in \Psi$ and $0 \leq s < t$,

$$\frac{d}{du} F[T_{u-s}\phi] = F[T_{u-s}A\phi] \quad \text{and}$$

$$F[T_{t-s}\phi] = F[\phi] + \int_s^t F[T_{u-s}A\phi]du$$

Next for some $q > 0$ $|F[T_{u-s}A\phi]| \leq \|F\|_{-q} \|T_{u-s}A\phi\|_q$. Define

$$G(\phi) = \sup_{s \leq u \leq t} \|T_{u-s}A\phi\|_q$$

Then by the continuity of T and A $G(\phi) < \infty \quad \forall \phi \in \Phi$ and it is a lower semicontinuous function of ϕ . Then by the Baire category argument there exist $\theta > 0$ and $r > 0$ s.t. $\forall u \in [s, t]$

$$\|T_{u-s}A\phi\|_q \leq \theta \|\phi\|_r \quad \forall \phi \in \Phi.$$

Hence, since F , T and A are continuous, by the dominated convergence theorem if $\phi_n \rightarrow \phi$ in Φ $\{\phi_n\} \subseteq \Psi$

$$\int_s^t F[T_{u-s} A \phi_n] du \xrightarrow{n \rightarrow \infty} \int_s^t F[T_{u-s} A \phi] du$$

The second equality in (3.5) follows in a similar way.

Q.E.D.

Proof of Theorem 3.1 Let Ω_1 be such that $P(\Omega_1) = 1$ and

$$\omega \in \Omega_1 \Rightarrow W(\omega) \in C(R_+; \Phi'_q).$$

Step 1. For $t > 0$ the map $\phi \rightarrow \int_0^t W_s [T_{t-s} A \phi] ds$ is continuous and linear on Φ :

Let $\phi \in \Phi$, then

$$\begin{aligned} \left| \int_0^t W_s [T_{t-s} A \phi] ds \right| &\leq \int_0^t \|W_s\|_{-q} \|T_{t-s} A \phi\|_q ds \\ &\leq \sup_{0 \leq s \leq t} \|W_s\|_{-q} \cdot R_t(\phi) \end{aligned} \quad (3.6)$$

where

$$R_t(\phi) = \int_0^t \|T_{t-s} A \phi\|_q ds.$$

Next $\phi_n \rightarrow \phi$ in Φ implies $\|T_{t-s} A \phi_n\|_q \rightarrow \|T_{t-s} A \phi\|_q$. Then by Fatou's lemma

$$R_t(\phi) \leq \underline{\lim} R_t(\phi_n)$$

i.e. $R_t(\phi)$ is a lower semicontinuous function. For $a \in R$ clearly we have

$$R_t(a\phi) = |a| R_t(\phi) \text{ and for } \phi_1, \phi_2 \in \Phi$$

$$R_t(\phi_1 + \phi_2) \leq R_t(\phi_1) + R_t(\phi_2).$$

Hence since $R_t(\phi) < \infty \forall \phi \in \Phi$, by the Baire category argument $R_t(\phi)$ is continuous in Φ and there exist $\theta_t > 0$ and $r_t > 0$ s.t.

$$R_t(\phi) \leq \theta_t \|\phi\|_{r_t} \quad \forall \phi \in \Phi.$$

Then from (3.6), for each fixed $t > 0$

$$\left| \int_0^t W_s [T_{t-s} A \phi] ds \right| \leq \theta_t \sup_{0 \leq s \leq t} \|W_s\|_{-q} \cdot \|\phi\|_{r_t} \quad \forall \phi \in \Phi$$

and thus the map $\phi \rightarrow \int_0^t W_s [T_{t-s} A \phi] ds$ is continuous and linear on Φ . Write

$$\xi_t^1 = \int_0^t W_s [T_{t-s} A \phi] ds.$$

For each $t \geq 0$ define

$$\xi_t(\omega) = \begin{cases} T_t' n(\omega) + \xi_t^1(\omega) + W_t(\omega) & \text{if } \omega \in \Omega_1 \\ 0 & \text{otherwise} \end{cases} \quad (3.7)$$

Step 2. We check that ξ_t is a solution of (3.2). Let $\omega \in \Omega_1$.

Using Lemma 3.1 with $F = W_s$ and $\phi \rightarrow A\phi$ we have

$$W_s(\omega) [T_{t-s} A \phi] = W_s(\omega) [A \phi] + \int_s^t W_s(\omega) [T_{u-s} A^2 \phi] du \quad (3.8)$$

Using again Lemma 3.2 with $F = n$ and $s = 0$ we obtain

$$n[T_t \phi] = n[\phi] + \int_0^t n[T_u A \phi] du \quad (3.9)$$

Substituting (3.8) and (3.9) in (3.4) we have

$$\begin{aligned} \xi_t(\omega) [\phi] &= n(\omega) [\phi] + \int_0^t n(\omega) [T_u A \phi] du + W_t(\omega) [\phi] \\ &\quad + \int_0^t W_s(\omega) [A \phi] + \int_0^t \int_s^t W_s(\omega) [T_{u-s} A^2 \phi] du ds \end{aligned}$$

Interchanging order of integration in the last term of the last expression and using (3.4) we have

$$\begin{aligned}
 \xi_t(\omega)[\phi] &= \eta(\omega)[\phi] + \int_0^t \eta(\omega)[T_u A \phi] du + W_t(\omega)[\phi] \\
 &\quad + \int_0^t W_u(\omega)[A \phi] du + \int_0^t \int_0^u W_s(\omega)[T_{u-s} A^2 \phi] ds du \\
 &= \eta(\omega)[\phi] + \int_0^t \eta(\omega)[T_u A \phi] du + W_t(\omega)[\phi] \\
 &\quad + \int_0^t \{W_u(\omega)[A \phi] + \int_0^u W_s(\omega)[T_{u-s} A^2 \phi] ds\} du
 \end{aligned}$$

i.e. $\forall \phi \in \Phi$ and $0 \leq t \leq T$

$$\xi_t(\omega)[\phi] = \eta(\omega)[\phi] + \int_0^t \xi_u(\omega)[A \phi] du + W_t(\omega)[\phi] \quad \omega \in \Omega_1.$$

Hence condition (iii) in Definition 3.1 is satisfied. Observe that

$(t, \omega) \rightarrow \xi_t(\omega)$ is $\mathcal{B}(\Phi')/\mathcal{B}(R_+) \times \mathcal{F}$ -measurable and for each $t > 0$ ξ_t is $\mathcal{F}_t^{W, \eta}$ -measurable where

$$\mathcal{F}_t^{W, \eta} = \sigma\{\eta[\phi], W_s[\phi]: s \leq t, \phi \in \Phi\} \vee \{P\text{-null sets}\}.$$

Next, from (3.4), the assumptions on W and η and Step 1 we have

$$\begin{aligned}
 E\left(\sup_{0 \leq t \leq T} (\xi_t[\phi])^2\right) &\leq 3E\left(\sup_{0 \leq t \leq T} (\eta[T_t \phi])^2\right) \\
 &\quad + 3E\left(\sup_{0 \leq t \leq T} \left(\int_0^t W_s[T_{t-s} A \phi] ds\right)^2\right) \\
 &\quad + 3E\left(\sup_{0 \leq t \leq T} (W_t[\phi])^2\right) \\
 &\leq c_T^2 \|\phi\|_{m_T}^2 < \infty \quad \forall \phi \in \Phi
 \end{aligned} \tag{3.10}$$

for some constants $c_T > 0$ and $m_T > 0$, which shows condition (iv) in Definition 3.1. In the next step we shall show that ξ_t satisfies (ii) in Definition 3.1.

Step 3 Let $p_T > m_T$ s.t. the map $\phi_{p_T} \hookrightarrow \phi_{m_T}$ is Hilbert-Schmidt and let $\{e_j\}_{j \geq 1} \subseteq \Phi$ be a CONS in ϕ_{p_T} with dual basis $\{\hat{e}_j\}_{j \geq 1}$ a CONS in ϕ'_{p_T} . Then using (3.10)

$$E\left(\sum_{j=1}^{\infty} \sup_{0 \leq t \leq T} (\xi_t[e_j])^2\right) \leq C_T \sum_{j=1}^{\infty} \|e_j\|_{m_T}^2 < \infty.$$

Let $\Omega_2 = \{\omega: \sum_{j=1}^{\infty} \sup_{0 \leq t \leq T} (\xi_t(\omega)[e_j])^2 < \infty\}$, then $P(\Omega_2) = 1$. Define

$$\tilde{\xi}_t(\omega) = \begin{cases} \sum_{j=1}^{\infty} \xi_t(\omega)[e_j] \hat{e}_j & \omega \in \Omega_2 \\ 0 & \omega \notin \Omega_2. \end{cases}$$

Then from (3.10)

$$E\left(\sup_{0 \leq t \leq T} \|\tilde{\xi}_t\|_{-p_T}^2\right) = E\left(\sup_{0 \leq t \leq T} \sum_{j=1}^{\infty} (\xi_t[e_j])^2\right) \leq C_T \sum_{j=1}^{\infty} \|e_j\|_{m_T}^2 < \infty$$

It remains to show that the series converges uniformly in $[0, T]$ in the ϕ'_{p_T} -norm and that $\tilde{\xi}_t(\omega) = \xi_t(\omega)$ a.s. in $0 \leq t \leq T$. Define

$$S_n(t, \omega) = \sum_{j=1}^n \xi_t(\omega)[e_j] \hat{e}_j$$

then

$$\sup_{0 \leq t \leq T} \|S_{n'}(t, \omega) - S_n(t, \omega)\|_{-p_T}^2 \rightarrow 0 \quad \text{as } n', n \rightarrow \infty$$

and $S_n(\cdot, \omega) \in C([0, T], \phi_{-p_T})$. But since $C([0, T], \phi_{-p_T})$ is a complete metric space there exists $\tau_t(\omega) \in C([0, T], \phi_{-p_T})$ s.t.

$$\sup_{0 \leq t \leq T} \|S_n(t, \omega) - \tau_t(\omega)\|_{-p_T} \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

and

$$\begin{aligned}\|\tau_t(\omega)\|_{-p_T}^2 &= \lim_n \|S_n(t, \omega)\|_{-p_T}^2 = \lim_{n \rightarrow \infty} \sum_{j=1}^n (\xi_t(\omega)[e_j])^2 \\ &= \|\tilde{\xi}_t(\omega)\|_{-p_T}^2.\end{aligned}$$

Hence

$$E\|\tilde{\xi}_t(\omega)\|_{-p_T}^2 = E\|\tau_t\|_{-p_T}^2.$$

Next

$$E|\tau_t[\phi] - S_n(t)[\phi]|^2 \leq E\|\tau_t - S_n(t)\|_{-p_T}^2 \|\phi\|_{p_T}^2 \rightarrow 0$$

then

$$E\tau_t[\phi] = \lim_n ES_n(t)[\phi] = \lim_{n \rightarrow \infty} \sum_{j=1}^n E\xi_t[e_j]\hat{e}_j[\phi]$$

$\therefore$

$$\tau_t[\phi] = \lim_{n \rightarrow \infty} S_n(t)[\phi] = \sum_{j=1}^{\infty} \xi_t[e_j]\hat{e}_j[\phi] = \tilde{\xi}_t[\phi] \text{ a.s.}$$

Then we have shown that for each $T > 0$ $\exists$ p_T s.t. the following set has probability one

$$\Omega_T = \{\omega: \tilde{\xi}_T(\omega) \in C([0, T]; \Phi'_{p_T})\}.$$

Then taking $T_n \uparrow \infty$ and $\overline{\Omega} = \bigcap_{n=1}^{\infty} \Omega_{T_n}$ we have that for $\omega \in \overline{\Omega}$

$$\tilde{\xi}(\omega) \in C(R_+; \Phi').$$

Step 4 Uniqueness.

Suppose that there exists a Φ' -valued process $\bar{\xi} = (\bar{\xi}_t)$ that is also a solution of (3.2). Then by Proposition 3.1 for each $T > 0$ there exists a set Ω_3 of probability one such that if $\omega \in \Omega_3$, $\bar{\xi}^T(\omega) \in C([0, T]; \Phi'_{n_T})$ for some $n_T > 0$ (WLOG take $p_T > n_T$) and

$$\bar{\xi}_t(\omega)[\phi] = \gamma(\omega)[\phi] + \int_0^t \bar{\xi}_s[A\phi]ds + W_t[\phi] \quad \forall \phi \in \Phi, \quad 0 \leq t \leq T.$$

Fix $\omega \in \Omega_2 \cap \Omega_3$. Then, suppressing ω in the following, from the above expression we have that for $0 \leq s \leq t \leq T$ and $\phi \in \Phi$

$$W_s[T_{t-s}A\phi] = \bar{\xi}_s[T_{t-s}A\phi] - n[T_{t-s}A\phi] - \int_0^s \bar{\xi}_u[AT_{t-s}A\phi]du. \quad (3.11)$$

Substituting the last expression in the second term of RHS of (3.4) we obtain

$$\begin{aligned} \xi_t[\phi] &= n[T_t\phi] + \int_0^t \bar{\xi}_s[T_{t-s}A\phi]ds - \int_0^t n[T_{t-s}A\phi]ds \\ &\quad - \int_0^t \int_0^s \bar{\xi}_u[AT_{t-s}A\phi]duds + W_t[\phi] \end{aligned}$$

and using Fubini's theorem

$$\begin{aligned} \xi_t[\phi] &= n[T_t\phi] + \int_0^t \bar{\xi}_s[T_{t-s}A\phi]ds - \int_0^t n[T_{t-s}A\phi]ds \\ &\quad - \int_0^t \int_u^t \bar{\xi}_u[AT_{t-s}A\phi]dsdu + W_t[\phi] \end{aligned} \quad (3.12)$$

Next, applying Lemma 3.1 to $F = \gamma$ we have

$$\int_0^t n[T_{t-s}A\phi]ds = n[T_t\phi] - n[\phi]. \quad (3.13)$$

Next since $AT_{t-s}\phi = T_{t-s}A\phi \quad \forall \phi \in \Phi$, applying Lemma 3.1 to $F = \bar{\xi}_u$ we have

$$\int_u^t \bar{\xi}_u[AT_{t-s}A\phi]ds = \int_u^t \bar{\xi}_u[T_{t-s}A^2\phi]ds = \bar{\xi}_u[T_{t-u}A\phi] - \bar{\xi}_u[A\phi] \quad (3.14)$$

Then using (3.14) and (3.13) in (3.12)

$$\begin{aligned}\xi_t[\phi] &= \eta[\phi] + \int_0^t \bar{\xi}_s[T_{t-s}A\phi]ds - \int_0^t \bar{\xi}_u[T_{t-u}A\phi]du \\ &\quad + \int_0^t \bar{\xi}_u[A\phi]du + W_t[\phi] = \eta[\phi] + \int_0^t \bar{\xi}_u[A\phi]du + W_t[\phi] = \bar{\xi}_u[\phi].\end{aligned}$$

Thus for each $T > 0$

$$\xi_t(\omega)[\phi] = \bar{\xi}_t(\omega)[\phi] \quad \forall \phi \in \Phi, \quad 0 \leq t \leq T \quad \omega \in \Omega_2 \cap \Omega_3$$

Hence we have shown that for each $T > 0$ there exists a set Ω_T of probability one given by

$$\Omega_T = \{\omega: \xi_t(\omega) = \bar{\xi}_t(\omega) \quad 0 \leq t \leq T\}.$$

Let $T_n \uparrow \infty$ and define $\bar{\Omega} = \bigcap_{n=1}^{\infty} \Omega_{T_n}$. Then $P(\bar{\Omega}) = 1$ and if $\omega \in \bar{\Omega}$

$$\xi_t(\omega) = \bar{\xi}_t(\omega) \quad \forall \quad t \geq 0$$

i.e.

$$P(\xi_t = \bar{\xi}_t \quad t \geq 0) = 1.$$

The proof of the Theorem is complete.

Q.E.D.

Remarks.

3.1 In the case of a special compatible family (Φ, H, T_t) (Example 1.2), for each $j \geq 1$ ξ_t^j is the one dimensional Ornstein-Uhlenbeck process satisfying

$$d\xi_t^j = -\lambda_j \xi_t^j dt + dW_t[\phi_j]$$

$$\xi_0^j = \eta_0[\phi_j].$$

3.2 In applications one usually deals with an SDE of the form

$$d\tau_t = (-L\tau_t + m)dt + dW_t = L\tau_t dt + dZ_t$$

where Z_t is a Φ' -valued Wiener process with parameters (m, Q) , with $m \in \Phi'$, i.e.

$$EZ_t[\phi] = tm[\phi].$$

In this case we have

$$|m[\phi]|^2 + Q(\phi, \phi) \leq \theta \|\phi\|_{r_2}^2 \quad \forall \phi \in \Phi$$

for some $\theta > 0$ and $r_2 > 0$. Hence in the case of a special compatible family

$$d\tau_t^j = (-\lambda_j \tau_t^j + m_j)dt + dW_t[\phi_j]$$

$$\begin{matrix} j \\ 0 \end{matrix} = \eta[\phi_j]$$

where $m_j = m[\phi_j]$, $\tau_t^j = \tau_t[\phi_j]$.

The following result can be shown in a similar way to Theorem 3.1

Theorem 3.2. Let $M = (M_t)_{t \geq 0}$ be a right continuous Φ' -valued martingale w.r.t. $(\mathcal{F}_t)$ such that

$$E(M_t[\phi])^2 < \infty \quad \forall t \text{ and } \forall \phi \in \Phi.$$

Then the SDE (3.1) has a unique solution given by

$$\xi_t = T_t' \eta + M_t + \int_0^t A' T_{t-s}' M_s ds \quad \text{a.s.}$$

i.e.

$$\xi_t[\phi] = \eta[T_t \phi] + M_t[\phi] + \int_0^t M_s[T_{t-s} A \phi] ds \quad (3.15)$$

with the following property:

a) For each $T > 0$ there exists $\Omega_T \in \mathcal{F}$ with $P(\Omega_T) = 1$ and $P_T > 0$ s.t.

$$\xi_{\bullet}^T(\omega) = \{\xi_t(\omega) : 0 \leq t \leq T\} \in D([0, T]; \Phi_{P_T}') \quad \forall \omega \in \Omega_T.$$

EXAMPLE 3.1 (Poisson driven OU-SDE).

Consider a special compatible family (Φ, H, T_t) as in Example 1.2 where $H = L^2(x, d\Gamma)$ for some $\mathfrak{X}$ and a σ -finite measure Γ . In neurophysiological applications (see [7]) $\mathfrak{X}$ represents the surface membrane of a neuron, e.g. $\mathfrak{X}$ is taken as in Example 1.3 to be $[0, b]$.

Define the Φ' -valued martingale

$$Y_t[\phi] = \int_0^t \int_{R \times \mathfrak{X}} a \phi(x) \tilde{N}(da \, dx \, ds) \quad (3.16)$$

where

$$i) \quad \tilde{N}(da \, d\eta \, ds) = N(da \, d\eta \, ds) - \mu(da \, d\eta) ds$$

and

ii) N is a poisson random measure, i.e. $N([0, t] \times A \times B)$ is a Poisson random variable with parameter $t \cdot \mu(A \times B)$ for $A \in B(R)$ and B is a measurable subset of $\mathfrak{X}$.

The interpretation is that $N([0, t] \times A \times B) =$ the number of voltage pulses of size $a \in A \subseteq R$ arriving at sites $x \in B$ at times $s \leq t$.

Then

$$\begin{aligned} E(Y_t[\phi]Y_t[\psi]) &= \int_0^t \int_{R \times \mathfrak{K}} a^2 \phi(x) \psi(x) \mu(da, dx) ds \\ &= t Q(\phi, \psi) \end{aligned}$$

where

$$Q(\phi, \phi) = \int_{R \times \mathfrak{K}} a^2 (\phi(x))^2 \mu(da \, dx) \quad (3.17)$$

In [7] the semigroup (T_t) represents the evolution semigroup describing the decay of the difference ξ_t between the actual voltage potential at the time $t \geq 0$ and the resting potential on $\mathfrak{K}$, and this difference is modeled as the Φ' -valued solution of the SDE

$$d\xi_t = -L'\xi_t dt + dY_t, \quad E\|\eta\|_{-r_0} < \infty \quad \text{some } r_0 > 0 \quad (3.18)$$

where $A = -L$ is the generator of T_t .

In this situation it can be shown that there exists $p > 0$ s.t. for each $T > 0$

$$\xi_\cdot^T \in D([0, T]; \Phi_p') \quad \text{a.s.}$$

(see [7]) and that $\sum_j \xi_t^j \phi_j$ converges uniformly in $[0, T]$ to $\xi_\cdot^T$ in the Φ_p' topology where

$$\xi_t^j = e^{-t\lambda_j} \xi_0[\phi_j] + \int_0^t e^{-\lambda_j(t-s)} dY_s[\phi_j] \quad (3.19)$$

and λ_j, ϕ_j satisfy (as in Example 1.2) $L\phi_j = \lambda_j \phi_j \quad j \geq 1$.

LECTURE IV

WEAK CONVERGENCE OF SOLUTIONS

At the end of the last lecture (Example 3.1) it was shown how the membrane voltage potential at time t ξ_t of a neuron can be modeled as a Φ' -valued stochastic differential equation driven by a stochastic process with stationary independent increments defined through a Poisson random measure. However, it is believed that the pulse sizes are quite small, making it reasonable to hope that they can be modeled by a Gaussian noise process. Let us now consider the weak convergence of the solutions of (3.14) to the corresponding SDE driven by a Gaussian noise. Most of the material in this lecture is taken from [2] and [7]. Sufficient conditions for the weak convergence of Φ' -valued stochastic processes are given in [11].

Let (Φ, H, T_t) be a compatible family and $\Lambda \in \mathcal{B}(\Phi')$. For each $n > 1$ let μ^n be a measure on $(R \times \Lambda, \mathcal{B}(R) \times \mathcal{B}(\Lambda))$ such that the positive definite bilinear form

$$Q^n(\phi, \psi) := \int_{R \times \Lambda} a^2 \eta[\phi] \eta[\psi] \mu^n(da, d\eta) \quad (4.1)$$

is continuous on $\Phi \times \Phi$, and let N^n be a Poisson random measure with intensity measure $\mu^n(da, d\eta)dt$. Define

$$\tilde{N}^n(da \, d\eta \, ds) := N^n(da \, d\eta \, ds) - \mu^n(da \, d\eta)ds \quad (4.2)$$

and

$$Y^n[\phi] := \int_0^t \int_{R \times \Lambda} a \eta[\phi] \tilde{N}^n(da \, d\eta \, ds) . \quad (4.3)$$

Y_t defined in Example 3.1 is a special case of the above for a large class of spaces $\mathcal{X}$.

For $n \geq 1$ let $m^n \in \Phi'$ and consider the Φ' -valued process ξ_t^n given by

$$\begin{aligned} d\xi_t^n &= A' \xi_t^n dt + m^n dt + dY_t^n \\ \xi_0^n &= \eta^n \end{aligned} \tag{4.4}$$

where η^n is $\mathcal{F}_0$ -measurable.

The following result is proved in [7].

Theorem 4.1 Assume the following six conditions hold:

1) There exists $r_2 > 0$ and $c > 0$ such that for $n \geq 1$

$$(m^n[\phi])^2 + Q^n(\phi, \phi) \leq c \|\phi\|_{r_2}^2 \quad \forall \phi \in \Phi.$$

2) $\lim_{n \rightarrow \infty} Q^n(\phi, \phi) = Q(\phi, \phi) \quad \forall \phi \in \Phi$ for some positive definite bilinear continuous form Q on $\Phi \times \Phi$.

3) $\lim_{n \rightarrow \infty} m^n[\phi] = m[\phi] \quad \forall \phi \in \Phi$ for some $m \in \Phi'$.

4) There exists $r_3 > 0$ such that

$$\sup_n \max\{\|E\eta^n\|_{-r_0}^2, \|E\eta\|_{-r_0}\} < \infty.$$

5) ξ_0^n converges in law to η on Φ'_{r_3} for some $\eta \in \Phi'_{r_0}$.

6) $\lim_{n \rightarrow \infty} \int_{R \times \Lambda} |a_n[\phi]|^3 \mu^n(da d\eta) = 0 \quad \forall \phi \in \Phi$.

Then for each $T > 0$ there exists $p_T > 0$ such that $\xi^{n,T}$ converge weakly to ξ^T on $D([0, T], \Phi_{p_T})$ where ξ is the unique solution of

$$\begin{aligned} d\xi_t &= A' \xi_t dt + m dt + dW_t \\ \xi_0 &= \eta \end{aligned} \quad (4.5)$$

and W is a centered Φ' -valued Wiener process with covariance functional Q .
Furthermore

$$\xi^T \in C([0, T]; \Phi'_T). \quad (4.6)$$

We conclude this lecture by discussing several important examples occurring in applications. They are discussed in [7], [15] and [16].

EXAMPLE 4.1 (White noise current injection at a single point x_0).

Let $H = L^2([0, b], dx), T_t, -L$ and Φ be as in Example 3.1, i.e.

$$\mathcal{X} = [0, b], b = \pi, H = L^2(x, r=Leb), L\phi = -\alpha\phi + \beta\Delta\phi,$$

$$\phi_j(x) = c_j(\cos(jx)), c_0 = 1/\sqrt{\pi}, c_j = \sqrt{2/\pi} \text{ and } \lambda_j = a + \beta(j)^2.$$

Assume that the impulses can arrive only at a single point $x_0 \in [0, b]$ with arrival rate measures μ^n of the form

$$\mu^n(A \times B) = \mu_1^n(A) \cdot 1_B(x_0) \quad A \in \mathcal{B}(R_+), B \in \mathcal{B}((0, b]) \quad (4.7)$$

where

$$\mu_1^n(A) = \sum_{k=1}^p f_e^{k,n} 1_A(a_e^{k,n}) + \sum_{\ell=1}^q f_i^{\ell,n} 1_A(-a_i^{\ell,n}) \quad (4.8)$$

and $a_e^{k,n} \in (0, \infty)$ are the possible sizes of "excitatory" pulse (positive) and $-a_i^{\ell,n}$ are the sizes of "inhibitory" pulses (negative), and $f_e^{k,n}, f_i^{\ell,n}$ are intensities of Poisson processes.

Write:

$$\sigma^2 := \sum_{k=1}^p f_e^{k,n} (a_e^{k,n})^2 + \sum_{\ell=1}^q f_i^{\ell,n} (a_i^{\ell,n})^2 \quad (4.9)$$

$$\gamma_n := \sum_{k=1}^p f_e^{k,n} a_e^{k,n} - \sum_{\ell=1}^q f_i^{\ell,n} a_i^{\ell,n} \quad (4.10)$$

and

$$X_t^n[\phi] = \gamma^n \phi(x_0) + Y_t^n[\phi] \quad \phi \in \Phi \quad (4.11)$$

where

$$\begin{aligned} Y_t^n[\phi] &= \sum_k a_e^{k,n} \int_0^t \int_{\mathbb{X}} \phi(x) \tilde{N}_e^{k,n}(dx, ds) \\ &\quad - \sum_{\ell} a_i^{\ell,n} \int_0^t \int_{\mathbb{X}} \phi(x) \tilde{N}_i^{\ell,n}(dx, ds) \end{aligned} \quad (4.12)$$

and $\tilde{N}_e^{k,n}, \tilde{N}_i^{\ell,n}$ are independent Poisson random measures with variance measures given by $f_e^{k,n} \nu(dx), f_i^{\ell,n} \nu(dx)$ with $\nu(B) = 1_B(x_0)$.

Hence from (4.1) we have that

$$Q^n(\phi, \psi) = \phi(x_0) \psi(x_0) \sigma_n^2 \quad (4.13)$$

It is worth to observe that $\{\phi_j(x)\}_{j \geq 1}$ does not diagonalize Q^n .

For each $n \geq 1$ consider the SDE

$$\begin{aligned} d\xi_t^n &= -L' \xi_t^n dt + dX_t^n \\ \xi_0^n &= \eta_0^n. \end{aligned} \quad (4.14)$$

An application of Theorem 4.1 gives the following weak convergence of solutions of (4.14).

Proposition 4.1. Assume the following four conditions hold:

- 1) $\lim_{n \rightarrow \infty} \max_{k, \ell} \{a_e^{k,n}, a_i^{\ell,n}\} = 0.$

$$2) \lim_{n \rightarrow \infty} \sigma_n^2 = \sigma^2 \text{ for some } 0 < \sigma^2 < \infty.$$

$$3) \lim_{n \rightarrow \infty} \gamma_n = \gamma \text{ for some } \gamma < \infty.$$

$$4) \eta_0^n \text{ converges in law to } \eta_0, \text{ where } \eta_0 \text{ is a } \Phi'_{r_0} \text{-valued gaussian random variable for some } r_0 > 0$$

Then $\lim Q^n(\phi, \psi) = Q(\phi, \psi) := \phi(x_0)\psi(x_0)\sigma^2$ and ξ^n converges weakly to ξ where ξ is the unique solution of the SDE

$$\begin{aligned} d\xi_t &= -L'\xi_t dt + dW_t \\ \xi_0 &= \eta_0 \end{aligned} \quad (4.15)$$

where W is a centered Φ' -valued Wiener process with covariance Q . Moreover using Example 1.3 and Theorem 3.1(c) we have that $W_\cdot \in C(R_+; \Phi'_p)$ for $p > 1/4$ and $\xi \in C(R_+; \Phi'_q)$ where $q > \max(1/4, r_0)$.

Furthermore $\xi^j := \xi_t[\phi_j]$ satisfies the real valued Ornstein-Uhlenbeck SDE

$$d\xi_t^j = [-\lambda_j \xi_t^j + \gamma \phi_j(x_0)]dt + \sigma |\phi_j(x_0)| dW_t^j \quad (4.16)$$

$$\xi_0^j = \xi_0[\phi_j]$$

where the one dimensional standard Wiener processes $W_t^j = W_t[\phi_j]$ $j \geq 1$ are not independent.

EXAMPLE 4.2 (White noise current uniformly distributed over $\mathbb{X}$).

Consider the previous example but now assuming that white noise injection can occur at any point in $[0, b]$ with arrival rate measure

$$\mu^n(A \times B) = \mu_1^n(A) \nu^n(B) \quad A \in \mathcal{B}(R_+), \quad B \in \mathcal{B}([0, \pi]) \quad (4.17)$$

where μ_1^n is defined in (4.8) and ν^n is a sequence of probability measures on $\mathcal{X} = [0, b]$ such that ν^n converge weakly to ν where ν is $1/b$ Lebesgue measure. Then for each $n \geq 1$

$$Q^n(\phi, \psi) = \sigma^2 \int_0^b \phi(x) \psi(x) \nu^n(dx) \quad \phi, \psi \in \Phi \quad (4.18)$$

In this case the corresponding weak convergence result is given by the following proposition.

Proposition 4.2. Assume that the conditions (1)-(4) of Proposition 4.1 are satisfied. Then

$$a) \quad \lim_{n \rightarrow \infty} Q^n(\phi, \psi) = Q(\phi, \psi) := \frac{\sigma^2}{b} \int_0^b \phi(x) \psi(x) dx.$$

$$b) \quad \xi_n \text{ converge weakly to } \xi.$$

where ξ_t is the unique solution of (4.15) satisfying the same conditions as in Proposition 4.1 with the difference that W is a Φ' -valued Wiener process with covariance functional Q given by (a).

The Wiener process W can, in this instance, be defined in terms of the centered 2-parameter Wiener process $W_{t,x}$ ($0 \leq t \leq T$, $0 \leq x \leq b$) with covariance

$$E(W_{t,x} W_{s,y}) = \frac{1}{b} \min(t, s) \min(x, y), \quad (4.19)$$

by

$$W_t[\phi] := \frac{1}{b} \int_0^b \phi(x) d_x W_{t,x}.$$

Further properties of the solution ξ_t in this case have been investigated in detail by J. Walsh [15].

Theorem 3.1(c) and Corollary 2.3 are examples of results for which in the case of a special compatible family one can obtain additional information on the spaces where some Φ' -valued processes lie. This information is given in terms of r_1 (which satisfies condition (1.3) in Example 1.2) and of r_2 given in Corollary 2.3 and related to the covariance functional Q . The following Lemma is useful in determining r_2 when Q is of the form presented in the previous examples.

Lemma 4.1 Let (Φ, H, T_t) be a special compatible family. Suppose there exists $r > 0$ such that

$$c_1 := \sup_x \sup_j |\phi_j(x)| (1 + \lambda_j)^{-r} < \infty \quad \forall \quad \phi \in \Phi$$

and

$$c_2 := \int_{R \times \mathcal{X}} a^2 \mu(da, dx) < \infty$$

Then

$$Q(\phi, \psi) = \int_{R \times \mathcal{X}} a^2 \phi(x) \psi(x) \mu(da, dx) \quad (4.20)$$

satisfies

$$Q(\phi, \phi) \leq \theta \|\phi\|_{r+r_1}^2 \quad (4.21)$$

where $(\lambda_j)_{j \geq 1}$ and r_1 are given in Example 1.2.

The proof of the Lemma is given in [7].

EXAMPLE 4.3

Let $\mathcal{X} = S^2$, the unit sphere in R^3 , with Lebesgue surface measure Γ . Let Δ_B be the Laplace-Beltrami operator

$$\Delta_B \phi = (\sin \theta)^{-1} \left[\frac{\partial}{\partial \theta} (\sin \theta) \frac{\partial \phi}{\partial \theta} + \frac{\partial}{\partial \eta} (\sin \theta)^{-1} \frac{\partial \phi}{\partial \eta} \right] \quad (4.22)$$

(where θ, η are the Euler angles on S^2). Let $L = -\beta + \delta \Delta_B$ for $\beta, \delta > 0$ and $H = L^2(\mathcal{X}, \Gamma)$. This time the eigenfunctions are the spherical harmonics $Y_{\ell m} (\ell=0,1,\dots; m = -\ell, \dots, \ell)$ with eigenvalues $\lambda = \beta + \delta \ell(\ell + 1)$ for L , $e^{-t\lambda}$ for T_t .

Write

$$\phi_j = Y_{\ell m}, \quad j = m + \ell(\ell + 1) \quad (4.23)$$

(i.e. $\ell = [\sqrt{j}]$, $m = j - \ell^2 - \ell$)

$$\lambda_j = \beta + \delta [\sqrt{j}]([\sqrt{j}] + 1) . \quad (4.24)$$

Then

$$\sum_j (1 + \lambda_j)^{-2r_1} = \sum_{\ell} (2\ell + 1) [1 + \beta + \delta(\ell^2 + \ell)]^{-2r_1} < \infty$$

if $r_1 > 1/2$. Hence condition (1.3) in Example 1.2 is satisfied and a special compatible family (ϕ, H, T_t) can be constructed. Furthermore

$$\sup_x |Y_{\ell m}(x)|^2 = \left(\frac{2\ell + 1}{4\pi} \right)$$

and therefore

$$\sup_j \sup_x |\phi_j(x)| (1 + \lambda_j)^{-1/4} = [4\pi^2 \min(\delta, 4\beta)]^{-1/4} < \infty .$$

Hence c_1 in Lemma 4.1 is finite for $r = 1/4$. Then by Lemma 4.1 for any measure μ on $R \times \mathfrak{X}$ such that

$$\int_{R \times \mathfrak{X}} a^2 \mu(da, dx) < \infty \quad (4.25)$$

we have that if Q is given by (4.26)

$$Q(\phi, \phi) \leq \theta \|\phi\|_{r+r_1}^2 \quad \forall \phi \in \Phi$$

where $r + r_1 > 3/4$, i.e. $r_2 = r + r_1$. Then if q is as in Corollary 2.3

$$q \geq r_1 + r_2 = \frac{1}{2} + \frac{3}{4}, \text{ i.e. } q \geq \frac{5}{4}.$$

A more general result is the following:

Proposition 4.3 Let $\mathfrak{X}$ be a smooth d -dimensional compact Riemannian manifold with smooth (possibly empty) boundary $\partial\mathfrak{X}$ and Riemannian volume element $d\Gamma$. Let L be a positive, self-adjoint operator on a domain $\mathcal{D} \subseteq H = L^2(\mathfrak{X}, d\Gamma)$ such that

- a) $C_0^\infty(\mathfrak{X}) \subseteq \mathcal{D}$ (if $\partial\mathfrak{X}$ is not empty, $\mathfrak{X} \cup \partial\mathfrak{X}$ is compact and $C_0^\infty(\) = C^\infty$ -functions on $\mathfrak{X}$ whose support lies in $\mathfrak{X}$. If $\partial\mathfrak{X}$ is empty, $\mathfrak{X}$ is compact and $C_0^\infty(\mathfrak{X}) = C^\infty(\mathfrak{X})$).
- b) $L_0 := L|_{C_0^\infty(\)}$ is a uniformly strongly elliptic differential operator of order $2m > 0$ with smooth coefficients.
- c) $\mathcal{D} \subseteq W_{2m}(\mathfrak{X})$, the Hilbert space of those elements in H with $2m$ weak derivatives in H .

Then L admits a CONS $\{\phi_j\}$ of eigenfunctions in H with eigenvalues $\{\lambda_j\}$ satisfying

- i) $L\phi_j = \lambda_j \phi_j$
- ii) $\phi_j \in C^\infty(\mathcal{X})$
- iii) $\sum_j (1 + \lambda_j)^{-2r_1} < \infty$ for $r_1 > \frac{d}{4m}$
- iv) $\sup_j \sup_x |\phi_j(x)| (1 + \lambda_j)^{-r} < \infty$ all $r > \frac{d}{4m}$

Hence by condition (iii) above and (1.3) we have that a special compatible family can be constructed in the manner of Example (1.2). Moreover from (iii) above if μ is any measure on $R \times \mathcal{X}$ satisfying (4.25), by Lemma 4.1 $r_2 > d/2m$. Then if q is as in Corollary 2.3 $q \geq r_1 + r_2$ i.e. $q \geq 3d/4m$.

Additional examples and applications of Φ' -valued processes can be seen in [7] and references therein.

LECTURE V

STOCHASTIC EVOLUTION EQUATIONS AND NONLINEAR STOCHASTIC DIFFERENTIAL EQUATIONS

In this lecture we will give an outline of recent works on stochastic evolution equations and nonlinear stochastic differential equations on the dual of a Countably Hilbert nuclear space.

Throughout this lecture $(\Omega, \mathcal{F}, P)$ will denote a complete probability space with a right continuous filtration $(\mathcal{F}_t)_{t \geq 0}$ and $(\Phi, \|\cdot\|_r, r \geq 0)$ will be a Countably Hilbert nuclear space.

1. STOCHASTIC EVOLUTION EQUATIONS

The material of this section is recent joint work with V. Perez-Abreu.

There are several possible extensions of the SDE (3.1). For example one may consider an evolution operator A_t instead of the infinitesimal generator $-L$ and/or a perturbation operator P_t . In this section we consider the SDE

$$d\xi_t = (A_t' \xi_t + P_t' \xi_t)dt + dW_t \quad (I)$$

$$\xi_0 = \gamma$$

where γ is a $\mathcal{F}_0$ -measurable Φ' -valued gaussian random variable s.t.

$E\|\gamma\|_{-r_0}^2 < \infty$ some $r_0 > 0$ and W_t is a Φ' -valued Wiener process with covariance Q . By Theorem 2.1. $W. \in C(R, \Phi_q')$ a.s. for some $q > 0$. The operators A_t and P_t from Φ to Φ are assumed to satisfy the following conditions:

Assumptions on A_t :

a) For each $t \geq 0$ $A_t: \Phi \rightarrow \Phi$ is a continuous linear operator.

b) For each $\phi \in \Phi$ the map $t \rightarrow A_t \phi$ is continuous.

c) A_t is the generator of a two parameter semigroup $T(s,t)$

$0 \leq s \leq t < \infty$ ($T(s,t) = T(s,t')T(t',t)$ $s < t' < t$, $T(t,t) = I$), i.e.

$$\frac{d}{dt} T(s,t)\phi = T(s,t)A_t \phi \quad \forall \phi \in \Phi \quad 0 \leq s \leq t$$

and

$$\frac{d}{ds} T(s,t)\phi = -A_s T(s,t)\phi \quad \forall \phi \in \Phi \quad 0 \leq s \leq t.$$

d) For $s \leq t$ $T(s,t): \Phi \rightarrow \Phi$ is a continuous linear operator,

e) $\lim_{t \rightarrow t_0} T(s,t)\phi = T(s,t_0)\phi$ in the Φ -topology for each s fixed and

$0 \leq s \leq t_0$, $\phi \in \Phi$, and $\lim_{s \rightarrow s_0} T(s,t)\phi = T(s_0,t)\phi$ for each t

fixed and $0 \leq s_0 \leq t$, $\phi \in \Phi$,

d) for each $T > 0$ and $n > 0$

$$\sup_{0 \leq s \leq t \leq T} \|T(s,t)\phi\|_n < \infty \quad \forall \phi \in \Phi.$$

Assumptions on P_t :

e) For each $t \geq 0$ $P_t: \Phi \rightarrow \Phi$ is a continuous linear operator

f) There exists a sequence of seminorms $\{\|\cdot\|_n; n \geq 0\}$ on Φ generating an equivalent topology as that given by the Hilbertian norms $\{\|\cdot\|_n; n \geq 0\}$ such that the following holds:

i) for each $T > 0$ there exists $m_T > 0$ s.t. for $m \geq m_T$ P_t has a continuous linear extension to $\Phi_{|m|}$ ($\|\cdot\|_m$ -completion of Φ) and the map $s \rightarrow P_s \phi$ is $\Phi_{|m|}$ -continuous,

ii) for each $T > 0$ there exists $m_T > 0$ s.t. for $m \geq m_T$ $\exists$
 $K(m, T) > 0$ and

$$\sup_{0 \leq s \leq t \leq T} \|P_s T(s, t) \phi\|_m \leq K(m, T) \|\phi\|_m \quad \forall \phi \in \Phi$$

Observe that condition f (ii) above can be obtained using f (i) if we assume that
 for each $T > 0$ and $m \geq 0$

$$\sup_{0 \leq s \leq t \leq T} \|T(s, t) \phi\|_m \leq D(m, T) \|\phi\|_m \quad \forall \phi \in \Phi$$

for some $D(m, T) > 0$.

Definition 5.1 We say that the SDE (I) has a Φ' -valued solution $\xi = (\xi_t)_{t \geq 0}$
 if the following four conditions hold:

- i) (ξ_t) is $\mathcal{F}_t$ -adapted and Φ' -valued.
- ii) $\xi_t[\phi] = \gamma[\phi] + \int_0^t \xi_s[A_s \phi] ds + \int_0^t \xi_s[P_s \phi] ds + W_t[\phi] \quad \forall \phi \in \Phi$
 a.s. $\forall t \geq 0$.
- iii) $\xi_\cdot \in C(\mathbb{R}_+; \Phi')$ a.s.
- iv) For each $T > 0$

$$E\left(\sup_{0 \leq t \leq T} |\xi_t[\phi]|^2\right) < \infty.$$

Proposition 5.1 If $(\xi_t)_{t \geq 0}$ is a solution of the SDE (I) then for each $T > 0$
 there exist $q_T > 0$ and a version of ξ (denoted also by ξ) such that

$$\xi_\cdot^T \in C([0, T]; \Phi'_{q_T}) \text{ a.s.}$$

and

$$\xi_t[\phi] = \gamma[\phi] + \int_0^t \xi_s[A_s \phi] ds + \int_0^t \xi_s[P_s \phi] ds + W_t[\phi]$$

$$\forall \phi \in \Phi, \quad 0 \leq t \leq T \quad \text{a.s.} \quad (5.1)$$

The proof of this proposition is done in a manner very similar to Proposition 3.1.

Remark 1. Condition (iv) in Definition 5.1 is implied by the following condition: For each $T > 0$

$$E \int_0^T (\xi_s[A_s \phi])^2 ds + E \int_0^T (\xi_s[P_s \phi])^2 ds < \infty \quad \forall \phi \in \Phi.$$

Theorem 5.1 Let γ and $W = (W_t)_{t \geq 0}$ be as in the beginning of this lecture and suppose that A_t and P_t satisfy the assumptions (a)-(d) and (e)-(f) respectively. Then the SDE (I) has a unique solution $\xi = (\xi_t)_{t \geq 0}$ such that for each $T > 0$ there exists $p_T > 0$ and

$$\xi_{\cdot}^T \in C([0, T]; \Phi'_{p_T}) \quad \text{a.s.}$$

and

$$E \left(\sup_{0 \leq t \leq T} \|\xi_t\|_{\Phi'_{p_T}}^2 \right) < \infty.$$

Moreover ξ_t is a Φ' -valued gaussian process.

Proof. The idea is first to show that the unperturbed SDE

$$d\eta_t = A_t' \eta_t dt + dW_t$$

$$\eta_0 = \gamma \quad (II)$$

has a unique solution. This is done a manner entirely similar to the argument in Theorem 3.1 using Lemma 5.1 below instead of Lemma 3.1, and using only the

assumptions on A_t . The solution is given by

$$\eta_t[\phi] = \gamma[T(0,t)\phi] + \int_0^t W_s[A_s T(s,t)\phi] ds + W_t[\phi] . \quad (5.2)$$

The next step is to show, by the method of successive approximations, that the stochastic equation

$$\xi_t = \int_0^t T'(s,t) P_t' \xi_s ds + \eta_t \quad (5.3)$$

has a unique solution on $C(R_+; \Phi')$: The process η_t given by (5.2) has the property that for each $T > 0$ there is a $q_T > 0$ s.t. $E(C_T)^2 < \infty$ where

$$C_T(\omega) := \sup_{0 \leq t \leq T} \|\eta_t(\omega)\|_{-q_T} \quad (5.4)$$

Next, using assumption (f) there exist constants $c_i = c_i(T, q_T)$ $i = 1, 2$ $m_T > 0$ and $n_T > 0$ s.t.

$$\|\phi\|_{q_T} \leq c_1 \|\phi\|_{m_T} \leq c_2 \|\phi\|_{n_T} \quad \forall \phi \in \Phi . \quad (5.5)$$

From (f) (ii) it then follows that

$$\sup_{0 \leq t \leq T} \|P_s T(s,t)\phi\|_{m_T} \leq K_T \|\phi\|_{m_T} \quad \forall \phi \in \Phi . \quad (5.6)$$

Let

$$\Omega_1 = \{\omega: C_T(\omega) < \infty\}$$

then $P(\Omega_1) = 1$. For $\omega \in \Omega_1$ and $0 \leq t \leq T$ define the sequence of successive approximations

$$\xi_t^0(\omega) = \eta_t(\omega)$$

$$\xi_t^1(\omega) = \int_0^t T'(s,t) P'_s \xi_s^0(\omega) ds + \eta_t(\omega)$$

⋮

⋮

$$\xi_t^n(\omega) = \int_0^t T'(s,t) P'_s \xi_s^{n-1}(\omega) ds + \eta_t(\omega).$$

Then for each $n \geq 0$ and $\phi \in \Phi$

$$\xi_t^0[\phi] = \eta_t[\phi]$$

$$\xi_t^1[\phi] = \int_0^t \eta_s[P_s T(s,t) \phi] ds + \eta_t[\phi]$$

⋮

⋮

⋮

$$\xi_t^n[\phi] = \int_0^t \xi_t^{n-1}[P_s T(s,t) \phi] ds + \eta_t[\phi]$$

$$= \int_0^t \int_0^{s_1} \dots \int_0^{s_{n-2}} \eta_{s_{n-1}}[P_{s_{n-1}} T(s_{n-1}, s_{n-2}) \dots$$

$$P_{s_1} T(s_1, t) \phi] ds_{n-1} \dots ds_1$$

$$+ \int_0^t \int_0^{s_1} \dots \int_0^{s_{n-3}} \eta_{s_{n-2}}[P_{s_{n-2}} T(s_{n-2}, s_{n-3}) \dots$$

$$P_{s_1} T(s_1, t) \phi] ds_{n-2} \dots ds_1$$

$$+ \dots + \eta_t[\phi].$$

Hence, using (5.5) and (5.6) one shows that the above integrals are well defined and moreover

$$|\xi_t^n(\omega)[\phi]| \leq C_T(\omega) c_1 c_2 \left(\sum_{k=0}^n \frac{(K_T T)^k}{k!} \right) \|\phi\|_{n_T} \quad \forall \phi \in \Phi \quad (5.7)$$

and

$$|\xi^n(\omega)[\phi] - \xi^m(\omega)[\phi]| \leq C_T(\omega) c_1 c_2 \left(\sum_{k=m+1}^n \frac{(K_T T)^k}{k!} \right) \|\phi\|_{n_T} \quad \forall \phi \in \Phi. \quad (5.8)$$

Then for each $\omega \in \Omega_1$, $0 \leq t \leq T$ and $\phi \in \Phi$ $\xi_t^n(\omega)$ is a Φ' -valued element for $n \geq 1$, $\xi_t(\omega)[\phi] = \lim \xi_t^n(\omega)[\phi]$ exists and

$$\sup_{0 \leq t \leq T} |\xi_t(\omega)[\phi]| \leq C_T(\omega) c_1 c_2 e^{K_T T} \|\phi\|_{n_T} \quad \forall \phi \in \Phi. \quad (5.9)$$

Hence $\xi_t(\omega) \in \Phi'$. Moreover let $\ell_T > n_T$ be such that the injection map $\Phi_{\ell_T} \hookrightarrow \Phi_{q_T}$ is Hilbert-Schmidt and let $\{\phi_j\}_{j \geq 1} \subseteq \Phi$ be a CONS for Φ_{ℓ_T} with dual basis $\{\phi_j\}_{j \geq 1}$ a CONS for Φ'_{ℓ_T} . Then from (5.9) we have

$$\sup_{0 \leq t \leq T} \sum_{j=1}^{\infty} |\xi_t(\omega)[\phi_j]|^2 \leq C_T^2(\omega) c_1^2 c_2^2 e^{2K_T T} \sum_{j=1}^{\infty} \|\phi_j\|_{n_T}^2 < \infty$$

and therefore for $0 \leq t \leq T$ and $\omega \in \Omega_1$

$$\tilde{\xi}_t(\omega) = \sum_{j=1}^{\infty} \xi_t(\omega)[\phi_j] \phi_j \quad (5.10)$$

is a well defined element of Φ'_{ℓ_T} and $\tilde{\xi}_t(\omega)[\phi] = \xi_t(\omega)[\phi] \quad \forall \phi \in \Phi$. From now on we will write ξ_t instead of $\tilde{\xi}_t$.

Next we shall show that ξ_t is a solution of (5.3): From (5.7) and assumption f(ii) we have (suppressing $\omega \in \Omega_1$ in the writing)

$$\begin{aligned} |\xi^{n-1}[P_S T(s,t)\phi]| &\leq \|\xi_S^{n-1}\|_{\mathcal{L}_T} \|P_S T(s,t)\phi\|_{\mathcal{L}_T} \\ &\leq C_T c_1 c_2 e^{TK_T} K_T \|\phi\|_{m_T} < \infty \end{aligned}$$

$$\forall \phi \in \Phi, n \geq 1, 0 \leq s \leq t \leq T.$$

Then since

$$\xi_t^n(\omega)[\phi] = \int_0^t \xi_s^{n-1}(\omega)[P_S T(s,t)\phi] ds + \eta_t(\omega)[\phi], \quad \omega \in \Omega_1,$$

by the dominated convergence theorem

$$\begin{aligned} \xi_t[\phi] &= \lim_{n \rightarrow \infty} \xi_t^n[\phi] = \lim_{n \rightarrow \infty} \int_0^t \xi_s^{n-1}[P_S T(s,t)\phi] ds + \eta_t[\phi] \\ &= \int_0^t \xi_s[P_S T(s,t)\phi] ds + \eta_t[\phi], \quad \forall \phi \in \Phi, 0 \leq t \leq T. \text{ a.s. } (5.11) \end{aligned}$$

which shows that ξ_t satisfies condition (ii) in Definition 5.1.

Next we shall show that $\xi_t \in C([0,T]; \Phi'_{p_T})$ a.s. for some $p_T > 0$. Let $t_0, t \in [0,T]$. Then using Lemma 5.1 below it is not difficult to show that for $\omega \in \Omega_1$

$$\left| \int_0^t \xi_u(\omega)[P_u T(u,t)\phi] du - \int_0^{t_0} \xi_u(\omega)[P_u T(u,t_0)\phi] du \right| \leq C_T(\omega) D_T \|\phi\|_{r_T} |t - t_0|$$

for some $D_T > 0$ and $r_T > 0$. Hence the process

$$Z_t[\phi] := \int_0^t \xi_u[P_u T(u,t)\phi] du$$

is a continuous process in $t \in [0,T]$ for each $\phi \in \Phi$. Hence from (5.11) we have that $\xi_t[\phi]$ is also a continuous function of t . Moreover,

$$\sup_{0 \leq t \leq T} |\xi_t(\omega)[\phi]| \leq \{C_T(\omega) D_T + C_T(\omega)\} \|\phi\|_{r_T} \quad \forall \phi \in \Phi, \omega \in \Omega_1. \quad (5.12)$$

Next let $p_T > r_T$ be such that the injection map $\Phi_{p_T} \hookrightarrow \Phi_{r_T}$ is Hilbert Schmidt and let $\{e_j\}_{j \geq 1} \subseteq \Phi$ be a CONS for Φ_{p_T} with dual basis $\{\hat{e}_j\}_{j \geq 1}$ a CONS for

Φ'_{p_T} . Then from (5.12)

$$\sup_{0 \leq t \leq T} \sum_{j=1}^{\infty} |\xi_t[e_j]|^2 \leq (C_T D_T + C_T)^2 \sum_{j=1}^{\infty} \|e_j\|_{r_T}^2 < \infty.$$

Hence for $\omega \in \Omega_1$ define $\tilde{\xi}_t(\omega) = \sum_{j=1}^{\infty} \xi_t(\omega)[e_j] \hat{e}_j$ which is an element in Φ'_{p_T} . Then

$$\tilde{\xi}_t(\omega)[\phi] = \xi_t(\omega)[\phi] \quad \forall \phi \in \Phi \quad 0 \leq t \leq T, \omega \in \Omega_1.$$

From now on we write ξ_t instead of $\tilde{\xi}_t$. Then by the dominated convergence theorem, since $\xi_t(\omega)[e_j]$ is continuous on t for each $j > 1$ we have

$$\begin{aligned} \lim_{t \rightarrow t_0} \|\xi_t - \xi_{t_0}\|_{-p_T}^2 &= \lim_{t \rightarrow t_0} \sum_{j=1}^{\infty} (\xi_t[e_j] - \xi_{t_0}[e_j])^2 \\ &= \sum_{j=1}^{\infty} \lim_{t \rightarrow t_0} (\xi_t[e_j] - \xi_{t_0}[e_j])^2 = 0 \quad t_0 \in [0, T] \end{aligned}$$

Then $\xi^T(\omega) \in C([0, T]; \Phi'_{p_T})$ for some $p_T > 0$ and $\omega \in \Omega_1$ where $P(\Omega_1) = 1$.

Also from (5.4) and (5.12) we have that for each $T > 0$

$$E\left(\sup_{0 \leq t \leq T} |\xi_t[\phi]|^2\right) < \infty \quad \forall \phi \in \Phi.$$

which shows condition (iv) in Definition 5.1. Moreover from (5.12) and since $E(C_T^2) < \infty$

$$E\left(\sup_{0 \leq t \leq T} \|\xi\|_{-p_T}^2\right) \leq E(C_T D_T + C_T)^2 \sum_{j=1}^{\infty} \|e_j\|_{r_T}^2 < \infty.$$

A similar argument to that at end of Step 3 in Theorem 3.1 gives condition (ii) in Definition 5.1, i.e.

$$\xi_{\cdot} \in C(R_+; \Phi') \quad \text{a.s.}$$

Hence, ξ is a solution of (5.3).

To show uniqueness let X be any solution of (5.3). For the present assume that X_t satisfies the following condition:

(*) For each $T > 0$ there exists $p_T^1 > 0$ s.t. $\chi_{\cdot}^T \in C([0, T]; \Phi_{p_T^1}^1)$ a.s.

W.L.O.G. let $p_T^1 > p_T$ and

$$\Omega_2 = \{\omega: \sup_{0 \leq t \leq T} \|\chi_t(\omega)\|_{-p_T} < \infty\}.$$

Then $P(\Omega_2) = 1$. Fix $\omega \in \Omega_1 \cap \Omega_2$ and let $0 \leq t \leq T$. Then for each $\phi \in \Phi$ (suppressing ω in the writing)

$$\chi_t[\phi] = \int_0^t \chi_s[P_s T(s, t) \phi] ds + \eta_t[\phi].$$

Next, if ξ_t^n is the sequence of successive approximations defined prior to (5.7) we have

$$\chi_t[\phi] - \xi_t^0[\phi] = \int_0^t \chi_s[P_s T(s, t) \phi] ds \quad (5.13)$$

$$\chi_t[\phi] - \xi_t^1[\phi] = \int_0^t \chi_s[P_s T(s, t) \phi] ds - \int_0^t \xi_s^0[P_s T(s, t) \phi] ds$$

⋮

$$\chi_t[\phi] - \xi_t^n[\phi] = \int_0^t (\chi_s[P_s T(s, t) \phi] - \xi_s^{n-1}[P_s T(s, t) \phi]) ds.$$

Hence

$$\begin{aligned} \chi_t[\phi] - \xi_t^n[\phi] &= \int_0^t \int_0^{s_1} \dots \int_0^{s_{n-1}} \{\chi_{s_n}[P_{s_n} T(s_n, s_{n-1}) \dots P_{s_1} T(s_1, t) \phi] \\ &\quad - \xi_{s_n}^1[P_{s_n} T(s_n, s_{n-1}) \dots P_{s_1} T(s_1, t) \phi]\} ds_n \dots ds_1 \\ (5.13) \quad &= \int_0^t \int_0^{s_1} \dots \int_0^{s_n} \chi_{s_{n+1}}[P_{s_{n+1}} T(s_{n+1}, s_n) \dots \\ &\quad P_{s_1} T(s_1, t) \phi] ds_{n+1} \dots ds_1 \end{aligned}$$

Then using inequalities similar to (5.5) and (5.6) it follows that

$$|X_t[\phi] - \xi_t^n[\phi]| \leq \sup_{0 \leq t \leq T} \|X_t\|^{-p_T} c_1 c_2 \frac{(K_T T)^n}{n!} \|\phi\|_m$$

for some positive constants c_1, c_2, K_T and m . Hence

$$\sup_{0 \leq t \leq T} |X_t[\phi] - \xi_t^n[\phi]| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus $P(X_t = \xi_t \quad 0 \leq t \leq T) = 1$ and a similar argument to that at the end of Step 4 in Theorem 3.1 gives that $P(X_t = \xi_t \quad t \geq 0) = 1$.

The next step is to show that the solution ξ of (5.3) obtained above is also a solution of (I). In order to do that we need the following Lemma.

Lemma 5.1 Suppose conditions (a)-(d) on A_t hold and let B be any continuous linear operator from Φ to Φ . Then for each $F \in \Phi'$ and $0 \leq u \leq t$

$$a) \quad F[BT(u,t)\phi] = F[B\phi] + \int_u^t F[BT(u,s)A_s\phi]ds \quad \forall \phi \in \Phi$$

$$b) \quad F[BT(u,t)\phi] = F[B\phi] + \int_u^t F[BA_sT(s,t)\phi]ds \quad \forall \phi \in \Phi.$$

The proof of the Lemma is similar to that of Lemma 3.1 using the Kolmogorov Forward and Backward equations:

$$\frac{d}{ds} T(u,s)\phi = T(u,s)A_s\phi \quad 0 \leq u \leq s, \quad \phi \in \Phi$$

$$\frac{d}{du} T(u,s) = -A_uT(u,s)\phi \quad 0 \leq u \leq s, \quad \phi \in \Phi.$$

End of the Proof of Theorem 5.1

We shall show that ξ is also a solution of (I) and that it is unique.

Let $\omega \in \Omega_1$ then from (5.11) (suppressing ω in the writing) we have

$$\xi_t[\phi] = \int_0^t \xi_s[P_s T(s,t)\phi]ds + \eta_t[\phi] \quad \forall \phi \in \Phi \quad 0 \leq t \leq T. \quad (5.14)$$

Next, applying Lemma 5.1(a) with $B = P_u$ and $F = \xi_u$ we have

$$\xi_u[P_u T(u,t)\phi] = \xi_u[P_u \phi] + \int_u^t \xi_u[P_u T(u,s)A_s \phi]ds$$

and therefore

$$\begin{aligned} \int_0^t \xi_u(\omega)[P_u T(u,t)\phi]du &= \int_0^t \xi_u(\omega)[P_u \phi]du + \int_0^t \int_u^t \xi_u(\omega)[P_u T(u,s)A_s \phi]dsdu \\ &= \int_0^t \xi_u(\omega)[P_u \phi]du + \int_0^t \int_0^s \xi_u(\omega)[P_u T(u,s)A_s \phi]duds. \end{aligned} \quad (5.15)$$

But from (5.14) since

$$\xi_s[A_s \phi] = \int_0^s \xi_u[P_u T(u,s)A_s \phi]du + \eta_s[A_s \phi],$$

using the above expression in the second term of (5.15) we obtain

$$\int_0^t \xi_u[P_u T(u,t)\phi]du = \int_0^t \xi_u[P_u \phi]du + \int_0^t \xi_s[A_s \phi]ds - \int_0^t \eta_s[A_s \phi]ds. \quad (5.16)$$

But also from (5.14) we obtain

$$\int_0^t \xi_u[P_u T(u,t)\phi]du = \xi_t[\phi] - \eta_t[\phi].$$

Hence from the above expression and (5.16),

$$\xi_t[\phi] - \eta_t[\phi] = \int_0^t \xi_s[P_s \phi]ds + \int_0^t \xi_s[A_s \phi]ds - \int_0^t \eta_s[A_s \phi]ds$$

i.e.

$$\xi_t[\phi] = \int_0^t \xi_s[P_s \phi]ds + \int_0^t \xi_s[A_s \phi]ds + \eta_t[\phi] - \int_0^t \eta_s[A_s \phi]ds$$

but since $\eta_t[\phi] - \int_0^t \eta_s[A_s \phi]ds = \gamma[\phi] + W_t[\phi],$

$$\xi_t[\phi] = \int_0^t \xi_s[P_s \phi] ds + \int_0^t \xi_s[A_s \phi] ds + \gamma[\phi] + W_t[\phi] \quad \forall \phi \in \Phi,$$

i.e.

$$d\xi_t = A_t' \xi_t dt + P_t' \xi_t dt + dW_t.$$

Finally we shall show the uniqueness of the solution (I) proving that any other solution $\bar{\xi}_t$ of (I) satisfies the SDE (5.3). Let Ω_3 be the set of probability one given by Proposition 5.1 s.t.

$$\bar{\xi}_\cdot^T \in C([0, T]; \Phi_{q_T}') \quad \text{a.s.}$$

and

$$\bar{\xi}_t[\phi] = \gamma[\phi] + \int_0^t \bar{\xi}_s[A_s \phi] ds + \int_0^t \bar{\xi}_s[P_s \phi] ds + W_t[\phi]$$

$$\forall \phi \in \Phi \quad 0 \leq t \leq T \quad \text{a.s.} \quad (5.17)$$

WLOG we can take $q_T > p_T$. Let $\omega \in \Omega_3 \cap \Omega_1$. Then (suppressing ω in the following) from (5.17) we have that for $0 \leq s \leq T$

$$\begin{aligned} W_s[A_s T(s, t) \phi] &= \bar{\xi}_s[A_s T(s, t) \phi] - \int_0^t \bar{\xi}_u[P_u A_s T(s, t) \phi] du - \\ &\quad - \int_0^t \bar{\xi}_u[A_u A_s T(s, t) \phi] du - \gamma[A_s T(s, t) \phi]. \end{aligned} \quad (5.18)$$

On the other hand from (5.3) and (5.2) if $0 \leq t \leq T$

$$\xi_t[\phi] - \int_0^t \xi_s[P_s T(s, t) \phi] ds = \int_0^t W_s[A_s T(s, t) \phi] ds + \gamma[T(0, t) \phi] + W_t[\phi]. \quad (5.19)$$

Hence using (5.18) in (5.19) we have

$$\begin{aligned}
 \xi_t[\phi] - \int_0^t \xi_s[P_s T(s,t)\phi]ds &= \int_0^t \bar{\xi}_s[A_s T(s,t)\phi]ds \\
 &\quad - \int_0^t \int_0^s \bar{\xi}_u[P_u A_s T(s,t)\phi]duds - \int_0^t \int_0^s \bar{\xi}_u[A_u A_s T(s,t)\phi]duds \\
 &\quad - \int_0^t \gamma[A_s T(s,t)\phi]ds + \gamma[T(0,t)\phi] + W_t[\phi] \quad . \quad (5.20)
 \end{aligned}$$

Next, using Lemma 5.1(b) with $F = \gamma$, $B = I$ we have

$$- \int_0^t \gamma[A_s T(s,t)\phi]ds + \gamma[T(0,t)\phi] = \gamma[\phi]. \quad (5.21)$$

Again, applying Lemma 5.1(b) with $F = \bar{\xi}_u$, $B = P_u$ and with $F = \bar{\xi}_u$ and $B = A_u$ we have the following two expressions

$$- \int_0^t \bar{\xi}_u[P_u A_s T(s,t)\phi]ds = \bar{\xi}_u[P_u \phi] - \bar{\xi}_u[P_u T(u,t)\phi] \quad (5.22)$$

$$- \int_0^t \bar{\xi}_u[A_u A_s T(s,t)\phi]ds = \bar{\xi}_u[A_u \phi] - \bar{\xi}_u[A_u T(u,t)\phi]. \quad (5.23)$$

Hence, using (5.21), (5.22) and (5.23) in (5.20) we have

$$\begin{aligned}
 \xi_t[\phi] - \int_0^t \xi_s[P_s T(s,t)\phi]ds &= \int_0^t \bar{\xi}_s[A_s T(s,t)\phi]ds + \int_0^t \bar{\xi}_u[P_u \phi]du \\
 &\quad - \int_0^t \bar{\xi}_u[P_u T(u,t)\phi]du + \int_0^t \bar{\xi}_u[A_u \phi]du - \int_0^t \bar{\xi}_u[A_u T(u,t)\phi]du \\
 &\quad + \gamma[\phi] + W_t[\phi],
 \end{aligned}$$

that is,

$$\begin{aligned}
 \xi_t[\phi] - \int_0^t \xi_s[P_s T(s,t)\phi]ds &= \int_0^t \bar{\xi}_u[P_u \phi]du + \int_0^t \bar{\xi}_u[A_u \phi]du \\
 &\quad + \gamma[\phi] + W_t[\phi] - \int_0^t \bar{\xi}_u[P_u T(u,t)\phi]du.
 \end{aligned}$$

Then using (5.17)

$$\bar{\xi}_t[\phi] - \int_0^t \bar{\xi}_s[P_s T(s,t)\phi]ds = \xi_t[\phi] - \int_0^t \xi_s[P_s T(s,t)\phi]ds = \eta_t[\phi],$$

i.e. $\bar{\xi}_t$ satisfies (5.3). The uniqueness now follows from the uniqueness of the solution of the SDE (5.3) using Proposition 5.1, which gives condition (*).

Finally the Gaussian property of the solution ξ follows from the fact that for each $\phi \in \Phi$ $\xi_t[\phi]$ is the a.s. limit of a sequence (the successive approximations) of gaussian random variables. Q.E.D.

Special case. Let $A = -L$ be the infinitesimal generator of a one parameter semigroup as in Lecture 3 and consider the SDE

$$d\xi_t = -L'\xi_t dt + P_t'\xi_t dt + dW_t$$

$$\xi_0 = \gamma$$

The unperturbed equation is a model used in neurophysiological applications (Example 3.1 and Lecture 4). However it is important to observe that in this field the kind of perturbation that occur are more likely to be nonlinear rather than linear.

Example 5.1. This example occurs in fluctuation theorems for interacting particle diffusions and has been considered by McKean [9], Hitsuda and Mitoma [4] and Mitoma [12].

For $n \geq 1$ let

$$\gamma^{(n)}(t) = (\gamma_1^{(n)}(t), \dots, \gamma_n^{(n)}(t))$$

be an n -particle diffusion given by the SDE

$$\begin{aligned} \gamma_k^n(t) = & \gamma_k + \frac{1}{n} \sum_{j=1}^n \int_0^t a(\gamma_k^{(n)}(s), \gamma_j^{(n)}(s)) dB_k(s) \\ & + \frac{1}{n} \sum_{j=1}^n \int_0^t b(\gamma_k^{(n)}(s), \gamma_j^{(n)}(s)) ds \quad (k = 1, \dots, n) \end{aligned}$$

where $(\gamma_k, B_k)_{k \geq 1}$ are independent copies of (γ, B) and γ is a random variable independent of the real valued Brownian motion B_t . The coefficients $a(x, y)$ and $b(x, y)$ are bounded C^∞ -functions in (x, y) .

For each $t > 0$ consider

$$U^{(n)}(t) = \frac{1}{n} \sum_{j=1}^n \delta_{\gamma_j^{(n)}(t)}$$

where δ_x is the unit mass at x . $U^{(n)}(t)$ is a measure valued process. McKean [9] has shown that

$$U^{(n)}(t) \xrightarrow{\text{a.s.}} u(dx, t)$$

where $u(dx, t)$ is the probability distribution of $X(t)$ that satisfies

$$dX(t) = \alpha(X(t), t) dB_t + \beta(X(t), t) dt$$

where

$$\alpha(x, t) := \int_{\mathbb{R}} a(x, y) u(dy, t)$$

$$\beta(x, t) := \int_{\mathbb{R}} b(x, y) u(dy, t).$$

Moreover, McKean [9] has also shown that $u(dy, t)$ has a density $u(x, t)$ and that $\alpha(x, t)$, $\beta(x, t)$ and $u(x, t)$ are C^∞ -functions in $\mathbb{R} \times \mathbb{R}_+$.

Let

$$S_n(t) = \sqrt{n} [U^{(n)}(t) - u(\cdot, t)].$$

Hitsuda and Mitoma [4] have shown that the measure valued processes $S_n(\cdot)$ converge weakly to the solution $\xi = (\xi_t)$ of the stochastic evolution equation

$$d\xi_t = A_t^1 \xi_t dt + P_t^1 \xi_t dt + dW_t \quad (5.24)$$

where for $\phi \in \Phi$

$$(A_t \phi)(x) = \frac{1}{2} \alpha(x, t)^2 \phi^{(2)}(x) + \beta(x, t) \phi^{(1)}(x) \quad (5.25)$$

$$\begin{aligned} (P_t \phi)(x) = & \int_{\mathbb{R}} b(y, x) \phi^{(1)}(y) u(y, t) dy \\ & + \int_{\mathbb{R}} \alpha(y, t) a(x, y) \phi^{(2)}(y) u(y, t) dy \end{aligned} \quad (5.26)$$

and W_t is a zero mean Φ' -valued gaussian process with independent increments, $w_0 = 0$. As pointed out in Mitoma [12], the nuclear space appropriate to the problem is given by the space Φ of real valued functions ϕ such that $\phi \in \Phi$ iff $\psi(x)\phi(x) \in \mathcal{S}$ where

$$\psi(x) = \int_{\mathbb{R}} e^{-|z|} \rho(x - z) dz$$

and ρ is the usual mollifier

$$\begin{aligned} \rho(x) = & c \cdot \exp(1/(1 - |x|^2)) & |x| < 1 \\ & 0 & |x| \geq 1. \end{aligned}$$

Notice that Φ is a modification of $\mathcal{S}$ with the following relations among the norms defining their corresponding topologies:

$$\|\phi\|_{n, \Phi} = \|\psi\phi\|_{n, \mathcal{S}} \quad (5.27)$$

$$\|\phi\|_{n, \Phi} = \|\psi\phi\|_{n, \mathcal{S}} \quad (5.28)$$

where

$$\|f\|_{n,\beta} = \sup_{0 \leq j \leq n} \sup_{x \in \mathbb{R}} |D^j f(x)| \quad n \geq 1 \quad (5.29)$$

$$\|f\|_{n,\beta}^2 = \sum_{k=0}^n \int_{\mathbb{R}} (1+x^2)^{2n} |D^k f(x)|^2 dx \quad n \geq 1. \quad (5.30)$$

It can be shown (see [12]) that under the above conditions the SDE (5.24) satisfies the assumptions of Theorem 5.1 and therefore $S_n(\cdot)$ converge weakly to the unique solution of the stochastic evolution equation (5.24).

The example just discussed is an instance where the two parameter evolution semigroup $T(s,t)$, its generator A_t and the perturbator P_t can all be defined directly on a countably Hilbertian nuclear space Φ so as to satisfy the assumptions stated at the beginning of this lecture. It is worth noting that, in many cases, these operators may be more naturally defined on a Hilbert or Banach space, as e.g., in the work of Kato and Tanabe [8, 14]. In such cases the problem of finding a Φ for which the assumptions concerning A_t and P_t are valid, has to be solved first before the results of this lecture can be applied.

2. Φ' -VALUED DIFFUSION STOCHASTIC DIFFERENTIAL EQUATIONS

A more general Φ' -valued SDE is given by

$$\begin{aligned} d\xi_t &= A(t, \xi_t)dt + B(t, \xi_t)dW_t \\ \xi_0 &= \gamma \end{aligned} \quad (5.31)$$

where the coefficient functions A, B are of the following type

$$a) \quad A: \mathbb{R}_+ \times \Phi' \rightarrow \Phi'$$

$$b) \quad B: R_+ \times \Phi' \rightarrow \mathcal{L}(\Phi', \Phi')$$

where $\mathcal{L}(\Phi', \Phi')$ denotes the class of all linear continuous operators from Φ' to Φ' , γ is a Φ' -valued random variable and W_t is a Φ' -valued Wiener process with covariance Q . In this case it is necessary to define a stochastic integral of the type

$$\int_0^t f_s(w) dW_s(w)$$

where $f_t(w) \in \mathcal{L}(\Phi', \Phi')$.

The following are conditions under which a unique solution to (5.31) exists:

For each $T > 0$ and sufficiently large $m \geq 0$, there exist numbers $r > 2$, $\theta > 0$ and $p \geq m$ such that A, B , the initial measure μ_0 for ξ_0 and the covariance functional Q satisfy the following properties

(IC) - Initial Condition

$$\int_{\Phi'} \|u\|_{-m}^r \mu_0(du) < \infty$$

(CC) Coercivity: Let j_m be the canonical isomorphism between Φ_m and Φ'_m .

For each $t \leq T$ and $u \in \Phi$

$$2A_t(j_m u)[u] + (r - 1) \|Q_{B_t^*(u)}\|_{-m, -m} \leq \theta(1 + \|j_m(u)\|_{-m}^2)$$

$\|\dots\|_{-m, -m}$ is the trace norm of the nuclear operator determined by the bilinear form $Q_{B_t^*(u)}$ where

$$Q_{B_t^*(u)}(\phi, \psi) = Q(B_t^*(u)\phi, B_t^*(u)\psi) \quad \phi, \psi \in \Phi.$$

and, $B_t^*(u) \in \mathcal{L}(\Phi, \Phi)$ is the adjoint operator defined by the relation.

$B_t(u) f[\phi] = f[B_t^*(u)\phi]$ for all $f \in \Phi'$ and all $\phi \in \Phi$

(LG) Linear Growth for each $t \leq T$ and $u \in \Phi'_m$

$$A_t(u) \in \Phi'_p$$

and

$$\|A_t(u)\|_{-p}^2 \leq \theta(1 + \|u\|_{-m}^2)$$

$$\|Q_{B_t^*(u)}\|_{-m, -m}^2 \leq \frac{1}{2} \theta^{2/r} (1 + \|u\|_{-m}^2)$$

(MC) Monotonicity. For each $t \leq T$ and $u, v \in \Phi'_m$

$$A_t(u), A_t(v) \in \Phi'_p$$

and

$$\begin{aligned} & 2(u - v, A_t(u) - A_t(v))_{-p} + (r - 1) \|Q_{B_t^*(u) - B_t^*(v)}\|_{-p, -p} \\ & \leq \theta \|u - v\|_{-p}^2 \end{aligned}$$

(JC) Joint Continuity

$$A: [0, T] \times \Phi' \rightarrow \Phi'$$

and

$$B: [0, T] \times \Phi' \rightarrow \mathcal{L}(\Phi', \Phi')$$

are each jointly continuous.

The proof of the existence of the solution will appear in a forthcoming paper by G. Kallianpur and R. Wolpert.

- [1] Christensen, S.K. (1985), "Linear stochastic differential equations on the dual of a countably Hilbert Nuclear space with applications of Neurophysiology", Tech. Rept. 104 Center for Stochastic Processes, The University of North Carolina at Chapel Hill.
- [2] Christensen, S.K. and G. Kallianpur (1985), "Stochastic differential equations for neuronal behavior", Tech. Rept. 103 Center for Stochastic Processes, The University of North Carolina at Chapel Hill.
- [3] Dellacherie, C. and P. Meyer (1982), "Probabilities and Potential B", North Holland, Amsterdam.
- [4] Hitsuda, H. and I. Mitoma, "Tightness problem and stochastic evolution equations arising from fluctuation phenomena for interacting diffusions", Preprint.
- [5] Itô, K. (1978), "Stochastic analysis in infinite dimensions", In. Friedman, A. Pinsky, M. (eds.), Stochastic Analysis, Academic Press, New York.
- [6] Itô, K. (1983), "Distribution-valued processes arising from independent Brownian motions", Math. Z. 182, 17-33.
- [7] Kallianpur, G. and R. Wolpert (1984), "Infinite dimensional stochastic differential equation models for spatially distributed neurons", Appl. Math. Optim. 12, 125-172.
- [8] Kato, T. (1970), "Linear evolution equations of hyperbolic type", J. Fac. Sci. Univ. Tokyo, Sec. I, 17, 241-258.
- [9] McKean, H.P. (1967), "Propagation of Chaos for a class of nonlinear parabolic equations", in Lecture Series in Differential Equations 2, pp. 177-193, Van Nostrand Math. Studies 19, New York.
- [10] Mitoma, I. (1981), "Martingales of random distributions", Memoirs Fac. Sci. Kyushu Univ. Ser. A 35, 1, 185-220.
- [11] Mitoma, I. (1983), "Tightness of probabilities on $C([0,1]; \cdot)$ and $D([0,1]; \cdot)$ ", Ann. Probab. 11, 989-999.
- [12] Mitoma, I. (1985), "An ∞ -Dimensional inhomogeneous Langevin equation", J. Funct. Anal. 61, 3, 342-359.
- [13] Perez-Abreu, V. (1985), "Product stochastic measures, multiple stochastic integrals and their extensions to nuclear space valued Wiener processes", Tech. Rept. 107 Center for Stochastic Processes, The University of North Carolina at Chapel Hill.
- [14] Tanabe, H. (1975), "Equations of Evolution", Monographs and Studies in Mathematics 6, Pitman, London.
- [15] Walsh, J.B. (1981), "A stochastic model of neuronal response", Adv. Appl. Prob. 13, 231-281.
- [16] Wan, F.Y.M. and H.C. Tuckwell (1979), "The response of a spatially distributed neuron to white noise current injection", Biol. Cybernetics 33, 39-55.