**ON THE ANALYSIS OF BRINE TRANSPORT
IN POROUS MEDIA**

By

C.J. van Duijn

L.A. Peletier

and

R.J. Schotting

IMA Preprint Series # 826

June 1991

On the Analysis of Brine Transport in Porous Media

C.J. van Duijn, L.A. Peletier & R.J. Schotting

1. Introduction

In this paper we analyse a model for the transport of brine through a porous medium. Brine is water which is strongly impregnated with salt, so much so that the volume of the fluid is affected. Therefore the corresponding fluid flow can no longer be considered as divergence free. This, in contrast to flows with low concentrations of salt, such as sea water, where one normally assumes that the fluid is incompressible.

We shall focuss on the aspect that the flow is not divergence free and we shall investigate how this influences the transport of the salt by the moving fluid. We do this for a two-dimensional flow through a homogeneous and isotropic porous medium. The fluid is assumed to have constant dynamic viscosity μ and a variable density ρ , and the porous medium is characterised by a constant porosity $\Phi \in (0, 1)$ and a constant permeability κ . We shall follow Hassanizadeh & Leijnse [HL] in their formulation of the flow of brine fluids. However, we deviate from their description in the sense that we shall use the classical form of Darcy's law for the momentum balance equation for the fluid and Fick's law for the dispersion of the salt. Moreover, in the equation of state, we shall ignore the influence of pressure variations. The equations describing the brine flow then result from the following basic laws:

mass balance of the fluid

$$\Phi \frac{\partial \rho}{\partial t} + \operatorname{div}(\rho \mathbf{q}) = 0 \quad (1.1)$$

mass balance of the salt

$$\Phi \frac{\partial \rho \omega}{\partial t} + \operatorname{div}(\rho \omega \mathbf{q} - D \rho \operatorname{grad} \omega) = 0, \quad (1.2)$$

Darcy's law

$$\frac{\mu}{\kappa} \mathbf{q} + \operatorname{grad} p - \rho \mathbf{g} = 0, \quad (1.3)$$

equation of state

$$\rho = \rho_f e^{\gamma \omega}. \quad (1.4)$$

In these equations $\mathbf{q} = (q_y, q_z)$ denotes the specific discharge of the fluid, p the fluid pressure and $\mathbf{g} = (g_y, g_z)$ the acceleration of gravity. In equation (1.2) and throughout the analysis we consider

$$D = \Phi D_{\text{mol}},$$

where D_{mol} denotes the molecular diffusion, which we assume to be constant. However, in the concluding section we shall also discuss the more realistic situation where D depends on the velocity (see for instance [B] or [HL]). In the equation of state, ρ_f denotes the mass

density of the fresh water, $\gamma = 0.6923$ is a constant and ω is the mass fraction of the salt in the brine, (i.e. $\omega = \text{mass concentration of the salt component of the brine} / \text{mass density } \rho \text{ of the brine}$).

We shall study two specific flow problems, which we shall refer to as Problems I and II.

Fig. 1. Initial salt distributions

In Problem I - see Figure 1(a) - the flow domain is unbounded above and below and initially, say at $t = 0$, the region above the plane $\{z = 0\}$ is filled with fresh water and the brine, with mass density $\rho_s > \rho_f$ fills the region below it. In Problem II - see Figure 1(b) - the flow domain consists of the upper half space $\{z > 0\}$ and is bounded below by an impermeable salt rock. Again, at $t = 0$, fresh water occupies the region $\{z > 0\}$ while the salt from the rock ensures that $\rho = \rho_s$ (mass density of saturated brine) along the boundary $\{z = 0\}$.

Since in both problems the y -coordinate ranges from $-\infty$ to $+\infty$ we may look for a density ρ and specific discharge q of the form

$$\rho = \rho(z, t) \quad \text{and} \quad q = q(z, t). \quad (1.5)$$

Under this assumption we obtain a linear relation between the y -component of the specific discharge and the fluid density (see also [dJdJvD]).

To derive the governing equations we first eliminate the pressure from Darcy's law by taking the curl. This results in

$$\frac{\partial}{\partial z} \left\{ q_y - \frac{\kappa}{\mu} \rho g_y \right\} - \frac{\partial}{\partial y} \left\{ q_z - \frac{\kappa}{\mu} \rho g_z \right\} = 0.$$

With (1.5) this gives

$$q_y + \frac{\kappa}{\mu} \rho g \sin \beta = \text{constant}, \quad (1.6)$$

where β is the inclination of the original fresh-brine interface in Problem I and the salt boundary in Problem II, and $g = |g|$. Next we impose the boundary conditions from

Figure 1 to determine the constant in (1.6). Since

$$q_y(z, t) \rightarrow q_f \quad \text{as } z \rightarrow \infty \quad \text{for all } t > 0, \quad (1.7)$$

in which $q_f \in \mathbf{R}$ is given, and

$$\rho(z, t) \rightarrow \rho_f \quad \text{as } z \rightarrow \infty \quad \text{for all } t > 0, \quad (1.8)$$

we obtain the relation

$$q_y = q_f - \frac{\kappa}{\mu}(\rho - \rho_f)g \sin \beta. \quad (1.9)$$

Using (1.5) in the equations (1.1) and (1.2) we obtain

$$\Phi \frac{\partial \rho}{\partial t} + \frac{\partial}{\partial z}(\rho q_z) = 0 \quad (1.10)$$

and

$$\Phi \frac{\partial \rho \omega}{\partial t} + \frac{\partial}{\partial z}(\rho \omega q_z - D \rho \frac{\partial \omega}{\partial z}) = 0.$$

Combining these equations yields

$$\Phi \rho \frac{\partial \omega}{\partial t} + \rho q_z \frac{\partial \omega}{\partial z} = \frac{\partial}{\partial z} \left(\rho D \frac{\partial \omega}{\partial z} \right)$$

and with (1.4)

$$\Phi \frac{\partial \rho}{\partial t} + q_z \frac{\partial \rho}{\partial z} = D \frac{\partial^2 \rho}{\partial z^2}. \quad (1.11)$$

In Problem I, equations (1.10) and (1.11) have to be solved for $(z, t) \in \mathbf{R} \times \mathbf{R}^+$ subject to the initial conditions

$$\rho(z, 0) = \begin{cases} \rho_f & z > 0 \\ \rho_s & z < 0. \end{cases} \quad (1.12a)$$

At one of the boundaries $z = -\infty$ or $z = +\infty$ we have to specify q_z . Here we choose

$$q_z(z, t) \rightarrow 0 \quad \text{as } z \rightarrow \infty \quad \text{for all } t > 0. \quad (1.12b)$$

In Problem II we solve the differential equations in the quarter plane $\mathbf{R}^+ \times \mathbf{R}^+$ subject to the initial condition

$$\rho(z, 0) = \rho_f, \quad z > 0 \quad (1.13a)$$

and the boundary condition

$$\rho(0, t) = \rho_s, \quad t > 0. \quad (1.13b)$$

In addition we now require (see [HL])

$$q_z(0, t) = -\frac{D}{\gamma \rho_s (1 - \omega_s)} \frac{\partial \rho}{\partial z}(0, t) \quad \text{for all } t > 0. \quad (1.13c)$$

Here $\rho_s = \rho_f \exp(\gamma \omega_s)$, where ω_s is the salt mass fraction of saturated brine.

At this point it is important to note that the differential equations (1.10) and (1.11) only involve as unknowns the velocity component q_z and the density ρ . Thus solving these equations subject to the appropriate boundary conditions and using ρ in (1.9) determines a solution $(\mathbf{q}, \rho)$ of the original problem.

To simplify the equations we introduce dimensionless variables, and set

$$\begin{aligned} \rho^* &= \frac{\rho - \rho_f}{\rho_s - \rho_f} & q_z^* &= \frac{q_z}{\frac{k}{\mu} \rho_f g} \\ z^* &= \frac{\frac{k}{\mu} \rho_f g}{D} z, & t^* &= \frac{(\frac{k}{\mu} \rho_f g)^2}{\Phi D} t \end{aligned} \quad (1.14)$$

and

$$\varepsilon = \frac{\rho_s - \rho_f}{\rho_f}. \quad (1.15)$$

In these new variables, (1.10) and (1.11) become (when we omit the asterisks again)

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial z}(\rho q_z) + \frac{1}{\varepsilon} \frac{\partial q_z}{\partial z} = 0 \quad \text{in } \Omega \times \mathbf{R}^+ \quad (1.16)$$

$$\frac{\partial \rho}{\partial t} + q_z \frac{\partial \rho}{\partial z} = \frac{\partial^2 \rho}{\partial z^2} \quad \text{in } \Omega \times \mathbf{R}^+, \quad (1.17)$$

where Ω is either $\mathbf{R}$ or $\mathbf{R}^+$.

In Problem I, the rescaled initial and boundary conditions become

$$\rho(z, 0) = \begin{cases} 0 & \text{if } z > 0 \\ 1 & \text{if } z < 0 \end{cases} \quad (1.18a)$$

and

$$q_z(z, t) \rightarrow 0 \quad \text{as } z \rightarrow \infty \quad \text{for all } t > 0. \quad (1.18b)$$

In Problem II we have

$$\rho(z, 0) = 0 \quad \text{for } z > 0 \quad (1.19a)$$

$$\rho(0, t) = 1 \quad \text{for } t > 0 \quad (1.19b)$$

and

$$q_z(0, t) = -K \varepsilon \frac{\partial \rho}{\partial z}(0, t) \quad \text{for } t > 0 \quad (1.19c)$$

in which

$$K = \frac{\rho_f}{\rho_s} \frac{1}{\gamma(1 - \omega_s)}.$$

Through ρ_s and ω_s , K is a function of ε ; it is given by

$$K(\varepsilon) = \frac{1}{(1 + \varepsilon)\{\gamma - \log(1 + \varepsilon)\}}, \quad 0 < \varepsilon < e^\gamma - 1. \quad (1.19d)$$

It is readily verified that K is strictly increasing, and that $K(\varepsilon) > 1/\gamma$ if $\varepsilon > 0$.

Remark. We note that in the formal limit as $\varepsilon \rightarrow 0$, equation (1.16) reduces to $\partial q_z / \partial z = 0$. Together with condition (1.18b), or with (1.19c), this implies that in this limit $q_z = 0$. For ρ then remains the linear heat equation. In the literature this is known as the Boussinesq approximation, in which the flow satisfies $\text{div} \mathbf{q} = 0$. For Problem I the Boussinesq approximation was studied in [dJdJvD], in particular when the dispersion depends on the velocity.

We solve Problems I and II by looking for self-similar solutions. We set

$$x = z/\sqrt{t} \quad (1.20)$$

and

$$\rho(z, t) = u(x) \quad \text{and} \quad q_z(z, t) = \frac{1}{\sqrt{t}} v(x). \quad (1.21)$$

This results in the differential equations

$$(uv)' + \frac{1}{\varepsilon} v' - \frac{1}{2} x u' = 0 \quad \text{in } \Omega \quad (1.22)$$

$$u'v - \frac{1}{2} x u' = u'' \quad \text{in } \Omega. \quad (1.23)$$

In Problem I the boundary conditions are

$$u(-\infty) = 1 \quad \text{and} \quad u(+\infty) = 0 \quad (1.24a)$$

and

$$v(\infty) = 0, \quad (1.24b)$$

and in Problem II

$$u(0) = 1 \quad \text{and} \quad u(+\infty) = 0 \quad (1.25a)$$

and

$$v(0) = -K\varepsilon u'(0). \quad (1.25b)$$

Throughout we shall restrict the analysis to solutions (u, v) such that $u > 0$ in Ω .

We analyze Problems I and II using ideas from a previous study of a fresh - salt ground water problem [vDP3], which also involved a third order differential equation.

In Sections 2 and 3 we study the self similar solutions related to Problem I. In Section 2 we prove that for every $\varepsilon > 0$, there exists a unique pair of smooth functions (u, v) which solves (1.22) - (1.24) (see Figures 2 and 3). The function u is strictly decreasing on $\mathbf{R}$ and has precisely one inflection point at some $x_0 > 0$, such that $u'' < 0$ on $(-\infty, x_0)$ and $u'' > 0$ on (x_0, ∞) . The function v is positive on $\mathbf{R}$, strictly increasing on $(-\infty, x_0)$ and strictly decreasing on (x_0, ∞) . Moreover v satisfies

$$v(-\infty) = \varepsilon^2 \int_0^1 \frac{y(u)}{(1 + \varepsilon u)^2} du, \quad (1.26)$$

where y is the solution of the boundary value problem (2.8), (2.9).

In Section 3 we investigate the Boussinesq limit as $\varepsilon \rightarrow 0$. Denoting for $\varepsilon > 0$ the solutions of (1.22)-(1.24) by $(u_\varepsilon, v_\varepsilon)$ we show that

$$v_\varepsilon(x) = -u'_0(x)\varepsilon + \{E_1(x) + o(1)\}\varepsilon^2 \quad (1.27a)$$

and

$$u_\varepsilon(x) = u_0(x) + \frac{1}{2}u_0(x)\{1 - u_0(x)\}\varepsilon + \{E_2(x) + o(1)\}\varepsilon^2 \quad (1.27b)$$

as $\varepsilon \rightarrow 0$, uniformly on $\mathbf{R}$.

Here u_0 is the Boussinesq limit

$$u_0(x) = \frac{1}{2} \operatorname{erfc}\left(\frac{x}{2}\right), \quad x \in \mathbf{R}, \quad (1.28)$$

which can be obtained formally by setting $v = 0$ in (1.23). The functions E_1 and E_2 can be explicitly calculated. This is outlined in Lemmas 3.6 and 3.7.

In Section 4 we turn to Problem II where we discuss for $0 < \varepsilon < e^\gamma - 1$ the existence and uniqueness of solutions (u, v) of (1.22), (1.23) and (1.25) (see also Figures 5 and 6). The upper bound for ε is a consequence of formula (1.19d) and is for all practical purposes no restriction. Brines with high salt concentrations have ε in the order of 0.2 (see [HL]). For salt water in coastal aquifers one often uses $\varepsilon = 0.025$ (see [B]).

Here we show that u_0 is strictly decreasing on $\mathbf{R}^+$ and has precisely one inflection point at some $x_0 > 0$: $u'' < 0$ on $(0, x_0)$ and $u'' > 0$ on (x_0, ∞) . The function v satisfies $v' > 0$ on $(0, x_0)$ and $v' < 0$ on (x_0, ∞) . Notice that

$$v(0) > 0 \quad \text{and} \quad v(\infty) = O(\varepsilon) \quad \text{as } \varepsilon \rightarrow 0 \quad (1.29a)$$

whereas we deduce from (1.26) that in Problem I

$$v(-\infty) = O(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0. \quad (1.29b)$$

In Section 5 we investigate again the limit $\varepsilon \rightarrow 0$. From the results obtained there we find

$$v_\varepsilon(x) = -u'_0(x)\varepsilon + O(\varepsilon^2) \quad (1.30a)$$

and

$$\begin{aligned} u_\varepsilon(x) = & u_0(x) + \left[-\frac{2(1-\gamma)}{\pi\gamma} \{u_0(x) + \sqrt{\pi}u'_0(x)\} \right. \\ & \left. + \frac{1}{2}u_0(x)\{1 - u_0(x)\} \right] \varepsilon + O(\varepsilon^2), \end{aligned} \quad (1.30b)$$

as $\varepsilon \rightarrow 0$, uniformly on $\mathbf{R}$. Here u_0 is the Boussinesq limit

$$u_0(x) = \operatorname{erfc}\left(\frac{x}{2}\right), \quad x \in \mathbf{R}^+. \quad (1.31)$$

In Problem I we have $v(-\infty) > 0$ for any $\varepsilon > 0$ and in Problem II $v(+\infty) > 0$ for ε sufficiently small. This implies for the velocity $\bar{q}(z, t)$ that

$$\lim_{t \downarrow 0} q_z(z, t) = \infty \quad \text{for each } z < 0 \text{ in Problem I}$$

and

$$\lim_{t \downarrow 0} q_z(z, t) = \infty \quad \text{for each } z > 0 \text{ in Problem II.}$$

These unbounded velocities at $t = 0^+$ are caused by the discontinuity in the initial density profile at $z = 0$ which gives rise to an unbounded density gradient. This problem will not occur when the profile is smooth: for instance, if we apply for any $\delta > 0$ the time shift $t \mapsto t + \delta$ and consider as initial distribution

$$\rho(z, 0) = \rho_f + (\rho_s - \rho_f)u(z/\sqrt{\delta}),$$

then the velocity will be bounded down to $t = 0$.

In Section 6 we present some numerical results. We computed solutions (u, v) of Problems I and II for the cases $\varepsilon = 0.025$ and $\varepsilon = 0.5$. The case $\varepsilon = 0.025$ corresponds to a relatively small density difference as encountered in coastal aquifers, where fresh and salt water are present. The second value $\varepsilon = 0.5$ is unrealistically high, but it is chosen here to show up in the effect of volume changes caused by high salt concentrations on the distribution of brine.

Finally, in the concluding Section 7, we make some observations about brine transport when the dispersion depends on velocity. We present a von Mises-type transformation which reduces equations (1.10) and (1.11) to a single nonlinear diffusion equation involving ρ only. This allows us to study more general initial value problems and leads to the conjecture that the self-similar solutions considered in this paper are the large time asymptotic profiles of solutions emanating from more general initial distributions. We leave the proof of this conjecture to a future paper.

2. Self-similar solutions on $\mathbb{R}$

In this section we shall study the self-similar solutions (u, v) of equations (1.22) and (1.23) which satisfy the required conditions at $x = \pm\infty$. Here we put the equations into the form

$$\begin{cases} u'' + (\frac{1}{2}x - v)u' = 0 \\ v' = -\frac{u''}{u + \frac{1}{\varepsilon}}, \quad u > 0 \end{cases} \quad -\infty < x < \infty \quad (2.1)$$

$$(2.2)$$

We recall that the boundary conditions are

$$u(-\infty) = 1 \quad \text{and} \quad u(+\infty) = 0 \quad (2.3)$$

and

$$v(+\infty) = 0. \quad (2.4)$$

As a preliminary observation we note that u and v' are invariant under shifts. Specifically, set

$$\bar{x} = x - a, \quad \bar{u}(\bar{x}) = u(x), \quad \bar{v}(\bar{x}) = v(x) - \frac{1}{2}a, \quad (2.5)$$

where a is some arbitrary number. Hence, if $(u(x), v(x))$ is a solution of (2.1) - (2.3), then so is $(\bar{u}(\bar{x}), \bar{v}(\bar{x}))$ but with x in (2.1) replaced by $\bar{x}$. Thus, if $(u(x), v(x))$ is a solution and $v(\pm\infty)$ exists, then by shifting x over a suitable distance a , we can reach any prescribed limiting value of $v(x)$, either at $x = -\infty$ or at $x = +\infty$. Thus it follows from (2.5) that by choosing $a = 2v(\infty)$, we can ensure that the boundary condition (2.4) is satisfied.

We begin by establishing a few qualitative properties of solutions of (2.1) - (2.4).

Lemma 2.1. *Let (u, v) be a solution of (2.1) - (2.4). Then*

$$(a) \quad u'(x) < 0 \quad \text{for all } x \in \mathbf{R}.$$

(b) *There exists a number $x_0 > 0$ such that*

$$\begin{aligned} u''(x) &> 0 \quad (<) \quad \text{if } x > x_0 \quad (<) \\ v'(x) &< 0 \quad (>) \quad \text{if } x > x_0 \quad (<). \end{aligned}$$

$$(c) \quad u'(x) \rightarrow 0 \quad \text{as } x \rightarrow \pm\infty.$$

Proof. (a) Suppose to the contrary that there exists a point $\xi \in \mathbf{R}$ such that $u'(\xi) = 0$. Then, by standard uniqueness theory applied to equation (2.1), $u'(x) = 0$ for all $x \in \mathbf{R}$ so that the conditions at $x = \pm\infty$ cannot both be satisfied.

(b) In view of the boundary conditions, u'' must have at least one zero. Let x_0 be such a zero. Then by the equations (2.1) and (2.2),

$$v(x_0) = \frac{1}{2}x_0 \quad \text{and} \quad v'(x_0) = 0. \quad (2.6)$$

Differentiating (2.1) we obtain

$$u'''(x_0) = -\left\{\frac{1}{2} - v'(x_0)\right\}u'(x_0) - \left\{\frac{1}{2} - v(x_0)\right\}u''(x_0)$$

and so

$$u'''(x_0) = -\frac{1}{2}u'(x_0) > 0.$$

Thus, at every zero, u'' passes from negative to positive, which means that there can be only one zero. Since $v(\infty) = 0$ and $v'(x) < 0$ for $x > x_0$, we conclude that $v(x_0) > 0$ which implies by (2.1) that $x_0 > 0$.

(c) Because $u'(x) < 0$ and $u''(x) > 0$ for x large enough, $\ell = \lim_{x \rightarrow \infty} u'(x)$ exists. Since $u(x) \rightarrow 0$, ℓ can only be zero. That $u'(x) \rightarrow 0$ as $x \rightarrow -\infty$ is proved the same way.

Remark. It is interesting to note that due to property (2.6), it is possible to shift the pair (u, v) by such an amount that we obtain

$$\bar{v}(0) = 0 \quad \text{as well as} \quad \bar{v}(\bar{x}) < 0 \quad \text{if} \quad \bar{x} \neq 0.$$

For the velocity field $\mathbf{q}$ this means that $q_z(0, t) = 0$ for all $t > 0$ so that there is no flux across the plane $\{z = 0\}$.

We now turn to proving existence and uniqueness of a solution (u, v) of (2.1) - (2.4). We proceed in two steps. First we replace the condition (2.4) on v at infinity by a third condition on u at the origin:

$$u(0) = \frac{1}{2} \tag{2.7}$$

and we establish existence and uniqueness for this modified problem (2.1) - (2.3), (2.7). We then show that the limit $v(\infty)$ exists. By means of the transformation (2.5) we finally obtain the desired solution of the original problem. Because the shift $a = 2v(\infty)$ is uniquely determined by the (unique) solution of the modified problem, the resulting solution is unique.

We transform the system (2.1), (2.2) to a single well known second order differential equation by introducing u as an independent variable. Since any solution u is strictly decreasing by Lemma 2.1, this is possible. Thus, we define

$$x = x(u) \quad \text{and} \quad y(u) = -u'(x(u)). \tag{2.8}$$

Then we find after an elementary computation that $y(u)$ is a solution of the equation

$$y\{(1 + \varepsilon u)y'\}' = -\frac{1}{2}(1 + \varepsilon u), \quad 0 < u < 1, \tag{2.9}$$

whilst the boundary conditions (2.3) become by Lemma 2.1

$$y(0) = 0 \quad \text{and} \quad y(1) = 0. \tag{2.10}$$

To bring (2.9) really into standard form we carry out one more transformation, setting

$$t = \frac{\log(1 + \varepsilon u)}{\log(1 + \varepsilon)} \quad \text{and} \quad z(t) = \frac{\varepsilon}{\log(1 + \varepsilon)} y(u). \tag{2.11}$$

This yields the problem

$$(P_1) \begin{cases} -zz'' = \frac{1}{2}e^{2t \log(1+\varepsilon)}, & z > 0 \quad \text{for} \quad 0 < t < 1 \\ z(0) = 0, & z(1) = 0. \end{cases}$$

Problem (P₁) has been studied in detail in [AP, vDGZ, vDP1, T]. It has a unique solution $z \in C^\infty(0, 1) \cap C([0, 1])$ which is concave and satisfies $z'(0^+) = \infty$ and $z'(1^-) = -\infty$. Plainly, this implies the existence, uniqueness and positivity of the solution $y(u)$ of (2.9), (2.10).

We return to the original variables x , u and v by means of

$$x(u) = \int_u^{\frac{1}{2}} \frac{ds}{y(s)}, \quad (2.12)$$

which follows from (2.8), and

$$v(x) = \frac{1}{2}x - y'(u(x)), \quad (2.13)$$

which follows from (2.1) and (2.8). The boundary conditions (2.3) follow from the fact that $1/y$ is not integrable near $u = 0$ and $u = 1$.

Lemma 2.2. *Let (u, v) be a solution of (2.1) - (2.3), (2.7). Then*

$$(a) \quad v(\pm\infty) = \lim_{x \rightarrow \pm\infty} v(x) \text{ exists}$$

and $v(-\infty)$ as well as $v(+\infty)$ are finite.

$$(b) \quad v(-\infty) > v(+\infty),$$

$$(c) \quad v(x) > v(+\infty) \text{ for all } x \in \mathbf{R}.$$

Proof. (a) That the limits exist follows at once from the monotonicity of v for large positive and large negative values of x . Thus, it remains to show that both limits are finite.

If we integrate (2.2) over $(0, x)$ we obtain

$$\begin{aligned} v(x) &= v(0) - \varepsilon \int_0^x \frac{u''(s)}{1 + \varepsilon u(s)} ds \\ &= v(0) - \varepsilon \frac{u'(s)}{1 + \varepsilon u(s)} \Big|_0^x - \varepsilon^2 \int_0^x \frac{u'^2(s)}{\{1 + \varepsilon u(s)\}^2} ds \\ &= v(0) + \varepsilon \frac{y(u(x))}{1 + \varepsilon u(x)} - \varepsilon \frac{y(\frac{1}{2})}{1 + \frac{\varepsilon}{2}} - \varepsilon^2 \int_{u(x)}^{\frac{1}{2}} \frac{y(u)}{(1 + \varepsilon u)^2} du. \end{aligned} \quad (2.14)$$

Thus, letting $x \rightarrow \infty$ we obtain

$$v(\infty) = v(0) - \varepsilon \frac{y(\frac{1}{2})}{1 + \frac{\varepsilon}{2}} + \varepsilon^2 \int_0^{\frac{1}{2}} \frac{y(u)}{(1 + \varepsilon u)^2} du. \quad (2.15)$$

That $v(-\infty)$ is finite can be proved in a similar way.

(b) and (c) We integrate (2.2) over (x, ∞) . This yields, after some manipulations similar to those carried out to derive (2.14),

$$v(x) = v(\infty) + \varepsilon \frac{y(u(x))}{1 + \varepsilon u(x)} + \varepsilon^2 \int_0^{u(x)} \frac{y(u)}{(1 + \varepsilon u)^2} du, \quad (2.16)$$

from which the assertions follow at once.

Remark. If we let $x \rightarrow -\infty$ we obtain a relation between the limits of v :

$$v(-\infty) - v(\infty) = \varepsilon^2 \int_0^1 \frac{y(u)}{(1 + \varepsilon u)^2} du, \quad (2.17)$$

Having proved that $v(\infty)$ exists and is finite, we can complete the proof of the existence and uniqueness of a solution of (2.1) - (2.4) by an appropriate shift, as indicated at the beginning of this section.

Theorem 2.3. *For each $\varepsilon > 0$, there exists one and only one pair of functions $(u(x), v(x))$ which satisfies (2.1) - (2.4). The function v is related to u through the relation*

$$v(x) = -\varepsilon \frac{u'(x)}{1 + \varepsilon u(x)} + \varepsilon^2 \int_x^\infty \frac{u'^2(s)}{\{1 + \varepsilon u(s)\}^2} ds. \quad (2.18)$$

We conclude this section with asymptotic estimates for $u(x)$ and $v(x)$ as $x \rightarrow -\infty$ and $x \rightarrow +\infty$.

Lemma 2.4. *Let $(u(x), v(x))$ be the solution of (2.1) - (2.4). Then there exist positive constants A_+ and A_- such that*

$$\begin{aligned} u(x) &< A_+ x^{-1} e^{-x^2/4} \\ v(x) &< A_+ \varepsilon x^{-1} e^{-x^2/4} \end{aligned} \quad \text{if } x > 0$$

and

$$\begin{aligned} 1 - u(x) &< A_- x^{-1} e^{-(x-a)^2/4} \\ v(x) - v(-\infty) &< A_- \varepsilon x^{-1} e^{-(x-a)^2/4} \end{aligned} \quad \text{if } x < 0,$$

where $a = 2v(-\infty)$.

Proof. For convenience we write

$$V(x) = \int_0^x v(s) ds.$$

Then equation (2.1) can be integrated once to yield

$$u'(x) = u'(0) \exp\left(-\frac{1}{4}x^2 + V(x)\right) \quad \text{for } x \in \mathbf{R}. \quad (2.19)$$

By Lemma 2.2 there exists a constant K such that

$$v(x) < K \quad \text{for } x \in \mathbf{R}. \quad (2.20)$$

Hence $V(x) < Kx$ for $x > 0$ and

$$|u'(x)| \leq |u'(0)| e^{K^2} e^{-(x-2K)^2/4} \quad \text{for } x > 0. \quad (2.21)$$

Using this bound in (2.18) we conclude that v is integrable on $(0, \infty)$ so that $V(\infty) < \infty$ and

$$|u'(x)| \leq |u'(0)| e^{V(\infty)} e^{-x^2/4} \quad \text{for } x > 0. \quad (2.22)$$

The estimate for $u(x)$ now follows upon integration; the one for $v(x)$ follows when we use (2.22) in (2.18).

Next we let $x \rightarrow -\infty$. Because $v(x) \rightarrow v(-\infty) > 0$ if $\varepsilon > 0$

$$V(x) = v(-\infty)x - \int_x^0 \{v(s) - v(-\infty)\} ds.$$

As before we find that the integral converges as $x \rightarrow -\infty$, so that

$$|u'(x)| \leq |u'(0)| e^{a^2/4} e^{-(x-a)^2/4} \quad \text{for } x < 0, \quad (2.23)$$

where $a = 2v(-\infty)$. The estimates for $x \rightarrow -\infty$ now follow as they did for $x \rightarrow +\infty$.

3. The limit $\varepsilon \rightarrow 0$

In this section we shall study the dependence of the solution of (2.1) - (2.4) on ε for small values of ε . Thus, to emphasize the dependence on ε , we shall denote the solution by $(u_\varepsilon(x), v_\varepsilon(x))$.

Lemma 3.1. *There exist positive constants C_1 and C_2 , which do not depend on ε such that*

$$|u'_\varepsilon(x)| < C_1 \quad \text{and} \quad 0 < v_\varepsilon(x) < C_2\varepsilon$$

for all $\varepsilon \in (0, 1)$.

Proof. Observe that

$$e^{2t \log(1+\varepsilon)} \leq 4 \quad \text{for } 0 \leq t \leq 1, \quad 0 < \varepsilon < 1.$$

Hence, by the monotone dependence of z_ε on the right hand side of its differential equation [vDP], it follows that

$$z_\varepsilon(t) < \bar{z}(t) \quad \text{for } 0 \leq t \leq 1,$$

where $\bar{z}$ is the positive solution of Problem (P) with right hand side equal to 2. Therefore z_ε , and thereby also y_ε , are uniformly bounded for $0 < \varepsilon < 1$. This means that $|u'_\varepsilon(x)|$ is uniformly bounded on $\mathbf{R}$.

The bound for v_ε follows at once from (2.16).

Corollary 3.2. (a) *The constants A_+ and A_- in Lemma 2.4 can be chosen independent of ε .*

(b) *We have*

$$u'_\varepsilon(x) \rightarrow 0 \quad \text{and} \quad u''_\varepsilon(x) \rightarrow 0 \quad \text{as } |x| \rightarrow \infty,$$

uniformly with respect to $\varepsilon \in (0, 1)$.

Proof. (a) By Lemma 3.1, the constant K in (2.20) can be chosen independent of ε if $\varepsilon < 1$. This implies, putting (2.21) into (2.18), that $V(\infty)$ can also be estimated independent of ε . Thus, using Lemma 3.1 again, the constant in (2.22) does not depend on ε and so A_+ does not depend on ε .

That A_- is independent of ε follows at once from (2.23) and Lemma 3.1.

Part (b) follows immediately from (2.19) and Lemma 3.1 as well as the differential equation (2.1).

Our first estimate states that u_ε converges to the limit profile u_0 which solves (2.1), (2.3) when we set $v = 0$. Thus

$$u_0(x) = \frac{1}{2} \operatorname{erfc}\left(\frac{x}{2}\right) = \frac{1}{\sqrt{\pi}} \int_{x/2}^{\infty} e^{-t^2} dt. \quad (3.1)$$

Lemma 3.3. *We have*

$$u_\varepsilon(x) \rightarrow u_0(x), \quad u'_\varepsilon(x) \rightarrow u'_0(x), \quad u''_\varepsilon(x) \rightarrow u''_0(x) \quad \text{as } \varepsilon \rightarrow 0$$

uniformly on $\mathbf{R}$.

Proof. By Lemma 3.1, u'_ε and v_ε are uniformly bounded. Hence by equations (2.1) and (2.2), u''_ε and v'_ε are uniformly bounded on bounded sets. Finally, by differentiating (2.1) we find that u'''_ε is also bounded on bounded sets. Therefore by Arzela-Ascoli's theorem there exists a sequence $\{\varepsilon_n\}$, $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$, and a function U such that

$$u_{\varepsilon_n} \rightarrow U, \quad u'_{\varepsilon_n} \rightarrow U', \quad u''_{\varepsilon_n} \rightarrow U'' \quad \text{as } \varepsilon_n \rightarrow 0$$

along the subsequence $\{\varepsilon_n\}$, uniformly on bounded intervals. Taking the limit in (2.1) and remembering that $v_{\varepsilon_n} = O(\varepsilon_n)$ as $\varepsilon_n \rightarrow 0$, we find that

$$U'' + \frac{1}{2}xU' = 0 \quad \text{for } x \in (-\infty, \infty).$$

By the asymptotic estimates of Lemma 2.4, which are uniform with respect to ε by Corollary 3.2(b), we conclude that

$$U(-\infty) = 1 \quad \text{and} \quad U(+\infty) = 0.$$

Therefore $U = u_0$ and the entire family $\{u_\varepsilon\}$ converges to u_0 as $\varepsilon \rightarrow 0$ uniformly on $\mathbf{R}$. The uniform convergence of the derivatives u'_ε and u''_ε follows from Corollary 3.2(b).

Lemma 3.4. *We have*

$$\lim_{\varepsilon \rightarrow 0} \frac{v_\varepsilon(x)}{\varepsilon} = -u'_0(x) \quad \text{uniformly on } \mathbf{R}.$$

Proof. By (2.18) we have

$$\frac{v_\varepsilon(x)}{\varepsilon} = -\frac{u'_\varepsilon(x)}{1 + \varepsilon u_\varepsilon(x)} + \varepsilon \int_0^{u_\varepsilon(x)} \frac{y_\varepsilon(u)}{(1 + \varepsilon u)^2} du. \quad (3.2)$$

Hence, because the integral is uniformly bounded, the limit follows from Lemma 3.3.

Lemma 3.5. *We have*

$$\lim_{\varepsilon \rightarrow 0} \frac{u_\varepsilon(x) - u_0(x)}{\varepsilon} = \frac{1}{2} u_0(x) \{1 - u_0(x)\} \quad \text{uniformly on } \mathbf{R}.$$

Proof. Set

$$w_\varepsilon = \frac{u_\varepsilon - u_0}{\varepsilon}.$$

Then, subtracting the equation for u_0 from the one for u_ε and dividing the difference by ε , we find that w_ε is a solution of the problem

$$\begin{cases} w'' + \frac{1}{2} x w' = -f_\varepsilon(x) & x \in (-\infty, \infty) \\ w(-\infty) = 0, & w(\infty) = 0, \end{cases}$$

where

$$f_\varepsilon(x) = -\frac{v_\varepsilon(x)}{\varepsilon} u'_\varepsilon(x). \quad (3.3)$$

Note that by Lemmas 3.3 and 3.4

$$f_0(x) \stackrel{\text{def}}{=} \lim_{\varepsilon \rightarrow 0} f_\varepsilon(x) = \{u'_0(x)\}^2 \quad \text{uniformly on } \mathbf{R}.$$

The solution w_ε can be written explicitly in terms of f_ε by means of

$$w_\varepsilon(x) = \int_x^\infty e^{-t^2/4} dt \int_0^t e^{s^2/4} f_\varepsilon(s) ds - A_\varepsilon \int_x^\infty e^{-t^2/4} dt \quad (3.5a)$$

where

$$A_\varepsilon = \frac{1}{2\sqrt{\pi}} \int_{-\infty}^{\infty} e^{-t^2/4} dt \int_0^t e^{s^2/4} f_\varepsilon(s) ds. \quad (3.5b)$$

By Lemma 2.4 and Corollary 3.2 there exists a constant C , which does not depend on ε such that

$$f_\varepsilon(x) \leq C e^{-x^2/4} \quad \text{for } -\infty < x < \infty.$$

Hence, we may let $\varepsilon \rightarrow 0$ in (3.5). Using the symmetry of $f_0(x)$ with respect to $x = 0$, we conclude that $\lim_{\varepsilon \rightarrow 0} A_\varepsilon = 0$ and that

$$\lim_{\varepsilon \rightarrow 0} w_\varepsilon(x) = w_0(x) \quad \text{uniformly on } \mathbf{R},$$

where

$$w_0(x) = \int_x^\infty e^{-t^2/4} dt \int_0^t e^{s^2/4} \{u'_0(s)\}^2 ds.$$

Remembering that $u_0(x)$ is given explicitly in (3.1) we can write $w_0(x)$ as

$$w_0(x) = \frac{1}{8} \left[1 - \left\{ \operatorname{erf} \left(\frac{x}{2} \right) \right\}^2 \right]. \quad (3.6)$$

Again, using the expression for u_0 we arrive at the desired result.

Next we estimate the second order terms in the expansions for u_ε and v_ε with respect to ε . For v_ε we find from (3.2) for $\varepsilon > 0$ and $x \in \mathbf{R}$

$$\begin{aligned} \frac{1}{2} \left\{ \frac{v_\varepsilon(x)}{\varepsilon} + u'_0(x) \right\} &= \frac{1}{2} \left\{ u'_0(x) - \frac{u'_\varepsilon(x)}{1 + \varepsilon u_\varepsilon(x)} \right\} + \int_0^{u_\varepsilon(x)} \frac{y_\varepsilon(u)}{(1 + \varepsilon u)^2} du \\ &= -w'_\varepsilon(x) + \frac{u_\varepsilon(x) u'_\varepsilon(x)}{1 + \varepsilon u_\varepsilon(x)} + \int_0^{u_\varepsilon(x)} \frac{y_\varepsilon(u)}{(1 + \varepsilon u)^2} du \end{aligned} \quad (3.7)$$

We inspect the convergence of the three terms on the right hand side of (3.7) in succession.

By differentiating (3.5a) we obtain

$$w_\varepsilon(x) = -e^{-x^2/4} \int_0^x e^{s^2/4} f_\varepsilon(s) ds - A_\varepsilon e^{-x^2/4}.$$

Using the arguments from the proof of Lemma 3.5 we conclude that

$$\lim_{\varepsilon \rightarrow 0} w'_\varepsilon(x) = -e^{-x^2/4} \int_0^x e^{s^2/4} f_0(s) ds = w'_0(x), \quad (3.8)$$

uniformly in $\mathbf{R}$.

From Lemma 3.3 we deduce that

$$\lim_{\varepsilon \rightarrow 0} \frac{u_\varepsilon(x) u'_\varepsilon(x)}{1 + \varepsilon u_\varepsilon(x)} = u_0(x) u'_0(x) \quad \text{uniformly in } \mathbf{R} \quad (3.9)$$

To pass to the limit in the third term, we first consider the convergence of the functions z_ε . Because the right hand side of the differential equation for z_ε varies monotonically with ε we have from [vDGZ] that

$$\varepsilon_1 > \varepsilon_2 > 0 \quad \Rightarrow \quad z_{\varepsilon_1}(t) > z_{\varepsilon_2}(t) > z_0(t) \quad \text{for } 0 < t < 1.$$

Hence there exists a function $\tilde{z}(t) \geq z_0(t)$ such that

$$\lim_{\varepsilon \rightarrow 0} z_\varepsilon(t) = \tilde{z}(t) \quad \text{pointwise in } [0, 1].$$

Again, using an argument from [vDGZ, proof of Proposition 2.4] we find that $\tilde{z}(t) = z_0(t)$ for $0 \leq t \leq 1$. Thus, we have the situation where a sequence of smooth functions $\{z_\varepsilon\}$ converges monotonically to a continuous function z_0 . Then Dini's theorem implies that

$$\lim_{\varepsilon \rightarrow 0} z_\varepsilon(t) = z_0(t) \quad \text{uniformly on } [0, 1],$$

and so, by (2.11),

$$\lim_{\varepsilon \rightarrow 0} y_\varepsilon(u) = y_0(u) \quad \text{uniformly on } [0, 1]. \quad (3.10)$$

Using (3.8) - (3.10) in (3.7) we obtain the desired limit

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \left\{ \frac{v_\varepsilon(x)}{\varepsilon} + u'_0(x) \right\} = E_1(x),$$

where

$$\begin{aligned} E_1(x) &= -w'_0(x) + u_0(x)u'_0(x) + \int_0^{u_0(x)} y_0(u) du \\ &= 2u_0(x)u'_0(x) - \frac{1}{2}u'_0(x) + \int_x^\infty \{u'_0(s)\}^2 ds. \end{aligned} \quad (3.11)$$

Thus we have proved

Lemma 3.6. *We have*

$$v_\varepsilon(x) = -u'_0(x)\varepsilon + \{E_1(x) + o(1)\}\varepsilon^2 \quad \text{as } \varepsilon \rightarrow 0,$$

where E_1 is given by (3.11), uniformly in $\mathbf{R}$.

To improve the estimate for u_ε given by Lemma 3.5, we investigate the limiting behaviour of the function

$$D_\varepsilon w \stackrel{\text{def}}{=} \frac{1}{\varepsilon}(w_\varepsilon - w_0) \quad \text{as } \varepsilon \rightarrow 0.$$

Thus, remembering (3.5) we need to investigate

$$D_\varepsilon f = \frac{1}{\varepsilon}(f_\varepsilon - f_0) \quad \text{as } \varepsilon \rightarrow 0.$$

By (3.3) and Lemma 3.6

$$\begin{aligned}
D_\varepsilon f(x) &= \frac{1}{\varepsilon} \left\{ -\frac{v_\varepsilon(x)}{\varepsilon} u'_\varepsilon(x) - \{u'_0(x)\}^2 \right\} \\
&= \frac{1}{\varepsilon} [\{u'_0(x) - (E_1(x) + o(1))\varepsilon\} u'_\varepsilon(x) - \{u'_0(x)\}^2] \\
&= u'_0(x) w'_\varepsilon(x) - (E_1(x) + o(1)) u'_\varepsilon(x)
\end{aligned}$$

Thus, by (3.8)

$$\lim_{\varepsilon \rightarrow 0} D_\varepsilon f(x) \text{ exists} = D_0 f(x) \quad \text{uniformly in } \mathbf{R},$$

where

$$D_0 f(x) = \{1 - 3u_0(x)\} \{u'_0(x)\}^2 - u'_0(x) \int_x^\infty \{u'_0(s)\}^2 ds. \quad (3.12)$$

Moreover, by Lemma 3.1 and Corollary 3.2 there exists a constant $C > 0$, which does not depend on ε , such that

$$D_\varepsilon f(x) \leq C e^{-x^2/4} \quad \text{for } -\infty < x < \infty.$$

Therefore we may pass to the limit in the expression for $D_\varepsilon w(x)$. This leads to

Lemma 3.7. *We have*

$$u_\varepsilon(x) = u_0(x) + \frac{1}{2} u_0(x) \{1 - u_0(x)\} \varepsilon + \{E_2(x) + o(1)\} \varepsilon^2 \quad \text{as } \varepsilon \rightarrow 0,$$

uniformly in $\mathbf{R}$. Here E_2 is given by (3.5) in which f_ε is replaced by $D_0 f$ which is defined in (3.12).

4. Self-similar solutions on $\mathbf{R}^+$

As in Sections 2 and 3, we study solutions (u, v) of the system of equations

$$\begin{cases} u'' + (\frac{1}{2}x - v)u' = 0 \end{cases} \quad 0 < x < \infty \quad (4.1)$$

$$\begin{cases} v' = -\frac{u''}{u + \frac{1}{\varepsilon}}, \end{cases} \quad (4.2)$$

but now subject to the boundary conditions

$$u(0) = 1 \quad \text{and} \quad u(\infty) = 0 \quad (4.3)$$

and

$$v(0) = -\varepsilon K(\varepsilon) u'(0), \quad (4.4)$$

where K is the positive constant which has been defined in (1.19d). Note that due to the boundary conditions at $x = 0$, the functions u and v' are no longer invariant under shifts.

We begin again with some qualitative properties of u and v .

Lemma 4.1. *Let (u, v) be a solution of (4.1) - (4.4). Then*

(a)
$$u'(x) < 0 \quad \text{for all } x \in [0, \infty).$$

(b) *There exists a number $x_0 > 0$ such that*

$$\begin{aligned} u''(x) &> 0 \quad (<) \quad \text{if } x > x_0 \quad (<) \\ v'(x) &< 0 \quad (>) \quad \text{if } x > x_0 \quad (<). \end{aligned}$$

(c)
$$u'(x) \rightarrow 0 \quad \text{as } x \rightarrow \infty.$$

Proof. We shall only prove Part (b); for the proofs of the other parts we refer to the proof of Lemma 2.1.

Because by Part (a) $u'(0) < 0$, we conclude from (4.4) that $v(0) > 0$. Using this in equation (4.1) we find that $u'' < 0$ in a neighbourhood of $x = 0$. By the boundary condition at infinity, u'' has to change sign at least once. Arguing as in the proof of Lemma 2.1(b) we show that this can happen no more than once.

To prove the existence and uniqueness of a solution (u, v) , we proceed as in Section 2 and introduce the variables y and z as in (2.8) and (2.11) respectively. This leads for z to a problem of the form

$$(P_2) \begin{cases} -zz'' = \frac{1}{2}e^{2t \log(1+\varepsilon)}, & z > 0 \quad \text{for } 0 < t < 1 \\ z(0) = 0, & z'(1) = -Lz(1), \end{cases}$$

in which L is a constant. If we choose

$$L = K(\varepsilon)(1 + \varepsilon) \log(1 + \varepsilon) = \frac{\log(1 + \varepsilon)}{\gamma - \log(1 + \varepsilon)} \quad (4.5)$$

then, returning to the original variables we find that $u(x)$ and $v(x)$ satisfy the boundary condition (4.4) at $x = 0$.

Theorem 4.2. *For any $L > -1$, Problem (P_2) has a unique solution $z \in C^\infty((0, 1]) \cap C([0, 1])$.*

Proof. Instead of Problem (P_2) we consider the problem

$$(P_\alpha) \begin{cases} -ww'' = \frac{1}{2}e^{2t \log(1+\varepsilon)} & w > 0 \quad \text{for } 0 < t < 1 \\ w(0) = 0, & w(1) = \alpha, \end{cases}$$

where $\alpha > 0$ is an arbitrary positive number.

From [vDGZ], where further references are given, we know that for each $\alpha > 0$, Problem (P_α) has a unique, concave solution $w_\alpha \in C^\infty((0, 1]) \cap C([0, 1])$.

It will be convenient to fix the boundary conditions. Thus we normalize w_α and write

$$r_\alpha = \alpha^{-1} w_\alpha.$$

The functions r_α then satisfy

$$-rr'' = \frac{1}{2\alpha^2} e^{2t \log(1+\epsilon)}, \quad r > 0 \quad \text{for} \quad 0 < t < 1 \quad (4.6)$$

and

$$r(0) = 0, \quad r(1) = 1.$$

For every $\alpha > 0$ we can define

$$\beta(\alpha) = r'_\alpha(1). \quad (4.7)$$

It then remains to show that the equation

$$\beta(\alpha) = -L \quad (4.8)$$

has one and only one solution if $L > -1$. In the following lemma we derive some properties of the function β which together ensure that this is indeed the case.

Lemma 4.3. *We have*

$$(a) \quad \alpha_1 < \alpha_2 \quad \Rightarrow \quad \beta(\alpha_1) < \beta(\alpha_2)$$

$$(b) \quad \beta \in C(0, \infty) \text{ and}$$

$$\lim_{\alpha \rightarrow \infty} \beta(\alpha) = 1 \quad \text{and} \quad \lim_{\alpha \rightarrow 0} \beta(\alpha) = -\infty.$$

Proof. (a) Because the right hand side of (4.6) decreases with increasing α , we apply again a comparison argument and find

$$\alpha_1 < \alpha_2 \quad \Rightarrow \quad r_{\alpha_1} > r_{\alpha_2} \quad \text{on} \quad (0, 1)$$

and so

$$r'_{\alpha_1}(1) \leq r'_{\alpha_2}(1). \quad (4.9)$$

In fact, using an argument from [vDP1, Theorem 4], we can show that strict inequality holds in (4.9):

$$\alpha_1 < \alpha_2 \quad \Rightarrow \quad r'_{\alpha_1}(1) < r'_{\alpha_2}(1).$$

To prove Part (b) we use a compactness argument. Let $\bar{\alpha} > 0$ and let α_n be an increasing sequence such that

$$\alpha_n \nearrow \bar{\alpha} \quad \text{as } n \rightarrow \infty.$$

Then $\{r_{\alpha_n}\}$ is a decreasing sequence, bounded below by $r_{\bar{\alpha}}$. Hence, there exists a function $\bar{r} \geq r_{\bar{\alpha}}$ such that for every $t \in [0, 1]$,

$$r_{\alpha_n}(t) \searrow \bar{r}(t) \quad \text{as } n \rightarrow \infty.$$

Because the sequence $\{r_{\alpha_n}\}$ is bounded away from zero on any interval $[\delta, 1]$, where $\delta \in (0, 1)$, the differential equation implies that $r'_{\alpha_n}, r''_{\alpha_n}$ and r'''_{α_n} are all bounded on $[\delta, 1]$, uniformly with respect to n . Hence, there exists a subsequence $\{r_{\alpha_{n'}}\}$ such that

$$r_{\alpha_{n'}} \rightarrow \bar{r}, \quad r'_{\alpha_{n'}} \rightarrow \bar{r}', \quad r''_{\alpha_{n'}} \rightarrow \bar{r}'' \quad n \rightarrow \infty, \quad (4.10)$$

uniformly on $[\delta, 1]$. By taking the limit in the equation we conclude that

$$-\bar{r}\bar{r}'' = \frac{1}{2\bar{\alpha}^2} e^{2t \log(1+\varepsilon)}, \quad \bar{r} > 0 \quad \text{for } 0 < t < 1,$$

because we may choose δ arbitrary small. Since

$$r_{\bar{\alpha}} \leq \bar{r} < r_{\alpha_n} \quad \text{for } 0 \leq t \leq 1,$$

we also have

$$\bar{r}(0) = 0 \quad \text{and} \quad \bar{r}(1) = 1.$$

Thus, $\bar{r} = r_{\bar{\alpha}}$ and the limit is uniquely determined by $\bar{\alpha}$. Therefore (4.10) holds, with $\bar{r}$ replaced by $r_{\bar{\alpha}}$, for the entire sequence. This means in particular that

$$r'_{\alpha_n}(1) \rightarrow r'_{\bar{\alpha}}(1) \quad \text{as } n \rightarrow \infty,$$

or

$$\beta(\alpha_n) \rightarrow \beta(\bar{\alpha}) \quad \text{as } n \rightarrow \infty.$$

A similar argument can be given for a decreasing sequence. This proves the continuity of β at $\bar{\alpha}$.

Thus, for any given $L > -1$, there exists a unique number α_L such that

$$\beta(\alpha_L) = -L$$

and (4.8) is satisfied. This completes the proof of Theorem 4.2.

We return to the original variables x , u and v by means of

$$x(u) = \int_u^1 \frac{ds}{y(s)} \quad \text{for } 0 < u \leq 1$$

and

$$v(x) = \frac{1}{2}x - y'(u(x)) \quad \text{for } 0 \leq x < \infty.$$

By an elementary computation we then verify that the pair of functions $(u(x), v(x))$ satisfies (4.1) - (4.4). Thus we have proved

Theorem 4.4. *For any $0 < \varepsilon < e^\gamma - 1$ there exists a unique pair of functions (u, v) which satisfies (4.1) - (4.4).*

In the next lemma we list some properties of $v(x)$.

Lemma 4.5. *Let (u, v) be the solution of (4.1) - (4.4). Then*

$$(a) \quad v(\infty) = \lim_{x \rightarrow \infty} v(x) \text{ exists}$$

and is finite;

$$(b) \quad v(x) > v(\infty) \quad \text{for } 0 \leq x < \infty.$$

$$(c) \quad v(x) = \varepsilon \left(K(\varepsilon) - \frac{1}{1 + \varepsilon} \right) y(1) + \varepsilon \frac{y(u(x))}{1 + \varepsilon u(x)} - \varepsilon^2 \int_{u(x)}^1 \frac{y(u)}{(1 + \varepsilon u)^2} du.$$

$$(d) \quad v(\infty) > 0 \quad \text{for } \varepsilon \text{ small enough.}$$

Proof. The proofs of Parts (a) - (c) use arguments very similar to those used in the proof of Lemma 2.2. We therefore omit them.

To prove Part (d) we observe that by (c),

$$v(\infty) = \varepsilon \left(K(\varepsilon) - \frac{1}{1 + \varepsilon} \right) y(1) - \varepsilon^2 \int_0^1 \frac{y(u)}{(1 + \varepsilon u)^2} du. \quad (4.11)$$

By (1.19d),

$$K(\varepsilon) - \frac{1}{1 + \varepsilon} = \frac{1}{1 + \varepsilon} \left(\frac{1}{\gamma - \log(1 + \varepsilon)} - 1 \right) > \frac{1}{\gamma} - 1 > 0 \quad (4.12)$$

and $y(1) = -u'(0) > 0$. Thus, since for small values of ε the first term on the right hand side of (4.11) dominates the second term, the assertion is proved.

In Problem I we required that $v(\infty) = 0$. Equation (2.16) then implies that $v(-\infty) > 0$ and that $v(-\infty) = O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$. Here we find that $v(\infty)$ is not so small.

Lemma 4.6. *We have*

$$v(\infty) = O(\varepsilon) \quad \text{as } \varepsilon \rightarrow 0.$$

Proof. Writing (4.11) in terms of z we obtain

$$v(\infty) = \log(1 + \varepsilon)z(1) \left\{ K(\varepsilon) - \frac{1}{1 + \varepsilon} - \log(1 + \varepsilon) \int_0^1 \frac{z(t)}{z(1)} e^{-t \log(1 + \varepsilon)} dt \right\}. \quad (4.13)$$

Thus, it remains to bound $z(t)$ from above and below, uniformly with respect to ε . To that end we introduce two auxiliary functions $\underline{z}$ and $\bar{z}$, which will serve as lower and upper bound respectively.

– For $\underline{z}$ we choose the solution of the problem

$$\begin{cases} -zz'' = \frac{1}{2}, & z > 0 & \text{for } 0 < t < 1 \\ z(0) = 0, & z'(1) = -Lz(1), \end{cases}$$

in which we choose $L = \Lambda(\varepsilon_0)$, where (see (4.11)),

$$\Lambda(\varepsilon) \stackrel{\text{def}}{=} (1 + \varepsilon) \log(1 + \varepsilon) K(\varepsilon) \quad \text{and} \quad \varepsilon_0 \in (0, e^\gamma - 1).$$

Note that Λ is an increasing function.

– For $\bar{z}$ we choose the solution of the problem

$$\begin{cases} -zz'' = \frac{1}{2}(1 + \varepsilon_0), & z > 0 & \text{for } 0 < t < 1 \\ z(0) = 0, & z'(1) = 0, \end{cases}$$

Lemma 4.7. *Let $z_\varepsilon(t)$ be the solution of Problem P_2 in which $L = \Lambda(\varepsilon)$, and let $\underline{z}$ and $\bar{z}$ be as defined above. Then for $\varepsilon \in (0, \varepsilon_0)$, where $0 < \varepsilon_0 < e^\gamma - 1$,*

$$\underline{z}(t) < z_\varepsilon(t) < \bar{z}(t) \quad \text{for } 0 < t < 1.$$

Proof. We shall only prove the lower bound; the upper bound is proved in the same way. We argue by contradiction. Thus, suppose there exist numbers $0 \leq a < b \leq 1$ such that

$$w(t) = z(t) - \underline{z}(t) < 0 \quad \text{for } a < t < b$$

and

$$w(a) = 0.$$

If $b \leq 1$ and $w(b) = 0$ we obtain a contradiction as in Theorem 2.1 of [vDGZ]. On the other hand, if $b = 1$ and $w(1) < 0$ it follows from the boundary condition at $t = 1$ that

$$w'(1) = -\Lambda(\varepsilon)z(1) + \Lambda(\varepsilon_0)\underline{z}(1) > \{\Lambda(\varepsilon_0) - \Lambda(\varepsilon)\}z(1) > 0, \quad (4.14)$$

where we have used the monotonicity of $\Lambda(\varepsilon)$. From the differential equations for z and $\underline{z}$ we finally deduce that

$$\begin{aligned} w''(t) &= \frac{1}{2\underline{z}} - \frac{1}{2z} e^{2t \log(1+\varepsilon)} \\ &< \frac{1}{2z} \left(1 - e^{2t \log(1+\varepsilon)}\right) < 0. \end{aligned} \quad (4.15)$$

It follows from (4.14) and (4.15) that

$$w'(t) > 0 \quad \text{for } a \leq t \leq 1.$$

Since $w(a) = 0$ this means that

$$w(t) > 0 \quad \text{for } a \leq t \leq 1$$

which contradicts our assumption. This proves the lower bound.

We conclude this section with the observation that the asymptotic behaviour of the solution $(u(x), v(x))$ as $x \rightarrow \infty$ is similar to what we found in Lemma 2.4. Here we have for $x > 0$

$$u(x) < A_+ x^{-1} e^{-x^2/4} \quad (4.16a)$$

and

$$v(x) - v(\infty) < A_+ x^{-1} e^{-x^2/4}, \quad (4.16b)$$

where A_+ is a positive constant, which does not depend on ε .

5. The limit $\varepsilon \rightarrow 0$ for Problem II

Having established the existence and uniqueness of a solution $(u_\varepsilon(x), v_\varepsilon(x))$ of (4.1)-(4.4), we now determine its limiting behaviour as $\varepsilon \rightarrow 0$.

Lemma 5.1. *There exist positive constants C_1 and C_2 , which do not depend on ε such that*

$$|u'_\varepsilon(x)| < C_1 \quad \text{and} \quad 0 < v_\varepsilon(x) < C_2 \varepsilon$$

for all $\varepsilon \in (0, \varepsilon_0)$.

Proof. By Lemma 4.7, $z_\varepsilon(t)$ is uniformly bounded and thus, so is $y_\varepsilon(u)$. This proves the first bound.

The second bound follows at once from Lemma 4.5(c).

Let u_0 be the solution of the problem

$$\begin{cases} u'' + \frac{1}{2}xu' = 0, & 0 < x < \infty \\ u(0) = 1, & u(\infty) = 0. \end{cases}$$

It can be written down explicitly and is found to be

$$u_0(x) = \operatorname{erfc}\left(\frac{x}{2}\right). \quad (5.1)$$

As in Section 3, we can establish the following limits.

Lemma 5.2. *We have*

$$u_\varepsilon(x) \rightarrow u_0(x), \quad u'_\varepsilon(x) \rightarrow u'_0(x), \quad u''_\varepsilon(x) \rightarrow u''_0(x) \quad \text{as } \varepsilon \rightarrow 0$$

uniformly on $[0, \infty)$.

To obtain an estimate for $v_\varepsilon(x)$ we use Lemma 4.5(c). With the limits from Lemma 5.2 we find that

$$\frac{v_\varepsilon(x)}{\varepsilon} \rightarrow -\{K(0) - 1\}u'_0(0) - u'_0(x) \quad \text{as } \varepsilon \rightarrow 0$$

uniformly on $[0, \infty)$. Remembering the definition (1.19d) of $K(\varepsilon)$ we arrive at the following limit

Lemma 5.3. *We have*

$$\lim_{\varepsilon \rightarrow 0} \frac{v_\varepsilon(x)}{\varepsilon} = \frac{1 - \gamma}{\sqrt{\pi} \gamma} - u'_0(x) \quad \text{uniformly on } [0, \infty).$$

To improve the estimate for $u_\varepsilon(x)$ we consider again the function

$$w_\varepsilon = \frac{u_\varepsilon - u_0}{\varepsilon}$$

which now is the solution of the problem

$$\begin{cases} w'' + \frac{1}{2}xw' = -f_\varepsilon(x) & x \in (0, \infty) \\ w(0) = 0, \quad w(\infty) = 0, \end{cases}$$

where

$$f_\varepsilon(x) = -\frac{v_\varepsilon(x)}{\varepsilon} u'_\varepsilon(x). \quad (5.2)$$

Note that by (4.16a) and the boundedness of $v_\varepsilon(x)$,

$$|f_\varepsilon(x)| \leq C e^{-x^2/4} \quad \text{for } x \geq 0, \quad (5.3)$$

where C may be chosen independent of ε . In addition, by Lemmas 5.1 and 5.2

$$\lim_{\varepsilon \rightarrow 0} f_\varepsilon(x) = -\frac{1 - \gamma}{\sqrt{\pi} \gamma} u'_0(x) + \{u'_0(x)\}^2 \quad \text{uniformly on } [0, \infty).$$

We find that

$$w_\varepsilon(x) = A_\varepsilon \int_0^x e^{-t^2/4} dt - \int_0^x e^{-t^2/4} dt \int_0^t e^{s^2/4} f_\varepsilon(s) ds \quad (5.4)$$

and the constant A_ε is chosen so that $w_\varepsilon(\infty) = 0$:

$$\sqrt{\pi} A_\varepsilon = \int_0^\infty e^{-t^2/4} dt \int_0^t e^{s^2/4} f_\varepsilon(s) ds. \quad (5.5)$$

It follows from (5.3) that the integral on the right hand side of (5.5) exists and is uniformly convergent with respect to ε . Therefore we may let $\varepsilon \rightarrow 0$ in (5.4) and (5.5) to obtain the required limit. In this manner we find:

Lemma 5.4. *We have*

$$\lim_{\varepsilon \rightarrow 0} \frac{u_\varepsilon(x) - u_0(x)}{\varepsilon} = -\frac{2(1-\gamma)}{\pi\gamma} \{u_0(x) + \sqrt{\pi} u'_0(x)\} + \frac{1}{2} u_0(x) \{1 - u_0(x)\}$$

uniformly with respect to $x \in [0, \infty)$.

Further terms in the asymptotic expansions for u_ε and v_ε as $\varepsilon \rightarrow 0$ may be obtained in exactly the same way as was done for Problem I in Section 3. We leave that to the interested reader.

6. Numerical results

Problem I

To solve equations (1.22) and (1.23) on $\mathbf{R}$ we first put them in the form of three first order equations by setting

$$p = u, \quad q = u' \quad \text{and} \quad r = v \quad (6.1)$$

This gives

$$p' = q \quad (6.2a)$$

$$q' = q(r - \frac{1}{2}x) \quad (6.2b)$$

$$r' = -\frac{q(r - \frac{1}{2}x)}{p + \frac{1}{\varepsilon}}, \quad (6.2c)$$

We solve this first order system with a Runge-Kutta method in the regions $\{x < 0\}$ and $\{x > 0\}$, subject to the initial conditions imposed on $x = 0$:

$$p(0) = u(0) = \frac{1}{2}, \quad q(0) = u'(0) = q_0, \quad r(0) = v(0) = r_0. \quad (6.3)$$

The values of q_0 and r_0 have to be selected so that

$$p(\infty) = u(\infty) = 0 \quad \text{and} \quad p(-\infty) = u(-\infty) = 1.$$

This is done by solving Problem (P₁) for z . This equation is discretised with $\Delta t = 10^{-3}$ and the nonlinear system is solved using a multi dimensional Newton method with an accuracy of 10^{-8} . In this manner we compute

$$y\left(\frac{1}{2}\right) = \frac{\log(1 + \varepsilon)}{\varepsilon} z\left(\frac{\log(1 + \frac{\varepsilon}{2})}{\log(1 + \varepsilon)}\right)$$

and

$$y'\left(\frac{1}{2}\right) = \frac{1}{1 + \frac{\varepsilon}{2}} z'\left(\frac{\log(1 + \frac{\varepsilon}{2})}{\log(1 + \varepsilon)}\right).$$

By (2.8) and (2.13) we have

$$q_0 = -y\left(\frac{1}{2}\right) \quad \text{and} \quad r_0 = -y'\left(\frac{1}{2}\right).$$

This yields a solution of (1.22), (1.23) and (1.24a) which satisfies $u(0) = \frac{1}{2}$. In particular it gives $v(\infty) = r(\infty)$, so that we can apply the shift (2.5) to obtain the desired solution.

Remark. In principle we could have constructed a solution (u, v) directly from the function $z(t)$ or $y(u)$, as was done theoretically in Section 2. However this does not yield smooth asymptotic behaviour near $u = 0$ as x becomes large and near $u = 0$ as x becomes small. Therefore we have chosen to work here with the Runge Kutta method for the system (6.2). This procedure gives smooth decay at large values of $|x|$. Moreover it serves as a check for the accuracy with which $y(\frac{1}{2})$ and $y'(\frac{1}{2})$ are determined.

We carried out the computations for a small ε ($\varepsilon = 0.025$) and for a large value of ε ($\varepsilon = 0.5$). The latter value was chosen in order to emphasize the differences. We find

$$\begin{array}{llll} \varepsilon = 0.025 & : & \text{shift } a = 2v(\infty) & = -1.84210^{-2}; \\ \varepsilon = 0.5 & : & \text{shift } a = 2v(\infty) & = -0.2198. \end{array}$$

Figure 2 shows the scaled density distribution u for the two values of ε . The difference between the Boussinesq limit u_0 , see expression (1.28), and the solution u for $\varepsilon = 0.025$ is so small that they would coincide in Figure 2.

Fig. 2. Computed scaled density distributions u on $\mathbf{R}$.

Figure 3 shows the scaled velocity $v = q_z \sqrt{t}$ induced by the density effects. Here the Boussinesq limit corresponds to $v = 0$.

Fig. 3. Computed scaled velocities $v = q_z \sqrt{t}$ on $\mathbf{R}$.

Observe that the results represented in both figures correspond qualitatively to the analysis in Sections 2 and 3.

Problem II

Here we solve the system (6.2) for $x > 0$, subject to the initial conditions

$$p(0) = u(0) = 1 \quad (6.4a)$$

$$q(0) = u'(0) = q_0 \quad (6.4b)$$

$$r(0) = v(0) = -\varepsilon K(\varepsilon)q_0 \quad (6.4c)$$

The special form of (6.4c) results from the boundary condition (1.25b). Again we apply a Runge Kutta procedure, this time with q_0 as a shooting parameter. Its value is obtained as follows:

(i) Using Newton's method we solve numerically, for each given $\varepsilon > 0$, the sequence of Problems (P_α) where we let α vary from 0 to a sufficiently large number $\bar{\alpha}$ in steps $\Delta\alpha = 10^{-3}$. This leads to graphs of the functions $\beta(\alpha)$, shown in Figure 4, with the properties of Lemma 4.3

Fig. 4. Representation of the function $\beta(\alpha)$ defined by (4.7).

(ii) Given the constant $L > 0$ from (4.5), we determine the value α_L from the graph $\beta(\alpha)$ such that

$$\beta(\alpha_L) = -L.$$

(iii) Having obtained $\alpha_L = z(1)$ in Problem (P_2) , we apply the transformation (2.10) to obtain

$$q_0 = -y(1) = -\frac{\log(1 + \varepsilon)}{\varepsilon} \alpha_L. \quad (6.5)$$

With this value of q_0 we solve the initial value problem (6.2), (6.4) and check if $r(x)$ is sufficiently close to zero for large x . For example, with $\varepsilon = 0.5$, we obtain from interpolation of points on the $\beta(\alpha)$ curve

$$\alpha_L = 0.4191 \quad \text{giving} \quad 0 < r(\infty) < 10^{-4}.$$

Figures 5 and 6 give the results of the computations for the functions u and v , respectively, using the values $\varepsilon = 0.025$ and $\varepsilon = 0.5$.

Fig. 5. Computed scaled density distributions on $\mathbb{R}^+$.

Fig. 6. Computed scaled velocities $v = q_z \sqrt{t}$ on $\mathbb{R}^+$.

Again observe that these results correspond to the qualitative behaviour as analysed in Sections 4 and 5.

From the analysis and the computations we conclude that the influence of q_z on the salt distribution is relatively small in all cases and even negligible for the value $\varepsilon = 0.025$ (as to be expected). Therefore in many applications one may disregard the influence of volume changes due to variations in the salt concentration and use the Boussinesq limit

$$\operatorname{div} \mathbf{q} = 0, \quad (6.6)$$

$$\Phi \frac{\partial \rho}{\partial t} + \operatorname{div}(\mathbf{q}\rho - D \operatorname{grad} \rho) = 0, \quad (6.7)$$

$$\frac{\mu}{\kappa} \mathbf{q} + \operatorname{grad} p - \rho \mathbf{g} = 0. \quad (6.8)$$

7. Velocity dependent dispersion

In a more realistic description of the transport of solutes through porous media, a velocity dependent dispersion matrix has to be introduced into equation (1.2). For the specific flow problems considered in this paper this means that, instead of equation (1.11), we are led to consider the transport equation

$$\Phi \frac{\partial \rho}{\partial t} + q_z \frac{\partial \rho}{\partial z} = \frac{\partial}{\partial z} \left(D(|\mathbf{q}|) \frac{\partial \rho}{\partial z} \right), \quad (7.1)$$

where

$$D(|\mathbf{q}|) = \Phi D_{\text{mol}} + a_T |\mathbf{q}| + (a_L - a_T) \frac{q_z^2}{|\mathbf{q}|}. \quad (7.2)$$

Here a_L and a_T are positive numbers, called the longitudinal and transversal dispersion length, respectively and by $|\mathbf{q}|$ we denote the length of the vector $\mathbf{q}$.

When $|q_z| \ll |q_y|$, the last term in (7.2) may be neglected and we can write D , with the help of (1.9) as

$$\begin{aligned} D &= \Phi D_{\text{mol}} + a_T |q_y| \\ &= \Phi D_{\text{mol}} + a_T \left| q_f - \frac{\kappa}{\mu} (\rho - \rho_f) g \sin \beta \right|. \end{aligned} \quad (7.3)$$

Thus, in this approximation, D is a function of ρ .

Scaling the variables in equations (1.10), (7.1) and (7.3) according to

$$\rho^* = \frac{\rho - \rho_f}{\rho_s - \rho_f}, \quad q_z^* = \frac{q_z}{\hat{q}},$$

$$z^* = \frac{z}{a_T}, \quad t^* = \frac{\hat{q}}{\Phi a_T} t \quad \text{and} \quad D^* = \frac{D}{a_T \hat{q}},$$

where $\hat{q} = \frac{\kappa}{\mu} (\rho_s - \rho_f) g \sin \beta$, and dropping the asterisks again, we find

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial z} (\rho q_z) + \frac{1}{\varepsilon} \frac{\partial q_z}{\partial z} = 0 \quad \text{in } \Omega \times \mathbf{R}^+ \quad (7.4)$$

$$\frac{\partial \rho}{\partial t} + q_z \frac{\partial \rho}{\partial z} = \frac{\partial}{\partial z} \left(D(\rho) \frac{\partial \rho}{\partial z} \right) \quad \text{in } \Omega \times \mathbf{R}^+ \quad (7.5)$$

in which

$$D(\rho) = \lambda + |U - \rho|. \quad (7.6)$$

The constant ε in (7.4) is given again by (1.15) and in (7.6) we have

$$\lambda = \frac{\Phi D_{\text{mol}}}{a_T \hat{q}} \quad \text{and} \quad U = \frac{qf}{\hat{q}}.$$

Observe that the scaling proposed here is different from the scaling applied in (1.14). It allows us to consider the limit of small molecular diffusion with respect to the transversal dispersion, i.e. $\lambda = 0$, a mathematically interesting limit because it may lead to degenerate diffusion at points where $\rho = U$.

Applying the similarity transformation (1.20) and (1.21) yields the equations

$$(uv)' + \frac{1}{\varepsilon}v' - \frac{1}{2}xu' = 0 \quad \text{in } \Omega \quad (7.7)$$

$$u'v - \frac{1}{2}xu' = (D(u)u')' \quad \text{in } \Omega \quad (7.8)$$

In the nondegenerate case, i.e. $\lambda > 0$, or when $\lambda = 0$ and either $U < 0$ or $U > 1$, the qualitative properties as obtained in Sections 2 and 4 can be obtained with minor modifications of the proofs. Only the degenerate case $\lambda = 0$ and $0 \leq U \leq 1$, requires extra attention. Also, for all choices of λ and U , the asymptotic analysis, as $\varepsilon \rightarrow 0$, becomes more involved and less explicit. We shall leave these questions to a future paper. Formally the $\varepsilon = 0$ limit gives the equations

$$v = 0 \quad \text{in } \Omega \quad (7.9)$$

$$(D(u)u')' + \frac{1}{2}xu' = 0 \quad \text{in } \Omega \quad (7.10)$$

Self similar solutions of (7.10) were studied in detail in [vDP1, dJdJvD].

The von Mises transformation

We return to equations (1.10) and (1.11). Absorbing the porosity into the time coordinate and confining ourselves to flow situations as described in Problem I, we obtain

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial z}(\rho q_z) = 0 \quad \text{in } \mathbf{R} \times \mathbf{R}^+ \quad (7.11)$$

$$\frac{\partial \rho}{\partial t} + q_z \frac{\partial \rho}{\partial z} = \frac{\partial}{\partial z} \left(D(\rho) \frac{\partial \rho}{\partial z} \right) \quad \text{in } \mathbf{R} \times \mathbf{R}^+ \quad (7.12)$$

with, for example, $D(\rho)$ given by (7.3).

We shall apply a coordinate transformation which is a variant of the *von Mises transformation*, well known in the theory of the Prandtl boundary layer equations, (see for instance [O,vMF]). Considering (7.11) as the divergence operator in the (t, z) -plane, acting on the vector $(\rho, \rho q_z)$, we introduce the *stream function* $\psi = \psi(z, t)$ which satisfies

$$\rho = \frac{\partial \psi}{\partial z} \quad \text{and} \quad \rho q_z = -\frac{\partial \psi}{\partial t}. \quad (7.13)$$

Next we set

$$\hat{\rho}(\psi, t) = \hat{\rho}(\psi(z, t), t) = \rho(z, t) \quad (7.14)$$

and obtain

$$\begin{aligned} \frac{\partial \rho}{\partial t} &= \frac{\partial \hat{\rho}}{\partial \psi} \cdot \frac{\partial \psi}{\partial t} + \frac{\partial \hat{\rho}}{\partial t} = -\rho q_z \frac{\partial \hat{\rho}}{\partial \psi} + \frac{\partial \hat{\rho}}{\partial t}, \\ \frac{\partial \rho}{\partial z} &= \frac{\partial \hat{\rho}}{\partial \psi} \cdot \frac{\partial \psi}{\partial z} = \rho \frac{\partial \hat{\rho}}{\partial \psi}. \end{aligned}$$

Hence

$$\frac{\partial \hat{\rho}}{\partial t} = \hat{\rho} \frac{\partial}{\partial \psi} \left(D(\hat{\rho}) \hat{\rho} \frac{d\hat{\rho}}{d\psi} \right). \quad (7.15)$$

In view of (7.13) we also consider this equation on the domain $\{(\psi, t) : -\infty < \psi < \infty, t > 0\}$.

Now suppose that the initial salt distribution, in the original coordinates, is given by

$$\rho(z, 0) = \rho_0(z), \quad -\infty < z < \infty, \quad (7.16)$$

where the function ρ_0 satisfies $\rho_f \leq \rho_0 \leq \rho_s$ such that

$$\lim_{z \rightarrow \infty} \rho_0(z) = \rho_f \quad \text{and} \quad \lim_{z \rightarrow -\infty} \rho_0(z) = \rho_s.$$

Then we define the function $\tilde{z} : \mathbf{R} \rightarrow \mathbf{R}$ by

$$\int_0^{\tilde{z}(\psi)} \rho_0(s) ds = \psi, \quad -\infty < \psi < \infty,$$

and we take as initial distribution in the (ψ, t) coordinate system

$$\hat{\rho}(\psi, 0) = \rho_0(\tilde{z}(\psi)) \quad , \quad -\infty < \psi < \infty. \quad (7.17)$$

After having obtained the solution $\hat{\rho} = \hat{\rho}(\psi, t)$, of the initial value problem (7.15), (7.17), we return to the original coordinates as follows. First define the *stream function* $\psi : \mathbf{R} \times \overline{\mathbf{R}}^+ \rightarrow \mathbf{R}$ by

$$z = \int_0^{\psi(z, t)} \frac{1}{\hat{\rho}(s, t)} ds + h(t), \quad (7.18)$$

where $h(t)$ will be chosen later in relation to the boundary conditions on q_z . Note that because by the maximum principle $0 < \rho_f \leq \hat{\rho}(\psi, t) \leq \rho_s$, $z \rightarrow \pm\infty$ implies that $\psi \rightarrow \pm\infty$. Next we define

$$\rho(z, t) = \hat{\rho}(\psi, t)$$

and finally

$$q_z(z, t) = h'(t) - \int_0^{\psi(z, t)} \frac{\hat{\rho}_t}{\hat{\rho}^2}(s, t) ds. \quad (7.19)$$

Then the boundary condition $q_z(\infty, t) = 0$ for all $t > 0$ will be satisfied if we choose

$$h(t) = \int_0^t \left\{ \int_0^\infty \frac{\hat{\rho}_t}{\hat{\rho}^2}(s, \tau) ds \right\} d\tau. \quad (7.20)$$

under the assumption that the integrals exist. In that case, q_z is given by

$$q_z(z, t) = \int_{\psi(z, t)}^\infty \frac{\hat{\rho}_t}{\hat{\rho}^2}(s, \tau) ds. \quad (7.21)$$

Another condition on q_z could have been: $q_z(0, t) = 0$. In that case we would have chosen $h(t) = 0$.

One easily verifies that the pair (ρ, q_z) solves (7.11), (7.12) and (7.16).

If we take as initial profile the piecewise constant function

$$\rho_0(z) = \begin{cases} \rho_f & z > 0 \\ \rho_s & z < 0 \end{cases}$$

then also

$$\hat{\rho}(\psi, 0) = \begin{cases} \rho_f & \psi > 0 \\ \rho_s & \psi < 0. \end{cases} \quad (7.22)$$

Consequently the solution of (7.15), (7.22) has self similar form:

$$\hat{\rho}(\psi, t) = f(\eta) \quad \text{with } \eta = \psi/\sqrt{t}.$$

Here f is the solution of the boundary value problem

$$\begin{cases} f\{D(f)ff'\}' + \frac{1}{2}\eta f' = 0, & -\infty < \eta < \infty \\ f(-\infty) = \rho_s, & f(\infty) = \rho_f. \end{cases}$$

Using the similarity in equation (7.18) we obtain

$$x = \frac{z}{\sqrt{t}} = \int_0^{\psi(z, t)/\sqrt{t}} \frac{d\xi}{f(\xi)} - \int_0^\infty \frac{\xi f'(\xi)}{f^2(\xi)} d\xi \quad (7.23)$$

This implies that $\psi/\sqrt{t}$ only depends on x , i.e.

$$\psi(z, t) = \phi(x)\sqrt{t},$$

and consequently that

$$\rho(z, t) = \hat{\rho}(\psi(z, t), t) = f\left(\frac{\psi(z, t)}{\sqrt{t}}\right) = f(\phi(x)) \stackrel{\text{def}}{=} u(x).$$

For q_z we obtain from (7.19)

$$q_z(z, t) = -\frac{1}{2\sqrt{t}} \int_{\phi(x)}^\infty \frac{\xi f'(\xi)}{f^2(\xi)} d\xi \stackrel{\text{def}}{=} \frac{1}{\sqrt{t}} v(x).$$

Much is known about the large time behaviour of solutions of the initial value problem (7.15), (7.17) for fairly general initial functions $\hat{\rho}(\psi, 0)$. In particular sharp estimates were obtained for the rate at which the solutions converge to the self-similar profile $f(\eta)$, (see for instance [vDP2,Z]). The translation of these convergence results to the original, physical, coordinates (z, t) has not yet been studied in detail. We intend to do this in a future study.

References

- [B] J. Bear, *Dynamics of fluids in porous media*, Elsevier, New York, 1972.
- [AP] F.V. Atkinson & L.A. Peletier, *Similarity solutions of nonlinear diffusion equations*, Arch. Rational Mech. Anal. **54** (1974) 373–392.
- [dJdJvD] G. de Josseling de Jong & C.J. van Duijn, *Transverse dispersion from an originally sharp fresh-salt interface caused by shear flow*, J. Hydrology **84** (1986) 55–79.
- [vDGZ] C.J. van Duijn, S.M. Gomez & Hongfei Zhang, *On a class of similarity solutions of the equation $u_t = (|u|^{m-1}u_x)_x$ with $m > -1$* , IMA J. Appl. Math. **41** (1988) 147–163.
- [vDP1] C.J. van Duijn & L.A. Peletier, *A class of similarity solutions of the nonlinear diffusion equation*, Nonlinear Anal. TMA **1** (1977) 223–233.
- [vDP2] C.J. van Duijn & L.A. Peletier, *Asymptotic behaviour of solutions of nonlinear diffusion equations*, Arch. Rational Mech. Anal. **65** (1977) 363–377.
- [vDP3] C.J. van Duijn & L.A. Peletier, *A boundary layer problem in fresh-salt groundwater flow*, Report Mathematical Institute, Leiden University W 90-07, 1990; to appear in Quart. J. Appl. Math. Mech.
- [HL] S.M. Hassanizadeh & T. Leijnse, *On the modelling of brine transport in porous media*, Water Resources Research **24** (1988) 321–330.
- [vMF] R. von Mises & K.O. Friedrichs, *Fluid dynamics*, Applied Mathematics Series 5, Springer, New York, 171.
- [O] O.A. Oleinik, *The Prandtl system of equations*, Sov. Math. **4** (1963) 383–386.
- [T] S.D. Taliaferro, *A nonlinear singular boundary value problem*, Nonlinear Anal. TMA **3** (1979) 897–904.
- [Z] Hongfei Zhang, *Large time behaviour of the maximal solution of the equation $u_t = (u^{m-1}u_x)_x$ with $-1 < m < 0$* , Submitted to SIAM J. Math. Anal.

Department of Mathematics
Delft University of Technology

Mathematical Institute
Leiden University

#	Author/s	Title
746	Chris Grant,	Solutions to evolution equations with near-equilibrium initial values
747	Mario Taboada and Yuncheng You,	Invariant manifolds for retarded semilinear wave equations
748	Peter Rejto and Mario Taboada,	Unique solvability of nonlinear Volterra equations in weighted spaces
749	Hi Jun Choe,	Holder regularity for the gradient of solutions of certain singular parabolic equations
750	Jack D. Dockery,	Existence of standing pulse solutions for an excitable activator-inhibitory system
751	Jack D. Dockery and Roger Lui,	Existence of travelling wave solutions for a bistable evolutionary ecology model
752	Giovanni Alberti, Luigi Ambrosio and Giuseppe Buttazzo,	Singular perturbation problems with a compact support semilinear term
753	Emad A. Fatemi,	Numerical schemes for constrained minimization problems
754	Y. Kuang and H.L. Smith,	Slowly oscillating periodic solutions of autonomous state-dependent delay equations
755	Emad A. Fatemi,	A new splitting method for scalar conservation laws with stiff source terms
756	Hi Jun Choe,	A regularity theory for a more general class of quasilinear parabolic partial differential equations and variational inequalities
757	Haitao Fan,	A vanishing viscosity approach on the dynamics of phase transitions in Van Der Waals fluids
758	T.A. Osborn and F.H. Molzahn,	The Wigner–Weyl transform on tori and connected graph propagator representations
759	Avner Friedman and Bei Hu,	A free boundary problem arising in superconductor modeling
760	Avner Friedman and Wenxiong Liu,	An augmented drift-diffusion model in semiconductor device
761	Avner Friedman and Miguel A. Herrero,	Extinction and positivity for a system of semilinear parabolic variational inequalities
762	David Dobson and Avner Friedman,	The time-harmonic Maxwell equations in a doubly periodic structure
763	Hi Jun Choe,	Interior behaviour of minimizers for certain functionals with nonstandard growth
764	Vincenzo M. Tortorelli and Epifanio G. Virga,	Axis-symmetric boundary-value problems for nematic liquid crystals with variable degree of orientation
765	Nikan B. Firoozye and Robert V. Kohn,	Geometric parameters and the relaxation of multiwell energies
766	Haitao Fan and Marshall Slemrod,	The Riemann problem for systems of conservation laws of mixed type
767	Joseph D. Fehribach,	Analysis and application of a continuation method for a self-similar coupled Stefan system
768	C. Foias, M.S. Jolly, I.G. Kevrekidis and E.S. Titi,	Dissipativity of numerical schemes
769	D.D. Joseph, T.Y.J. Liao and J.-C. Saut,	Kelvin–Helmholtz mechanism for side branching in the displacement of light with heavy fluid under gravity
770	Chris Grant,	Solutions to evolution equations with near-equilibrium initial values
771	B. Cockburn, F. Coquel, Ph. LeFloch and C.W. Shu,	Convergence of finite volume methods
772	N.G. Lloyd and J.M. Pearson,	Computing centre conditions for certain cubic systems
773	João Palhoto Matos,	Young measures and the absence of fine microstructures in the $\alpha - \beta$ quartz phase transition
774	L.A. Peletier & W.C. Troy,	Self-similar solutions for infiltration of dopant into semiconductors
775	H. Scott Dumas and James A. Ellison,	Nekhoroshev’s theorem, ergodicity, and the motion of energetic charged particles in crystals
776	Stathis Filippas and Robert V. Kohn,	Refined asymptotics for the blowup of $u_t - \Delta u = u^p$.
777	Patricia Bauman, Nicholas C. Owen and Daniel Phillips,	Maximum principles and a priori estimates for an incompressible material in nonlinear elasticity
778	Patricia Bauman, Nicholas C. Owen and Daniel Phillips,	Maximal smoothness of solutions to certain Euler–Lagrange equations from nonlinear elasticity
779	Jack Carr and Robert Pego,	Self-similarity in a coarsening model in one dimension
780	J.M. Greenberg,	The shock generation problem for a discrete gas with short range repulsive forces
781	George R. Sell and Mario Taboada,	Local dissipativity and attractors for the Kuramoto–Sivashinsky equation in thin 2D domains
782	T. Subba Rao,	Analysis of nonlinear time series (and chaos) by bispectral methods
783	Nicholas Baumann, Daniel D. Joseph, Paul Mohr and Yuriko Renardy,	Vortex rings of one fluid in another free fall
784	Oscar Bruno, Avner Friedman and Fernando Reitich,	Asymptotic behavior for a coalescence problem
785	Johannes C.C. Nitsche,	Periodic surfaces which are extremal for energy functionals containing curvature functions
786	F. Abergel and J.L. Bona,	A mathematical theory for viscous, free-surface flows over a perturbed plane
787	Gunduz Caginalp and Xinfu Chen,	Phase field equations in the singular limit of sharp interface problems
788	Robert P. Gilbert and Yongzhi Xu,	An inverse problem for harmonic acoustics in stratified oceans
789	Roger Fosdick and Eric Volkmann,	Normality and convexity of the yield surface in nonlinear plasticity
790	H.S. Brown, I.G. Kevrekidis and M.S. Jolly,	A minimal model for spatio-temporal patterns in thin film flow

- 791 Chao-Nien Chen, On the uniqueness of solutions of some second order differential equations
 792 Xinfu Chen and Avner Friedman, The thermistor problem for conductivity which vanishes at large temperature
 793 Xinfu Chen and Avner Friedman, The thermistor problem with one-zero conductivity
 794 E.G. Kalnins and W. Miller, Jr., Separation of variables for the Dirac equation in Kerr Newman space time
 795 E. Knobloch, M.R.E. Proctor and N.O. Weiss, Finite-dimensional description of doubly diffusive convection
 796 V.V. Pukhnachov, Mathematical model of natural convection under low gravity
 797 M.C. Knaap, Existence and non-existence for quasi-linear elliptic equations with the p-laplacian involving
 critical Sobolev exponents
 798 Stathis Filippas and Wenxiong Liu, On the blowup of multidimensional semilinear heat equations
 799 A.M. Meirmanov, The Stefan problem with surface tension in the three dimensional case with spherical
 symmetry: non-existence of the classical solution
 800 Bo Guan and Joel Spruck, Interior gradient estimates for solutions of prescribed curvature equations
 of parabolic type
 801 Hi Jun Choe, Regularity for solutions of nonlinear variational inequalities with gradient constraints
 802 Peter Shi and Yongzhi Xu, Quasistatic linear thermoelasticity on the unit disk
 803 Satyanad Kichenassamy and Peter J. Olver, Existence and non-existence of solitary wave solutions to higher
 order model evolution equations
 804 Dening Li, Regularity of solutions for a two-phase degenerate Stefan Problem
 805 Marek Fila, Bernhard Kawohl and Howard A. Levine, Quenching for quasilinear equations
 806 Yoshikazu Giga, Shun'ichi Goto and Hitoshi Ishii, Global existence of weak solutions for interface equations
 coupled with diffusion equations
 807 Mark J. Friedman and Eusebius J. Doedel, Computational methods for global analysis of homoclinic and hetero-
 clinic orbits: a case study
 808 Mark J. Friedman, Numerical analysis and accurate computation of heteroclinic orbits in the case of center
 manifolds
 809 Peter W. Bates and Songmu Zheng, Inertial manifolds and inertial sets for the phase-field equations
 810 J. López Gómez, V. Márquez and N. Wolanski, Global behavior of positive solutions to a semilinear
 equation with a nonlinear flux condition
 811 Xinfu Chen and Fahuai Yi, Regularity of the free boundary of a continuous casting problem
 812 Eden, A., Foias, C., Nicolaenko, B. and Temam, R., Inertial sets for dissipative evolution equations
 Part I: Construction and applications
 813 Jose-Francisco Rodrigues and Boris Zaltzman, On classical solutions of the two-phase steady-state Stefan
 problem in strips
 814 Viorel Barbu and Srdjan Stojanovic, Controlling the free boundary of elliptic variational inequalities on
 a variable domain
 815 Viorel Barbu and Srdjan Stojanovic, A variational approach to a free boundary problem arising in electro-
 photography
 816 B.H. Gilding and R. Kersner, Diffusion-convection-reaction, free boundaries, and an integral equation
 817 Shoshana Kamin, Lambertus A. Peletier and Juan Luis Vazquez, On the Barenblatt equation of elasto-
 plastic filtration
 818 Avner Friedman and Bei Hu, The Stefan problem with kinetic condition at the free boundary
 819 M.A. Grinfeld, The stress driven instabilities in crystals: mathematical models and physical
 manifestations
 820 Bei Hu and Lihe Wang, A free boundary problem arising in electrophotography: solutions with connected
 toner region
 821 Yongzhi Xu, T. Craig Poling, and Trent Brundage, Direct and inverse scattering of time harmonic
 acoustic waves in an inhomogeneous shallow ocean
 822 Steven J. Altschuler, Singularities of the curve shrinking flow for space curves
 823 Steven J. Altschuler and Matthew A. Grayson, Shortening space curves and flow through singularities
 824 Tong Li, On the Riemann problem of a combustion model
 825 L.A. Peletier & W.C. Troy, Self-similar solutions for diffusion in semiconductors
 826 C.J. van Duijn, L.A. Peletier & R.J. Schotting, On the analysis of brine transport in porous media
 827 Minkyu Kwak, Finite dimensional description of convective reaction-diffusion equations
 828 Minkyu Kwak, Finite dimensional inertial forms for the 2D Navier-Stokes equations
 829 Victor A. Galaktionov and Sergey A. Posashkov, On some monotonicity in time properties for a quasilinear
 parabolic equation with source
 830 Victor A. Galaktionov, Remark on the fast diffusion equation in a ball
 831 Hi Jun Choe and Lihe Wang, A regularity theory for degenerate vector valued variational inequalities
 832 Vladimir I. Olikar and Nina N. Uraltseva, Evolution of nonparametric surfaces with speed depending on
 curvature, II. The mean curvature case.
 833 S. Kamin and W. Liu, Large time behavior of a nonlinear diffusion equation with a source
 834 Shoshana Kamin and Juan Luis Vazquez, Singular solutions of some nonlinear parabolic equations
 835 Bernhard Kawohl and Robert Kersner, On degenerate diffusion with very strong absorption