q-SERIES AND ORTHOGONAL POLYNOMIALS ASSOCIATED WITH BARNES' FIRST LEMMA

E.G. KALNINS[†] AND WILLARD MILLER, JR.^{*†}

Abstract. We exploit symmetry (recurrence relation) techniques for the derivation of properties associated with families of basic hypergeometric functions. Similar methods have been used by Nikiforov, Suslov and Uvarov. Here we apply these ideas to find new proofs of Barnes' First Lemma and some of its q -analogs. We show that these integrals correspond to the weight functions determining the orthogonality relations for Hahn, q -Hahn, and big q -Jacobi polynomials. As another example of our method we introduce a biorthogonal system of rational functions whose weight function corresponds to the q -analog of Kummer's Theorem

Key words. basic hypergeometric functions, orthogonal polynomials, Barnes' Lemma, biorthogonal functions

AMS(MOS) subject classifications. 33A65, 33A75, 39A10

1. Introduction. In papers Agarwal et.al. (1987), Kalnins and Miller (1987), Miller (1988) the authors have advocated the exploitation of symmetry (recurrence relation) techniques for the derivation of properties associated with families of basic hypergeometric functions, in analogy with the local Lie theory techniques for ordinary hypergeometric functions. In particular, we have used these ideas to give simple derivations of the orthogonality relations for the Askey-Wilson and Wilson polynomials (Askey and Wilson (1985), Wilson (1980)), and, in particular, simple evaluations of the weight function integrals that determine the normalizations of these polynomials. Similar techniques have been employed by Nikiforov, Suslov and Uvarov (1985) and Nikiforov and Suslov (1986), but they have apparently not applied them to the computation of contour integrals and summation formulas.

In Section 2 we use recurrence relations obeyed by a family of q -Hahn polynomials to derive the complex orthogonality of these polynomials and several q -analogs of Barnes' First Lemma, including those of Watson (1910) and Askey and Roy, (1987, eqn. 2.8). These integrals correspond to the square of the norm of the constant polynomial 1. Expanding one of these contour integrals by residues we obtain the real orthogonality relations for the Big q -Jacobi polynomials.

In Section 3 we carry out the corresponding computations for the limiting case $q \rightarrow 1$ — and obtain the classical Barnes' Lemma, which we now see is associated with the orthogonality of the Hahn polynomials.

[†]Mathematics Department University of Waikato, Hamilton, New Zealand

^{*}Supported in part by the National Science Foundation under grant DMS 86-00372

[‡]School of Mathematics, University of Minnesota, Minneapolis, MN 55455

In Section 4 we work out a simple but nontrivial example of the use of these ideas to derive biorthogonality relations for rational basic hypergeometric functions. The associated summation formula is the q -analog of Kummer's Theorem, originally due to Andrews (1973).

This link between recurrence relations obeyed by families of special functions, orthogonality relations for the functions, associated contour integrals and summation formulas appears capable of extensive generalization.

Most of the computations in the following sections were checked with SMP. One of the authors (W.M.) wishes to thank Dick Askey and Dennis Stanton for consultation on this work.

2. q -analogs of Barnes' Lemma. We are concerned with the q -Hahn polynomials

$$(2.1) \quad \Phi_n^{a,b,c,d}(z) = {}_3\varphi_2 \left(\begin{matrix} q^{-n}, & q^{n-1}abcd, & az \\ ac, & ad \end{matrix} ; q \right)$$

$$n = 0, 1, 2, \dots$$

where the basic hypergeometric functions ${}_{p+1}\varphi_p$ are defined as usual by

$${}_{p+1}\varphi_p \left(\begin{matrix} a_1, \dots, & a_{p+1} \\ b_1, \dots, & b_p \end{matrix} ; x \right) = \sum_{m=0}^{\infty} \frac{(a_1; q)_n \dots (a_{p+1}; q)_m x^m}{(b_1; q)_m \dots (b_p; q)_m (q; q)_m}$$

and

$$(a; q)_0 = 1$$

$$(a; q)_m = (1 - a)(1 - aq) \dots (1 - aq^{m-1}), \quad m \geq 1.$$

Initially we require $0 < |q| < 1$, $acd \neq 0$ and $|a|, |b|, |c|, |d| < 1$, but some of these conditions can be relaxed later. The polynomials obey the fundamental recurrence relations

$$(2.2A) \quad \mu^{(a,b,c,d)} \Phi_n^{(a,b,c,d)} = \frac{q^{\frac{1}{2}}}{d} \left(1 - \frac{ad}{q} \right) \Phi_n^{(aq^{-\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{-\frac{1}{2}})}$$

$$(2.2B) \quad \tau^{(a,b,c,d)} \Phi_n^{(a,b,c,d)} = \frac{a(1 - q^{-n})(1 - q^{n-1}abcd)}{q^{-\frac{1}{2}}(1 - ad)(1 - ac)} \Phi_{n-1}^{(aq^{\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{\frac{1}{2}})}$$

where

$$\mu^{(a,b,c,d)} = \frac{1}{z} \left[(1 - azq^{-\frac{1}{2}}) E_z^{\frac{1}{2}} - \left(1 - \frac{zq^{\frac{1}{2}}}{d} \right) E_z^{\frac{1}{2}} \right]$$

$$\tau^{(a,b,c,d)} = \frac{1}{z} [E_z^{\frac{1}{2}} - E_z^{-\frac{1}{2}}]$$

and $E_z^\alpha f(z) = f(q^\alpha z)$. These relations follow from

$$\begin{aligned}\mu(za; q)_n &= \frac{q^{\frac{1}{2}}}{d}(1 - adq^{n-1})(zaq^{\frac{1}{2}}; q)_n \\ \tau(za; q)_n &= \frac{a}{q^{\frac{1}{2}}}(1 - q^n)(zaq^{\frac{1}{2}}; q)_{n-1}.\end{aligned}$$

The existence of μ suggests the existence of a recurrence taking $\Phi_n^{(aq^{-\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{-\frac{1}{2}})$ to $\Phi_n^{(a,b,c,d)}$. Indeed, the appropriate operator is

$$\begin{aligned}(2.2c) \quad \mu^* &= -\frac{q^{\frac{1}{2}}}{\rho} \mu^{(bq^{\frac{1}{2}}, aq^{-\frac{1}{2}}, dq^{-\frac{1}{2}}, cq^{\frac{1}{2}}) : \\ &\quad \mu^* \Phi_n^{(aq^{-\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{-\frac{1}{2}})} \\ &= \frac{q^{\frac{1}{2}}}{\rho} \frac{(\frac{ad}{q} - q^{-n})(1 - bcq^n)}{c(1 - \frac{ad}{q})} \Phi_n^{(a,b,c,d)}, \quad \rho \neq 0,\end{aligned}$$

which follows from

$$\frac{\rho}{q^{\frac{1}{2}}} \mu^*(azq^{-\frac{1}{2}}; q)_n = (az; q)_n (bcq^n - 1) \frac{q^{-n}}{c} + (az; q)_{n-1} (acq^{n-1} - 1)(q^n - 1) \frac{q^{-n}}{c}.$$

Let $w_{a,b,c,d}(z)$ be a (complex-valued) weight function and $S_{a,b,c,d}$ be the indefinite inner product space of polynomials $f(z)$ with respect to the inner product

$$(2.3) \quad (f_1, f_2)_{a,b,c,d} = \frac{1}{2\pi i} \oint_C f_1(z) f_2(z) w_{a,b,c,d}(z) \frac{dz}{z}$$

where C is a deformation of the unit circle $|z| = 1$. Now consider $\mu = \mu^{(a,b,c,d)}$ and $\mu^* = -\frac{q^{\frac{1}{2}}}{\rho} \mu^{(bq^{\frac{1}{2}}, aq^{-\frac{1}{2}}, dq^{-\frac{1}{2}}, cq^{\frac{1}{2}})$ as mappings

$$\begin{aligned}(2.4) \quad \mu &: S_{a,b,c,d} \rightarrow S_{aq^{-\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{-\frac{1}{2}}} \\ \mu^* &: S_{aq^{-\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{-\frac{1}{2}}} \rightarrow S_{a,b,c,d}\end{aligned}$$

and determine $w_{a,b,c,d}$ so that

$$(2.5) \quad (\mu f, g)_{aq^{-\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{-\frac{1}{2}}} = (f, \mu^* g)_{a,b,c,d}$$

for all $f \in S_{a,b,c,d}$, $g \in S_{aq^{-\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{-\frac{1}{2}}}$. Condition (2.5) yields a q -difference equation for $w_{a,b,c,d}$ with solution

$$(2.6) \quad w_{a,b,c,d}(z) = \frac{(\frac{\rho z}{d}; q)_\infty (\frac{qd}{z\rho}; q)_\infty (\frac{\rho c}{z}; q)_\infty (\frac{qz}{c\rho}; q)_\infty}{(az; q)_\infty (bz; q)_\infty (\frac{c}{z}; q)_\infty (\frac{d}{z}; q)_\infty}$$

where

$$(x; q)_\infty = \lim_{n \rightarrow \infty} (x; q)_n.$$

It follows immediately that $\mu^* \mu$ is a self-adjoint operator

$$\mu^* \mu : S_{a,b,c,d} \rightarrow S_{a,b,c,d}$$

and from the recurrence relations (2.2) we have

$$(2.7) \quad \mu^* \mu \Phi_n^{(a,b,c,d)} = \lambda_n \Phi_n^{(a,b,c,d)}, \quad \lambda_n = \frac{q}{cd\rho} \left(\frac{ad}{q} - q^n \right) (1 - bcq^n).$$

Since eigenfunctions corresponding to distinct eigenvalues are orthogonal we have

$$(2.8) \quad (\Phi_n^{(a,b,c,d)}, \Phi_m^{(a,b,c,d)})_{a,b,c,d} = 0 \text{ for } m \neq n.$$

Relation (2.5) for $f = g \equiv 1$ yields

$$(2.9) \quad \|1\|_{aq^{-\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{-\frac{1}{2}}}^2 = -\frac{d}{\rho c} \frac{1 - bc}{1 - ad} \|1\|_{a,b,c,d}^2$$

where

$$\|1\|_{a,b,c,d}^2 = (1, 1)_{a,b,c,d}.$$

The symmetry of the weight function in (a, b) yields an additional relation of the form (2.9). Furthermore the obvious relation

$$(\Phi_1^{(a,b,c,d)}, \Phi_0^{(a,b,c,d)})_{a,b,c,d} = 0,$$

the explicit expression (2.1) and the property $(1, p_n)_{a,b,c,d} = \|1\|_{aq^n, b, c, d}^2$ for $p_n(z) = (az; q)_n$, yield the relation

$$(2.10) \quad \|1\|_{aq, b, c, d}^2 = \frac{(1 - ad)(1 - ac)}{(1 - abcd)} \|1\|_{a, b, c, d}^2.$$

Again, the symmetry in (a, b) gives an additional relation. The solution of these q -difference equations is

$$(2.11) \quad \|1\|_{a,b,c,d}^2 = \frac{(abcd; q)_\infty \left(\frac{c\rho}{d}; q\right)_\infty \left(\frac{qd}{c\rho}; q\right)_\infty}{(ad; q)_\infty (ac; q)_\infty (ac; q)_\infty (bc; q)_\infty (bd; q)_\infty} \mathcal{K}(\rho, q)$$

Where $\mathcal{K}(\rho, q)$ is to be determined. In the special case $a = \rho/d$, $b = q/c\rho$ we can compute the (trivial) integral directly: $\|1\|_{\rho/d, q/c\rho, c, d}^2 = 1$. Hence $\mathcal{K}(\rho, q) = (\rho; q)_\infty (q/\rho; q)_\infty / (q; q)_\infty$ and we have

$$\begin{aligned}
\|1\|_{a, b, c, d}^2 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{(\frac{\rho}{d}e^{i\theta}; q)_\infty (\frac{qd}{\rho}e^{-i\theta}; q)_\infty (\rho ce^{i\theta}; q)_\infty}{(ae^{i\theta}; q)_\infty (be^{i\theta}; q)_\infty (ce^{i\theta}; q)_\infty} \\
&\quad \cdot \frac{(\frac{q}{c\rho}e^{i\theta}; q)_\infty}{(de^{-i\theta}; q)_\infty} d\theta \\
(2.12) \quad &= \frac{(abcd; q)_\infty (\frac{c\rho}{d}; q)_\infty (\frac{qd}{c\rho}; q)_\infty (\rho; q)_\infty (\frac{q}{\rho}; q)_\infty}{(ad; q)_\infty (ac; q)_\infty (bc; q)_\infty (bd; q)_\infty (q; q)_\infty}
\end{aligned}$$

in agreement with Askey and Roy (1986).

Now we consider the recurrence

$$\tau^{(a, b, c, d)} : S_{a, b, c, d} \rightarrow S_{aq^{\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{\frac{1}{2}}},$$

(2.2B) and compute the adjoint $\tau^* \equiv \tau^{*(aq^{\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{\frac{1}{2}})}$ such that

$$(2.13) \quad (\tau f, g)_{aq^{\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{\frac{1}{2}}} = (f, \tau^* g)_{a, b, c, d}$$

for all $f \in S_{a, b, c, d}$, $g \in S_{aq^{\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{\frac{1}{2}}}$. A straightforward computation yields

$$(2.14) \quad \tau^{*(aq^{\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{\frac{1}{2}})} = \frac{1}{q^{\frac{1}{2}}z} [-(1-az)(1-bz)E_z^{\frac{1}{2}} + (1-\frac{z}{c})(1-\frac{z}{d})E_z^{\frac{1}{2}}].$$

It follows that $\tau^* \tau : S_{a, b, c, d} \rightarrow S_{a, b, c, d}$ is self-adjoint. Moreover, the action of τ^* on the polynomial basis is

$$(2.15) \quad \tau^* \Phi_{n-1}^{(aq^{\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{\frac{1}{2}})} = \frac{(1-ac)(1-ad)}{q^{\frac{1}{2}}acd} \Phi_n^{(a, b, c, d)}.$$

This follows from

$$\tau^*(aq^{\frac{1}{2}}z; q)_k = \frac{q^{-k-\frac{1}{2}}}{acd} (1-acq^k)(1-adq^k)(az; q)_k - \frac{q^{-k-\frac{1}{2}}}{acd} (1-abcdq^k)(az; q)_{k+1}.$$

Thus

$$(2.16) \quad \tau^* \tau \Phi_n^{(a, b, c, d)} = \frac{1}{cd} (1-q^{-n})(1-q^{n-1}abcd) \Phi_n^{(a, b, c, d)}.$$

Setting $f = \Phi_n^{(a, b, c, d)}$, $g = \Phi_{n-1}^{(aq^{\frac{1}{2}}, bq^{\frac{1}{2}}, cq^{\frac{1}{2}}, dq^{\frac{1}{2}})}$ in (2.13) we obtain the recurrence

$$(2.17) \quad \|\Phi_n^{(a, b, c, d)}\|_{a, b, c, d}^2 = \frac{qa^2cd(1-q^{-n})(1-q^{n-1}abcd)}{(1-ad)^2(1-ac)^2} \|\Phi_{n-1}^{(aq^{\frac{1}{2}}, \dots, dq^{\frac{1}{2}})}\|_{aq^{\frac{1}{2}}, \dots, dq^{\frac{1}{2}}}^2$$

which permits us to compute the norms $\|\Phi_n^{(a,\dots,d)}\|_{a,\dots,d}^2$ recursively from $\|1\|_{a,\dots,d}^2$. Note that the norms are all nonzero. We have shown that the q -Hahn polynomials $\{\Phi_n^{(a,b,c,d)}\}$ are uniquely characterized by their orthogonality with respect to the complex weight function $w_{a,b,c,d}$.

Since the weight function is symmetric in $\{a,b\}$ the orthogonal polynomials satisfies the transformation rule

$$(2.18.) \quad {}_3\varphi_2 \left(\begin{matrix} q^{-n}, & q^{n-1}abcd, & bz \\ bc, & dc & \end{matrix} ; q \right) = \left(\frac{b}{a} \right)_n \frac{(ac;q)_n(ad;q)_n}{(bc;q)_n(bd;q)_n} {}_3\varphi_2 \left(\begin{matrix} q^{-n}, & q^{n-1}abcd, & az \\ ac, & ad & \end{matrix} ; q \right)$$

Also, the action of $\tau^{*(aq^{\frac{1}{2}}, \dots, dq^{\frac{1}{2}})}$ determines a Rodrigues formula.

The complex orthogonality (2.8) for the q -Hahn polynomials leads to real discrete orthogonality for the Big q -Jacobi polynomials

$$(2.19) \quad {}_3\varphi_2 \left(\begin{matrix} q^{-n}, & q^{n-1+\alpha+\beta+\gamma+\delta}, & q^{x+1} \\ q^{\alpha+\gamma}, & q^{\alpha+\delta} & \end{matrix} ; q \right)$$

(We thank Dennis Stanton for pointing out this fact.) The polynomials (2.19) are orthogonal with respect to the discrete measure with mass points and corresponding weights (Ismail and Wilson (1982)):

$$(2.20) \quad \begin{aligned} x = \alpha + \delta + k - 1 & \quad \frac{(q^{\delta-\gamma+k+1}; q)_\infty (q^{k+1}; q)_\infty}{(q^{\alpha+\delta+k}; q)_\infty (q^{\beta+\delta+k}; q)_\infty} q^k q^{\alpha+\delta-1} (1-q) \\ x = \alpha + \gamma + k - 1 & \quad \frac{(q^{\gamma-\delta+k+1}; q)_\infty (q^{k+1}; q)_\infty}{(q^{\alpha+\gamma+k}; q)_\infty (q^{\beta+\gamma+k}; q)_\infty} q^k q^{\alpha+\gamma-1} (1-q), \end{aligned}$$

$k = 0, 1, 2, \dots$. To obtain this result from (2.6), (2.8) we first set $a = q^\alpha, \dots, d = q^\delta$. If $\text{Re}(\alpha, \beta, \gamma, \delta) > 0$ and there are no double poles we can expand the integral $(\Phi_n, \Phi_m)_{a,b,c,d}$ by residues, using the simple poles of $w_{a,b,c,d}(z)$, (2.6), at $z = q^{\gamma+k}, z = q^{\delta+k}$. The result of this expansion is (2.20) with $z = q^x$. In particular, the dependence on ρ cancels out.

Our approach relating orthogonal polynomials to integrals of weight functions can be used to evaluate other important integrals. One class of such integrals can be conveniently studied through the change of variables given in the preceding paragraph:

$$(2.21) \quad a = q^\alpha, b = q^\beta, c = q^\gamma, d = q^\delta, z = q^x.$$

This change reduces q -difference equations for the weight function to ordinary difference equations. Indeed the operators (2.2) now take the form

$$(2.22) \quad \begin{aligned} \mu^{(\alpha,\beta,\gamma,\delta)} &= q^{-x} [(1 - q^{\alpha+x-\frac{1}{2}}) \mathcal{E}_x^{\frac{1}{2}} - (1 - q^{x-\delta+\frac{1}{2}}) \mathcal{E}_x^{-\frac{1}{2}}] \\ \tau^{(\alpha,\beta,\gamma,\delta)} &= q^{-x} [\mathcal{E}_x^{\frac{1}{2}} - \mathcal{E}_x^{-\frac{1}{2}}] \\ \mu^* &= -q^{-x} [(1 - q^{\beta+x}) \mathcal{E}_x^{\frac{1}{2}} - (1 - q^{x-\gamma}) \mathcal{E}_x^{-\frac{1}{2}}]. \end{aligned}$$

where $\mathcal{E}_x^s g(x) = g(x + s)$.

The orthogonal functions are now polynomials in q^x . We require that the inner product take the form

$$(2.23) \quad (f_1, f_2)_{\alpha, \beta, \gamma, \delta} = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} f_1(q^x) f_2(q^x) w_{\alpha, \beta, \gamma, \delta}(x) dx$$

where the contour in the complex x -plane will run from $-i\infty$ to $+i\infty$ so that decreasing sequences of poles for w lie on the left and increasing sequences of poles lie on the right. The condition that μ^* be the adjoint of μ now becomes

$$(2.24) \quad (\mu f, g)_{\alpha - \frac{1}{2}, \beta + \frac{1}{2}, \gamma + \frac{1}{2}, \delta - \frac{1}{2}} = (f, \mu^* g)_{\alpha, \beta, \gamma, \delta}$$

with $f \in S_{\alpha, \beta, \gamma, \delta}$, $g \in S_{\alpha - \frac{1}{2}, \beta + \frac{1}{2}, \gamma + \frac{1}{2}, \delta - \frac{1}{2}}$, where $S_{\alpha, \beta, \gamma, \delta}$ is the space of polynomials in q^x with inner product (2.23). The nonunique solution for w is

$$(2.25) \quad w_{\alpha, \beta, \gamma, \delta}(x) = \frac{(q^{x-\gamma+1}; q)_{\infty} (q^{x-\delta+1}; q)_{\infty}}{(q^{x+\alpha}; q)_{\infty} (q^{x+\beta}; q)_{\infty}} q^x H(\alpha, \beta, \gamma, \delta, x)$$

where H is an analytic function of its variables satisfying the periodicity properties

$$(2.26) \quad \begin{aligned} H(\alpha, \beta, \gamma, \delta, x) &= H(\alpha, \beta, \gamma, \delta, x + 1) \\ &= H(\alpha - \frac{1}{2}, \beta - \frac{1}{2}, \gamma \pm \frac{1}{2}, \delta \mp \frac{1}{2}, x + \frac{1}{2}) \\ &= H(\alpha - \frac{1}{2}, \beta - \frac{1}{2}, \gamma \pm \frac{1}{2}, \delta \mp \frac{1}{2}, x + \frac{1}{2}) \\ &= H(\alpha - \frac{1}{2}, \beta + \frac{1}{2}, \gamma + \frac{1}{2}, \delta + \frac{1}{2}, x + \frac{1}{2}). \end{aligned}$$

One solution of (2.26) is

$$H(\alpha, \beta, \gamma, \delta, x) = \frac{\sin \pi(\gamma - \delta)}{\sin \pi(\gamma - x) \sin \pi(\delta - x)}$$

so that the weight function becomes

$$(2.27) \quad w_{\alpha, \beta, \gamma, \delta}(x) = \frac{(q^{x-\gamma+1}; q)_{\infty} (q^{x-\delta+1}; q)_{\infty} q^x \sin \pi(\gamma - \delta)}{(q^{x+\alpha}; q)_{\infty} (q^{x+\beta}; q)_{\infty} \sin \pi(\gamma - x) \sin \pi(\delta - x)}.$$

Then the polynomials are

$$(2.28) \quad \Phi_n^{(\alpha, \beta, \gamma, \delta)}(q^x) = {}_3\varphi_2 \left(\begin{matrix} q^{-n}, & q^{n+\alpha+\beta+\gamma+\delta-1}, & q^{\alpha+x} \\ q^{\alpha+\gamma}, & q^{\alpha+\delta} \end{matrix}; q \right), n = 0, 1, 2, \dots$$

and the eigenvalue equation is

$$\mu^* \mu \Phi_n^{\alpha, \beta, \gamma, \delta} = q^{\frac{1}{2} - \gamma - \delta} (q^{\alpha + \delta - 1} - q^{-n}) (1 - q^{\beta + \gamma + n}) \Phi_n^{\alpha, \beta, \gamma, \delta}.$$

We have immediately

$$(\Phi_n^{(\alpha, \dots, \delta)}, \Phi_m^{(\alpha, \dots, \delta)})_{\alpha, \dots, \delta} = 0, \quad m \neq n$$

and

$$(2.29) \quad \|1\|_{\alpha - \frac{1}{2}, \beta + \frac{1}{2}, \gamma + \frac{1}{2}, \delta - \frac{1}{2}}^2 = q^{-\frac{1}{2} + \delta - \gamma} \frac{(1 - q^{\beta + \gamma})}{(1 - q^{\alpha + \delta - 1})} \|1\|_{\alpha, \beta, \gamma, \delta}^2.$$

The symmetry of w in (α, β) leads to a similar recurrence. Also, the skew-symmetry of w in γ, δ yields a new recurrence. Finally, the orthogonality

$$(\Phi_1^{(a, \dots, \delta)}, \Phi_0^{(\alpha, \dots, \delta)}) = 0$$

leads to the recurrences

$$\|1\|_{\alpha+1, \beta, \gamma, \delta}^2 = \frac{(1 - q^{\alpha + \gamma})(1 - q^{\alpha + \delta})}{(1 - q^{\alpha + \beta + \gamma + \delta})} \|1\|_{\alpha, \beta, \gamma, \delta}^2$$

and

$$\|1\|_{\alpha, \beta, \gamma+1, \delta}^2 = -\frac{q^{\delta - \gamma}(1 - q^{\alpha + \gamma})(1 - q^{\beta + \gamma})}{(1 - q^{\alpha + \beta + \gamma + \delta})} \|1\|_{\alpha, \beta, \gamma, \delta}^2$$

with the solution

$$(2.30) \quad \|1\|_{\alpha, \beta, \gamma, \delta}^2 = \frac{(q^{\alpha + \beta + \gamma + \delta}; q)_\infty (q^{\delta - \gamma}; q)_\infty (q^{1 + \gamma - \delta}; q)_\infty q^\gamma}{(q^{\alpha + \delta}; q)_\infty (q^{\alpha + \gamma}; q)_\infty (q^{\beta + \gamma}; q)_\infty (q^{\beta + \delta}; q)_\infty} M(\alpha, \beta, \gamma, \delta)$$

where M is an analytic function of its arguments, symmetric in (α, β) and in (γ, δ) , satisfying the periodicity relations

$$(2.31) \quad \begin{aligned} M(\alpha, \beta, \gamma, \delta) &= M(\alpha + 1, \beta, \gamma, \delta) = M(\alpha, \beta, \gamma + 1, \delta) \\ &= M(\alpha - \frac{1}{2}, \beta + \frac{1}{2}, \gamma + \frac{1}{2}, \delta - \frac{1}{2}). \end{aligned}$$

To evaluate M we replace α by $\alpha + k$, k a positive integer, and rewrite the integral for $\|1\|_{\alpha+k, \beta, \gamma, \delta}^2$ in the form ($x = iy$)

$$(2.32) \quad \begin{aligned} &\frac{1}{2\pi} \int_{-\infty}^{\infty} \left[\frac{(q^{\alpha+k+\delta}; q)_\infty (q^{\alpha+k+\delta}; q)_\infty}{(q^{\alpha+\beta+\gamma+\delta+k}; q)_\infty (q^{iy+\alpha+k}; q)_\infty} \right] \frac{(q^{iy-\gamma+1}; q)_\infty}{(q^{iy+\beta}; q)_\infty} \\ &\quad \cdot \frac{(q^{iy-\delta+1}; q)_\infty q^{iy} dy}{\sin \pi(\gamma - iy) \sin \pi(\delta - iy)} \\ &= \frac{q^\gamma (q^{\delta - \gamma}; q)_\infty (q^{1 + \gamma - \delta}; q)_\infty}{(q^{\beta + \gamma}; q)_\infty (q^{\beta + \delta}; q)_\infty} M(\alpha, \beta, \gamma, \delta). \end{aligned}$$

Notice that the right-hand side of (2.32) is independent of k and that the bracketed quantity on the left goes to 1 as $k \rightarrow +\infty$. From the Lebesgue dominated convergence theorem we conclude that

$$\begin{aligned} & \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{q^{iy-\gamma+1}; q)_{\infty} (q^{iy-\delta+1}; q)_{\infty} q^{iy}}{(q^{iy+\beta}; q)_{\infty} \sin \pi(\gamma - iy) \sin \pi(\delta - iy)} dy \\ &= \frac{q^{\delta} (q^{\delta-\gamma}; q)_{\infty} (q^{1+\gamma-\delta}; q)_{\infty}}{(q^{\beta+\gamma}; q)_{\infty} (q^{\beta+\delta}; q)_{\infty}} M(\alpha, \beta, \gamma, \delta). \end{aligned}$$

It follows immediately that M is independent of α and β . Now in (2.32) set $k = 0$, $\alpha = 1 - \gamma$, $\beta = 1 - \delta$:

$$(2.33) \quad \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{q^{iy} \sin \pi(\gamma - \delta) dy}{\sin \pi(\gamma - iy) \sin \pi(\delta - iy)} = \frac{M(\gamma, \delta) q^{\gamma} (1 - q^{\delta-\gamma})}{(1 - q)(q; q)_{\infty}}.$$

The rather elementary integral on the left-hand side of (2.33) can be easily evaluated by residues and the resulting geometric series summed to yield $M(\gamma, \delta) = (q; q)_{\infty} / \pi$. Thus

$$(2.34) \quad \begin{aligned} & \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{(q^{iy-\gamma+1}; q)_{\infty} (q^{iy-\delta+1}; q)_{\infty} q^{iy} \sin \pi(\gamma - \delta)}{(q^{iy+\alpha}; q)_{\infty} (q^{iy+\beta}; q)_{\infty} \sin \pi(\gamma - iy) \sin \pi(\delta - iy)} dy \\ &= \frac{(q; q)_{\infty} q^{\gamma} (q^{\alpha+\beta+\gamma+\delta}; q)_{\infty} (q^{\delta-\gamma}; q)_{\infty} (q^{1+\gamma-\delta}; q)_{\infty}}{\pi (q^{\alpha+\delta}; q)_{\infty} (q^{\alpha+\gamma}; q)_{\infty} (q^{\beta+\gamma}; q)_{\infty} (q^{\beta+\delta}; q)_{\infty}} \end{aligned}$$

which is Watson's q -analogue of Barnes' First Lemma, Watson (1910).

By choosing other solutions H of relations (2.26) we can evaluate other integrals in the form (2.30), (2.31). For example, if $H = \cos^{-2} \pi x$ then $M = (q; q)_{\infty} q^{\frac{1}{4}} (1 - q^{\frac{1}{2}}) / 2\pi (q^{\gamma} - q^{\delta})$. However, in general one cannot evaluate M by the simple method we used for (2.34).

3. The Classical Barnes' Lemma. To see the relationship between our results and Barnes' Lemma we could, with care, let $q \rightarrow 1-$ in expression (2.12) (i.e., at the end of our construction). However, it is more instructive to take the limit $q \rightarrow 1-$ immediately and then proceed step-by-step through the argument of the proceeding section. From this point of view the functions to be considered are the Hahn polynomials

$$(3.1) \quad \Phi_n^{(\alpha, \beta, \gamma, \delta)}(x) = {}_3F_2 \left(\begin{matrix} -n, & n + \alpha + \beta + \gamma + \delta - 1, & \alpha + x \\ \alpha + \gamma, & \alpha + \delta \end{matrix}; 1 \right)$$

$$n = 0, 1, 2, \dots, \quad \alpha, \beta, \gamma, \delta > 0$$

where ${}_3F_2$ is a generalized hypergeometric function:

$${}_{p+1}F_p \left(\begin{matrix} \alpha, \dots, \alpha_{p+1} \\ \beta_1, \dots, \beta_p \end{matrix}; z \right) = \sum_{m=0}^{\infty} \frac{(\alpha_1)_m \dots (\alpha_{p+1})_m z^m}{(\beta_1)_m \dots (\beta_p)_m m!},$$

$$(\alpha)_m = \begin{cases} 1 & \text{if } m = 0 \\ \alpha(\alpha+1)\dots(\alpha+m-1) & \text{if } m \geq 1. \end{cases}$$

The recurrence relations are

$$(3.2A) \quad \mu^{(\alpha,\beta,\gamma,\delta)} \Phi_n^{(\alpha,\beta,\gamma,\delta)} = (\alpha + \delta - 1) \Phi_n^{(\alpha-\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta-\frac{1}{2})}$$

$$(3.2B) \quad \mu^{*(\alpha-\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta-\frac{1}{2})} \Phi_n^{(\alpha-\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta-\frac{1}{2})} = \frac{(n+\alpha+\delta-1)(n+\beta+\gamma)}{(\alpha+\delta-1)} \Phi_n^{(\alpha,\beta,\gamma,\delta)}$$

$$(3.2C) \quad \tau^{(\alpha,\beta,\gamma,\delta)} \Phi_n^{(\alpha,\beta,\gamma,\delta)} = \frac{-n(n+\alpha+\beta+\gamma+\delta-1)}{(\alpha+\delta)(\alpha+\gamma)} \Phi_{n-1}^{(\alpha+\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta+\frac{1}{2})}$$

$$(3.2D) \quad \tau^{*(\alpha+\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta+\frac{1}{2})} \Phi_{n-1}^{(\alpha+\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta+\frac{1}{2})} = (\alpha+\delta)(\alpha+\gamma) \Phi_n^{(\alpha,\beta,\gamma,\delta)}$$

where

$$\begin{aligned} \mu^{(\alpha,\beta,\gamma,\delta)} &= (\alpha + x - \frac{1}{2}) \mathcal{E}_x^{\frac{1}{2}} + (\delta - x - \frac{1}{2}) \mathcal{E}_x^{-\frac{1}{2}}, \\ \mu^{*(\alpha-\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta-\frac{1}{2})} &= (\beta + x) \mathcal{E}_x^{\frac{1}{2}} + (\gamma - 1) \mathcal{E}_x^{-\frac{1}{2}}, \\ \tau^{(\alpha,\beta,\gamma,\delta)} &= \mathcal{E}_x^{\frac{1}{2}} - \mathcal{E}_x^{-\frac{1}{2}} \\ \tau^{*(\alpha+\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta+\frac{1}{2})} &= (\gamma - x)(\delta - x) \mathcal{E}_x^{-\frac{1}{2}} - (\alpha + x)(\beta + x) \mathcal{E}_x^{\frac{1}{2}} \end{aligned}$$

and $\mathcal{E}_x^s g(x) = g(x+s)$.

We define a complex inner product by

$$(3.3) \quad (g_1, g_2)_{\alpha,\beta,\gamma,\delta} = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} g_1(x) g_2(x) w_{\alpha,\beta,\gamma,\delta}(x) dx$$

for polynomials g_1, g_2 where the integration path is the imaginary axis in the complex x -plane. Let $S_{\alpha,\beta,\gamma,\delta}$ be the space of polynomials with this inner product. We require that

$$(3.4) \quad (\mu f, g)_{\alpha-\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta-\frac{1}{2}} = (f, \mu^* g)_{\alpha,\beta,\gamma,\delta}$$

for all $f \in S_{\alpha,\beta,\gamma,\delta}$, $g \in S_{\alpha-\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta-\frac{1}{2}}$. In order that (3.4) hold, the weight function must satisfy a difference equation whose solution is (essentially)

$$(3.5) \quad w_{\alpha,\beta,\gamma,\delta}(x) = \Gamma(\alpha+x)\Gamma(\beta+x)\Gamma(\gamma-x)\Gamma(\delta-x)$$

where Γ is the gamma function, Whittaker and Watson (1958, Chapter XII). Here we are using the fundamental recurrence for the gamma function

$$\Gamma(z+1) = z\Gamma(z).$$

It now follows that $\mu^*\mu : S_{\alpha,\beta,\gamma,\delta} \rightarrow S_{\alpha,\beta,\gamma,\delta}$ is self-adjoint with respect to this inner product and has eigenfunctions $\Phi_n^{\alpha,\beta,\gamma,\delta}$:

$$(3.6) \quad \mu^*\mu\Phi_n^{\alpha,\beta,\gamma,\delta} = (n+\alpha+\delta-1)(n+\beta+\gamma)\Phi_n^{\alpha,\beta,\gamma,\delta}$$

Since eigenfunctions corresponding to distinct eigenvalues are orthogonal we have

$$(3.7) \quad (\Phi_n^{(\alpha,\beta,\gamma,\delta)}, \Phi_m^{(\alpha,\beta,\gamma,\delta)})_{\alpha,\beta,\gamma,\delta} = 0, n \neq m.$$

A similar computation gives

$$(3.8) \quad (\tau f, g)_{\alpha+\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta+\frac{1}{2}} = (f, \tau^*g)_{\alpha,\beta,\gamma,\delta}$$

for the same weight function and all

$$g \in S_{\alpha+\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta+\frac{1}{2}}, f \in S_{\alpha,\beta,\gamma,\delta}.$$

Thus $\tau^*\tau$ is self-adjoint on $S_{\alpha,\beta,\gamma,\delta}$ and

$$(3.9) \quad \tau^*\tau\Phi_n^{(\alpha,\beta,\gamma,\delta)} = -n(n+\alpha+\beta+\gamma+\delta-1)\Phi_n^{(\alpha,\beta,\gamma,\delta)}.$$

Setting $f = g = 1$ in (3.4) we find

$$(3.10) \quad \|1\|_{\alpha-\frac{1}{2},\beta+\frac{1}{2},\gamma+\frac{1}{2},\delta-\frac{1}{2}}^2 = \frac{(\beta+\gamma)}{(\alpha+\delta-1)} \|1\|_{\alpha,\beta,\gamma,\delta}^2.$$

Symmetry of the weight function in (α, β) and in (γ, δ) gives three more such relations. Furthermore

$$(\Phi_1^{(\alpha,\beta,\gamma,\delta)}, \Phi_0^{(\alpha,\beta,\gamma,\delta)})_{\alpha,\beta,\gamma,\delta} = 0,$$

which, from (3.1) and (3.5), implies

$$(3.11) \quad \|1\|_{\alpha+1,\beta,\gamma,\delta}^2 = \frac{(\alpha+\gamma)(\alpha+\delta)}{(\alpha+\beta+\gamma+\delta)} \|1\|_{\alpha,\beta,\gamma,\delta}^2$$

and also

$$(3.12) \quad \|1\|_{\alpha,\beta,\gamma+1,\delta}^2 = \frac{(\gamma+\alpha)(\gamma+\beta)}{(\alpha+\beta+\gamma+\delta)} \|1\|_{\alpha,\beta,\gamma,\delta}^2.$$

The symmetry of the weight function in (α, β) and in (γ, δ) gives two more such relations. It follows that

$$(3.13) \quad \|1\|_{\alpha, \beta, \gamma, \delta}^2 = \frac{\Gamma(\alpha + \gamma)\Gamma(\alpha + \delta)\Gamma(\beta + \gamma)\Gamma(\beta + \delta)}{\Gamma(\alpha + \beta + \gamma + \delta)} M(\alpha, \beta, \gamma, \delta)$$

where M is symmetric in (α, β) and in (γ, δ) , and satisfies the periodicity properties

$$(3.14) \quad \begin{aligned} M(\alpha + 1, \beta, \gamma, \delta) &= M(\alpha - \frac{1}{2}, \beta + \frac{1}{2}, \gamma + \frac{1}{2}, \delta - \frac{1}{2}) \\ &= M(\alpha, \beta, \gamma + 1, \delta) = M(\alpha, \beta, \gamma, \delta). \end{aligned}$$

To evaluate M we replace α by $\alpha + k$ and γ by $\gamma + k$, k a positive integer, and write the expression (3.13) in the form

$$(3.15) \quad \begin{aligned} \frac{1}{2\pi} \int_{-\infty}^{\infty} &\left(\frac{\Gamma(\alpha + \beta + \gamma + \delta + 2k)\Gamma(\alpha + k + iy)\Gamma(\gamma + k - iy)}{\Gamma(\alpha + \gamma + 2k)\Gamma(\alpha + \delta + k)\Gamma(\beta + \gamma + k)} \right) \\ &\cdot \Gamma(\beta + iy)\Gamma(\delta - iy)dy = \Gamma(\beta + \delta)M(\alpha, \beta, \gamma, \delta). \end{aligned}$$

From Stirling's formula, Whittaker and Watson (1958, Chapter XII), we have

$$\lim_{k \rightarrow +\infty} \frac{\Gamma(\alpha + \beta + \gamma + \delta + 2k)\Gamma(\alpha + k + iy)\Gamma(\gamma + k - iy)}{\Gamma(\alpha + \gamma + 2k)\Gamma(\alpha + \delta + k)\Gamma(\beta + \gamma + k)} = 2^{\beta + \delta}$$

and it follows easily that

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} \Gamma(\beta + iy)\Gamma(\delta - iy)dy = 2^{-(\beta + \delta)}\Gamma(\beta + \delta)M.$$

Hence, M is a constant, independent of $\alpha, \beta, \gamma, \delta$. To evaluate the constant we set $\alpha = \beta + \gamma = \delta = \frac{1}{2}$ in (3.13) and use the reflection formula for gamma functions:

$$\|1\|_{\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}}^2 = M = \pi \int_0^{\infty} \frac{dy}{\cosh^2 \pi y} = 1.$$

Thus,

$$(3.16) \quad \begin{aligned} \|1\|_{\alpha, \beta, \gamma, \delta}^2 &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \Gamma(\alpha + iy)\Gamma(\beta + iy)\Gamma(\gamma - iy)\Gamma(\delta - iy)dy \\ &= \frac{\Gamma(\alpha + \gamma)\Gamma(\alpha + \delta)\Gamma(\beta + \gamma)\Gamma(\beta + \delta)}{\Gamma(\alpha + \beta + \gamma + \delta)} \end{aligned}$$

This is Barnes' First Lemma, Bailey (1935, p.6), Slater (1966, p. 109).

In relation (3.8) we set $f = \Phi_n^{(\alpha, \beta, \gamma, \delta)}$, $g = \Phi_{n-1}^{(\alpha+\frac{1}{2}, \beta+\frac{1}{2}, \gamma+\frac{1}{2}, \delta+\frac{1}{2})}$ to obtain the recurrence

$$(3.17) \quad \|\Phi_n^{(\alpha, \beta, \gamma, \delta)}\|_{\alpha, \beta, \gamma, \delta}^2 = \frac{-n(n + \alpha + \beta + \gamma + \delta - 1)}{(\alpha + \gamma)^2(\alpha + \delta)^2} \cdot \|\Phi_{n-1}^{(\alpha+\frac{1}{2}, \beta+\frac{1}{2}, \gamma+\frac{1}{2}, \delta+\frac{1}{2})}\|_{\alpha+\frac{1}{2}, \beta+\frac{1}{2}, \gamma+\frac{1}{2}, \delta+\frac{1}{2}}^2.$$

It follows that the norms of the orthogonal polynomials are nonzero and can be computed recursively from $\|1\|_{\alpha, \beta, \gamma, \delta}^2$. Thus these polynomials are defined uniquely by their orthogonality with respect to the weight function w .

The symmetry of the weight function in (α, β) implies the identity

$$\begin{aligned} & {}_3F_2 \left(\begin{matrix} -n, & n + \alpha + \beta + \gamma + \delta - 1, & x + \beta \\ \beta + \gamma, & \beta + \delta \end{matrix} ; 1 \right) \\ &= \frac{(\alpha + \gamma)_n(\alpha + \delta)_n}{(\beta + \gamma)_n(\beta + \delta)_n} {}_3F_2 \left(\begin{matrix} -n, & n + \alpha + \beta + \gamma + \delta - 1, & x + \alpha \\ \alpha + \gamma, & \alpha + \delta \end{matrix} ; 1 \right). \end{aligned}$$

Furthermore, the symmetry of the weight function with respect to the interchanges $x \leftrightarrow -x$, $\alpha \leftrightarrow \delta$, $\beta \leftrightarrow \gamma$ implies

$$\begin{aligned} & {}_3F_2 \left(\begin{matrix} -n, & n + \alpha + \beta + \gamma + \delta - 1, & -x + \alpha \\ \alpha + \gamma, & \alpha + \delta \end{matrix} ; 1 \right) = \\ & (-1)^n \frac{(\beta + \delta)_n}{(\alpha + \gamma)_n} {}_3F_2 \left(\begin{matrix} -n, & n + \alpha + \beta + \gamma + \delta - 1, & x + \delta \\ \alpha + \delta, & \beta + \delta \end{matrix} ; 1 \right). \end{aligned}$$

4. Biorthogonality Relations. We will extend the ideas of the previous sections by considering rational functions rather than polynomials. Thus the basic object of study will be the rational function of z

$$(4.1) \quad p_n^{a,b}(z) = \frac{(az; q)_n}{(bz; q)_n}$$

rather than the polynomial $(az; q)_n$ of Section 2. Two fundamental recurrences are

$$(4.2A) \quad \mu_1 p_{n-1}^{a,bq} = \frac{(a - \frac{b\rho}{a} q^{-n+1})}{a - b} p_{n-2}^{a,b} + \frac{b(-b + \rho q^{-n+1})}{a} \frac{1}{a - b} p_n^{a,b}$$

$$(4.2B) \quad \mu_2 p_n^{a,b} = (a - b)(1 - q^n) p_{n-1}^{a,bq}$$

where

$$\mu_1 = \frac{(1 - \frac{b\rho z}{aq})}{(1 - \frac{az}{q})} E_z^{-1} + \frac{b}{a} \frac{1 - \rho z}{1 - bz} I_z$$

$$\mu_2 = \frac{q}{z}(1 - bq^{-1}z)(1 - bz)[I_z - E_z^{-1}]$$

and

$$E_z^s f(z) = f(q^s z), \quad I_z f(z) = f(z).$$

It is not difficult to verify that the eigenvalue equation

$$(4.3) \quad \mu_1 \mu_2 \Psi(z) = \lambda \Psi(z)$$

has the solutions

$$(4.4) \quad \Psi_\ell^{a,b,\rho}(z) = \sum_{k=0}^{\ell} \frac{(q^{-2\ell}; q^2)_k (\frac{b}{\rho} q^{2\ell-1}; q^2)_k (az; q)_{2k}}{(q^2; q^2)_k (\frac{a^2 q}{b\rho}; q^2)_k (bz; q)_{2k}} q^{2k}$$

corresponding to the eigenvalues

$$(4.5) \quad \lambda_\ell = \frac{b}{a}(1 - q^{-2\ell})(bq^{2\ell} - \rho q), \quad \ell = 0, 1, 2, \dots$$

Let $S_e^{a,b,\rho}$ be the complex vector space of all finite linear combinations of the functions $\{\Psi_\ell^{a,b,\rho}\}$. Consider the bilinear form

$$\langle f, g \rangle_{a,b,\rho}^e = \oint_C f(z)g(z)w_{a,b,\rho}^e(z)\frac{dz}{z}$$

where $g \in S_e^{a,b,\rho}$, w is a weight function, C is a positively oriented closed curve in the complex z -plane and $f \in \hat{S}_e^{a,b,\rho}$ (a space to be determined). We interpret $\mu_1 \mu_2$ as the map

$$\mu_1 \mu_2 : S_e^{a,b,\rho} \rightarrow S_e^{a,b,\rho}$$

and try to determine w, C , and $\hat{S}_e^{a,b,\rho}$ such that the adjoint eigenvalue equation

$$(4.7) \quad (\mu_1 \mu_2)^* \hat{\Psi}_\ell^{a,b,\rho} = \lambda_\ell \hat{\Psi}_\ell^{a,b,\rho}, \quad \ell = 0, 1, 2, \dots$$

has hypergeometric solutions $\hat{\Psi}_\ell^{a,b,\rho}$ where

$$(\mu_1 \mu_2)^* : \hat{S}_e^{a,b,\rho} \rightarrow \hat{S}_e^{a,b,\rho}.$$

An evident solution is

$$\begin{aligned} \hat{\Psi}_\ell^{a,b,\rho}(z) &= \Psi_\ell^{a,b,\rho}\left(\frac{q}{b\rho z}\right) \\ &= \sum_{k=0}^{\ell} \frac{(q^{-2\ell}; q^2)_k (\frac{b}{\rho} q^{2\ell-1}; q^2)_k (\frac{aq}{b\rho z}; q)_{2k}}{(q^2; q^2)_k (\frac{a^2 q}{b\rho}; q^2)_k (\frac{q}{\rho z}; q)_{2k}} q^{2k} \end{aligned}$$

with $\hat{S}_e^{a,b,\rho}$ as the space of all finite linear combinations of the $\{\hat{\Psi}_\ell^{a,b,\rho}\}$. The weight function must satisfy the recurrence

$$(4.9) \quad \frac{w_{a,b,\rho}^e(qz)}{w_{a,b,\rho}^e(z)} = -\frac{(1-az)(1-\frac{1}{\rho z})}{(1-\frac{a}{b\rho z})(q-bz)}.$$

This recurrence has many solutions, depending on our choice of the zeros and the poles of w in the z -plane. One of the solutions with the simplest pole structure is

$$(4.10) \quad w_{a,b,\rho}^e(z) = \frac{(bz;q)_\infty(-\rho z;q)_\infty(-\frac{q}{\rho z};q)_\infty}{(az;q)_\infty(\frac{qa}{\rho bz};q)_\infty(\rho z;q)_\infty}$$

where we assume

$$(4.11) \quad |q| < |\rho| < 1, |qa| < |\rho b|.$$

For C we take the unit circle: $|z| = 1$. We will adopt solution (4.10) in the computations to follow.

Note that

$$(4.12) \quad \mu_2 \Psi_\ell^{a,b,\rho} = \frac{q^2(a-b)(1-q^{-2\ell})(1-\frac{b}{\rho}q^{2\ell}-1)}{(1-\frac{a^2q}{b\rho})} \Theta_{\ell-1}^{a,b,\rho}$$

where

$$\begin{aligned} \Theta_{-1}^{a,b,\rho} &= 0, \\ \Theta_\ell^{a,b,\rho} &= \sum_{k=0}^{\ell} \frac{(q^{-2\ell}; q^2)_k (\frac{b}{\rho}q^{2\ell+3}; q^2)_k (az;q)_{2k+1} q^{2k}}{(q^2; q^2)_k (\frac{a^2q}{b\rho}; q^2)_k (bzq;q)_{2k+1}} \\ \ell &= 0, 1, \dots \end{aligned}$$

It follows from (4.3)-(4.5) that

$$(4.14) \quad \mu_1 \Theta_{\ell-1}^{a,b,\rho} = \frac{(a^2q - \rho b)}{qa(a-b)} \Psi_\ell^{a,b,\rho}, \quad \ell = 1, 2, \dots$$

Let $S_o^{a,b,\rho}$ be the space of all finite linear combinations of the $\{\Theta_\ell^{a,b,\rho}\}$. We have the interpretation

$$\mu_1 : S_o^{a,b,\rho} \rightarrow S_e^{a,b,\rho}, \mu_2 : S_e^{a,b,\rho} \rightarrow S_o^{a,b,\rho}.$$

Furthermore, $(\mu_1\mu_2)^*$ factors as $(\mu_1\mu_2)^* = \mu_2^*\mu_1^*$ where

$$\begin{aligned} \mu_1^* &= \frac{q^2}{\rho^2 z} (1 - \rho z) (1 - \frac{\rho z}{q}) (I_z - E_z^1), \\ \mu_2^* &= \frac{b^2 q^2}{qa^2} \frac{(1-az)}{(1-\frac{b\rho z}{a})} E_z^1 + \frac{b\rho^2}{qa} \frac{(1-\frac{bz}{q})}{(1-\frac{\rho z}{q})} I_z \end{aligned}$$

and

$$\begin{aligned}
\mu_1^* \hat{\Psi}_\ell^{a,b,\rho} &= \frac{\frac{q^3}{\rho}(\frac{a}{b} - 1)}{(1 - \frac{a^2 q}{b\rho})} (1 - q^{-2\ell}) (1 - \frac{b}{\rho} q^{2\ell-1}) \hat{\Theta}_{\ell-1}^{a,b,\rho} \\
\mu_2^* \hat{\Theta}_{\ell-1}^{a,b,\rho} &= -\frac{b^2 \rho^2 (1 - \frac{a^2 q}{b\rho})}{a q^2 (a - b)} \hat{\Psi}_\ell^{a,b,\rho}
\end{aligned}
\tag{4.15}$$

with

$$\begin{aligned}
\hat{\Theta}_{-1}^{a,b,\rho}(z) &= 0, \\
\hat{\Theta}_\ell^{a,b,\rho}(z) &= \sum_{k=0}^{\ell} \frac{(q^{-2\ell}; q^2)_k (\frac{b}{\rho} q^{2\ell+3}; q^2)_k (\frac{a q}{b \rho z}; q)_{2k+1}}{(q^2; q^2)_k (\frac{a^2 q^3}{b \rho}; q^2)_k (\frac{q^2}{\rho z}; q)_{2k+1}} q^{2k}, \\
\ell &= 0, 1, \dots
\end{aligned}
\tag{4.16}$$

Let $\hat{S}_o^{a,b,\rho}$ be the space of all finite linear combinations of the functions $\{\hat{\Theta}_\ell^{a,b,\rho}\}$. Then we have the interpretations

$$\mu_1^* : \hat{S}_e^{a,b,\ell} \rightarrow \hat{S}_o^{a,b,\rho}, \quad \mu_2^* : \hat{S}_o^{a,b,\rho} \rightarrow \hat{S}_e^{a,b,\rho}.$$

We now try to determine a weight function $w_{a,b,\rho}^o(z)$ such that the adjoint relation

$$< \mu_1^* f, g >_{a,b,\rho}^o = < f, \mu_1 g >_{a,b,\rho}^e \tag{4.17}$$

holds for all $f \in \hat{S}_e^{a,b,\rho}$, $g \in \hat{S}_o^{a,b,\rho}$, where

$$< g_1, g_2 >_{a,b,\rho}^o = \oint_C g_1(z) g_2(z) w_{a,b,\rho}^o(z) \frac{dz}{z}. \tag{4.18}$$

A straightforward computation yields

$$\begin{aligned}
w_{a,b,\rho}^o(z) &= \frac{\rho^2 b z w_{a,b,\rho}^e(z)}{a q^2 (1 - b z) (1 - \frac{\rho z}{q})} \\
&= \frac{\rho^2 b z (b q z; q)_\infty (-\rho z; q)_\infty (-\frac{q}{\rho z}; q)_\infty}{a q^2 (a z; q)_\infty (\frac{q a}{\rho b z}; q)_\infty (\frac{\rho z}{q}; q)_\infty}.
\end{aligned}
\tag{4.19}$$

We can similarly verify that the adjoint relation

$$< \mu_2^* g, f >_{a,b,\rho}^e = < g, \mu_2 f >_{a,b,\rho}^o \tag{4.20}$$

holds for all $f \in \hat{S}_e^{a,b,\rho}$, $g \in \hat{S}_o^{a,b,\rho}$.

It follows immediately from these adjoint relations and the eigenvalue equations (4.3) and (4.7) that the biorthogonality relations

$$(4.21A) \quad \langle \hat{\Psi}_\ell^{a,b,\rho}, \Psi_{\ell'}^{a,b,\rho} \rangle_{a,b,\rho}^e = 0,$$

$$(4.21B) \quad \langle \hat{\Theta}_\ell^{a,b,\rho}, \Theta_{\ell'}^{a,b,\rho} \rangle_{a,b,\rho}^o = 0,$$

hold for all $\ell \neq \ell'$. Our remaining problem is to compute the left-hand sides of expressions (4.21A), (4.21B) for $\ell = \ell'$.

One appropriate operator for this problem is

$$(4.22) \quad \xi = \frac{q}{z}(1 - bqz)\left(1 - \frac{b^2\rho z}{a^2}\right)I_z - \frac{q}{z} \frac{(1 - bz)(1 - bqz)\left(1 - \frac{b\rho z}{aq}\right)}{\left(1 - \frac{az}{q}\right)}E_z^{-1}$$

which satisfies

$$(4.23) \quad \begin{aligned} \xi p_{2k+1}^{a,bq} &= (a - bq) \frac{b\rho}{a^2} \left(1 - \frac{q^{2k+1}a^2}{b\rho}\right) p_{2k}^{a,bq^2}, \\ k &= 0, 1, 2, \dots \end{aligned}$$

where the basis functions $p_n^{a,b}$ are defined by (4.1). It follows that

$$\begin{aligned} \xi \Theta_\ell^{a,b,\rho} &= -q(a - bq) \left(1 - \frac{b\rho}{a^2q}\right) \Psi_\ell^{a,bq^2,\rho q^{-2}} \\ \ell &= 0, 1, \dots \end{aligned}$$

and that ξ has the interpretation

$$\xi : S_o^{a,b,\rho} \rightarrow S_e^{a,bq^2,\rho q^{-2}}.$$

We define the adjoint operator

$$\xi^* : \hat{S}_e^{a,bq^2,\rho q^{-2}} \rightarrow \hat{S}_o^{a,b,\rho}$$

by

$$(4.25) \quad \langle g, \xi f \rangle_{a,bq^2,\rho q^{-2}}^e = \langle \xi^* g, f \rangle_{a,b,\rho}^o$$

where $g \in \hat{S}_e^{a,bq^2,\rho q^{-2}}$, $f \in S_o^{a,b,\rho}$. A straightforward computation yields

$$(4.26) \quad \xi^* = qE_z^1 + \frac{a}{b} \frac{\left(1 - \frac{b^2\rho z}{a^2}\right)}{\left(1 - \frac{\rho z}{q^2}\right)} I_z$$

and

$$(4.27) \quad \frac{(a-bq)}{q} \xi^* \hat{\Psi}_\ell^{a,bq^2,pq^{-2}} = -\frac{q^2}{a} \frac{(1 - \frac{a^2 q^{2\ell+1}}{b\rho})}{(1 - \frac{a^2 q}{b\rho})} (b^2 - a^2 q^{-2\ell-2}) \hat{\Theta}_\ell^{a,b,\rho}.$$

Set

$$(4.28) \quad \begin{aligned} \|\Psi_\ell^{a,b,\rho}\|_e^2 &= \langle \hat{\Psi}_\ell^{a,b,\rho}, \Psi_\ell^{a,b,\rho} \rangle_{a,b,\rho}^e \\ \|\Theta_\ell^{a,b,\rho}\|_o^2 &= \langle \hat{\Theta}_\ell^{a,b,\rho}, \Theta_\ell^{a,b,\rho} \rangle_{a,b,\rho}^o \\ \ell &= 0, 1, 2, \dots \end{aligned}$$

From (4.14), (4.15) and (4.17) with $f = \hat{\Psi}_\ell^{a,b,\rho}$, $g = \Theta_{\ell-1}^{a,b,\rho}$ we have

$$(4.29) \quad \|\Psi_\ell^{a,b,\rho}\|_e^2 = -\frac{a\rho q^2(a-b)^2(1-q^{-2\ell})(1-\frac{b}{\rho}q^{2\ell-1})}{(b\rho-a^2q)^2} \|\Psi_{\ell-1}^{a,b,\rho}\|_o^2.$$

Relation (4.20) yields the same recurrence. Expressions (4.24), (4.25) and (4.27) with $f = \Theta_\ell^{a,b,\rho}$, $g = \hat{\Psi}_\ell^{a,bq^2,\rho q^{-2}}$ produce

$$(4.30) \quad \|\Psi_\ell^{a,bq^2,\rho q^{-2}}\|_e^2 = -\frac{a^2 q^3(1 - \frac{a^2 q^{2\ell+1}}{b\rho})(b^2 - a^2 q^{-2\ell-2})}{b\rho(a-bq)^2(1 - \frac{a^2 q}{b\rho})^2} \|\Theta_\ell^{a,b,\rho}\|_o^2.$$

It follows from these results that if we know $\|1^{a,b,\rho}\|_e^2$ for all a, b, ρ then we can compute recursively expressions (4.28) for all ℓ . In general these norms will be nonzero.

Replacing ρz by z in (4.6), (4.10) we have

$$(4.31) \quad \|1^{a,b,\rho}\|_e^2 = G(u, v, q) = \frac{1}{2\pi} \oint_C \frac{(\frac{uz}{v}; q)_\infty (-z; q)_\infty (-\frac{q}{z}; q)_\infty}{(uz; q)_\infty (\frac{qv}{z}; q)_\infty (z; q)_\infty} \frac{dz}{z}$$

where $u = a/\rho$, $v = a/b$. From the relation

$$\langle \hat{\Psi}_0^{a,b,\rho}, \Psi_1^{a,b,\rho} \rangle_e = 0$$

it is straightforward to compute the recurrence

$$\|1^{aq^2,bq^2,\rho}\|_e^2 = \frac{(1 - \frac{a^2 q}{b\rho})}{(1 - \frac{bq}{\rho})} \|1^{a,b,\rho}\|_e^2$$

or $G(q^2u, v, q) = (1 - uvq)G(u, v, q)/(1 - uq/v)$, which implies

$$(4.32) \quad G(u, v, q) = \frac{(\frac{uq}{v}; q^2)_\infty}{(uvq; q^2)_\infty} \mathcal{G}(v, q).$$

To finish the computation of (4.31) we utilize the recurrence operator

$$(4.33) \quad \eta = \frac{(aq - b)}{(1 - az)} \left[\left(1 - \frac{bz}{q}\right) E_z^{-1} + \frac{b}{aq} \left(1 - \frac{a^2 qz}{b}\right) I_z \right].$$

Here,

$$\eta p_n^{a,b} = aq(1 - q^{-n}) p_{n-2}^{a,b} + \frac{(-b^2 + a^2 q^{2-n})}{aq} p_n^{a,b},$$

hence

$$(4.34) \quad \eta \Psi_\ell^{a,b,\rho} = - \frac{q^{-2\ell} (b - \frac{a^2}{\rho} q^{2\ell+1})}{abq(1 - \frac{a^2 q}{b\rho})} (-a^2 q^2 + b^2 q^{2\ell}) \Psi_\ell^{a,b,\rho}.$$

Interpreting $\eta : S_e^{a,b,\rho} \rightarrow S_e^{aq,b,\rho}$ we compute the adjoint $\eta^* : \hat{S}_e^{aq,b,\rho} \rightarrow \hat{S}_e^{a,b,\rho}$ with the result

$$(4.35) \quad \eta^* = (aq - b) \left[\frac{(1 - \rho z)}{\rho z} (1 - az) E_z^1 + \frac{b}{aq} \left(1 - \frac{qa}{\rho bz}\right) \left(1 - \frac{a^2 qz}{b}\right) I_z \right].$$

This leads to the recurrence

$$(4.36) \quad \eta^* \hat{\Psi}_\ell^{aq,b,\rho} = - \frac{(aq - b)^2}{aq} \left(1 - \frac{a^2 q}{b\rho}\right) \hat{\Psi}_\ell^{a,b,\rho}.$$

Thus the relation

$$\langle \eta^* 1, 1 \rangle_{a,b,\rho}^e = \langle 1, \eta 1 \rangle_{aq,b,\rho}^e$$

implies

$$\|1^{aq,b,\rho}\|_e^2 = \left(1 - \frac{a^2 q}{bm}\right) \left(\frac{b - aq}{b + aq}\right) \|1^{a,b,\rho}\|_e^2$$

or $\mathcal{G}(qv, q) = (1 - qv)\mathcal{G}(v, q)/(1 + qv)$. We conclude that $\mathcal{G}(v, q) = (-vq; q)_\infty \mathcal{K}(q)/(vq; q)_\infty$ where $\mathcal{K}(q)$ is to be determined. Setting $u = 0, v = 1$ in (4.31) we find $G(0, -1, q) = 1 = (q; q)_\infty \mathcal{K}(q)/(-q; q)_\infty$, so

$$(4.37) \quad \|1^{a,b,\rho}\|_e^2 = \frac{(-q; q)_\infty (\frac{bq}{\rho}; q^2)_\infty (-\frac{aq}{b}; q)_\infty}{(q; q)_\infty (\frac{a^2 q}{b\rho}; q^2)_\infty (\frac{aq}{b}; q)_\infty}.$$

Expanding the original integral $\|1^{a,b,\rho}\|_e^2$ by residues inside the circle C , we see that our result is equivalent to the summation formula

$$(4.38) \quad {}_2\varphi_1 \left(\begin{matrix} vq, & uvq \\ uq \end{matrix}; -\frac{1}{v} \right) = \frac{(-q; q)_\infty (\frac{uq}{v}; q^2)_\infty (uvq^2; q^2)_\infty}{(uq; q)_\infty (-\frac{1}{v}; q)_\infty}.$$

This is a q -analog of Kummer's Theorem, first proved by Andrews (1973), see also Andrews (1977, page 20).

REFERENCES

- [1] A.K. AGARWAL, E.G. KALNINS AND W. MILLER, *Canonical equations and symmetry techniques for q -series*, SIAM J. Math. Anal., (to appear) (),.
- [2] G.E. ANDREWS, *On the q -analogue of Kummer's theorem and applications*, Duke Math. J., 40 (1973), 525–528.
- [3] G.E. ANDREWS, “The Theory of Partitions,” Addison-Wesley, Reading, Mass, 1977.
- [4] R. ASKEY AND R. ROY, *More q -beta integrals*, Rocky Mtn. J. Math, 26 (1986), 365–372.
- [5] R. ASKEY AND J. WILSON, *Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials*, Memoirs of the AMS, no. 319 (1985),.
- [6] W.N. BAILEY, “Generalized Hypergeometric Series,” Cambridge University Press, 1935, Reprinted by Stechert-Hafner, New York, 1964.
- [7] M.E.-H. ISMAIL AND J.A. WILSON, *Asymptotic and generating relations for the q -Jacobi and ${}_4\varphi_3$ polynomials*, J. Approx. Th., 36 (1982), 43–54.
- [8] E.G. KALNINS AND W. MILLER, *Symmetry techniques for q -series: Askey-Wilson polynomials*, Rocky Mtn. J. Math., (to appear) (),.
- [9] W. MILLER, *A note on Wilson polynomials*, SIAM J. Math. Anal, (to appear) (),.
- [10] A.F. NIKIFOROV AND S.K. SUSLOV, *Classical orthogonal polynomials of a discrete variable on nonuniform lattices*, Letters in Mathematical Physics, 11 (1986), 27–34.
- [11] A. F. NIKIFOROV, S.K. SUSLOV AND V.B. UVAROV, “Classical Orthogonal Polynomials of a Discrete Variable,” Nauka, Moscow (in Russian), 1985.
- [12] L.J. SLATER, “Generalized Hypergeometric Functions,” Cambridge University Press, Cambridge, 1966.
- [13] G.N. WATSON, *The continuation of functions defined by generalized hypergeometric series*, Trans Cambridge Phil. Soc, 21 (1910), 281–299.
- [14] E.T. WHITTAKER AND G.N. WATSON, “A Course in Modern Analysis,” Cambridge University Press, Cambridge, 1958.
- [15] J.A. WILSON, *Some hypergeometric orthogonal polynomials*, SIAM J. Math. Anal, 11 (1980), 690–701. ■

MAIN SHOP ☒		Job N ^o 89561	
Sub. Acct.	Fund	Dept. No.	Budget
	0624	5381	02
Department I.M.A.			
Date of Order 4/14		Date Wanted 4/28	AM ☐ PM ☐
For further questions contact: Patricia			Phone: 46869
Call when ready for pickup: ☐			
Phone: Name:			

BINDERY ONLY ☐ OFFSET ☒	Deliver to:
STOCK ONLY ☐ COMPOSITION ☐	545 Vincent Hall

Job Title: Reprint Series # 306
Quantity & Description: 200-20 + covers
INK/covers: Rembrandt Brown
Run Size 8 1/2 x 11 Trim Size x Finish Size 8 1/2 x 11 One Side ☐ Two Sides ☒

PAPER PRINT CARRIERS	Body	White 20# Sulphite 8 1/2 x 11	Other:
		White 60# OFFSET 8 1/2 x 11	
	Cover	Kind Sundance Weight 65# Color Desert Bronze Size	
		Elec. Masters ☒ Remarks:	
		Megas ☐	
		Metal Plts. ☐	

BINDERY ☒
COLLATE ☒
BINDING: GBC Binding ☐ Perfect ☐ Tape ☐ Shrink/Plastic Wrap ☐ Banding ☐
STITCHING: Left Corner ☐ 2 on Side ☒ Saddle ☐
FOLDING: For #10's ☐ 1/2 Fold ☐ OTHER:
OTHER: Drill ☐ 2-hole ☐ 3-hole ☐ Perforate ☐ Score ☐ Count in 100's ☐
Cutting ☐ Trimming ☐ Pad: NCR ☐ Glue ☐ Pads of sheets/sets

Additional Instructions: Carol W. - (Punch) - left 54

q -SERIES AND ORTHOGONAL POLYNOMIALS
ASSOCIATED WITH BARNES' FIRST LEMMA

BY

E.G. KALNINS

AND

WILLARD MILLER, JR.

IMA Preprint Series # 306

April 1987

INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
UNIVERSITY OF MINNESOTA
514 Vincent Hall
206 Church Street S.E.
Minneapolis, Minnesota 55455

- Recent IMA Preprints (continued)
#	Author(s)	Title
165	D. Arnold and R.S. Falk,	Continuous Dependence on the Elastic Coefficients for a Class of Anisotropic Materials
166	I.J. Bakelman,	The Boundary Value Problems for Non-Linear Elliptic Equation and the Maximum Principle for Euler-Lagrange Equations
167	Ingo Müller,	Gases and Rubbers
168	Ingo Müller,	Pseudoelasticity in Shape Memory Alloys - an Extreme Case of Thermoelasticity
169	Luigi Magalhães,	Persistence and Smoothness of Hyperbolic Invariant Manifolds for Functional Differential Equations
170	A. Damiljan and M. Vogelius,	Homogenization Limits of the Equations of Elasticity in Thin Domains
171	H.C. Simpson and S.J. Spector,	On Hadamard Stability in Finite Elasticity
172	J.L. Vazquez and C. Yaver,	Isolated Singularities of the Solutions of the Schrödinger Equation with a Radial Potential
173	G. Dal Maso and U. Mosco,	Wiener's Criterion and Γ -Convergence
174	John H. Maddocks,	Stability and Folds
175	R. Hardt and D. Kinderlehrer,	Existence and Partial Regularity of Static Liquid Crystal Configurations
176	M. Marzuke,	Construction of Smooth Ergodic Cocycles for Systems with Fast Periodic Approximations
177	J.L. Ericksen,	Stable Equilibrium Configurations of Elastic Crystals
178	Patricio Aviles,	Local Behavior of Solutions of Some Elliptic Equations
179	S.-M. Chow and R. Lauterbach,	A Bifurcation Theorem for Critical Points of Variational Problems
180	R. Pego,	Phase Transitions: Stability and Admissibility in One Dimensional Nonlinear Viscoelasticity
181	Mariano Giletti,	Quadratic Functions and Partial Regularity
182	J. Bone,	Fully Discrete Galerkin Methods for the Korteweg De Vries Equation
183	J. Maddocks and J. Keller,	Mechanics of Robes
184	F. Bernis,	Qualitative Properties for some nonlinear higher order Nonlinear Higher Order Parabolic Equations with Absorption
185	S. Reichelstein and S. Reiter,	Game Forms with Minimal Strategy Spaces
186	J. Rubinstein and R. Mouril,	Dispersion and Convection in Periodic Media
187	W.H. Fleming and P.E. Souganidis,	Asymptotic Series and the Method of Vanishing Viscosity
188	H. Bofroo Da Veiga,	Existence and Asymptotic Behavior for Strong Solutions of Navier-Stokes Equations in the Whole Space
189	L.A. Caffarelli, J.L. Vazquez, and N.I. Wolanski,	Lipschitz Continuity of Solutions and Interfaces of the N-Dimensional Porous Medium Equation
190	R. Johnson,	m -functions and Floquet Exponents for Linear Differential Systems
191	F.V. Atkinson and L.A. Peletier,	Ground States and Dirichlet Problems for $- \Delta = F(U)$ in R^n
192	G. Dal Maso, U. Mosco,	The Wiener Modulus of a Radial Measure
193	H. A. Levine and H.F. Weinberger,	Inequalities between Dirichlet and Neumann Eigenvalues
194	J. Rubinstein,	On the Macroscopic Description of Slow Viscous Flow Past a Dirichlet Problem
195	G. Dal Maso and U. Mosco,	Wiener Criteria and Energy Decay for Relaxed Dirichlet Problems
196	V. Olliker and P. Wiltman,	On the Monge-Ampere Equation Arising in the Reflect Mapping Problem
197	M. Chipot, D. Kinderlehrer and L. Caffarelli,	Some Smoothness Properties of Linear Laminae
198	Y. Iijima and R. Kohn,	Characterizing Blow-up Using Similarity Variables
199	P. Cannarsa and H. M. Soner,	On the Singularities of the Viscosity Solutions to Hamilton-Jacobi-Bellman Equations
200	Andrew Majda,	Nonlinear Geometric Optics for Hyperbolic Systems of Conservation Laws
201	G. Buttazzo, G. Dal Maso and U. Mosco,	A Derivation Theorem for Capacities with Respect to a Radon Measure
202	S. Cowin, M. Mehraiebi,	On the Identification of Material Symmetry for Anisotropic Elastic Materials
203	R.M.R. Darling,	Constructing Nonhomomorphic Stochastic Flows.
204	M. Chipot,	On the Reynolds Lubrication Equation
205	R.V. Kohn and G.M. Wiltman,	On Bounding the Effective Conductivity of Anisotropic Composites
206	I.J. Bakelman,	Notes Concerning the Torsion of Hardening Rods and Its N-Dimensional Generalizations
207	I.J. Bakelman,	The Boundary Value Problems for Non-Linear Elliptic Equations
208	Guanglu Gong and Mingpin Qian,	On the Large Deviation Functions of Markov Chains
209	Arlo Leizerowitz,	Control Problems with Random and Progressively Known Targets
210	R.M.R. Darling,	Ergodicity of a Measure-Valued Markov Chain Induced by Random Transformations
211	G. Gong, M. Qian & Zhongxin Zhao,	Killed Diffusions and Its Conditioning
212	Arlo Leizerowitz,	Controlling Diffusion Processes on Infinite Horizon with the Overtaking Criterion
213	Willard Beatty,	The Poisson Function of Finite Elasticity
214	David Terman,	Travelling Wave Solutions Arising From a Combustion Model
215	Yuh-Jia Lee,	Sharp Inequalities and Regularity of Heat Semi-Group on Infinite Dimensional Spaces
216	D. Stroock,	Lecture Notes
217	Claudio Canuto,	Spectral Methods and Maximum Principle
218	Thomas O'Brien,	A Two Parameter Family of Pension Contribution Functions and Stochastic Optimization
219	Takeyuki Hida,	Analysis of Brownian Functionals
220	Leonid Murty,	On Informational Decentralization and Efficiency of Resource Allocation Mechanisms
221	E.B. Fabes and D.M. Stroock,	A New Proof of Moser's Parabolic Harnack Inequality via the Old Ideas of Nash
222	Minoru Murata,	Structure of Positive Solution to $(- \Delta + V)u = 0$ in R^n
223	Paul Dupuis,	Large Deviations Analysis of Reflected Diffusion
224	F. Bernis,	Existence Results for Doubly Nonlinear Higher Order Parabolic Equations on Unbounded Domains.
225	S. Grey and S. Pelikan,	Large Deviations Principles for Stationary Processes
226	R. Gulliver and S. Hildebrandt,	Boundary Configurations Spanning Continuous of Minimal Surfaces.
227	J. Baxter, G. Dal Maso, U. Mosco,	Stopping Times and Γ -Convergence.
228	Julio Bouilliet,	Self-Similar Solutions, Having Jumps and Intervals of Constancy of a Diffusion-heat Conduction Equation
229	R. Hardt, D. Kinderlehrer & F.-H. Lin,	A Remark About the Stability of Smooth Equilibrium Configurations of Static Liquid Crystal
230	M. Chipot and M. Luskin,	The Compressible Reynolds Lubrication Equation
231	J.H. Maddocks,	A Model for Disclinations in Nematic Liquid Crystal
232	C. Foles, G.R. Sell and R. Temam,	Inertial Manifolds for Nonlinear Evolutionary Equations
233	P.L. Chow,	Expectation Functionals Associated with Some Stochastic Evolution
234	Giuseppe Buttazzo,	Reinforcement by a Thin Layer with Oscillating Thickness
235	W.H. Fleming, S.J. Sheu and H.M. Soner,	On Existence of the Dominant Eigenfunction and Its Application to the Large Deviation Properties of Ergodic Markov Processes
236	R. Jensen and P.E. Souganidis,	A Regularity Result for Viscosity Solutions of Hamilton-Jacobi Equations in one Space Dimension
237	B. Boczar-Karakl, J.L. Bone and D.L. Cohen,	Interaction of Shallow-Water Waves and Bottom Topography
238	F. Colonius and W. Klemann,	Infinite Time Optimal Control and Periodic Dirichlet Problems

#	Author(s)	Title	Recent IMA Preprints (continued)	#	Author(s)	Title
2241	Herry Kesten,	Scaling Relations for 2D-Percolation	277	Cockburn, B.	The Quasi-Monotone Schemes for Scalar Conservation Laws Part II	
2242	A. Leizorowicz,	Infinite Horizon Optimization for Markov Process with Finite States Spaces	278	M. Chipot, T. Sideris,	On the Abelian Higgs Model	
2243	Lois H.Y. Chen,	The Rate of Convergence in A Central Limit Theorem for Dependent Random Variables with Arbitrary Index Set	279	C. Fofas, B. Nicolaenko, G.R.Sell, R. Temam,	Inertial Manifolds for the Kuramoto-Sivashinsky Equation and an Estimate of their Lowest Dimension	
2244	G. Kallianpur,	Stochastic Differential Equations in Duals of Nuclear Spaces with some Applications	280	R. Duran,	On the Approximation of Miscible Displacement in Porous Media by a Method of Characteristics Combined with a Mixed Method	
2245	Tzuu-Shuh Chiang, Yunshong Chow and Yuh-Jia Lee,	Evaluation of Certain Functional Integrals	281	H. Aixiang, Zhang Bo,	The Convergence for Nodal Expansion Method	
2246	L. Karp and M. Pinsky,	The First Eigenvalue of a Small Geodesic Ball in a Riemannian Manifold	282	V. Tverskiy,	Dispersive Bulk Parameters for Coherent Propagation in Correlated Random Distributions	
2247	Chi-Sing Man,	Towards An Acoustoelastic Theory for Measurement of Residual Stress	283	W.-T.F. den Hollander,	Mixing Properties for Random Walk in Random Scenery	
2248	Andreas Stoll,	Invariance Principles for Brownian Intersection	284	H.-R. Jausslin,	Nondifferentiable Potentials for Nonequilibrium Steady States	
2249	R.W.R. Darling,	Local Time and Polymer Measures	285	K. Meyer, G.R.Sell,	Homoclinic Orbits and Bernoulli Bundles in Almost Period Systems	
2250	R. Cohen, R. Hardt, D. Kinderlehrer, S.Y.Lin, M. Luskin,	Minimum Energy for Liquid Crystals: Computational Results	286	J. Douglas, Jr., Y. Yuan,	Finite Difference Methods for the Transient Behavior of a Semiconductor Device	
2251	Suzanne M. Lenhart,	Viscosity Solutions for Weakly Coupled Systems of First Order PDEs	287	Li Kaitai, Yan Ningning,	The Extrapolation for Boundary Finite Elements	
2252	M. Cranston, E. Fabes, Z. Zhao,	Condition Gauge and Potential Theory for the Schrödinger Operator	288	R. Durrett, R.H. Schonmann,	Stochastic Growth Models	
2253	H. Brezis, J.-M.Coron, E.H. Lieb,	Harmonic Maps with Defects	289	David Kinderlehrer,	Remarks about Equilibrium Configurations of Crystals	
2254	A. Carverhill,	Flows of Stochastic Dynamical Systems: Nontriviality of the Lyapunov Spectrum	290	D.G. Aronson, J.L. Vazquez,	Eventual C^∞ -Regularity and Concavity for Flows in One-Dimensional Porous Media	
2255	A. Carverhill,	Conditioning A 'Lifted' Stochastic System in a Product Case	291	L.R. Scott, J.M. Boyle, B. Bagheri,	Distributed Data Structures for Scientific Computation	
2256	R.J. Williams,	Local Time and Excursions of Reflected Brownian Motion	292	J. Douglas, Jr., P.-J.Paes Leme, T. Arbogast, T. Schmitt,	Simulation of Flow in Naturally Fractured Petroleum Reservoirs	
2257	H. Föllmer, S. Grey,	Large Deviations for the Empirical Field of a Gibbs Measure	293	D.G. Aronson, L.A. Caffarelli,	Optimal Regularity for One-Dimensional Porous Medium Flow	
2258	A. Leizorowicz,	Characterization of Optimal Trajectories On an Infinite Horizon	294	Haim Brezis,	Liquid Crystals and Energy Estimates for S^2 -Valued Maps	
2259	Y.Giga, T. Miyakawa, H. Osada,	Two Dimensional Navier Stokes Flow with Measures As Initial Vorticity	295	Todd Arbogast,	Analysis of the Simulation of Single Phase Flow Through a Naturally Fractured Reservoir	
2260	M. Chipot, V. Ollier,	Sur Une Propriété Des Fonctions Propres De L'Opérateur De Laplace Beltrami	296	He Yiminan, Li Kaitai,	The Coupling Method of Finite Elements and Boundary Elements for Radiation Problems	
2261	V. Perez-Abreu,	Decompositions of Semimartingales On Duals of Countably Nuclear Spaces	297	T. Cazenave, A. Harauz, L. Vazquez, F.B. Weissler,	Nonlinear Effects in the Wave Equation with a Cubic Restoring Force	
2262	J.M. Ball,	Does Rank-One Convexity Imply Quasiconvexity	298	M. Chipot, F.B. Weissler,	Some Blow Up Results for a Nonlinear Parabolic Parabolic Equation with a Gradient Term	
2263	B. Cockburn,	The Quasi-Monotone Schemes for Scalar Conservation Laws. Part I	299	Li Kaitai,	Perturbation Solutions of Simple and Double Bifurcation Problems for Navier-Stokes Equations	
2264	K.A. Perlick-Spector,	On Radially Symmetric Simple Waves in Elasticity	300	Chen Zhongxin, Li Kaitai,	The Convergence on the Multigrid Algorithm for Navier-Stokes Equations	
2265	P.N. Shivakumar, Chi-Sing Men, Simon W. Rabkin,	Modelling of the Heart and Pericardium at End-Diastole	301	A. Gerardi, G. Nappo,	Martingale Approach for Modelling DNA Synthesis	
2266	Jose-Luis Menaldi,	Probabilistic View of Estimates for Finite Difference Methods	302	D.N. Arnold, L. Ridgway Scott Michael Vogelius,	Regular Inversion of the Divergence Operator with Dirichlet Boundary Conditions on a Polygon	
2267	Robert Hardt, Harold Rosenberg,	Open Book Structures and Unicity of Minimal Submanifolds	303	Duran, Ricardo G.	Error Analysis in L^p , $1 \leq p < \infty$, for Mixed Definite Element Methods for Linear and Quasi-Linear Elliptic Problems	
2268	Bernardo Cockburn,	The Quasi-Monotone Schemes for Scalar Conservation Laws	304	Nochetto, Ricardo H., Verdi, Claudio,	An Efficient Linear Scheme to Approximate Parabolic Free Boundary Problems: Error Estimates and Implementation	
2269	H.R. Jausslin, M. Zimmermann, Jr	Dynamics of a Model for an Ac Josephson Effect in Superfluid ^4He	305	K.A. Perlick-Spector, Scott J. Spector,	Nonuniqueness for a Hyperbolic System	
2270	A.K. Kapla,	Introductory Lecture on Reacting Flows	306	E.G. Kalnins, Willard Miller, Jr.,	q-Series and Orthogonal Polynomials Associated with Barnes' First Lemma	
2271	J.C. Taylor,	Do Minimal Solutions of Heat Equations Characterize Diffusions?	307	D.N. Arnold, Richard S. Falk,	A Uniformly Accurate Finite Element Method for Mindlin-Reissner Plate	
2272	J.C. Taylor,	The Minimal Eigenfunctions Characterize the Ornstein-Uhlenbeck Process	308	Chi-Wang Shu,	TVD Properties of a Class of Modified Eno Schemes for Scalar Conservation Laws	
2273	Chi-Sing Man, Quan-Xin Sun,	On the Significance of Normal Stress Effects in the Flow of Glaciers	309	Edmund Dikow, Ulrich Horning,	A Random Boundary Value Problem Modeling Spatial Variability in Porous Media Flow	
2274	Omar Hijab,	On Partially Observed Control of Markov Processes	310	Jack K. Hale,	Compact Attractors and Singular Perturbations	
2275	Lawrence Gray,	The Behavior of Processes with Statistical Mechanical Properties	311	Bourgeat, A. Cockburn, B.	The TVD-Projection Method for Solving Implicit Numerical Schemes for Scalar Conservation Laws: A Numerical Study of a Simple Case	
2276	R. Hardt, D. Kinderlehrer, M. Luskin,	Remarks About the Mathematical Theory of Liquid Crystals				