**ON THE SPREAD OF CONTINUOUS-TIME
LINEAR SYSTEMS**

By

Avner Friedman

and

Michael L. Honig

IMA Preprint Series # 412

April 1988

ON THE SPREAD OF CONTINUOUS-TIME LINEAR SYSTEMS*

AVNER FRIEDMAN† AND MICHAEL L. HONIG‡

Abstract. Given the impulse response h of a linear time invariant system, we consider signals $y = h * u$ with inputs u subject to $|u(t)| \leq 1$ and ask: for a given $\tau > 0$ and $y(t_0)$, what is the set of all the possible values (the "spread") of $y(t_0 + \tau)$? We characterize this set, study its properties, and compute it for some functions h .

§1. Basic definitions and results. Let $h(t)$ be a prescribed continuous function defined for $0 \leq t < \infty$ and belonging to $L^1(0, \infty)$; we refer to it as a *an impulse response*. Let $u(t)$ be any measurable function for $0 \leq t < \infty$ satisfying $|u(t)| \leq 1$; we refer to it as an *input*. The function $y(t)$, defined

$$y(t) = \int_0^t h(t-s)u(s) \, ds,$$

is called the *output* or the *signal*.

Given $\alpha \in \mathbb{R}$, $t_0 \geq 0$, $\tau > 0$, we would like to estimate the range of the output $y(t)$ at time $t = t_0 + \tau$, given that $y(t_0) = \alpha$. More quantitatively, we wish to bound the numbers

$$\begin{aligned}\tilde{\sigma}^+(\alpha, \tau, t_0) &= \sup_u \{y(t_0 + \tau); \text{ given } y(t_0) = \alpha\}, \\ \tilde{\sigma}^-(\alpha, \tau, t_0) &= \inf_u \{y(t_0 + \tau); \text{ given } y(t_0) = \alpha\}.\end{aligned}$$

Introduce the class of control functions

$$(1.1) \quad K_{\tau, \alpha} = \{u \in L^\infty(-\infty, \tau), \quad -1 \leq u(s) \leq 1, \int_{-\infty}^0 h(-s)u(s) \, ds = \alpha\}$$

and the functional

$$(1.2) \quad J_\tau(u) = \int_{-\infty}^{\tau} h(\tau-s)u(s) \, ds,$$

*This work is partially supported by National Science Foundation Grant DMS-86-12880 and Air Force Grant AFOSR-86-0134

†Institute for Mathematics and its Applications, University of Minnesota, Minneapolis, Minnesota 55455

‡Bell Communications Research, Morristown, New Jersey 07960

and define

$$(1.3) \quad \sigma^+(\tau, \alpha) = \sup_{u \in K_{\tau, \alpha}} J_{\tau}(u) ,$$

$$(1.4) \quad \sigma^-(\tau, \alpha) = \inf_{u \in K_{\tau, \alpha}} J_{\tau}(u) .$$

Definition 1.1. The function $\sigma(\tau, \alpha) = \sigma^+(\tau, \alpha) - \sigma^-(\tau, \alpha)$ is called the *spread of the linear system*.

The motivation comes from the following theorem.

THEOREM 1.1. For any $\alpha \in \mathbb{R}, \tau > 0$,

$$(1.5) \quad \sup_{t_0 \geq 0} \tilde{\sigma}^+(\tau, \alpha, t_0) = \sigma^+(\tau, \alpha),$$

$$(1.6) \quad \inf_{t_0 \geq 0} \tilde{\sigma}^-(\tau, \alpha, t_0) = \sigma^-(\tau, \alpha)$$

and, consequently,

$$(1.7) \quad \sup_{t_0 \geq 0} \tilde{\sigma}^+(\tau, \alpha, t_0) - \inf_{t_0 \geq 0} \tilde{\sigma}^-(\tau, \alpha, t_0) = \sigma(\tau, \alpha) .$$

Proof. The condition $y(t_0) = \alpha$ means that

$$(1.8) \quad \int_0^{t_0} h(t_0 - s)u(s) ds = \alpha.$$

Writing

$$y(t_0 + \tau) = \int_0^{t_0 + \tau} h(t_0 + \tau - s)u(s) ds$$

and substituting $t_0 - s = s'$, $u(t_0 - s') = v(s')$ we get

$$y(t_0 + \tau) = \int_{-t_0}^{\tau} h(\tau - s')v(s') ds' .$$

The same substitution applied to (1.8) gives

$$\int_{-t_0}^0 h(-s')v(s') ds' = \alpha.$$

Hence

$$\tilde{\sigma}^+(\tau, \alpha, t_0) = \sup \left\{ \int_{-t_0}^{\tau} h(\tau - s') v(s') ds' ; v \text{ satisfies} \right. \\ \left. |v(s')| \leq 1, \quad \int_{-t_0}^0 h(-s') v(s') ds' = \alpha \right\}.$$

Extending $v(s')$ to $s' < -t_0$ by zero we see that $\sigma^+(\tau, \alpha, t_0)$ is $\sup J_{\tau}(v)$ when v is restricted to a subset say K_{τ, α, t_0} , of $K_{\tau, \alpha}$; hence

$$\tilde{\sigma}^+(\tau, \alpha, t_0) \leq \sigma^+(\tau, \alpha).$$

As $t_0 \rightarrow \infty$ the subsets K_{τ, α, t_0} increase and every $u \in K_{\tau, \alpha}$ restricted to a bounded interval is a function in $\bigcup_{t_0 > 0} K_{\tau, \alpha, t_0}$ restricted to the same interval; this implies the equality in (1.5). The proof of (1.6) is similar.

THEOREM 1.2. *For any $\alpha \in \mathbf{R}$, $\tau > 0$ there exist admissible functions $u_{\tau, \alpha}^+$, $u_{\tau, \alpha}^-$ in $K_{\tau, \alpha}$ such that*

$$(1.9) \quad J_{\tau}(u_{\tau, \alpha}^+) = \sup_{u \in K_{\tau, \alpha}} J_{\tau}(u) = \sigma^+(\tau, \alpha),$$

$$(1.10) \quad J_{\tau}(u_{\tau, \alpha}^-) = \inf_{u \in K_{\tau, \alpha}} J_{\tau}(u) = \sigma^-(\tau, \alpha).$$

Indeed, taking a maximizing sequence u_j , we can extract a subsequence which is weakly convergent in L_{loc}^1 to a function u_0 . It is easy to check that u_0 is a maximizer for J_{τ} , i.e., u_0 is the asserted $u_{\tau, \alpha}^+$. The proof of (1.10) is similar.

In this paper we study the structure of $u_{\tau, \alpha}^{\pm}$ and this enables us to compute the spread of some linear systems of interest. In §2 we solve a general maximization problem, which is then used (in §3) to analyze the structure of $u_{\tau, \alpha}^{\pm}$. In §4 we establish various properties of $\sigma(\tau, \alpha)$ and, in §5, we compute $\sigma(\tau, \alpha)$ for some examples. Finally in §6 we show that all the results can be extended to the case where $y(t_0)$, $y(t_0 + \tau_1), \dots, y(t_0 + \tau_{N-1})$ are prescribed and the range of $y(t_0 + \tau_N)$ is sought; here $0 < \tau_1 < \tau_2 < \dots < \tau_N$.

Motivation for studying the function $\sigma^{\pm}(\tau, \alpha)$ comes from the following problem posed in [1] and [2]. For any $d > 0$, $T > 0$ and impulse response $h(\cdot)$, denote by $N_{\max}(T, d)$ the maximum number of inputs $u_j(s)$ such that the corresponding outputs $y_j(t)$ satisfy:

$$\max_{0 < t \leq T} |y_i(t) - y_j(t)| \geq d \quad \forall i \neq j.$$

We define

$$(1.11) \quad MCT(d) = \lim_{T \rightarrow \infty} \frac{\log N_{\max}(T, d)}{T} \quad \text{bits/sec},$$

and would like to obtain bounds on the $MCT(d)$ for any $h(\cdot)$. Set

$$\tau^* = \inf\{\tau \mid \sigma(\tau, 0) = d\}.$$

Work in progress [3] indicates that

$$(1.12) \quad MCT(d) \leq \frac{1}{\tau^*}$$

for any $h(\cdot)$ that satisfies

$$\int_{\left\{\frac{h(\tau-s)}{h(-s)} \geq 1\right\}} |h(-s)| \, ds \leq \int_{\left\{\frac{h(\tau-s)}{h(-s)} \leq 1\right\}} |h(-s)| \, ds$$

for all $\tau \leq \tau^*$; the arguments used depend on results derived in this paper. Results obtained here (in §6) for the N constraint problem, in which N output values are specified, can be used to tighten the upper bound given by (1.12) (see [3]).

The problem of computing spread for a discrete-time linear system with impulse response h_i , $i = 0, 1, 2, \dots$, is considered in [2]. This computation is equivalent to solving a linear program with bounded variables and one equality constraint. Here we show how the spread can be computed for a continuous-time linear system. Two examples of special interest are presented in which the spread can be computed by finding a solution to a transcendental equation.

A general optimization problem. Let $f(s)$, $g(s)$ be continuous functions in $-\infty < s \leq 0$, which belong to $L^1(-\infty, 0)$, and assume:

$$(2.1) \quad f \neq 0 \quad \text{a.e.,}$$

$$(2.2) \quad \text{meas} \left\{ \frac{g}{f} = \mu \right\} = 0 \quad \text{for any } \mu \in \mathbb{R}.$$

Let

$$K = \left\{ u(s) \text{ measurable for } -\infty < s < 0, \quad |u(s)| \leq 1, \right. \\ \left. \int_{-\infty}^0 f(s)u(s) \, ds = \alpha \right\}$$

for some fixed $\alpha \in \mathbb{R}$, and

$$J(u) = \int_{-\infty}^0 g(s)u(s) ds.$$

As in the proof of Theorem 1.2 one can show that there exists a function $u_0 \in K$ such that

$$(2.3) \quad J(u_0) = \max_{u \in K} J(u).$$

THEOREM 2.1. *For any solution $u_0 \in K$ of (2.3) there exists a number $\lambda \in \mathbb{R}$ such that almost everywhere*

$$(2.4) \quad u_0(s) = \begin{cases} \operatorname{sgn} f(s) & \text{if } \frac{g(s)}{f(s)} > \lambda \\ -\operatorname{sgn} f(s) & \text{if } \frac{g(s)}{f(s)} < \lambda. \end{cases}$$

Notice that (2.4) is equivalent to

$$u_0(s) = \operatorname{sgn} [g(s) - \lambda f(s)].$$

Proof. We begin by proving that $u_0 = 1$ a.e. If the assertion is not true then the set $G_0 = \{|u_0| < 1\}$ has positive measure. Denote by G the subset of G_0 consisting of all points t of G_0 -density equal to 1, such that also $f(t) \neq 0$. Then $\operatorname{meas} G = \operatorname{meas} G_0 > 0$.

Take t_1, t_2 in G ($t_1 \neq t_2$) and let G_i be a subset of G contained in δ_0 -neighborhood of t_i , such that $\sup_{G_i} |u| < 1$,

$$G_1 \cap G_2 = \emptyset, \quad \operatorname{meas} G_1 = \operatorname{meas} G_2 = \delta > 0.$$

For any real numbers A_1, A_2 , if ϵ is positive and small enough then the function

$$(2.5) \quad \tilde{u} = u_0 + A_1 \frac{\epsilon}{\delta} \chi_{G_1} + A_2 \frac{\epsilon}{\delta} \chi_{G_2}$$

satisfies $|\tilde{u}| \leq 1$. Further, if

$$(2.6) \quad A_1 \int_{G_1} f(s) ds + A_2 \int_{G_2} f(s) ds = 0$$

then $\int_{-\infty}^0 f(s)\tilde{u}(s) ds = \alpha$, so that $\tilde{u} \in K$. Notice that (2.6) is equivalent to

$$(2.7) \quad A_1 f(t_1) + A_2 f(t_2) = \sigma_1(\delta_0)$$

for some $\sigma_1(\delta_0)$ such that $\sigma_1(\delta_0) \rightarrow 0$ if $\delta_0 \rightarrow 0$.

From the maximality of u_0 it follows that (2.6), or (2.7), implies $J(\tilde{u}) \leq J(u_0)$, that is

$$(2.8) \quad A_1 \int_{G_1} g(s) ds + A_2 \int_{G_2} g(s) ds \leq 0 ,$$

i.e.,

$$(2.9) \quad A_1 g(t_1) + A_2 g(t_2) \leq \sigma_2(\delta_0)$$

for some $\sigma_2(\delta_0)$ such that $\sigma_2(\delta_0) \rightarrow 0$ if $\delta_0 \rightarrow 0$.

Choosing

$$(2.10) \quad A_1 = -A_2 \frac{f(t_2)}{f(t_1)} + \frac{\sigma_1(\delta_0)}{f(t_1)}$$

so that (2.7) is satisfied, (2.9) must then hold and, upon letting $\delta_0 \rightarrow 0$, we get

$$(2.11) \quad A_2 \left[-\frac{g(t_1)f(t_2)}{f(t_1)} + g(t_2) \right] \leq 0 .$$

Since A_2 is arbitrary, it follows that the expression in brackets must vanish. Thus

$$\frac{g(t_1)}{f(t_1)} = \frac{g(t_2)}{f(t_2)}$$

for all t_1, t_2 in G . Since G has a positive measure, this is a contradiction to (2.2).

Denote by D the set of all points t such that $f(t) \neq 0$ and t is a Lebesgue point of u_0 . Thus almost all t in $(-\infty, 0)$ belong to D . Take any t_1, t_2 in D with

$$(2.12) \quad \frac{g(t_1)}{f(t_1)} > \frac{g(t_2)}{f(t_2)} .$$

We shall prove that a.e.

$$(2.13) \quad u_0(t_2) = \operatorname{sgn} f(t_2) \quad \text{implies} \quad u_0(t_1) = \operatorname{sgn} f(t_1)$$

and

$$(2.14) \quad u_0(t_1) = -\operatorname{sgn} f(t_1) \quad \text{implies} \quad u_0(t_2) = -\operatorname{sgn} f(t_2) .$$

These two statements clearly imply the assertion (2.4).

To prove (2.13) suppose the assertion is not true. Then the set $\tilde{G}$ of pair (t_1, t_2) for which (2.13) is not true has positive measure. Choose t_1, t_2 at which $\tilde{G}$ has density 1. Then for any $\delta_0 > 0$ we can find sets G_1, G_2 such that G_i is contained in δ_0 -neighborhood of t_i , $\text{meas } G_1 = \text{meas } G_2 = \delta > 0$, and

$$\begin{aligned} u_0(t) &= \text{sgn } f(t) & \text{for all } t \in G_2, \\ u_0(t) &= -\text{sgn } f(t) & \text{for all } t \in G_1. \end{aligned}$$

We again form the function (2.5). If

$$(2.15) \quad A_2 \text{sgn } f(t_2) < 0, \quad A_1 \text{sgn } f(t_1) > 0$$

then $|\tilde{u}| \leq 1$ if ϵ is sufficiently small.

If we can further choose A_1, A_2 such that (2.6) (or (2.7)) holds, then (2.8) (or 2.9)) must be satisfied. The condition (2.7) is satisfied by the choice (2.10) of A_1 , and if $A_2 \text{sgn } f(t_2) < 0$ then clearly also $A_1 \text{sgn } f(t_1) > 0$ provided δ_0 is sufficiently small. We conclude, after letting $\delta_0 \rightarrow 0$, that (2.11) must hold provided $A_2 \text{sgn } f(t_2) < 0$. Dividing (2.11) by $A_2 f(t_2)$ we arrive at the inequality

$$-\frac{g(t_1)}{f(t_1)} + \frac{g(t_2)}{f(t_2)} \geq 0$$

which is a contradiction to (2.12). This completes the proof of (2.13); the proof of (2.14) is similar.

From Theorem 2.1 we immediately get

COROLLARY 2.2. *The constant λ in Theorem 2.1 is uniquely determined by*

$$(2.16) \quad \int_{\{\frac{g(s)}{f(s)} > \lambda\}} |f(s)| ds - \int_{\{\frac{g(s)}{f(s)} < \lambda\}} |f(s)| ds = \alpha;$$

consequently the maximizer u_0 is also uniquely determined. As α decreases from $\int_{-\infty}^0 |f(s)| ds$ to $-\int_{-\infty}^0 |f(s)| ds$, $\lambda = \lambda(\alpha)$ increases monotonically from

$$\inf_{s < 0} \{g(s)/f(s)\} \quad \text{to} \quad \sup_{s < 0} \{g(s)/f(s)\}.$$

§3. The structure of $u_{\tau, \alpha}^\pm$. Choose $h(t)$ as in §1, i.e.,

$$(3.1) \quad h \in L^1(0, \infty) \cap C^0[0, \infty)$$

and assume further that

$$(3.2) \quad h(t) \neq 0 \quad \text{a.e.}$$

$$(3.3) \quad \text{meas}\{0 < t < \infty; \frac{h(t+\tau)}{h(t)} = \lambda\} = 0 \quad \forall \tau > 0, \lambda \in \mathbb{R}.$$

Taking $f(t) = h(-t)$, $g(t) = h(\tau - t)$ in Theorem 2.1 and Corollary 2.2, we get:

THEOREM 3.1. *There exists a unique solution $u_{\tau,\alpha}^+$ of (1.9) and it is given by*

$$(3.4) \quad u_{\tau,\alpha}^+(s) = \begin{cases} \operatorname{sgn} h(-s) & \text{if } \frac{h(\tau-s)}{h(-s)} > \lambda^+ \\ -\operatorname{sgn} h(-s) & \text{if } \frac{h(\tau-s)}{h(-s)} < \lambda^+ \end{cases}$$

where λ^+ is determined by

$$(3.5) \quad \int_{\{\frac{h(\tau-s)}{h(-s)} > \lambda^+\}} |h(-s)| \, ds - \int_{\{\frac{h(\tau-s)}{h(-s)} < \lambda^+\}} |h(-s)| \, ds = \alpha .$$

Clearly also $u_{\tau,\alpha}^+(s) = \operatorname{sgn} h(\tau-s)$ if $0 < s < \tau$.

We now consider a special case.

THEOREM 3.2. *If $h \in L^1(0, \infty)$, $h > 0$, $d^2(\log h)/dt^2 > 0$, then there is a unique solution of (1.9), given by*

$$(3.6) \quad u_{\tau,\alpha}^+(s) = \begin{cases} 1 & \text{if } -\infty < s < \mu, \\ -1 & \text{if } \mu < s < 0 \end{cases}$$

and $u_{\tau,\alpha}^+(s) = 1$ if $0 < s < \tau$, where μ is determined by

$$(3.7) \quad \int_{-\infty}^{\mu} h(-s) \, ds - \int_{\mu}^0 h(-s) \, ds = \alpha .$$

Proof. By assumption

$$\frac{h'(s)}{h(s)} \quad \text{is strictly increasing,}$$

hence

$$\frac{h'(\tau+s)}{h(\tau+s)} > \frac{h'(s)}{h(s)} .$$

This means that

$$\frac{d}{ds} \frac{h(\tau+s)}{h(s)} > 0$$

and thus

$$\frac{h(\tau-s)}{h(-s)} \quad \text{is strictly decreasing in } s.$$

Now apply Theorem 3.1 to complete the proof.

REMARK 3.1. If $\log h$ is convex (but not satisfying $d^2(\log h)/dt^2 > 0$) then we can approximate it by a smooth function h_n with $d^2(\log h_n)/dt^2 > 0$. Applying Theorem 3.2 to the corresponding maximizers $u_{\tau,\alpha}^{h_n}$ we deduce that there is a maximizer $u_{\tau,\alpha}$ (for h) having the form (3.6), (3.7). There may be other maximizers; for instance if $h(t) = e^{-t}$ then every $u \in K_{\tau,\alpha}$ is a maximizer. (Notice that (3.3) does not hold for $h(t) = e^{-t}$.)

THEOREM 3.3. If $h \in L^1(0, \infty)$, $h > 0$ and $d^2(\log h)/dt^2 < 0$ then there is a unique solution of (1.9) given by

$$(3.8) \quad u_{\tau,\alpha}^+(s) = \begin{cases} -1 & \text{if } -\infty < s < \tilde{\mu} \\ 1 & \text{if } \tilde{\mu} < s < 0 \end{cases}$$

and $u_{\tau,\alpha}^+(s) = 1$ if $0 < s < \tau$, where $\tilde{\mu}$ is determined by

$$(3.9) \quad - \int_{-\infty}^{\tilde{\mu}} h(-s) ds + \int_{\tilde{\mu}}^0 h(-s) ds = \alpha.$$

Notice that $d\mu/d\alpha > 0$, $d\tilde{\mu}/d\alpha < 0$ where $\mu = \mu(\alpha)$ and $\tilde{\mu} = \tilde{\mu}(\alpha)$ are defined by (3.7) and (3.9) respectively.

§4. Properties of the spread.

Theorem 3.1 implies that

$$(4.1) \quad \begin{aligned} \sigma^+(\tau, \alpha) &= \int_{\left\{ \frac{h(\tau-s)}{h(-s)} > \lambda^+ \right\}} [\operatorname{sgn} h(-s)] h(\tau-s) ds \\ &\quad - \int_{\left\{ \frac{h(\tau-s)}{h(-s)} < \lambda^+ \right\}} [\operatorname{sgn} h(-s)] h(\tau-s) ds + \int_0^\tau |h(\tau-s)| ds \\ &= \int_{\left\{ \frac{h(\tau-s)}{h(-s)} > \lambda^+ \right\}} \frac{h(\tau-s)}{h(-s)} |h(-s)| ds \\ &\quad - \int_{\left\{ \frac{h(\tau-s)}{h(-s)} < \lambda^+ \right\}} \frac{h(\tau-s)}{h(-s)} |h(-s)| ds + \int_0^\tau |h(\tau-s)| ds, \end{aligned}$$

where λ^+ is determined by (3.5). Similarly, one can show that

$$(4.2) \quad \begin{aligned} \sigma^-(\tau, \alpha) &= - \int_{\left\{ \frac{h(\tau-s)}{h(-s)} > \lambda^- \right\}} \frac{h(\tau-s)}{h(-s)} |h(-s)| ds + \int_{\left\{ \frac{h(\tau-s)}{h(-s)} < \lambda^- \right\}} |h(-s)| ds \\ &\quad - \int_0^\tau |h(\tau-s)| ds, \end{aligned}$$

where λ^- is determined by

$$(4.3) \quad - \int_{\left\{ \frac{h(\tau-s)}{h(-s)} > \lambda^- \right\}} |h(-s)| ds + \int_{\left\{ \frac{h(\tau-s)}{h(-s)} < \lambda^- \right\}} |h(-s)| ds = \alpha .$$

As α decreases from $\int_0^\infty |h(s)| ds$ to $-\int_0^\infty |h(s)| ds$, $\lambda^-(\alpha)$ decreases monotonically from $\sup_{s < 0} \{h(\tau-s)/h(-s)\}$ to $\inf_{s < 0} \{h(\tau-s)/h(-s)\}$. Also, $\lambda^-(0) = \lambda^+(0)$.

Combining (4.1) and (4.2) gives the spread

$$(4.4) \quad \begin{aligned} \frac{1}{2} \sigma(\tau, \alpha) &= \frac{1}{2} [\sigma^+(\tau, \alpha) - \sigma^-(\tau, \alpha)] \\ &= \int_{\left\{ \frac{h(\tau-s)}{h(-s)} > \lambda_M \right\}} \frac{h(\tau-s)}{h(-s)} |h(-s)| ds - \int_{\left\{ \frac{h(\tau-s)}{h(-s)} < \lambda_m \right\}} \frac{h(\tau-s)}{h(-s)} |h(-s)| ds \\ &\quad + \int_0^\tau |h(\tau-s)| ds \end{aligned}$$

where $\lambda_M = \max(\lambda^-, \lambda^+)$ and $\lambda_m = \min(\lambda^-, \lambda^+)$,

THEOREM 4.1. *There holds:*

$$(4.5) \quad \frac{\partial \sigma^\pm(\tau, \alpha)}{\partial \alpha} = \lambda^\pm .$$

Proof. Since $\lambda^+(\alpha)$ is a monotonically decreasing function of α , we can write

$$(4.6) \quad \int_{\left\{ \frac{h(\tau-s)}{h(-s)} > \lambda^+ + \Delta\lambda \right\}} |h(-s)| ds - \int_{\left\{ \frac{h(\tau-s)}{h(-s)} < \lambda^+ + \Delta\lambda \right\}} |h(-s)| ds = \alpha - \Delta\alpha$$

where $\Delta\lambda, \Delta\alpha > 0$. Subtracting (4.6) from (4.1) gives

$$(4.7) \quad 2 \int_{\left\{ \lambda^+ < \frac{h(\tau-s)}{h(-s)} < \lambda^+ + \Delta\lambda \right\}} |h(-s)| ds = \Delta\alpha .$$

From (4.1), (4.6), and (4.7),

$$\begin{aligned} \sigma^+(\tau, \alpha) - \sigma^+(\tau, \alpha - \Delta\alpha) &= 2 \int_{\left\{ \lambda^+ < \frac{h(\tau-s)}{h(-s)} < \lambda^+ + \Delta\lambda \right\}} \frac{h(\tau-s)}{h(-s)} |h(-s)| ds \\ &= [\lambda^+ + \epsilon(\Delta\lambda)] \Delta\alpha \end{aligned}$$

where $\epsilon(\Delta\lambda) \rightarrow 0$ as $\Delta\lambda \rightarrow 0$. Letting $\Delta\alpha \rightarrow 0$ while observing (4.7) gives $\partial\sigma^+/\partial\alpha = \lambda^+$. A similar argument shows that $\partial\sigma^-/\partial\alpha = \lambda^-$.

THEOREM 4.2. (i) $\sigma^+(\tau, \alpha)$ is concave in α , $\sigma^-(\tau, \alpha)$ is convex in α , and thus $\sigma(\tau, \alpha)$ is concave in α ;

(ii) $\sigma^\pm(\tau, \alpha) = -\sigma^\pm(\tau, -\alpha)$ and therefore $\sigma(\tau, \alpha) = \sigma(\tau, -\alpha)$;

(iii) $\frac{\partial \sigma(\tau, \alpha)}{\partial \alpha} \leq 0$ if $\alpha > 0$.

Proof. Assertions (i) and (ii) follow immediately from Theorem 4.1 and the fact that $\partial \lambda^+ / \partial \alpha$ ($\partial \lambda^- / \partial \alpha$) is negative (positive) for all α . The assertion (ii) is obvious from the definition of $\sigma^\pm$. Finally, since $\sigma(\tau, \alpha)$ is concave in α (by (i)) and $\partial \sigma(\tau, \alpha) / \partial \alpha = 0$ at $\alpha = 0$ (by (ii)), (iii) follows.

We now specialize to the case where either $\log h$ is convex, so that

$$(4.8) \quad \sigma^+(\tau, \alpha) = \int_{-\infty}^{\mu} h(\tau - s) ds - \int_{\mu}^0 h(\tau - s) ds + \int_0^{\tau} h(s') ds'$$

where μ is determined by (3.7), or $\log h$ is concave so that

$$(4.9) \quad \sigma^+(\tau, \alpha) = - \int_{-\infty}^{\tilde{\mu}} h(\tau - s) ds + \int_{\tilde{\mu}}^0 h(\tau - s) ds + \int_0^{\tau} h(s') ds'$$

where $\tilde{\mu}$ is determined by (3.9).

THEOREM 4.3. If $h' < 0$ and $\log h$ is convex or concave then

$$(4.10) \quad \frac{\partial \sigma^\pm(\tau, \alpha)}{\partial \tau} > 0.$$

Proof. If $\log h$ is convex then from (4.8) we get,

$$\begin{aligned} \frac{\partial \sigma^+(\tau, \alpha)}{\partial \tau} &= - \int_{-\infty}^{\mu} \frac{d}{ds} h(\tau - s) ds + \int_{\mu}^0 \frac{d}{ds} h(\tau - s) ds + h(\tau) \\ &= 2h(\tau) - 2h(\tau - \mu) > 0. \end{aligned}$$

Similarly, if $\log h$ is concave then

$$\begin{aligned} \frac{\partial \sigma^+(\tau, \alpha)}{\partial \tau} &= \int_{-\infty}^{\tilde{\mu}} \frac{d}{ds} h(\tau - s) ds - \int_{\tilde{\mu}}^0 \frac{d}{ds} h(\tau - s) ds + h(\tau) \\ &= 2h(\tau - \tilde{\mu}) > 0. \end{aligned}$$

Finally, the second inequality in (4.10) follows from the first inequality and Theorem 4.2 (ii).

§5. Examples.

If $h(t) = \exp\{-k(t)\}$ where $k(t) \rightarrow \infty$, k convex (k concave) then $\log h$ is concave (convex). For $h(t) = (t+a)^b$ where $a > 0$, $b > 0$, $\log h$ is convex.

We shall now consider two functions $h(t)$ of special interest.

THEOREM 5.1. Let

$$(5.1) \quad h(t) = \sum_{i=1}^N a_i e^{-\beta_i t} \quad (a_i > 0, \beta_i > 0).$$

Then $d^2 \log h / dt^2 > 0$.

Proof. As in the proof of Theorem 3.2, the assertion is equivalent to showing that

$$\frac{d}{ds} \frac{h(\tau-s)}{h(-s)} = \frac{h(-s) \sum a_i \beta_i e^{-\beta_i(\tau-s)} - h(\tau-s) \sum a_i \beta_i e^{\beta_i s}}{h^2(-s)}$$

is negative for any $\tau > 0$. But the numerator is equal to

$$\begin{aligned} & \sum \sum a_i \beta_i a_j (e^{\beta_i(\tau-s)+\beta_j s} - e^{-\beta_j(\tau-s)+\beta_i s}) \\ &= \sum \sum a_i a_j \beta_i e^{s(\beta_i+\beta_j)} (e^{-\beta_i \tau} - e^{-\beta_j \tau}) \\ &= \frac{1}{2} \sum \sum a_i a_j e^{s(\beta_i+\beta_j)} \left[\beta_i (e^{-\beta_i \tau} - e^{-\beta_j \tau}) + \beta_j (e^{-\beta_j \tau} - e^{-\beta_i \tau}) \right] \\ &= \frac{1}{2} \sum \sum a_i a_j e^{s(\beta_i+\beta_j)} (\beta_i - \beta_j) (e^{-\beta_i \tau} - e^{-\beta_j \tau}) \end{aligned}$$

and each term in the last sum is negative if $\beta_i \neq \beta_j$.

For the function (5.1), the μ determined by (3.7) is given by

$$\sum_{i=1}^N \frac{a_i}{\beta_i} (2 e^{\beta_i \mu} - 1) = \alpha.$$

The next example is

$$(5.2) \quad h(t) = e^{-\beta t} \cos \omega t \quad (\beta > 0, \omega > 0).$$

Since

$$\frac{h(\tau-s)}{h(-s)} = e^{-\beta \tau} (\cos \omega \tau + \sin \omega \tau \tan \omega s),$$

one can check that the optimal solution $u_{\tau, \alpha}^+$, which for simplicity we shall denote by u_0 , satisfies:

$$u_0(s) = \begin{cases} \operatorname{sgn} h(-s) & \text{if } \gamma - n\pi < \omega s < -\frac{(2n+1)\pi}{2} \\ -\operatorname{sgn} h(-s) & \text{if } -\frac{(2n+3)\pi}{2} < \omega s < \gamma - n\pi \end{cases}$$

if $n = 0, 1, 2, \dots$, and

$$u_0(s) = \begin{cases} -\operatorname{sgn} h(-s) & \text{if } -\frac{\pi}{2} < \omega s < \min(\gamma + \pi, 0) \\ \operatorname{sgn} h(-s) & \text{if } \min(\gamma + \pi, 0) < \omega s \leq 0 \end{cases}$$

where $\gamma \in [-\frac{3\pi}{2}, -\frac{\pi}{2}]$ is to be selected such that

$$(5.3) \quad \int_{-\infty}^0 h(-s) u_0(s) ds = \alpha.$$

Recalling (5.2) one can check that

$$u_0(s) = \begin{cases} -1 & \text{if } \gamma - 2n\pi < \omega s < \gamma - (2n-1)\pi \\ 1 & \text{if } \gamma - (2n-1)\pi < \omega s < \gamma - 2(n-1)\pi \end{cases}$$

for $n = 1, 2, \dots$, and

$$u_0(s) = \begin{cases} -1 & \text{if } \gamma < \omega s < \min(\gamma + \pi, 0) \\ 1 & \text{if } \min(\gamma + \pi, 0) < \omega s < 0. \end{cases}$$

Setting $\gamma' = \min(\gamma + \pi, 0)$ and using the formula

$$\int_a^b h(-s) ds = -\operatorname{Re} \left\{ \frac{1}{\beta + i\omega} [e^{-(\beta + i\omega)b} - e^{-(\beta + i\omega)a}] \right\},$$

we can compute

$$\begin{aligned} \int_{-\infty}^0 h(-s) u_0(s) ds &= \sum_{n=1}^{\infty} \left[- \int_{(\gamma - 2n\pi)/\omega}^{(\gamma - (2n-1)\pi)/\omega} h(-s) ds \right. \\ &\quad \left. + \int_{(\gamma - (2n-1)\pi)/\omega}^{(\gamma - 2(n-1)\pi)/\omega} h(-s) ds \right] \\ &\quad - \int_{\gamma/\omega}^{\gamma'/\omega} h(-s) ds + \int_{\gamma'/\omega}^0 h(-s) ds. \end{aligned}$$

After somewhat lengthy calculations we get the expression

$$(5.4) \quad \operatorname{Re} \left\{ \frac{\beta - i\omega}{\beta^2 + \omega^2} e^{\beta + i\omega} \gamma/\omega \frac{1 + e^{-\beta\pi/\omega}}{1 - e^{-\beta\pi/\omega}} \right. \\ \left. + \frac{\beta - i\omega}{\beta^2 + \omega^2} \left[1 - 2 e^{(\beta + i\omega)\gamma'/\omega} + e^{(\beta + i\omega)\gamma/\omega} \right] \right\},$$

or

$$\frac{\beta}{\beta^2 + \omega^2} + \frac{2e^{\beta\gamma/\omega}}{(\beta^2 + \omega^2)(1 - e^{-\beta\pi/\omega})} (\beta \cos \gamma + \omega \sin \gamma) - \frac{2e^{\beta\gamma'/\omega}}{\beta^2 + \omega^2} (\beta \cos \gamma' + \omega \sin \gamma') .$$

Hence (5.3) determines γ by the following formulas:

$$(5.5) \quad \begin{aligned} \frac{2e^{\beta\gamma\omega}}{1 - e^{-\beta\pi/\omega}} (\beta \cos \gamma + \omega \sin \gamma) &= \alpha(\omega^2 + \beta^2) + \beta \quad \text{if } -\pi < \gamma < 0, \\ \left[\frac{2e^{\beta\gamma/\omega}}{1 - e^{-\beta\pi/\omega}} + 2e^{\beta(\gamma+\pi)/\omega} \right] (\beta \cos \gamma + \omega \sin \gamma) \\ &= \alpha(\omega^2 + \beta^2) - \beta \quad \text{if } -\frac{3\pi}{2} < \gamma < -\pi. \end{aligned}$$

Since

$$\begin{aligned} \int_{-\infty}^0 h(\tau - s) u_0(s) ds &= \operatorname{Re} \left\{ \int_{-\infty}^0 e^{-(\beta + i\omega)(\tau - s)} u_0(s) ds \right\} \\ &= \operatorname{Re} \left\{ e^{-(\beta + i\omega)\tau} \int_{-\infty}^0 h_0(-s) u_0(s) ds \right\} \end{aligned}$$

and the last integral is equal to the expression in braces in (5.4), we find that

$$\begin{aligned} \sigma^+(\tau, \alpha) &= \frac{2e^{\beta(\frac{\pi}{\omega} - \tau)}}{(\beta^2 + \omega^2)(1 - e^{-\beta\pi/\omega})} [\beta \cos(\gamma - \omega\tau) + \omega \sin(\gamma - \omega\tau)] \\ &\quad - \frac{e^{-\beta\tau}}{\beta^2 + \omega^2} (\beta \cos \omega\tau - \omega \sin \omega\tau) + \int_0^\tau |h(s)| ds \quad \text{if } -\pi < \gamma < 0, \\ \sigma^+(\tau, \alpha) &= \frac{2e^{\beta(\frac{\pi+\pi}{\omega} - \tau)}}{(\beta^2 + \omega^2)(1 - e^{-\beta\pi/\omega})} [\beta \cos(\gamma - \omega\tau) + \omega \sin(\gamma - \omega\tau)] \\ &\quad + \frac{e^{-\beta\tau}}{\beta^2 + \omega^2} (\beta \cos \omega\tau - \omega \sin \omega\tau) + \int_0^\tau |h(s)| ds \\ &\quad \text{if } -\frac{3\pi}{2} < \gamma < -\pi. \end{aligned}$$

§6. Several constraints. The results of the previous sections can be extended to the case of several constraints. In fact it all hinges upon generalizing Theorem 2.1 to the problem:

$$(6.1) \quad \max_{u \in K_\alpha} \int_{-\infty}^0 g(s) u(s) ds$$

where K_α is the set of all measurable functions $u(s)$ satisfying:

$$(6.2) \quad -1 \leq u(s) \leq 1 \quad \text{for} \quad -\infty < s \leq 0,$$

$$(6.3) \quad \int_{-\infty}^0 f_i(s)u(s) ds = \alpha_i \quad (i = 1, 2, \dots, N).$$

Here g and f_i are given functions in $L^1(-\infty, 0) \cap C^0(-\infty, 0]$ and α_i are given real numbers.

THEOREM 6.1. Assume that $f_1 \neq 0$ a.e. and that, for any real numbers $\mu_1, \dots, \mu_N$,

$$\text{measure}\{g = \sum_{i=1}^N \mu_i f_i\} = 0.$$

Then there exist sequences $u_m, \lambda_{i,m}, \alpha_{i,m}$ with $u_m \rightarrow u_0$ weakly in L^1_{loc} , $\alpha_{1,m} = \alpha_1$, $\alpha_{i,m} \rightarrow \alpha_i$ for $2 \leq i \leq N$, where u_0 is a maximizer of (6.1), and

$$(6.4) \quad u_m(s) = \text{sgn} \left[g(s) - \sum_{i=1}^N \lambda_{i,m} f_i(s) \right],$$

$$(6.5) \quad \int_{-\infty}^0 f_i(s)u_m(s) ds = \alpha_{i,m} \quad (i = 1, 2, \dots, N).$$

Thus, in order to evaluate (6.1) we need to analyze the u_m from (6.4), (6.5) and then compute $\int_{-\infty}^0 g u_m$, noting that

$$\int_{-\infty}^0 g(s)u_m(s) ds \rightarrow \int_{-\infty}^0 g(s)u_0(s) ds = \max_{u \in K_\alpha} \int_{-\infty}^0 g(s)u(s) ds.$$

Proof. For any small $\eta > 0$ introduce the "penalized" functional

$$(6.6) \quad J_\eta(u) = \int_{-\infty}^0 g(s)u(s) ds - \frac{1}{\eta} \sum_{i=2}^N \left[\int_{-\infty}^0 f_i(s)u(s) ds - \alpha_i \right]^2$$

and consider the problem:

$$(6.7) \quad \text{maximize } J_\eta(u) \quad \text{for } u \in K$$

where K consists of all functions u satisfying:

$$-1 \leq u(s) \leq 1, \quad \int_{-\infty}^0 f_1(s) ds = \alpha_1.$$

Proceeding as in the proof of Theorem 2.1 we deduce that if $|\tilde{u}| \leq 1$ where $\tilde{u}$ is defined by (2.6) with $u_0 = u_\eta$, then (2.6) implies

$$A_1 g(t_1) + A_2 g(t_2) - \frac{2}{\eta} \sum_{i=2}^N \left(\int_{-\infty}^0 f_i u_\eta ds - \alpha_i \right) (A_1 f_i(t_1) + A_2 f_i(t_2)) \leq \sigma_2(\delta_0)$$

where u_η is a solution of (6.7) and $\sigma_2(\delta_0) \rightarrow 0$ if $\delta_0 \rightarrow 0$. Taking $\delta_0 \rightarrow 0$ we get the inequality

$$\begin{aligned} A_1 [g(t_1) - \sum_{i=2}^N \lambda_{i,\eta} f_i(t_1)] \\ + A_2 [g(t_2) - \sum_{i=2}^N \lambda_{i,\eta} f_i(t_2)] \leq 0 \end{aligned}$$

for some scalars $\lambda_{i,\eta}$. We can now proceed as in §2 to deduce that $\text{meas}\{|u_\eta| < 1\} = 0$; furthermore,

$$(6.8) \quad u_\eta(s) = \text{sgn} \left[g(s) - \sum_{i=1}^N \lambda_{i,\eta} f_i(s) \right].$$

We note that

$$J_\eta(u_\eta) \geq J_\eta(\hat{u}) \quad \forall \hat{u} \in K;$$

From the inequality $J_\eta(u_\eta) \geq J_\eta(\hat{u})$ for some $\hat{u} \in K_\eta$ it follows that

$$\frac{1}{\eta} \sum_{i=2}^N \left[\int_{-\infty}^0 f_i(s) u_\eta(s) ds - \alpha_i \right]^2 \leq C, \quad C \text{ independent of } \eta.$$

Hence, as $\eta \rightarrow 0$,

$$(6.9) \quad \alpha_{i,\eta} \equiv \int_{-\infty}^0 f_i(s) u_\eta(s) ds \rightarrow \alpha_i \quad (2 \leq i \leq N).$$

It is also easy to verify that for any convergent subsequence u_{η_m} (weakly in L^1_{loc}), the limit u_0 is a solution to problem (6.1). Indeed

$$(6.10) \quad J_\eta(u_\eta) \leq \int_{-\infty}^0 g u_\eta \, ds \leq \max_{u \in K_{\alpha_\eta}} \int_{-\infty}^0 g u \, ds = \int_{-\infty}^0 g \hat{u}_\eta \, ds$$

where K_{α_η} is defined as K_α but with

$$\alpha_2 = \alpha_{2,\eta}, \dots, \alpha_N = \alpha_{N,\eta}.$$

Since $\alpha_{j,\eta} \rightarrow \alpha_j$, if we take η to vary in a subsequence of η_m such that $\hat{u}_\eta \rightarrow \hat{u}$ weakly in L^1_{loc} , then $\int_{-\infty}^0 g \hat{u}_\eta \rightarrow \int_{-\infty}^0 g \hat{u}$ and $\hat{u} \in K_\alpha$ (i.e., $\hat{u}$ satisfies (6.2), (6.3)). Denoting by u_1 any solution of (6.1) – (6.3) we then have

$$\int_{-\infty}^0 g \hat{u} \, ds \leq \int_{-\infty}^0 g u_1 \, ds;$$

also, by maximality of u_η (see (6.7))

$$\int_{-\infty}^0 g u_1 \, ds = J_\eta(u_1) \leq J_\eta(u_\eta).$$

Using these relations in (6.10) and noting that

$$\int_{-\infty}^0 g u_\eta \, ds \rightarrow \int_{-\infty}^0 g u_0 \, ds,$$

we conclude that

$$\int_{-\infty}^0 g u_0 \, ds = \int_{-\infty}^0 g u_1 \, ds = \max_{u \in K_\alpha} \int_{-\infty}^0 g u \, ds.$$

Thus u_0 is a solution to (6.1)–(6.3). Recalling also (6.8), (6.9), the proof of Theorem 6.1 is complete.

REMARK 6.1. The $\lambda_{i,m}$ satisfy

$$\int_{\{g > \Sigma \lambda_{j,m} f_j\}} f_i(s) \, ds - \int_{\{g < \Sigma \lambda_{j,m} f_j\}} f_i(s) \, ds = \alpha_i \quad (1 \leq i \leq N).$$

From these equations one should be able to determine the $\lambda_{j,m}$, at least in some relatively simple examples, and show that $\lambda_{j,m} \rightarrow \lambda_j$ (λ_j finite) as $m \rightarrow \infty$; consequently

$$u_0 = \operatorname{sgn}\left[g(s) - \sum_{i=1}^N \lambda_i f_i(s)\right].$$

$$\text{and } \int_{-\infty}^0 f_j(s) u_0(s) ds = \alpha_j \quad \text{for } 1 \leq j \leq N.$$

REMARK 6.2. Since uniqueness has not been established for (6.1), it is worth noting that any solution u_0 of (6.1) can be recovered as a weak limit of maximizers u_η provided we modify the definitions of $J_\eta(u)$ in (6.6), adding the term

$$- \int_{-\infty}^0 \frac{(u - u_0)^2}{1 + s^2} ds$$

to the right-hand side. One can then show that

$$\int_{-\infty}^0 \frac{(u_\eta - u_0)^2}{1 + s^2} ds \rightarrow 0 \quad \text{if } \eta \rightarrow 0.$$

The only differences occurs in (6.8): In the first inequality we have to replace $\lambda_{1,\eta} f_1(s)$ by $\lambda_{1,\eta} f_1(s) + \gamma(\eta)$ and in the second inequality we have to replace $\lambda_{1,\eta} f_1(s)$ by $\lambda_{1,\eta} f_1(s) - \gamma(\eta)$, for some function $\gamma(\eta) \rightarrow 0$ if $\eta \rightarrow 0$; furthermore, (6.8) holds a.e. except for set of measure $\leq \tilde{\gamma}(\eta)$ where $\tilde{\gamma}(\eta) \rightarrow 0$ if $\eta \rightarrow 0$.

REMARK 6.3. Theorem 2.1 can actually also be proved using the penalized functional

$$\int_{-\infty}^0 g u ds - \frac{1}{\eta} \left(\int_{-\infty}^0 f u ds - \alpha \right)^2 - \int_{-\infty}^0 \frac{(u - u_0)^2}{1 + s^2} ds.$$

REMARK 6.4. Consider the problem

$$(6.11) \quad \max_{u \in K} J_\tau(u)$$

where K is the set of all inputs u that satisfy

$$(6.12a) \quad \int_{-\infty}^{t_j} h(t_j - s) u(s) ds = \alpha_j, \quad j = 1, \dots, N,$$

and where $J_\tau(u)$ is defined by (1.2) and $0 = t_1 < t_2 < \dots < t_{N-1} < t_N \equiv \tau$.

Set

$$(6.13a) \quad \sigma^+(\tau; t_1, \alpha_1, \dots, t_N, \alpha_N) = \max_{u \in K} J_\tau(u),$$

$$(6.13b) \quad \sigma^-(\tau; t_1, \alpha_1, \dots, t_N, \alpha_N) = \min_{u \in K} J_\tau(u).$$

Then there exists a solution to (6.11) if and only if

$$(6.14) \quad \begin{cases} |\alpha_1| \leq \int_0^\infty |h(s)| ds, \\ \sigma^-(t_j; t_1, \alpha_1, \dots, t_{j-1}, \alpha_{j-1}) \leq \alpha_j \leq \sigma^+(t_j; t_1, \alpha_1, \dots, t_{j-1}, \alpha_{j-1}), \\ 1 < j \leq N. \end{cases}$$

Assuming $h(s) = 0$ for $s < 0$ (6.12a) can be written as

$$(6.15) \quad \int_{-\infty}^{\tau} h(t_j - s)u(s) ds = \alpha_j, \quad j = 1, \dots, N,$$

so that Theorem 6.1 implies that

$$(6.16) \quad u(s) = \begin{cases} \operatorname{sgn} [h(\tau - s) - \sum_{i=1}^j \lambda_i h(t_i - s)], & t_j < s < t_{j+1} \\ \operatorname{sgn} [h(\tau - s) - \sum_{i=1}^N \lambda_i h(t_i - s)], & s < t_1, \end{cases}$$

where $\lambda_1, \dots, \lambda_N$ are chosen to satisfy (6.12a). In general, it is necessary to resort to numerical techniques to compute $\lambda_1, \dots, \lambda_N$.

REFERENCES

- [1] M.L. HONIG, S. BOYD AND B. GOPINATH, *On optimum signal sets for digital communications with finite precision and amplitude constraints*, Proc. IEEE Globecom. Conference, Tokyo, Japan, Nov. 1987.
- [2] M.L. HONIG, K. STEIGLITZ AND B. GOPINATH, *Bounds on Maximum Throughput for Digital Communications with Finite-Precision and Amplitude Constraints*, ICASSP, April 1988, New York City.
- [3] M.L. HONIG, K. STEIGLITZ, S. BOYD, AND B. GOPINATH, *Bounds on Maximum Throughput for Digital Communications with Finite-Precision and Amplitude Constraints*, in preparation.