**SOME NECESSARY CONDITIONS AT AN INTERNAL
BOUNDARY FOR MINIMIZERS IN FINITE ELASTICITY**

By

Henry C. Simpson

and

Scott J. Spector

IMA Preprint Series # 521

April 1989

1. Introduction	1
2. Notations	5
3. The Constitutive Relation	6
4. The Energy. Minimizers	9
5. Necessary Conditions	13
6. Sufficient Conditions	18
7. An Internal Boundary Between Two Isotropic Materials	20

This work was supported in part by the National Science Foundation (DMS 86-00281, DMS 86-02201, and DMS 88-10653). It was also supported by the Air Force Office of Scientific Research (AFOSR 86-0184 and AFOSR 88-0200).

SOME NECESSARY CONDITIONS AT AN INTERNAL BOUNDARY FOR MINIMIZERS IN FINITE ELASTICITY

by

Henry C. Simpson* & Scott J. Spector**

*Department of Mathematics, University of Tennessee,
Knoxville, TN 37996-1300, USA.

**Department of Mathematics, Southern Illinois University,
Carbondale, IL 62901-4408, USA.

1. Introduction

Consider a deformed body that contains surfaces across which the material properties are not continuous such as a twin boundary in a crystal, a boundary between shear bands in a metal or polymer, or a boundary at which two distinct materials have been joined. Assuming that the body is elastic at equilibrium it is of interest to determine constitutive restrictions upon the stored-energy density that are consequences of the observed stability of the body.

In this paper we address the problem of finding pointwise algebraic conditions upon the elasticity tensors $\underline{C}^+(\underline{x}_0, \underline{\nabla} \underline{f}^+(\underline{x}_0))$ and $\underline{C}^-(\underline{x}_0, \underline{\nabla} \underline{f}^-(\underline{x}_0))$ that are necessary for a continuous, piecewise smooth deformation $\underline{f}$ to be a weak relative minimizer, i.e.; a local minimizer of the energy in the C^1 -topology. In particular we show that the Legendre-Hadamard condition, Agmon's condition,

and two new conditions: if for some vector $\underline{e}$

$$\underline{e} \otimes \underline{n}_0 \cdot \underline{C}_0^+[\underline{e} \otimes \underline{n}_0] = \underline{e} \otimes \underline{n}_0 \cdot \underline{C}_0^-[\underline{e} \otimes \underline{n}_0] = 0$$

then

$$\underline{C}_0^+[\underline{e} \otimes \underline{n}_0] = \underline{C}_0^-[\underline{e} \otimes \underline{n}_0];$$

and if for some vector $\underline{a}$

$$\underline{a} \otimes \underline{t}_0 \cdot \underline{C}_0^+[\underline{a} \otimes \underline{t}_0] = \underline{a} \otimes \underline{t}_0 \cdot \underline{C}_0^-[\underline{a} \otimes \underline{t}_0] = 0$$

then

$$\underline{C}_0^+[\underline{a} \otimes \underline{t}_0] \underline{n}_0 = \underline{C}_0^-[\underline{a} \otimes \underline{t}_0] \underline{n}_0$$

are consequences of the infinitesimal stability¹ of the deformation $\underline{f}$. Here $\underline{n}_0 = \underline{n}(\underline{x}_0)$ is the unit normal and $\underline{t}_0$ is a tangent vector to a surface $\mathcal{S}$ across which the elasticity tensor and/or deformation gradient suffer a discontinuity while $\underline{\nabla f}^\pm(\underline{x}_0)$ are the limits of the deformation gradients and $\underline{C}_0^\pm = \underline{C}^\pm(\underline{x}_0, \underline{\nabla f}^\pm(\underline{x}_0))$ are the limits of the elasticity tensors as the point $\underline{x}_0$ is approached from opposite sides of the surface $\mathcal{S}$.

Remark. Agmon's condition (cf. (3.1)-(3.2)) can be interpreted as a dynamic stability requirement for linear elastodynamics² since the failure of this condition yields a nontrivial, continuous $\underline{u}: \mathbb{R}^n \times \mathbb{R}^+ \rightarrow \mathbb{R}^n$ that satisfies

¹That is, the nonnegativity of the second variation of the energy.

²Hayes & Rivlin [7] have noted that the Legendre-Hadamard condition can also be viewed as a dynamic stability requirement for linear elastodynamics. See also Rivlin [11], Truesdell & Noll [13, Section 73] and Simpson & Spector [12].

$$\begin{aligned} \operatorname{div} C_{\approx 0}^{\pm}[\nabla u] &= u_{\approx tt} \text{ in } \mathcal{H}_{\pm}, \\ C_{\approx 0}^{+}[\nabla u^{+}]n_{\approx 0} &= C_{\approx 0}^{-}[\nabla u^{-}]n_{\approx 0} \text{ on } \partial\mathcal{H}_{\pm}, \end{aligned}$$

where

$$u(\underline{x}, t) = \exp(\alpha t) \operatorname{Re}\{z(\underline{x} \cdot \underline{n}_{\approx 0}) \exp(i \underline{x} \cdot \underline{t}_{\approx 0})\},$$

$\alpha > 0$, z is bounded, and $\underline{t}_{\approx 0}$ is tangent and $\underline{n}_{\approx 0}$ is normal to the boundary of the two complementary half-spaces $\mathcal{H}_{+}$ and $\mathcal{H}_{-}$. Thus the failure of Agmon's condition implies that the rest state, $u = 0$, is unstable since there are initial data, arbitrarily close to 0 that give rise to a solution growing exponentially in time.

In Section 7 we apply our results to a boundary at which two linearly elastic isotropic materials have been joined. Here the elasticity tensors are given by

$$C_{\approx 0}^{\pm}[\underline{H}] = \mu^{\pm}(\underline{H} + \underline{H}^T) + \lambda^{\pm}(\operatorname{trace} \underline{H})\underline{I}.$$

In particular, we show that given the strong ellipticity of $C_{\approx 0}^{+}$ and $C_{\approx 0}^{-}$ (i.e., given $\mu^{\pm} > 0$ and $2\mu^{\pm} + \lambda^{\pm} > 0$)

$$\mu^{-} \geq \frac{-\mu^{+}(\mu^{+} + \lambda^{+})}{3\mu^{+} + \lambda^{+}}, \quad \mu^{+} \geq \frac{-\mu^{-}(\mu^{-} + \lambda^{-})}{3\mu^{-} + \lambda^{-}}$$

are necessary conditions for the reference configuration to be infinitesimally stable.

James [8] and Gurtin [6] have considered the problem of determining constitutive restrictions upon the stored-energy density that are necessary for a continuous, piecewise smooth deformation to be a strong relative minimizer, i.e., a local

minimizer of the energy in the C^0 -topology. In particular, James has noted that for a smooth stored-energy density the Maxwell relation must be satisfied at every point on the surface where the deformation gradient is discontinuous while Gurtin has obtained a new integral inequality³ (see (4.4)) that the stored-energy density must satisfy. Our results make crucial use of Gurtin's inequality.

We note that the analysis in this paper is essentially the same as the analysis of Simpson & Spector [12] who derive pointwise algebraic conditions on the elasticity tensor at an (external) boundary that are necessary for a deformation to be a weak relative minimizer. Both the mechanics of internal boundaries and the precise results obtained are sufficiently different to lead us to believe that the results in this paper are of interest.

Finally, we note that we have not been able to determine if the four necessary conditions that we have obtained are also sufficient for the infinitesimal stability of even a continuous, piecewise homogeneous deformation of a piecewise homogeneous body. In particular, we have been unable to find algebraic conditions that are necessary and sufficient for

$$\int_{\mathcal{H}\mathcal{B}_+} \nabla u \cdot C_0^+[\nabla u] dx + \int_{\mathcal{H}\mathcal{B}_-} \nabla u \cdot C_0^-[\nabla u] dx \geq 0$$

whenever $u \in C_0^1(\mathcal{B}, \mathbb{R}^3)$. (Here $\mathcal{H}\mathcal{B}_+$ and $\mathcal{H}\mathcal{B}_-$ are two complementary

³This inequality is similar to Morrey's [10] quasiconvexity condition.

half-balls that comprise the unit ball $\mathcal{B}$.) However in the special case when $\underline{C}_0^+$ and $\underline{C}_0^-$ are strongly-elliptic Agmon's condition alone is necessary and sufficient for the above inequality.

2. Notations

We let

$\text{Lin} :=$ space of all linear transformations from $\mathbb{R}^n$ into $\mathbb{R}^n$ with inner product and norm

$$\underline{G} \cdot \underline{H} := \text{trace } (\underline{G} \underline{H}^T), \quad |\underline{G}|^2 := \underline{G} \cdot \underline{G},$$

where $\underline{H}^T$ denotes the transpose of $\underline{H}$. We write

$$\text{Lin}^+ := \{\underline{H} \in \text{Lin} : \det \underline{H} > 0\},$$

where $\det$ denotes the determinant. Given any $\underline{M} \in \text{Lin}$ we extend its domain of definition to $\mathbb{C}^n$ and write $\underline{M}z := \underline{M}x + i \underline{M}y$ where $x := (z + \bar{z})/2$ and $y := -i(z - \bar{z})/2$. We denote by $\underline{a} \otimes \underline{b}$ the tensor product of any two vectors $\underline{a}, \underline{b} \in \mathbb{C}^n$; in components

$$(\underline{a} \otimes \underline{b})_{ij} = a_i b_j.$$

We write ∇ and div for the gradient and divergence operators in $\mathbb{R}^n$: for a vector field $\underline{u}$, $\nabla \underline{u}$ is the tensor field with components $(\nabla \underline{u})_{ij} = \partial u_i / \partial x_j$; for a tensor field $\underline{S}$, $\text{div } \underline{S}$ is the vector field with components $\sum_j \partial S_{ij} / \partial x_j$. Given any function $\Phi(\underline{a}, \underline{b}, \dots, \underline{c})$ with vector or tensor arguments we write, e.g., $\partial \Phi / \partial \underline{a}$ for the partial Fréchet derivative of Φ with respect to $\underline{a}$ holding the remaining arguments fixed.

Throughout this paper $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) will denote a bounded open region with C^1 boundary, $\partial\Omega$. Further we assume that

$$\Omega = \Omega_+ \cup \mathcal{S} \cup \Omega_-, \quad \Omega_+ \cap \Omega_- = \emptyset,$$

where Ω_+ and Ω_- are open regions with C^1 boundaries and

$$\mathcal{S} = \bar{\Omega}_+ \cap \bar{\Omega}_- \cap \Omega$$

is a C^1 surface that separates the subregions Ω_+ and Ω_- . Further we assume that

$$\partial\Omega = \mathcal{D} \cup \Gamma, \quad (2.1)$$

where $\mathcal{D}$ and Γ are disjoint and Γ is relatively open.

Given any $\underline{x}_0 \in \mathbb{R}^n$ and $r > 0$ we write

$$\mathcal{B} := B_r(\underline{x}_0) := \{\underline{y} \in \mathbb{R}^n : |\underline{y} - \underline{x}_0| < r\}$$

for the ball of radius r centered at $\underline{x}_0$. We denote by $L^2(\mathcal{B}, \mathbb{R}^n)$ the space (of equivalence classes) of square-integrable vector-valued functions on $\mathcal{B}$ and define

$$H_0^1(\mathcal{B}, \mathbb{R}^n) := \text{completion of } \{\underline{u} \in C_0^\infty(\mathcal{B}, \mathbb{R}^n) : \|\underline{u}\|_1 < \infty\}$$

with respect to the norm

$$\|\underline{u}\|_1^2 := \int_{\mathcal{B}} |\underline{u}(\underline{x})|^2 d\underline{x} + \int_{\mathcal{B}} |\nabla \underline{u}(\underline{x})|^2 d\underline{x}.$$

3. The Constitutive Relation

We consider a body that, for convenience, we identify with the region $\bar{\Omega}$ that it occupies in a fixed reference configuration. A (two-phase) deformation of the body is a member of the space

$$\text{Def} := \{\underline{f} \in C^0(\bar{\Omega}, \mathbb{R}^n) : \underline{f}^\pm \in C^2(\bar{\Omega}_\pm, \mathbb{R}^n), \det \nabla \underline{f}^\pm > 0\},$$

where

$$\underset{\sim}{f}^+ : \bar{\Omega}_+ \rightarrow \mathbb{R}^n, \quad \underset{\sim}{f}^- : \bar{\Omega}_- \rightarrow \mathbb{R}^n$$

denote the restrictions of $\underset{\sim}{f}$ to $\bar{\Omega}_+$ and $\bar{\Omega}_-$, respectively.

We assume that the subbodies $\bar{\Omega}_-$ and $\bar{\Omega}_+$ are each hyperelastic with continuous response functions $W^\pm : \bar{\Omega}_\pm \times \text{Lin} \rightarrow \mathbb{R} \cup \{+\infty\}$. $W^\pm$ gives the stored energy

$$W^\pm(\underset{\sim}{x}, \underset{\sim}{\nabla f}^\pm(\underset{\sim}{x}))$$

at any point $\underset{\sim}{x} \in \bar{\Omega}_\pm$ when the body is deformed by $\underset{\sim}{f} \in \text{Def}$.

We further assume that $W^\pm$ restricted to $\bar{\Omega}_\pm \times \text{Lin}^+$ are C^2 and finite-valued. The derivatives

$$\underset{\sim}{S}^\pm(\underset{\sim}{x}, \underset{\sim}{F}) = \frac{\partial}{\partial \underset{\sim}{F}} \underset{\sim}{W}^\pm(\underset{\sim}{x}, \underset{\sim}{F}), \quad \underset{\sim}{C}^\pm(\underset{\sim}{x}, \underset{\sim}{F}) := \frac{\partial^2}{\partial \underset{\sim}{F}^2} \underset{\sim}{W}^\pm(\underset{\sim}{x}, \underset{\sim}{F})$$

are called the (Piola-Kirchhoff) stress and the elasticity tensor, respectively. To simplify the notation we will

occasionally drop the superscripts + and -; thus $\underset{\sim}{W}$, $\underset{\sim}{S}$, and $\underset{\sim}{C}$ will denote the stored energy, stress, and the elasticity tensor, respectively.

Given $\underset{\sim}{x}_0 \in \bar{\Omega}$ and $\underset{\sim}{f} \in \text{Def}$ we write $\underset{\sim}{C}_0^\pm := \underset{\sim}{C}^\pm(\underset{\sim}{x}_0, \underset{\sim}{\nabla f}^\pm(\underset{\sim}{x}_0))$. We say that $\underset{\sim}{C}_0$ satisfies the Legendre-Hadamard condition provided that

$$\underset{\sim}{a} \otimes \underset{\sim}{b} \cdot \underset{\sim}{C}_0[\underset{\sim}{a} \otimes \underset{\sim}{b}] \geq 0$$

whenever $\underset{\sim}{a}, \underset{\sim}{b} \in \mathbb{R}^n$. We say that $\underset{\sim}{C}_0$ is strongly elliptic provided that

$$\underset{\sim}{a} \otimes \underset{\sim}{b} \cdot \underset{\sim}{C}_0[\underset{\sim}{a} \otimes \underset{\sim}{b}] > 0$$

whenever $\underset{\sim}{a}, \underset{\sim}{b} \in \mathbb{R}^n$ with $\underset{\sim}{a} \neq 0$ and $\underset{\sim}{b} \neq 0$.

For $\underline{x}_0 \in \mathcal{L}$ we write $\underline{n}_0 = \underline{n}(\underline{x}_0)$ for the unit normal to $\mathcal{L}$ that points into Ω_- and $\mathcal{H}_\pm$ for the half-spaces

$$\mathcal{H}_+ := \{\underline{y} \in \mathbb{R}^n : (\underline{x}_0 - \underline{y}) \cdot \underline{n}_0 > 0\}, \quad \mathcal{H}_- := \{\underline{y} \in \mathbb{R}^n : (\underline{x}_0 - \underline{y}) \cdot \underline{n}_0 < 0\}.$$

The remark in the introduction then motivates us to consider the problem: Find $\underline{w} \in C^0(\mathbb{R}^n, \mathbb{C}^n) \cap C^2(\overline{\mathcal{H}}_+, \mathbb{C}^n) \cap C^2(\overline{\mathcal{H}}_-, \mathbb{C}^n)$ that satisfies

$$\begin{aligned} \operatorname{div} C_0^\pm[\nabla \underline{w}] &= \alpha^2 \underline{w} \text{ in } \mathcal{H}_\pm, \\ C_0^+[\nabla \underline{w}^+] \underline{n}_0 &= C_0^-[\nabla \underline{w}^-] \underline{n}_0 \text{ on } \partial \mathcal{H}, \end{aligned} \quad (3.1)$$

where $\alpha > 0$ and $\underline{w}^+$ and $\underline{w}^-$ are the restrictions of $\underline{w}$ to $\mathcal{H}_+$ and $\mathcal{H}_-$, respectively.

We are specifically interested in whether or not there are solutions of (3.1) that are bounded exponentials, that is, functions

$$\underline{w}(\underline{y}) = \underline{z}((\underline{x}_0 - \underline{y}) \cdot \underline{n}_0) \exp(i(\underline{y} - \underline{x}_0) \cdot \underline{t}) \quad (3.2)$$

for some $\underline{t} \in \mathbb{R}^n$ that is tangent to $\partial \mathcal{H}$ ($\underline{t} \cdot \underline{n}_0 = 0$) and some $\underline{z} \in C^0(\mathbb{R}, \mathbb{C}^n) \cap C^2([0, \infty), \mathbb{C}^n) \cap C^2((-\infty, 0], \mathbb{C}^n)$ satisfying

$$\sup_s |z(s)| < +\infty.$$

We say that the triple $(C_0^+, C_0^-, \underline{n}_0)$ satisfies Agmon's condition⁴ at an internal boundary provided that for every real $\alpha > 0$ the only bounded exponential solution of (3.1) is $\underline{w} = 0$.

Remark. Suppose that (3.1) has a bounded exponential solution $\underline{w}_\alpha$ for some $\alpha > 0$. Then (3.1) has such a solution for all $\alpha > 0$.

⁴Cf. Agmon [1, p. 123].

The required solution is given by

$$\underline{\underline{w}}_{\underline{\underline{\beta}}}(\underline{\underline{x}}) = \underline{\underline{w}}_{\underline{\underline{\alpha}}}(\frac{\underline{\underline{\beta}}}{\underline{\underline{\alpha}}}(\underline{\underline{x}} - \underline{\underline{x}}_0) + \underline{\underline{x}}_0).$$

4. The Energy. Minimizers

We assume that in each deformation $\underline{\underline{f}}$ the environment exerts a body force $\underline{\underline{b}}_{\underline{\underline{f}}}(\underline{\underline{x}})$ on the points $\underline{\underline{x}} \in \underline{\underline{\Omega}}$ and a surface force $\underline{\underline{s}}_{\underline{\underline{f}}}(\underline{\underline{x}})$ on the points $\underline{\underline{x}} \in \underline{\underline{\Gamma}}$ (cf. (2.1)). We assume further that there are potentials $\underline{\underline{\beta}} \in C^2(\underline{\underline{\Omega}} \times \mathbb{R}^n, \mathbb{R})$ and $\underline{\underline{\sigma}} \in C^2(\underline{\underline{\Gamma}} \times \mathbb{R}^n, \mathbb{R})$ such that

$$\underline{\underline{b}}_{\underline{\underline{f}}}(\underline{\underline{x}}) = \frac{\partial}{\partial \underline{\underline{f}}} \underline{\underline{\beta}}(\underline{\underline{x}}, \underline{\underline{f}}(\underline{\underline{x}})), \quad \underline{\underline{s}}_{\underline{\underline{f}}}(\underline{\underline{x}}) = \frac{\partial}{\partial \underline{\underline{f}}} \underline{\underline{\sigma}}(\underline{\underline{x}}, \underline{\underline{f}}(\underline{\underline{x}}))$$

for any $\underline{\underline{f}} \in \text{Def}$. We let

$$E(\underline{\underline{f}}) := \int_{\underline{\underline{\Omega}}_+} W^+(\nabla \underline{\underline{f}}^+) + \int_{\underline{\underline{\Omega}}_-} W^-(\nabla \underline{\underline{f}}^-) - \int_{\underline{\underline{\Omega}}} \underline{\underline{\beta}}(\underline{\underline{f}}) - \int_{\underline{\underline{\Gamma}}} \underline{\underline{\sigma}}(\underline{\underline{f}}) \quad (4.1)$$

denote the total energy when the body is deformed by $\underline{\underline{f}}$.

We are interested in a deformation $\underline{\underline{f}}$ that minimizes the total energy among certain classes of competing deformations, i.e.,

$$E(\underline{\underline{f}}) \leq E(\underline{\underline{f}} + \underline{\underline{u}}) \quad (4.2)$$

for certain variations

$$\underline{\underline{u}} \in \text{Var} := \{ \underline{\underline{u}} \in C^1(\underline{\underline{\Omega}}, \mathbb{R}^n) : \underline{\underline{u}} = 0 \text{ on } \underline{\underline{\mathcal{D}}} \}.$$

Definition. Let $\underline{\underline{f}} \in C^0(\underline{\underline{\Omega}}, \mathbb{R}^n)$ be such that $\underline{\underline{f}}^\pm \in C^1(\underline{\underline{\Omega}}_\pm, \mathbb{R}^n)$ and $E(\underline{\underline{f}}) < +\infty$. We say that $\underline{\underline{f}}$ is a

(i) strong local minimizer provided that there is an $\epsilon > 0$ such that (4.2) is satisfied by all $\underline{\underline{u}} \in \text{Var}$ with

$$\sup_{\underline{\underline{x}} \in \underline{\underline{\Omega}}} |\underline{\underline{u}}(\underline{\underline{x}})| < \epsilon;$$

(ii) weak local minimizer provided that there is an $\epsilon > 0$ such that (4.2) is satisfied by all $u \in \text{Var}$ with

$$\sup_{\substack{x \in \Omega \\ \approx}} |u(x)| + \sup_{\substack{x \in \Omega \\ \approx}} |\nabla u(x)| < \epsilon.$$

Given any $\underline{x}_0 \in \Omega \setminus \mathcal{L}$, $f \in \text{Def}$, and $r > 0$ we define $\underline{F}_0 := \nabla f(\underline{x}_0)$.

$$\underline{f}_0(\underline{y}) := \underline{F}_0(\underline{y} - \underline{x}_0) + f(\underline{x}_0), \quad \underline{W}_0(\underline{G}) := \underline{W}(\underline{x}_0, \underline{G})$$

for any $\underline{G} \in \text{Lin}^+$ and

$$\underline{y} \in \mathcal{B} := \{\underline{y} \in \mathbb{R}^n : |\underline{y} - \underline{x}_0| < r\}.$$

We interpret $\underline{f}_0$ as a homogeneous deformation of a homogeneous body that occupies the region $\mathcal{B}$ in a homogeneous reference configuration.

If $\underline{x}_0 \in \mathcal{L}$ we write $\underline{F}_0^+ := \nabla f^+(\underline{x}_0)$, $\underline{F}_0^- := \nabla f^-(\underline{x}_0)$.

$$\begin{aligned} \underline{f}_0^+(\underline{y}) &:= \underline{F}_0^+(\underline{y} - \underline{x}_0) + f(\underline{x}_0), & \underline{W}_0^+(\underline{G}) &:= \underline{W}^+(\underline{x}_0, \underline{G}), \\ \underline{f}_0^-(\underline{z}) &:= \underline{F}_0^-(\underline{z} - \underline{x}_0) + f(\underline{x}_0), & \underline{W}_0^-(\underline{G}) &:= \underline{W}^-(\underline{x}_0, \underline{G}) \end{aligned}$$

for any $\underline{G} \in \text{Lin}^+$,

$$\begin{aligned} \underline{y} \in \mathcal{B}_+ &:= \{\underline{y} \in \mathcal{B} : (\underline{x}_0 - \underline{y}) \cdot \underline{n}_0 > 0\}, \\ \underline{z} \in \mathcal{B}_- &:= \{\underline{z} \in \mathcal{B} : (\underline{x}_0 - \underline{z}) \cdot \underline{n}_0 < 0\}. \end{aligned}$$

We interpret the pair $(\underline{f}_0^+, \underline{f}_0^-)$ as a piecewise homogeneous deformation of the piecewise homogeneous body that occupies the region $\mathcal{B}$ in a piecewise homogeneous reference configuration.

Proposition 4.1. Let f be a strong local minimizer. Then,⁵ for every $x_0 \in \Omega \setminus \mathcal{L}$,

$$\int_{\mathcal{B}} W_0(F_0) dy \leq \int_{\mathcal{B}} W_0(F_0 + \nabla u(y)) dy \quad (4.3)^6$$

for all $u \in C_0^1(\mathcal{B}, \mathbb{R}^n)$ such that $W_0(F_0 + \nabla u(\cdot))$ is uniformly bounded on $\mathcal{B}$. Moreover,⁷ for every $x_0 \in \mathcal{L}$,

$$\left[\begin{array}{c} \int_{\mathcal{H}\mathcal{B}_+} W_0^+(F_0^+) dy \\ + \\ \int_{\mathcal{H}\mathcal{B}_-} W_0^-(F_0^-) dy \end{array} \right] \leq \left[\begin{array}{c} \int_{\mathcal{H}\mathcal{B}_+} W_0^+(F_0^+ + \nabla u(y)) dy \\ + \\ \int_{\mathcal{H}\mathcal{B}_-} W_0^-(F_0^- + \nabla u(y)) dy \end{array} \right] \quad (4.4)$$

for all $u \in C_0^1(\mathcal{B}, \mathbb{R}^n)$ such that $W_0^\pm(F_0^\pm + \nabla u(\cdot))$ are uniformly bounded on $\mathcal{H}\mathcal{B}_\pm$, respectively.

In other words, in order for f to be a strong local minimizer each of the homogeneous deformations $f_0(y) = [\nabla f(x_0)](y - x_0) + f(x_0)$, $x_0 \in \Omega \setminus \mathcal{L}$, must be a global minimizer for the pure displacement problem on a ball composed of the material at x_0 and each of the piecewise homogeneous deformations (f_0^+, f_0^-)

⁵This result is essentially due to Meyers [9, pp. 128-131]. See Ball [2, Theorem 2.1] or Gurtin [6, pp. 9-10] for a direct proof when W is finite-valued and Ball & Marsden [3] for a proof under our hypotheses.

⁶This inequality is Morrey's [10] quasiconvexity condition.

⁷This result is due to Gurtin [6, pp. 9-10]. See Ball & Marsden [3] for a proof under our hypotheses.

$(f_{\underline{\underline{0}}}^+(y) = [\nabla f_{\underline{\underline{0}}}^+(x_0)](\underline{\underline{y}} - \underline{\underline{x}}_0) + f_{\underline{\underline{0}}}^+(x_0), f_{\underline{\underline{0}}}^-(y) = [\nabla f_{\underline{\underline{0}}}^-(x_0)](\underline{\underline{y}} - \underline{\underline{x}}_0) + f_{\underline{\underline{0}}}^-(x_0)),$
 $\underline{\underline{x}}_0 \in \mathcal{A}$, must be a global minimizer for the pure displacement problem on a pair of half-balls each of which is composed of one of the two materials at $\underline{\underline{x}}_0$.

We now suppose that $\underline{\underline{f}} \in \text{Def}$ is a weak local minimizer. Then, for every $\underline{\underline{u}} \in \text{Var}$

$$0 \leq \varphi(t) := E(\underline{\underline{f}} + t\underline{\underline{u}}) - E(\underline{\underline{f}})$$

provided t is sufficiently small. Thus $\dot{\varphi}(0) = 0$ and $\ddot{\varphi}(0) \geq 0$.

If $\underline{\underline{f}}$ is C^2 on $\Omega^+ \cup \Omega^-$ then one can show that $\underline{\underline{f}}$ is equilibrated, that is

$$\begin{aligned} \text{div } S_{\underline{\underline{f}}}(\nabla f) + b_{\underline{\underline{f}}} &= 0 \text{ in } \Omega^+ \cup \Omega^-, \\ S_{\underline{\underline{f}}}(\nabla f)N &= s_{\underline{\underline{f}}} \text{ on } \Gamma, \\ S_{\underline{\underline{f}}}^+(\nabla f^+)n &= S_{\underline{\underline{f}}}^-(\nabla f^-)n \text{ on } \mathcal{A}, \end{aligned}$$

where $N = N(x)$ is the outward unit normal to $\partial\Omega$ at x . In addition (whether or not $\underline{\underline{f}}$ is C^2) the second variation of E at $\underline{\underline{f}}$ must be nonnegative, i.e.,

$$0 \leq \int_{\Omega^+} \nabla u \cdot C^+[\nabla u] + \int_{\Omega^-} \nabla u \cdot C^-[\nabla u] - \int_{\Omega} u \cdot Bu - \int_{\Gamma} u \cdot \Sigma u \quad (4.5)$$

for all $\underline{\underline{u}} \in \text{Var}$, where

$$B_{\underline{\underline{f}}}(\underline{\underline{x}}, f(\underline{\underline{x}})) := \frac{\partial^2}{\partial f^2} \beta(\underline{\underline{x}}, f(\underline{\underline{x}})), \quad \Sigma_{\underline{\underline{f}}}(\underline{\underline{x}}, f(\underline{\underline{x}})) := \frac{\partial^2}{\partial f^2} \sigma(\underline{\underline{x}}, f(\underline{\underline{x}})).$$

If we now define

$$\begin{aligned} \hat{W}_{\underline{\underline{f}}}^{\pm}(\underline{\underline{x}}, U) &:= U \cdot C_{\underline{\underline{f}}}^{\pm}(\underline{\underline{x}}, \nabla f^{\pm}(\underline{\underline{x}})) [U], \\ \hat{\beta}_{\underline{\underline{f}}}(\underline{\underline{x}}, v) &:= v \cdot B_{\underline{\underline{f}}}(\underline{\underline{x}}, f(\underline{\underline{x}})) v, \quad \hat{\sigma}_{\underline{\underline{f}}}(\underline{\underline{x}}, v) := v \cdot \Sigma_{\underline{\underline{f}}}(\underline{\underline{x}}, f(\underline{\underline{x}})) v, \\ \hat{E}_{\underline{\underline{f}}}(\underline{\underline{u}}) &:= \int_{\Omega^+} \hat{W}_{\underline{\underline{f}}}^+(\nabla u) + \int_{\Omega^-} \hat{W}_{\underline{\underline{f}}}^-(\nabla u) - \int_{\Omega} \hat{\beta}_{\underline{\underline{f}}}(\underline{\underline{u}}) - \int_{\Gamma} \hat{\sigma}_{\underline{\underline{f}}}(\underline{\underline{u}}), \end{aligned}$$

we find that (4.5) is equivalent to $\hat{E}(\underline{0}) \leq \hat{E}(\underline{u})$ for all $\underline{u} \in \text{Var}$. Thus the function $\underline{0}$ is a strong local minimizer of $\hat{E}$. Therefore by Proposition 4.1, the density of $C_0^1(\mathcal{B}, \mathbb{R}^n)$ in $H_0^1(\mathcal{B}, \mathbb{R}^n)$, and the continuity of the resulting quadratic form on $H_0^1(\mathcal{B}, \mathbb{R}^n)$ we arrive at the following result.

Proposition 4.2. Let $f \in \text{Def}$ be a weak local minimizer. Then, for every $\underline{x}_0 \in \mathcal{L}$,

$$0 \leq \int_{\mathcal{H}\mathcal{B}_+} \nabla u(\underline{y}) \cdot C_{\underline{0}}^+[\nabla u(\underline{y})] d\underline{y} + \int_{\mathcal{H}\mathcal{B}_-} \nabla u(\underline{y}) \cdot C_{\underline{0}}^-[\nabla u(\underline{y})] d\underline{y} \quad (4.6)$$

for all $\underline{u} \in H_0^1(\mathcal{B}, \mathbb{R}^n)$. Here $C_{\underline{0}}^\pm := C^\pm(\underline{x}_0, \nabla f^\pm(\underline{x}_0))$.

5. Necessary Conditions

Theorem 1. Let $f \in \text{Def}$ be a weak local minimizer. Then, for every $\underline{x}_0 \in \mathcal{L}$ ($= \bar{\Omega}_+ \cap \bar{\Omega}_- \cap \Omega$),

- (i) $C_{\underline{0}}^+$ and $C_{\underline{0}}^-$ satisfy the Legendre-Hadamard condition;
- (ii) $(C_{\underline{0}}^+, C_{\underline{0}}^-, n_{\underline{0}})$ satisfies Agmon's condition;
- (iii) if, for some $\underline{a} \in \mathbb{R}^n$,

$$\underline{a} \otimes \underline{n}_{\underline{0}} \cdot C_{\underline{0}}^+[\underline{a} \otimes \underline{n}_{\underline{0}}] = \underline{a} \otimes \underline{n}_{\underline{0}} \cdot C_{\underline{0}}^-[\underline{a} \otimes \underline{n}_{\underline{0}}] = 0$$

then

$$C_{\underline{0}}^+[\underline{a} \otimes \underline{n}_{\underline{0}}] = C_{\underline{0}}^-[\underline{a} \otimes \underline{n}_{\underline{0}}];$$

- (iv) if, for some $\underline{a} \in \mathbb{R}^n$,

$$\underline{a} \otimes \underline{t}_{\underline{0}} \cdot C_{\underline{0}}^+[\underline{a} \otimes \underline{t}_{\underline{0}}] = \underline{a} \otimes \underline{t}_{\underline{0}} \cdot C_{\underline{0}}^-[\underline{a} \otimes \underline{t}_{\underline{0}}] = 0$$

then

$$C_{\underline{0}}^+[\underline{a} \otimes \underline{t}_{\underline{0}}]_{n_{\underline{0}}} = C_{\underline{0}}^-[\underline{a} \otimes \underline{t}_{\underline{0}}]_{n_{\underline{0}}}.$$

Here, $C_{\underline{0}}^\pm := C^\pm(\underline{x}_0, \nabla f^\pm(\underline{x}_0))$ ($C^\pm = \partial^2 W^\pm / \partial (\nabla f^\pm)^2$), $n_{\underline{0}} := n(\underline{x}_0)$ the

unit normal to $\mathcal{X}$ that points into Ω_- , and $\underline{t}_0$ is tangent to $\mathcal{X}$ at $\underline{x}_0$.

Proof.⁸ We first apply Proposition 4.2 to conclude that (4.6) is satisfied at every $\underline{x}_0 \in \mathcal{X}$. The Legendre-Hadamard condition is then a well-known consequence of (4.6).⁹ We show that (ii)-(iv) also follow from (4.6). In order to simplify the notation we first change coordinates so that $\underline{x}_0 = \underline{0}$.

(4.6) $\Rightarrow$ (ii). Let $b \in C^\infty(\mathbb{R}, \mathbb{R})$ satisfy $b(s) = 1$ for $s \leq r/3$ and $b(s) = 0$ for $s \geq 2r/3$. Define $\varphi(\underline{x}) := b(|\underline{x}|)$. Then given $\underline{w} \in C^0(\mathbb{R}^n, \mathbb{C}^n) \cap C^2(\mathcal{H}_+, \mathbb{C}^n) \cap C^2(\mathcal{H}_-, \mathbb{C}^n)$ the functions $\underline{u} := \operatorname{Re}\{\varphi \underline{w}\}$ and $\underline{v} := \operatorname{Im}\{\varphi \underline{w}\}$ are contained in $H_0^1(\mathcal{B}, \mathbb{R}^n)$ and a straightforward computation (using the symmetry of $\underline{C}_0^\pm$) shows that

$$\begin{aligned} \int_{\mathcal{H}_\pm} (\underline{v} \underline{u} \cdot \underline{C}_0^\pm[\underline{v} \underline{u}] + \underline{v} \underline{v} \cdot \underline{C}_0^\pm[\underline{v} \underline{v}]) &= \int_{\mathcal{H}_\pm} \underline{v}(\varphi \underline{w}) \cdot \underline{C}_0^\pm[\underline{v}(\varphi \underline{w})] \\ &= \int_{\mathcal{H}_\pm} (\underline{w} \otimes \underline{v} \varphi \cdot \underline{C}_0^\pm[\underline{w} \otimes \underline{v} \varphi] + \operatorname{Re}\{\underline{v}(\varphi^2 \underline{w}) \cdot \underline{C}_0^\pm[\underline{v} \underline{w}]\}) \end{aligned} \quad (5.1)$$

and, by the divergence theorem,

$$\int_{\mathcal{H}_\pm} \underline{v}(\varphi^2 \underline{w}) \cdot \underline{C}_0^\pm[\underline{v} \underline{w}] = - \int_{\mathcal{H}_\pm} \varphi^2 \underline{w} \cdot \operatorname{div} \underline{C}_0^\pm[\underline{v} \underline{w}] + \int_{\partial(\mathcal{H}_\pm)} \varphi^2 \underline{w} \cdot \underline{C}_0^\pm[\underline{v} \underline{w}] \underline{N}^\pm, \quad (5.2)$$

where $\underline{N}^\pm$ is the outward unit normal to $\partial(\mathcal{H}_\pm)$.

⁸This proof is almost identical to the proof of Theorem 2 in [12] where necessary conditions at an (external) boundary for a minimizer are obtained.

⁹Cf., e.g., Truesdell & Noll [13, Section 68bis].

We now suppose that Agmon's condition is not satisfied and let $w_{\underline{1}}$ be a bounded exponential solution of (3.1) with $\alpha = 1$ (see the remark following the definition of Agmon's condition). Then $w_{\underline{\alpha}}(x) := w_{\underline{1}}(\alpha x)$ will be a bounded exponential solution of (3.1) for any $\alpha > 0$. Moreover, $|w_{\underline{\alpha}}(x)|$ is uniformly bounded independently of α .

We note that $N_{\underline{0}}^+ = n_{\underline{0}}$ and $N_{\underline{0}}^- = -n_{\underline{0}}$ on $\partial\mathcal{H}$. Thus by (3.1), (5.1), and (5.2) we find that there is a constant $k > 0$ such that

$$\begin{aligned} & \int_{\mathcal{H}\mathcal{B}_+} (\nabla u_{\underline{\alpha}} \cdot C_{\underline{0}}^+[\nabla u_{\underline{\alpha}}] + \nabla v_{\underline{\alpha}} \cdot C_{\underline{0}}^+[\nabla v_{\underline{\alpha}}]) + \int_{\mathcal{H}\mathcal{B}_-} (\nabla u_{\underline{\alpha}} \cdot C_{\underline{0}}^-[\nabla u_{\underline{\alpha}}] + \nabla v_{\underline{\alpha}} \cdot C_{\underline{0}}^-[\nabla v_{\underline{\alpha}}]) \\ & \leq k - \alpha^2 \int_{\mathcal{B}} \varphi^2 |w_{\underline{\alpha}}|^2 \leq k - \alpha^2 \int_{B_{r/3}} |w_{\underline{\alpha}}|^2, \end{aligned} \quad (5.3)$$

where $B_{r/3}$ is the ball of radius $r/3$ centered at $x_{\underline{0}}$.

We next note that $w_{\underline{\alpha}}$ is of the form (3.2) and hence $|w_{\underline{\alpha}}(x)| = |z_{\underline{\alpha}}(s)|$. Thus

$$\begin{aligned} \int_{B_{r/3}} |w_{\underline{\alpha}}(x)|^2 dx & \geq (2r/3n)^{n-1} \int_{-r/3n}^{r/3n} |z_{\underline{\alpha}}(s)|^2 ds \\ & = (2r/3n)^{n-1} \int_{-r/3n}^{r/3n} |z_{\underline{1}}(\alpha s)|^2 ds \\ & = \alpha^{-1} (2r/3n)^{n-1} \int_{-\alpha r/3n}^{\alpha r/3n} |z_{\underline{1}}(\sigma)|^2 d\sigma. \end{aligned}$$

If we combine the last inequality with (5.3) we conclude that, for $\alpha \geq 3n$,

$$\left[\begin{aligned} & \int_{\mathcal{H}\mathcal{B}_+} \{\nabla u_{\underline{\alpha}} \cdot C_{\underline{0}}^+[\nabla u_{\underline{\alpha}}] + \nabla v_{\underline{\alpha}} \cdot C_{\underline{0}}^+[\nabla v_{\underline{\alpha}}]\} \\ & + \int_{\mathcal{H}\mathcal{B}_-} \{\nabla u_{\underline{\alpha}} \cdot C_{\underline{0}}^-[\nabla u_{\underline{\alpha}}] + \nabla v_{\underline{\alpha}} \cdot C_{\underline{0}}^-[\nabla v_{\underline{\alpha}}]\} \end{aligned} \right] \leq k - \alpha (2r/3n)^{n-1} \int_{-r}^r |z_{\underline{1}}(\sigma)|^2 d\sigma.$$

Therefore, by choosing α sufficiently large, we can find a

function in $H_0^1(\mathcal{B}, \mathbb{R}^n)$ for which (4.6) is not satisfied.

(4.6) $\Rightarrow$ (iv). Suppose that $\underline{a} \in \mathbb{R}^n$ satisfies

$$\underline{a} \otimes \underline{t}_0 \cdot \underline{C}_{\underline{n}_0}^+[\underline{a} \otimes \underline{t}_0] = \underline{a} \otimes \underline{t}_0 \cdot \underline{C}_{\underline{n}_0}^-[\underline{a} \otimes \underline{t}_0] = 0, \quad (5.4)$$

where $\underline{t}_0 \perp \underline{n}_0$ and $|\underline{t}_0| = 1$. Let $\underline{b} \in \mathbb{R}^n$. Choose coordinates so that $\underline{n}_0 = (-1, 0, 0, \dots, 0)$, $\underline{t}_0 = (0, 1, 0, 0, \dots, 0)$ and write $\underline{x} = (s, t, z)$, where $s, t \in \mathbb{R}$ and $\underline{z} \in \mathbb{R}^{n-2}$. Let $\varphi, \psi \in C^\infty(\mathbb{R}, \mathbb{R})$ and $\underline{\rho} \in C^\infty(\mathbb{R}^{n-2}, \mathbb{R})$ satisfy (recall that r is the radius of the ball $\mathcal{B}$)

$$\left. \begin{aligned} \int_{-1}^1 |\dot{\psi}(t)|^2 dt &= 1, & \psi(t) &= 0 & \text{for } |t| \geq 1, \\ \varphi(0) &= 1, & \varphi(s) &= 0 & \text{for } |s| \geq 1, \\ \int_{|\underline{z}| \leq r} |\underline{\rho}(\underline{z})|^2 d\underline{z} &= 1, & \underline{\rho}(\underline{z}) &= 0 & \text{for } |\underline{z}| \geq r/3. \end{aligned} \right\} \quad (5.5)$$

For $\epsilon > 0$, $\delta > 0$, and $\beta > 0$ define

$$\underline{u}_{\delta, \epsilon}^\beta(s, t, \underline{z}) = \underline{\rho}(\underline{z}) [\underline{a} \delta^{-1} \varphi(\frac{\epsilon s}{\beta}) \psi(\frac{t}{\beta}) + \underline{b} \delta \varphi(\frac{s}{\beta}) \dot{\psi}(\frac{t}{\beta})]$$

Then $\underline{u}_{\delta, \epsilon}^\beta \in C_0^\infty(\overline{\mathcal{B}}, \mathbb{R}^n)$ (provided $\beta < r$ and $\epsilon > \beta/r$) and

$$\underline{\nabla} \underline{u}_{\delta, \epsilon}^\beta = \begin{bmatrix} \beta^{-1} \underline{\rho}(\underline{a} \otimes \underline{n}_0 \delta^{-1} \epsilon \dot{\varphi} \psi + \underline{b} \otimes \underline{n}_0 \delta \dot{\varphi} \dot{\psi}) \\ + \beta^{-1} \underline{\rho}(\underline{a} \otimes \underline{t}_0 \delta^{-1} \varphi \dot{\psi} + \underline{b} \otimes \underline{t}_0 \delta \varphi \ddot{\psi}) \\ + (\underline{a} \delta^{-1} \varphi \psi + \underline{b} \delta \varphi \dot{\psi}) \otimes \underline{\nabla} \underline{\rho} \end{bmatrix}. \quad (5.6)$$

Now, by (4.6),

$$0 \leq \int_{|\underline{z}| \leq r} d\underline{z} \int_{-\beta}^{\beta} dt \left[\int_0^\beta \underline{\nabla} \underline{u}_{\delta, \epsilon}^\beta \cdot \underline{C}_{\underline{n}_0}^+[\underline{\nabla} \underline{u}_{\delta, \epsilon}^\beta] ds + \int_{-\beta}^0 \underline{\nabla} \underline{u}_{\delta, \epsilon}^\beta \cdot \underline{C}_{\underline{n}_0}^-[\underline{\nabla} \underline{u}_{\delta, \epsilon}^\beta] ds \right].$$

We make the change of variables $\sigma = s/\beta$ and $\tau = t/\beta$ (and hence $ds dt = \beta^2 d\sigma d\tau$) and then let $\beta \rightarrow 0^+$ in the last inequality to

conclude, with the aid of (5.4), (5.5), and (5.6), that (assuming $\epsilon < 1$)

$$\begin{aligned}
& \underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{n}_0 \cdot \underset{\approx}{C}_0^+[\underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{n}_0] \delta^2 \int_0^1 \dot{\varphi}^2 + \underset{\approx}{a}\underset{\approx}{\theta}\underset{\approx}{n}_0 \cdot \underset{\approx}{C}_0^+[\underset{\approx}{a}\underset{\approx}{\theta}\underset{\approx}{n}_0] \delta^{-2} \epsilon^2 \int_0^{1/\epsilon} |\dot{\varphi}(\sigma\epsilon)|^2 d\sigma \int_{-1}^1 \psi^2 \\
& + \underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{n}_0 \cdot \underset{\approx}{C}_0^-[\underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{n}_0] \delta^2 \int_{-1}^0 \dot{\varphi}^2 + \underset{\approx}{a}\underset{\approx}{\theta}\underset{\approx}{n}_0 \cdot \underset{\approx}{C}_0^-[\underset{\approx}{a}\underset{\approx}{\theta}\underset{\approx}{n}_0] \delta^{-2} \epsilon^2 \int_{-1/\epsilon}^0 |\dot{\varphi}(\sigma\epsilon)|^2 d\sigma \int_{-1}^1 \psi^2 \\
& + \underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{t}_0 \cdot \underset{\approx}{C}_0^+[\underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{t}_0] \delta^2 \int_0^1 \varphi^2 \int_{-1}^1 \ddot{\varphi}^2 + 2\underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{n}_0 \cdot \underset{\approx}{C}_0^+[\underset{\approx}{a}\underset{\approx}{\theta}\underset{\approx}{t}_0] \int_0^1 \dot{\varphi}(\sigma) \varphi(\sigma\epsilon) d\sigma \\
& + \underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{t}_0 \cdot \underset{\approx}{C}_0^-[\underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{t}_0] \delta^2 \int_{-1}^0 \varphi^2 \int_{-1}^1 \ddot{\varphi}^2 + 2\underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{n}_0 \cdot \underset{\approx}{C}_0^-[\underset{\approx}{a}\underset{\approx}{\theta}\underset{\approx}{t}_0] \int_{-1}^0 \dot{\varphi}(\sigma) \varphi(\sigma\epsilon) d\sigma \\
& - 2\underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{t}_0 \cdot \underset{\approx}{C}_0^+[\underset{\approx}{a}\underset{\approx}{\theta}\underset{\approx}{n}_0] \epsilon \int_0^1 \dot{\varphi}(\sigma\epsilon) \varphi(\sigma) d\sigma \\
& - 2\underset{\approx}{b}\underset{\approx}{\theta}\underset{\approx}{t}_0 \cdot \underset{\approx}{C}_0^-[\underset{\approx}{a}\underset{\approx}{\theta}\underset{\approx}{n}_0] \epsilon \int_{-1}^0 \dot{\varphi}(\sigma\epsilon) \varphi(\sigma) d\sigma \geq 0.
\end{aligned} \tag{5.7}$$

In deriving the above inequality we have made use of the identities

$$\int_{-1}^1 \psi(\tau) \dot{\varphi}(\tau) d\tau = \int_{-1}^1 \dot{\varphi}(\tau) \ddot{\varphi}(\tau) d\tau = 0, \quad \int_{-1}^1 \psi(\tau) \ddot{\varphi}(\tau) d\tau = -1,$$

which are consequences of (5.5)₁.

We note that the change of variables $y = \sigma\epsilon$ yields

$$\begin{aligned}
\epsilon^2 \int_0^{1/\epsilon} |\dot{\varphi}(\sigma\epsilon)|^2 d\sigma &= \epsilon \int_0^1 |\dot{\varphi}(y)|^2 dy, \\
\epsilon^2 \int_{-1/\epsilon}^0 |\dot{\varphi}(\sigma\epsilon)|^2 d\sigma &= \epsilon \int_{-1}^0 |\dot{\varphi}(y)|^2 dy,
\end{aligned} \tag{5.8}$$

and by the bounded convergence theorem and (5.5)₂ we find that

$$\begin{aligned}
\lim_{\epsilon \rightarrow 0^+} \int_0^1 \dot{\varphi}(\sigma) \varphi(\sigma\epsilon) d\sigma &= \varphi(0) \int_0^1 \dot{\varphi}(y) dy = -1, \\
\lim_{\epsilon \rightarrow 0^+} \int_{-1}^0 \dot{\varphi}(\sigma) \varphi(\sigma\epsilon) d\sigma &= \varphi(0) \int_{-1}^0 \dot{\varphi}(y) dy = +1.
\end{aligned} \tag{5.9}$$

Finally, we note that φ and $\dot{\varphi}$ are bounded and let $\epsilon \rightarrow 0^+$ followed by $\delta \rightarrow 0^+$ in (5.7) to conclude, with the aid of (5.8)

and (5.9), that

$$2b_{\theta n_0} \cdot C_0^-[a_{\theta t_0}] - 2b_{\theta n_0} \cdot C_0^+[a_{\theta t_0}] \geq 0$$

and hence that

$$b \cdot (C_0^-[a_{\theta t_0}]_{n_0} - C_0^+[a_{\theta t_0}]_{n_0}) \geq 0.$$

Since b is arbitrary we find that

$$C_0^+[a_{\theta t_0}]_{n_0} = C_0^-[a_{\theta t_0}]_{n_0}.$$

(4.6) $\Rightarrow$ (iii). The variations $u_{\delta, \epsilon}^\beta$ that were used to prove that (iv) is a consequence of (4.6) will also yield this implication upon letting $\beta \rightarrow 0^+$ followed by $\epsilon \rightarrow +\infty$ and then $\delta \rightarrow 0^+$. See the proof of Theorem 2 in [12] for the details. $\square$

6. Sufficient Conditions

In this section we briefly consider the problem of finding algebraic conditions on C_0^+ and C_0^- that are sufficient for the nonnegativity of the second variation of the energy of a piecewise homogeneous body that occupies the region $\mathcal{B}$ in a piecewise homogeneous reference configuration; i.e., conditions to ensure that

$$0 \leq \int_{\mathcal{B}_+} \nabla u(y) \cdot C_0^+[\nabla u(y)] dy + \int_{\mathcal{B}_-} \nabla u(y) \cdot C_0^-[\nabla u(y)] dy \quad (6.1)$$

whenever $u \in H_0^1(\mathcal{B}, \mathbb{R}^n)$.

We do not know if conditions (i)-(iv) of Theorem 1 are sufficient for (6.1) to be satisfied. However we would like to

note that the techniques of [4, 12] can be used to show that when $C_{\approx 0}^+$ and $C_{\approx 0}^-$ are strongly elliptic Agmon's condition alone implies that (6.1) is satisfied by all $u \in H_0^1(\mathcal{B}, \mathbb{R}^n)$.

Theorem 2. Suppose that $C_{\approx 0}^+$ and $C_{\approx 0}^-$ are strongly elliptic. Then the following are equivalent:

- (a) (6.1) is satisfied by all $u \in H_0^1(\mathcal{B}, \mathbb{R}^n)$;
- (b) $(C_{\approx 0}^+, C_{\approx 0}^-, n)$ satisfies Agmon's condition;
- (c) If $t \perp n_{\approx 0}$ and $z \in C^0(\mathbb{R}, \mathbb{C}^n) \cap C^2([0, \infty), \mathbb{C}^n) \cap C^2((-\infty, 0], \mathbb{C}^n)$

is any bounded solution of

$$\left. \begin{aligned} -M_{\approx}^{+} \ddot{z} + i[N_{\approx t}^{+} + (N_{\approx t}^{+})^T] \dot{z} + P_{\approx t}^{+} z &= 0 \text{ on } (0, \infty), \\ -M_{\approx}^{-} \ddot{z} + i[N_{\approx t}^{-} + (N_{\approx t}^{-})^T] \dot{z} + P_{\approx t}^{-} z &= 0 \text{ on } (-\infty, 0) \end{aligned} \right\} \quad (6.2)$$

then

$$\bar{z}(0) \cdot [(M_{\approx}^{+} \dot{z}(0^{+}) - iN_{\approx t}^{+} z(0)) - (M_{\approx}^{-} \dot{z}(0^{-}) - iN_{\approx t}^{-} z(0))] \leq 0. \quad (6.3)$$

Here

$$\begin{aligned} M_{\approx}^{\pm} a &:= C_{\approx 0}^{\pm} [a \otimes n]_{n_{\approx 0}}, \quad N_{\approx t}^{\pm} a := C_{\approx 0}^{\pm} [a \otimes t]_{n_{\approx 0}}, \\ P_{\approx t}^{\pm} a &:= C_{\approx 0}^{\pm} [a \otimes t]_t. \end{aligned} \quad (6.4)$$

Remark. Agmon's condition can be difficult to verify for particular constitutive relations. We have found that condition (c) of Theorem 2 is sometimes simpler to check.

7. An Internal Boundary Between Two Isotropic Materials

We let

$$C_{\approx 0}^{\pm}[H] := \mu^{\pm}(\underset{\approx}{H} + \underset{\approx}{H}^T) + \lambda^{\pm}(\underset{\approx}{H} \cdot \underset{\approx}{I}) \underset{\approx}{I} \quad (7.1)$$

and consider the problem of finding necessary and sufficient conditions on $\mu^{\pm}$ and $\lambda^{\pm}$ so that (6.1) is satisfied whenever

$$\underset{\approx}{u} \in H_0^1(\mathcal{B}, \mathbb{R}^n).$$

Lemma 7.1. Let (6.1) be satisfied by all $\underset{\approx}{u} \in H_0^1(\mathcal{B}, \mathbb{R}^n)$ where $C_{\approx 0}^{\pm}$ are given by (7.1). Then

$$\mu^{\pm} \geq 0, \quad 2\mu^{\pm} + \lambda^{\pm} \geq 0. \quad (7.2)$$

Proof. We let

$$W^{\pm}(\underset{\approx}{x}, \underset{\approx}{U}) := \underset{\approx}{U} \cdot C_{\approx 0}^{\pm}[\underset{\approx}{U}]$$

and apply Theorem 1 to conclude that $C_{\approx 0}^{+}$ and $C_{\approx 0}^{-}$ each satisfy the Legendre-Hadamard condition. Thus

$$\mu^{\pm} |\underset{\approx}{a}|^2 |\underset{\approx}{b}|^2 + (\mu^{\pm} + \lambda^{\pm})(\underset{\approx}{a} \cdot \underset{\approx}{b})^2 \geq 0$$

for every $\underset{\approx}{a}, \underset{\approx}{b} \in \mathbb{R}^n$. In particular we let $\underset{\approx}{a} \neq 0$ and $\underset{\approx}{b} \neq 0$ satisfy $\underset{\approx}{a} \cdot \underset{\approx}{b} = 0$ to get (7.2)₁, while the choice $\underset{\approx}{a} = \underset{\approx}{b} \neq 0$ yields (7.2)₂. $\square$

Proposition 7.2. Let $C_{\approx 0}^{\pm}$ be given by (7.1). Suppose that

$$\mu^{\pm} > 0, \quad 2\mu^{\pm} + \lambda^{\pm} > 0. \quad (7.3)$$

Then necessary and sufficient conditions for (6.1) to be satisfied for all $\underset{\approx}{u} \in H_0^1(\mathcal{B}, \mathbb{R}^n)$ are

$$\mu^{-} \geq \frac{-\mu^{+}(\mu^{+} + \lambda^{+})}{3\mu^{+} + \lambda^{+}}, \quad \mu^{+} \geq \frac{-\mu^{-}(\mu^{-} + \lambda^{-})}{3\mu^{-} + \lambda^{-}}. \quad (7.4)$$

Proof. A standard result¹⁰ shows that (7.3) is equivalent to the strong ellipticity of $\underline{C}_0^\pm$. We will show that (7.4) is equivalent to condition (c) of Theorem 2 and the desired result will follow from Theorem 2.

For simplicity we suppose that $n = 3$ and choose coordinates so that $\underline{n} = (-1, 0, 0)$ and $\underline{t} = (0, \tau, 0)$ where $\tau \geq 0$. Then (6.4) and (7.1) imply that

$$\left. \begin{aligned} \underline{M} \underline{a} &= \mu \underline{a} + (\mu + \lambda) (\underline{a} \cdot \underline{n}) \underline{n}, \\ \underline{N}_{\underline{t}} \underline{a} &= \mu (\underline{a} \cdot \underline{n}) \underline{t} + \lambda (\underline{a} \cdot \underline{t}) \underline{n}, \\ \underline{P}_{\underline{t}} \underline{a} &= \mu \tau^2 \underline{a} + (\mu + \lambda) (\underline{a} \cdot \underline{t}) \underline{t}, \end{aligned} \right\} \quad (7.5)$$

and hence (6.2) reduces to

$$\left. \begin{aligned} -(2\mu + \lambda) \ddot{z}_1 - i\tau(\mu + \lambda) \dot{z}_2 + \mu \tau^2 z_1 &= 0, \\ -\mu \ddot{z}_2 - i\tau(\mu + \lambda) \dot{z}_1 + (2\mu + \lambda) \tau^2 z_2 &= 0, \\ -\ddot{z}_3 + \tau^2 z_3 &= 0. \end{aligned} \right\} \quad (7.6)$$

We first note that if $\tau = 0$ then the general bounded solution to (7.6) is $\underline{z}(s) = \underline{c}$ for all $s \in \mathbb{R}$ and hence we conclude, with the aid of (7.5)₂, that (6.3) is satisfied.

We next note that when $\mu + \lambda = 0$ (and $\tau > 0$) the general bounded solution to (7.6) is

$$\underline{z}(s) = \underline{c} \exp(-\tau |s|), \quad (7.7)$$

while when $\mu + \lambda \neq 0$ (and $\tau > 0$) the general bounded solution to

¹⁰ Cf., e.g., Gurtin [5, p. 86].

(7.6) is

$$z_{\pm}^{\pm}(s) = \begin{cases} c_{1\pm}^+ z_{\pm}^1(s) + c_{2\pm}^+ z_{\pm}^2(s) + c_{3\pm}^+ z_{\pm}^3(s), & s \geq 0 \\ c_{1\pm}^- z_{\pm}^1(s) + c_{2\pm}^- z_{\pm}^2(s) + c_{3\pm}^- z_{\pm}^3(s), & s \leq 0 \end{cases} \quad (7.8)$$

$$\begin{aligned} z_{\pm}^1(s) &= (\pm i, 1, 0) \exp(\mp \tau s), \quad z_{\pm}^3(s) = (0, 0, 1) \exp(-\tau |s|), \\ z_{\pm}^2(s) &= \mp (\mu^{\pm} + \lambda^{\pm}) \tau s \, z_{\pm}^1(s) + (\mp i \mu^{\pm}, 2\mu^{\pm} + \lambda^{\pm}, 0) \exp(\mp \tau s). \end{aligned} \quad (7.9)$$

Case I. $\mu^+ + \lambda^+ = \mu^- + \lambda^- = 0$. Then (7.5) and (7.7) imply that

$$\begin{aligned} M_{\pm}^{\pm} z_{\pm}(0^{\pm}) &= \mp \mu^{\pm} \tau c_{\pm}, \\ N_{\pm}^{\pm} z_{\pm}(0) &= -\mu^{\pm} [c_{1\pm} t + c_{2\pm} \tau n], \end{aligned}$$

and hence (6.3) reduces to

$$(\mu^+ + \mu^-) |c_{\pm}|^2 \geq i(\mu^+ - \mu^-)(c_1 \bar{c}_2 - c_2 \bar{c}_1).$$

Since μ^+ and μ^- are positive it is easy to show that the above inequality is always satisfied.

Main Case. $\mu^+ + \lambda^+ \neq 0$ and $\mu^- + \lambda^- \neq 0$. Then (7.5) and (7.8)-(7.9) imply that

$$\begin{aligned} z_{\pm}(0^{\pm}) &= D_{\pm}^{\pm} c_{\pm}^{\pm}, \quad M_{\pm}^{\pm} z_{\pm}(0^{\pm}) = -E_{\pm}^{\pm} c_{\pm}^{\pm}, \\ D_{\pm}^{\pm} &:= \begin{bmatrix} \pm i & \mp i \mu^{\pm} & 0 \\ 1 & 2\mu^{\pm} + \lambda^{\pm} & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad E_{\pm}^{\pm} := \tau \begin{bmatrix} i(2\mu^{\pm} + \lambda^{\pm}) & i\lambda^{\pm}(2\mu^{\pm} + \lambda^{\pm}) & 0 \\ \pm \mu^{\pm} & \pm \mu^{\pm}(3\mu^{\pm} + 2\lambda^{\pm}) & 0 \\ 0 & 0 & \pm \mu^{\pm} \end{bmatrix}. \end{aligned}$$

We note that $z_{\pm}$ is continuous at $s = 0$. Therefore if we let

$$c_{\pm}^+ := (D_{\pm}^+)^{-1} z_{\pm}(0), \quad c_{\pm}^- := (D_{\pm}^-)^{-1} z_{\pm}(0)$$

we find that (6.3) is equivalent to

$$\begin{aligned} \bar{z}(0) \cdot [G^+ - G^-] z(0) &\geq 0, \\ G^\pm &:= [E^\pm (D^\pm)^{-1} + iN_t^\pm]. \end{aligned} \quad (7.10)$$

Next, a simple calculation shows that

$$(D^\pm)^{-1} = \frac{1}{i(3\mu^\pm + \lambda^\pm)} \begin{bmatrix} \pm(2\mu^\pm + \lambda^\pm) & i\mu^\pm & 0 \\ \mp 1 & i & 0 \\ 0 & 0 & i(3\mu^\pm + \lambda^\pm) \end{bmatrix}$$

and hence that

$$G^\pm = \frac{\tau\mu^\pm}{3\mu^\pm + \lambda^\pm} \begin{bmatrix} \pm 2(2\mu^\pm + \lambda^\pm) & 2\mu^\pm i & 0 \\ -2\mu^\pm i & \pm 2(2\mu^\pm + \lambda^\pm) & 0 \\ 0 & 0 & \pm(3\mu^\pm + \lambda^\pm) \end{bmatrix}.$$

Since (7.3) implies that $(G^+ - G^-)_{jj} > 0$ ($j = 1, 2, 3$), (7.10) will be satisfied for all initial values $z(0)$ if and only if

$$\det(G^+ - G^-) \geq 0. \quad (7.11)$$

However

$$\det(G^+ - G^-) = 4\tau^3(\mu^+ + \mu^-) \left[\mu^+ + \frac{\mu^-(\mu^- + \lambda^-)}{3\mu^- + \lambda^-} \right] \left[\mu^- + \frac{\mu^+(\mu^+ + \lambda^+)}{3\mu^+ + \lambda^+} \right]$$

and hence we conclude, with the aid of (7.3), that (7.11) is equivalent to (7.4).

Case III. $\mu^+ + \lambda^+ = 0$ and $\mu^- + \lambda^- \neq 0$. We let D^- , E^- , and G^- be given by the previous case. If we use (7.7) and the fact that $\lambda^+ = -\mu^+$ we find that

$$D^+ = I, \quad E^+ = \tau\mu^+ I,$$

and hence that

$$\underset{\sim}{G}^+ = \tau \mu^+ \begin{bmatrix} 1 & i & 0 \\ -i & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Clearly $(\underset{\sim}{G}^+ - \underset{\sim}{G}^-)_{jj} > 0$ ($j = 1, 2, 3$) and

$$\det(\underset{\sim}{G}^+ - \underset{\sim}{G}^-) = 4\tau^3 \mu^- (\mu^+ + \mu^-) \left[\mu^+ + \frac{\mu^- (\mu^- + \lambda^-)}{3\mu^- + \lambda^-} \right].$$

Thus the desired result follows by the reasoning used in the previous case.

Case IV. $\mu^+ + \lambda^+ \neq 0$ and $\mu^- + \lambda^- = 0$. Same as Case III. □

Acknowledgement ~~~~~

We thank Professor R. Fosdick for his useful suggestions and comments. This work was supported in part by the Institute for Mathematics and its Applications, University of Minnesota. It was also supported by the National Science Foundation (DMS 86-00281 and DMS 86-02201) and the Air Force Office of Scientific Research (USAFOSR 86-0184).

References

1. Agmon, S., On the eigenfunctions and on the eigenvalues of general elliptic boundary value problems. Comm. Pure Appl. Math. 15 (1962), 119-147.
2. Ball, J. M., Constitutive inequalities and existence theorems in nonlinear elastostatics. Nonlinear Analysis and Mechanics Vol. I (R. J. Knops, ed.). London: Pitman, 1977.
3. Ball, J. M., & J. E. Marsden, Quasiconvexity at the boundary, positivity of the second variation, and elastic stability. Arch Rational Mech. Anal. 86 (1984), 251-277.
4. De Figueiredo, D. G., The coerciveness problem for forms over vector valued functions. Comm. Pure Appl. Math. 16 (1963), 63-94.
5. Gurtin, M. E., The linear theory of elasticity. Handbuch der Physik VIa/2 (C. Truesdell, ed.), Berlin: Springer-Verlag, 1972.
6. Gurtin, M. E., Two-phase deformations of elastic solids. Arch. Rational Mech. Anal. 84 (1983), 1-29.
7. Hayes, M., & R. S. Rivlin, Propagation of a plane wave in an isotropic elastic material subjected to pure homogeneous deformation. Arch. Rational Mech. Anal. 8 (1961), 15-22.
8. James, R. D., Finite deformation by mechanical twinning. Arch. Rational Mech. Anal. 77 (1981), 143-176.
9. Meyers, N. G., Quasi-convexity and lower semi-continuity of multiple variational integrals of any order. Trans. Amer. Math. Soc. 119 (1965), 125-149.

10. Morrey, C. B., Quasi-convexity and the lower semicontinuity of multiple integrals. Pacific J. Math. 2 (1952), 25-53.
11. Rivlin, R. S., Some reflections on material stability. Mechanics Today, vol. 5, Oxford: Pergamon, 1980.
12. Simpson, H. C., & S. J. Spector, Necessary conditions at the boundary for minimizers in finite elasticity. To appear: Arch. Rational Mech. Anal.
13. Truesdell, C., & W. Noll, The Non-linear Field Theories of Mechanics. Handbuch der Physik III/3 (S. Flügge, ed.). Berlin: Springer-Verlag, 1965.

#	Author/s	Title
441	Laurent Habsieger and Dennis Stanton	More Zeros of Krawtchouk Polynomials
442	K.M. Ramachandran	Nearly Optimal Control of Queues in Heavy Traffic with Heterogeneous Servers
443	Y.M. Zhu and G. Yin	A New Algorithm for Constrained Adaptive Array Processing
444	G. Yin and K.M. Ramachandran	A Differential Delay Equation with Wideband Noise Perturbations
445	Chaitan P. Gupta	Integral Type Asymptotic Conditions for the Solvability of a Periodic Fourth Order Boundary Value Problem
446	Chaitan P. Gupta	A Two-Point Boundary Value Problem of Dirichlet Type with Resonance at Infinitely Many Eigenvalues
447	Jong-Shenq Guo	On the Quenching Behavior of a Semilinear Parabolic Equation
448	Bei Hu	A Quasi-Variational Inequality Arising in Elastohydrodynamics
449	E. Somersalo, G. Beylkin, R. Burridge and M. Cheney	Inverse Scattering Problem for the Schrödinger Equation in Three Dimensions: Connections Between Exact and Approximate Methods
450	J.H. Dinitz and D.R. Stinson	Some New Perfect One-Factorizations from Starters in Finite Fields
451	Albert Fässler and René Jeanneret	Optimum Filter for DC/AC-Converter in Electronics
452	Paul Lemke	On the Question of Obtaining Optimal Partitions of Point Sets in E^d with Hyperplane Cuts
453	Paul Lemke and Michael Werman	On the Complexity of Inverting the Autocorrelation Function of a Finite Integer Sequence, and the Problem of Locating n Points on a Line, Given the $\binom{n}{2}$ Unlabelled Distances Between Them
454	Chris J. Budd, Avner Friedman, Bryce McLeod and Adam A. Wheeler	The Space Charge Problem
455	Jerrold R. Griggs, Daniel J. Kleitman and Aditya Shastri	Spanning Trees With Many Leaves in Cubic Graphs
456	Akos Seress	On λ -designs with $\lambda = 2P$
457	Chjan C. Lim	Quasi-periodic Dynamics of Desingularized Vortex Models
458	Chjan C. Lim	On Singular Hamiltonians: The Existence of Quasi-periodic Solutions and Nonlinear Stability
459	Eugene Fabes, Mitchell Luskin and George R. Sell	Construction of Inertial Manifolds by Elliptic Regularization
460	Matthew Witten	A Quantitative Model for Lifespan Curves
461	Jay A. Wood	Self-Orthogonal Codes and The topology of Spinor Group
462	Avner Friedman and Miguel A. Herrero	A Nonlinear Nonlocal Wave Equation Arising in Combustion Theory
463	Avner Friedman and Victor Isakov	On the Uniqueness in the Inverse Conductivity Problem with One Measurement
464	Yisong Yang	Existence, Regularity, and Asymptotic Behavior of the Solutions to the Ginzburg-Landau Equations on $\mathbb{R}^3$
465	Chjan. C. Lim	On Symplectic Tree Graphs
466	Wilhelm I. Fushchich, Ivan Krivsky and Vladimir Simulik	On Vector and Pseudovector Lagrangians for Electromagnetic Field
467	Wilhelm I. Fushchich	Exact Solutions of Multidimensional Nonlinear Dirac's and Schrödinger's Equations
468	Wilhelm I. Fushchich and Renat Zhdanov	On Some New Exact Solutions of Nonlinear D'Allembert and Hamilton Equations
469	Brian A. Coomes	The Lorenz System Does Not Have a Polynomial Flow
470	J.W. Helton and N.J. Young	Approximation of Hankel Operators: Truncation Error in an H^∞ Design Method
471	Gregory Ammar and Paul Gader	A Variant of the Gohberg-Semencul Formula Involving Circulant Matrices
472	R.L. Fosdick and G.P. MacSithigh	Minimization in Nonlinear Elasticity Theory for Bodies Reinforced with Inextensible Cords
473	Fernando Reitich	Rapidly Stretching Plastic Jets: The Linearized Problem
474	Francisco Bernis and Avner Friedman	Higher Order Nonlinear Degenerate Parabolic Equations
475	Xinfu Chen and Avner Friedman	Maxwell's Equations in a Periodic Structure
476	Avner Friedman and Michael Vogelius	Determining Cracks by Boundary Measurements
477	Yuji Kodama and John Gibbons	A Method for Solving the Dispersionless KP Hierarchy and its Exact Solutions II
478	Yuji Kodama	Exact Solutions of Hydrodynamic Type Equations Having Infinitely Many Conserved Densities
479	Robert Carroll	Some Forced Nonlinear Equations and the Time Evolution of Spectral Data
480	Chjan. C. Lim	Spanning Binary Trees, Symplectic Matrices, and Canonical Transformations for Classical N-body Problems
481	E.F. Assmus, Jr. and J.D. Key	Translation Planes and Derivation Sets
482	Matthew Witten	Mathematical Modeling and Computer Simulation of the Aging-Cancer Interface

#	Author/s	Title
483	Matthew Witten and Caleb E. Finch,	Re-Examining The Gompertzian Model of Aging
484	Bei Hu,	A Free Boundary Problem for a Hamilton-Jacobi Equation Arising in Ions Etching
485	T.C. Hu, Victor Klee and David Larman,	Optimization of Globally Convex Functions
486	Pierre Goossens,	Shellings of Tilings
487	D. David, D. D. Holm, and M.V. Tratnik,	Integrable and Chaotic Polarization Dynamics in Nonlinear Optical Beams
488	D. David, D.D. Holm and M.V. Tratnik,	Horseshoe Chaos in a Periodically Perturbed Polarized Optical Beam
489	Laurent Habsieger,	Linear Recurrent Sequences and Irrationality Measures
490	Laurent Habsieger,	MacDonald Conjectures and The Selberg Integral
491	David Kinderlehrer and Giorgio Vergara-Caffarelli,	The Relaxation of Functionals with Surface Energies
492	Richard James and David Kinderlehrer,	Theory of Diffusionless Phase Transitions
493	David Kinderlehrer,	Recent Developments in Liquid Crystal Theory
494	Niky Kamran and Peter J. Olver,	Equivalence of Higher Order Lagrangians 1. Formulation and Reduction
495	Lucas Hsu, Niky Kamran and Peter J. Olver,	Equivalence of Higher Order Lagrangians II. The Cartan Form for Particle Lagrangians
496	D.J. Kaup and Peter J. Olver,	Quantization of BiHamiltonian Systems
497	Metin Arik, Fahrünisa Neyzi, Yavuz Nutku, Peter J. Olver and John M. Verosky	Multi-Hamiltonian Structure of the Born-Infeld Equation
498	David H. Wagner,	Detonation Waves and Deflagration Waves in the One Dimensional ZND Model for High Mach Number Combustion
499	Jerrold R. Griggs and Daniel J. Kleitman,	Minimum Cutsets for an Element of a Boolean Lattice
500	Dieter Jungnickel,	On Affine Difference Sets
501	Pierre Leroux,	Reduced Matrices and q-log Concavity Properties of q-Stirling Numbers
502	A. Narain and Y. Kizilyalli,	The Flow of Pure Vapor Undergoing Film Condensation Between Parallel Plates
503	Donald A. French,	On the Convergence of Finite Element Approximations of a Relaxed Variational Problem
504	Yisong Yang,	Computation, Dimensionality, and Zero Dissipation Limit of the Ginzburg-Landau Wave Equation
505	Jürgen Sprekels,	One-Dimensional Thermomechanical Phase Transitions with Non-Convex Potentials of Ginzburg-Landau Type
506	Yisong Yang,	A Note On Nonabelian Vortices
507	Yisong Yang,	On the Abelian Higgs Models with Sources
508	Chjan. C. Lim,	Existence of Kam Tori in the Phase Space of Vortex Systems
509	John Weiss,	Bäcklund Transformations and the Painlevé Property
510	Pu Fu-cho and D.H. Sattinger,	The Yang-Baxter Equation for Integrable Systems
511	E. Bruce Pitman and David G. Schaeffer,	Instability and Ill-Posedness in Granular Flow
512	Brian A. Coomes,	Polynomial Flows on C^n
513	Bernardo Cockburn, Suchung Hou and Chi-Wang Shu,	The Runge-Kutta Local Projection Discontinuous Galerkin Finite Element Method for Conservation Laws IV: The Multidimensional Case
514	Peter J. Olver,	Invariant Theory, Equivalence Problems, and the Calculus of Variations
515	Daniel D. Joseph and Thomas S. Lundgren with an appendix by R. Jackson and D.A. Saville,	Ensemble Averaged and Mixture Theory Equations
516	P. Singh, Ph. Caussignac, A. Fortes, D.D. Joseph and T. Lundgren,	Stability of Periodic Arrays of Cylinders Across the Stream by Direct Simulation
517	Daniel D. Joseph,	Generalization of the Foscolo-Gibilario Analysis of Dynamic Waves
518	A. Narain and D.D. Joseph,	Note on the Balance of Energy at a Phase Change Interface
519	Daniel D. Joseph,	Remarks on inertial radii, persistent normal stresses, secondary motions , and non-elastic extensional viscosities
520	D. D. Joseph,	Mathematical Problems Associated with the Elasticity of Liquids
521	Henry C. Simpson and Scott J. Spector,	Some Necessary Conditions at an Internal Boundary for Minimizers in Finite Elasticity
522	Peter Gritzmann and Victor Klee,	On the 0-1 Maximization of Positive Definite Quadratic Forms