



BEDROCK GEOLOGY

By  
John H. Mossler


2012



CORRELATION OF MAP UNITS



MAP SYMBOLS



Digital base modified from the Minnesota Department of Transportation BaseMap data; digital base annotation by the Minnesota Geological Survey. Elevation contours were derived from the U.S. Geological Survey 30-meter Digital Elevation Model (DEM) by the Minnesota Geological Survey. Universal Transverse Mercator Projection, grid zone 15 1983 North American Datum GIS compilation by R.S. Lively Edited by Lori Robinson







Figure 1. Stratigraphic column depicting the lithology, thickness, and vertical succession of rock units shown on the map. Includes the approximate age in millions of years for the rocks in Minnesota. The gamma log is a compilation of the following gamma logs from Anoka and Hennepin Counties: 44789, 20292, 20665, 578957, and 225599. A higher gamma count typically reflects increased shale or clay content.




Figure 2. Acromagnetic map for the metropolitan area showing anomalies related to geologic structures shown on the bedrock map and on Figure 3. Shown in color shaded relief format with false illumination from the northwest at 45° inclination (Chandler, 2007).




Figure 3. Map of Anoka County contoured at the stratigraphic top of the Wenocese Sandstone showing geologic structure, scale: 1:300,000. Contour interval is 50 feet (15 meters). The approximate areas where some of all of the Wenocese Sandstone is missing because of erosion, and where the Wenocese Sandstone, Eau Claire Formation, or Mt. Simon Sandstone are first bedrock is shaded; contours in those areas are inferred from projection using the estimated full thickness of the formations. The inferred traces of faults and folds are shown.

REFERENCES

Berg, R.R., 1954. Franconia Formation of Minnesota and Wisconsin. Geological Society of America Bulletin, v. 66, p. 857-882.  
Chandler, V.W., 2007. Upgrade of aeromagnetic databases and processing systems at the Minnesota Geological Survey. Minnesota Geological Survey Open File Report 07-6.  
Chandler, V.W., Lively, R.S., and Wahl, T.E., 2004. Gravity and aeromagnetic data grids of Minnesota: On file at the Minnesota Geological Survey.  
Kohls, D.W., 1958. The geology of the Prescott quadrangle. Minneapolis, University of Minnesota, M.S. thesis, 138 p.  
Matsch, C.L., 1962. Pleistocene geology of the St. Paul Park and Prescott quadrangles. Minneapolis, University of Minnesota, M.S. thesis, 49 p.  
Meyer, G.N., and Paterson, C.J., 1999. Surficial geology of the Anoka 30 x 60 minute quadrangle, east-central Minnesota. Minnesota Geological Survey Miscellaneous Map M-97, scale 1:100,000.  
Mossler, J.H., 1972. Paleozoic structure and stratigraphy of the Twin City region. In Sims, P.K., and Morey, G.B., eds., Geology of Minnesota: A centennial volume. Minnesota Geological Survey, p. 485-497.  
———1992. Sedimentary rocks of Dresbachian age (Late Cambrian). Hollandale embayment, southeastern Minnesota. Minnesota Geological Survey Report of Investigations 40, 71 p.  
———2005. Bedrock geology of the Hudson quadrangle, Washington County, Minnesota. Minnesota Geological Survey Miscellaneous Map M-154, scale 1:24,000.  
———2006. Bedrock geology of the Prescott quadrangle, Washington and Dakota Counties, Minnesota. Minnesota Geological Survey Miscellaneous Map M-167, scale 1:24,000.  
———2007. Bedrock geology of the Coon Rapids quadrangle, Anoka and Hennepin Counties, Minnesota. Minnesota Geological Survey Miscellaneous Map M-175, scale 1:24,000.  
———2008. Paleozoic stratigraphic nomenclature for Minnesota. Minnesota Geological Survey Report of Investigations 65, 76 p., 1 pl.  
———2009a. Bedrock geology, pl. 2 of Bauer, E.J., project manager. Geologic atlas of Carver County, Minnesota. Minnesota Geological Survey County Atlas C-21, 6 pls., scale 1:100,000.  
———2009b. Bedrock geology of the Anoka quadrangle, Anoka and Hennepin Counties, Minnesota. Minnesota Geological Survey Miscellaneous Map M-184, scale 1:24,000.  
Mossler, J.H., Runkel, A.C., and Bauer, E.J., 2006. Submap of the St. Lawrence Formation in the seven-county Twin Cities metropolitan area. Minnesota Geological Survey Open-File Report 06-5, 1 pl., scale 1:100,000.  
Mossler, J.H., and Tipping, R.G., 2000. Bedrock geology and structure of the seven-county Twin Cities metropolitan area, Minnesota. Minnesota Geological Survey Miscellaneous Map M-104, scale 1:250,000.  
Quaschnick, R.K., 1959. The geology of the Marine quadrangle and the Falls Creek area. Minneapolis, University of Minnesota, M.S. thesis, 89 p.  
Runkel, A.C., 1994. Revised stratigraphic nomenclature for the Upper Cambrian (St. Croix) Jordan Sandstone, southeastern Minnesota. In Southwick, D.L., ed., Short contributions to the geology of Minnesota. Minnesota Geological Survey Report of Investigations 43, p. 60-71.  
Runkel, A.C., McKay, R.M., and Palmer, A.R., 1998. Origin of a classic carbonate sheet sandstone: Stratigraphy across the Sauk II-Sauk III boundary in the upper Mississippi valley. Geological Society of America Bulletin, v. 110, p. 188-210.  
Runkel, A.C., Miller, J.F., McKay, R.M., Shaw, T.H., and Bassett, D.J., 1999. Cambrian-Ordovician boundary strata in the central midcontinent of the St. Lawrence Formation. Acta Universitatis Carolinae—Geologica, v. 43, p. 17-20.  
Runkel, A.C., and Mossler, J.H., 2006. Bedrock geology, pl. 2 of Setherholm, D.R., project manager. Geologic atlas of Scott County, Minnesota. Minnesota Geological Survey County Atlas C-17, 6 pls., scale 1:100,000.  
Runkel, A.C., Mossler, J.H., Tipping, R.G., and Bauer, E.J., 2006. A hydrologic and mapping investigation of the St. Lawrence Formation in the Twin Cities metropolitan area. Minnesota Geological Survey Open-File Report 06-4, 20 p.  
Schwartz, G.M., 1936. The geology of the Minneapolis-St. Paul metropolitan area. Minnesota Geological Survey Bulletin 27, 267 p., 7 pls.  
Sims, P.K., and Zeitz, I., 1967. Aeromagnetic and inferred Precambrian paleogeographic map of eastern Minnesota and part of Wisconsin. U.S. Geological Survey Geophysical Investigations Map GP-563, 6 p., scale 1:250,000.  
Sloan, R.E., and Dunes, Z.F., 1962. A geologic and gravity survey of the Belle Plaine area, Minnesota. Minnesota Academy of Science Proceedings, v. 30, no. 1, p. 49-52.

SOURCES USED TO COMPILE THE GEOLOGIC MAP

Mapping relied almost entirely on subsurface data, including drillers' logs for water wells, exploratory test holes, and monitoring wells. Cutting samples and down-hole geophysical logs were available for some of the boreholes. Some data were obtained from engineering test borings completed for highway and bridge construction by the Minnesota Department of Transportation, and borings completed for an Anoka regional landfill remedial investigation.

INTRODUCTION

This geologic map shows the bedrock formations that lie beneath unconsolidated Quaternary deposits of variable thickness (Fig. 1). The county is almost completely covered by Quaternary glacial till, lake sediments, outwash deposits, swamp deposits (peat), and recent alluvium along the Mississippi and Rum Rivers and minor streams (see Plate 3, *Surficial Geology*; Meyer and Patterson, 1999). The only outcrops are in the "panhandle" of the county in Columbia Heights. The unconsolidated Quaternary and recent deposits generally range from less than 50 (15 meters) to more than 400 feet (122 meters) in thickness (see Plate 5, *Depth to Bedrock*).

Buried stream channels that are incised into several rock formations are a primary influence on the map distribution of bedrock geologic units. The distribution of units is also affected by the Douglas and Pine fault zones (Fig. 2) along the western boundary of the Mesoproterozoic Midcontinent rift (Sims and Zeitz, 1967). Although the faults developed during the Mesoproterozoic era they apparently were reactivated during the early Paleozoic era. Evidence for reactivation includes thickening of finer-grained units, such as the Lone Rock Formation, along down-thrown sides of faults throughout the Twin Cities area (Mossler, 1972).

Because bedrock is concealed beneath Quaternary and recent sediments, distribution of the bedrock units and the pattern of faults and folds are interpreted from subsurface data including water wells and engineering borings, supported by aeromagnetic and gravity data that reflect underlying Mesoproterozoic geology (Sims and Zeitz, 1967; Chandler and others, 2004).

Prior to collection of large amounts of subsurface data, there was little evidence for faults in Paleozoic rocks of the metropolitan area, with the exception of some that are exposed in southern and northeastern Washington County (Schwartz, 1936; Kohls, 1958; Quaschnick, 1959; Matsch, 1962). In addition, faulting had been inferred in the Belle Plaine area of Scott County from gravity data and water well records (Sloan and Dunes, 1962). Recent study utilizing a more extensive subsurface data base (primarily gamma logs, well cuttings, and driller's logs) has indicated that there are faults and folds that displace Paleozoic rocks along the western side of the Twin Cities basin (Mossler and others, 2006; Runkel and Mossler, 2006; Mossler, 2009a), where none were inferred before, just as there are along the southeastern side of the basin where earlier mapping (Kohls, 1958; Quaschnick, 1959; Matsch, 1962; Mossler, 2005, 2006) revealed extensive faulting. In some instances, mostly in rural areas, where data are insufficient to support the presence of faults along the western side of the Twin Cities basin, there are enough data to indicate localized, open folding of Paleozoic rocks along zones inferred from aeromagnetic and gravity data to be underlain by Mesoproterozoic faults. There may be faults along these zones but data are insufficient to map them. In rural areas most wells are shallow domestic wells that end in Quaternary drift or penetrate only a few feet into bedrock. There are numerous folds and faults along the trace of the Mesoproterozoic Douglas fault that Sims and Zeitz (1967) mapped using aeromagnetic data (Fig. 2). Subsurface data southeast of the Douglas fault zone indicate that there is a monoclinical fold and faulting along the trace of the Mesoproterozoic Pine fault zone that Sims and Zeitz (1967) mapped. There are also northwest-trending folds and faults to the northwest of the Douglas fault identified by Mossler (2007, 2009b, this report). The faulting and folding in Paleozoic rocks caused by the reactivation of Mesoproterozoic structures, combined with deep pre-glacial erosion of the bedrock surface, created a subpattern in the bedrock units that was not displayed in earlier maps such as Mossler and Tipping (2000). Specifically, older formations are commonly exposed along upthrown sides of faults and younger units are preserved along downthrown sides. Deep incision by pre-glacial streams combined with the regional dip of bedrock formations toward the southeast, shown in Figure 3, has caused progressively older formations to subcrop toward the north and west across the county.

The rise in altitude of stratigraphic tops of formations across the Anoka (Mossler, 2009b) and Coon Rapids (Mossler, 2007) quadrangles (see Index to 7.5-minute quadrangles) is not as marked as it is on the eastern side of the Twin Cities basin. It is only about half the rise observed, for a comparable distance, across quadrangles mapped on the eastern side of the Twin Cities basin in southern Washington County across the Hudson-Afton horst (see Index to 7.5-minute Quadrangles for location of hors).

The cross sections show bedrock at depth and illustrate stratigraphic relationships. The geologic formations are thin in relation to their areal extent and would only be as thick as shown in the cross sections if no vertical exaggeration was used. The exaggeration needed to show the thin rock formations also gives the appearance of steeper slopes on the land surface topography, buried bedrock surface, and geologic contacts between rock formations. Deeper formations such as the lower part of the Mt. Simon Sandstone (unit Cm) and underlying Mesoproterozoic formations are not shown on the cross sections because of scarcity of data. In addition, configuration of the bedrock contacts for shallower geologic units and their thicknesses are poorly constrained along parts of the cross sections because of paucity of data. For example, few wells penetrate the entire Tunnel City Group (unit Cc), and many of those that do penetrate it do not contain the upper part and the contact with the overlying St. Lawrence Formation (Op) because these intervals have eroded.

DESCRIPTION OF MAP UNITS

Nomenclature has been revised for some Paleozoic formations in Minnesota (Mossler, 2008) and some formation names formerly in use at the Minnesota Geological Survey have been replaced by

names widely accepted elsewhere in the region. Rocks formerly included in the Franconia Formation are now assigned to the Tunnel City Group, which is subdivided into the Mazomanie Formation and the Lone Rock Formation. The Lone Rock Formation includes the three members formerly included in the Franconia Formation: the Reno, Tomah, and Birkmore Members, which were described and named by Berg (1954). The interval formerly referred to as the Ironton-Galeville Sandstone is now named the Wenocese Sandstone. More detailed discussion of these revisions is given in Mossler (2008).

**Op Platteville and Glenwood Formations (Upper Ordovician)**—The Platteville Formation is dominantly limestone and dolomite. The Glenwood Formation is dominantly shale. Together they are about 25 to 34 feet (8 to 10 meters), but they generally are thinner because the upper part of the Platteville Formation has been removed by erosion. The formations are exposed in Columbia Heights and are present in the subsurface nearby. The Platteville Formation is 25 to 30 feet (8 to 9 meters) thick. It is composed of yellowish-gray to light brown-gray, thick, to medium-bedded dolomite overlying yellowish-gray to light gray, thin, to medium-bedded limestone. There is a thin bed of sandy, phosphatic dolomite at the base of the formation. The Glenwood Formation is grayish-green to brownish-gray, calcareous, sandy, phosphatic shale. It is typically from 3 to 5 feet (1 to 1.5 meters) thick.

**Os St. Peter Sandstone (Middle to Upper Ordovician)**—The St. Peter Sandstone generally is 145 to 155 feet (44 to 47 meters) thick in the Twin Cities area where it overlies by the Glenwood and Platteville Formations. It is present in the subsurface mainly in the panhandle of the county in Fridley, Hilltop, and Columbia Heights, and in southeastern Anoka County southeast of Lino Lakes. The upper part is white to light gray, medium- to fine-grained, quartzose sandstone. Thick beds characterize this part; there is some cross stratification in outcrops near the top of the unit. The upper part of the formation is very friable. The basal part is light to medium gray, fine- to coarse-grained, and poorly sorted quartzose sandstone that is more indurated. It is interbedded with shale and feldspathic siltstone of varied colors. The upper part of the formation is exposed at the Northtown railroad yard. The basal contact of the formation with the underlying Prairie du Chien Group dolomite (unit Op) is a major unconformity (Smith and others, 1993).

**Op Prairie du Chien Group (Lower Ordovician)**—The Prairie du Chien Group is generally from 125 to 140 feet (38 to 43 meters) thick in water wells in Anoka County and adjoining areas where covered by St. Peter Sandstone. It can be much thinner where it is first bedrock and has been subjected to erosion. It is present in the subsurface mainly in the panhandle of the county and in the southeastern part of the county near Centerville and Lino Lakes. The Prairie du Chien Group is not exposed in Anoka County, but is extensively exposed elsewhere in the Twin Cities metropolitan area along bluffs on the Mississippi, Minnesota, and St. Croix Rivers as well as along smaller streams. The Prairie du Chien Group is typically divided into two formations: the upper Shakopee Formation and lower Onota Dolomite; however, this subdivision is not always practical in the subsurface because the lack of data locally precludes subdivision. The Shakopee Formation is a heterolithic unit that contains dolomite, sandy dolomite, and sandstone. It is grayish-orange to yellowish-gray, thinly bedded, and oolitic and sandy in its lower part. The Onota Dolomite is yellowish-gray to pale brown dolomite, typically in medium to thick beds. It is less sandy than the Shakopee Formation except near the base. Contacts between the Shakopee Formation and the Onota Dolomite and between the Onota Dolomite and the underlying Jordan Sandstone (unit Cj) are unconformable (Runkel and others, 1999).

**Cj Jordan Sandstone (Upper Cambrian)**—Dominantly light gray sandstone characterized by coarsening-upward sequences consisting of two intertuffaceous facies (Runkel, 1994). Although not shown separately on the geologic map, they are apparent on Figure 1. They are medium- to coarse-grained, cross-stratified, generally friable, quartz sandstone; and very fine-grained, commonly bioturbated, feldspathic sandstone and lenses of siltstone and shale. The major part of the very fine-grained facies forms a regionally continuous interval that gradually overlies the St. Lawrence Formation (unit Cc), although there are lithologically similar intervals intercalated with the medium- to coarse-grained facies at higher stratigraphic intervals. The Jordan Sandstone is generally 85 to 100 feet (26 to 30 meters) thick where uneroded. However, throughout much of Anoka County in the areas where the overlying Prairie du Chien Group is missing because of erosion, only the basal part to as much as three quarters (40 to 80 feet [12 to 24 meters]) of the formation is present.

**Cc St. Lawrence Formation (Upper Cambrian)**—The St. Lawrence Formation is principally light gray to yellowish-gray and pale yellowish-green, dolomitic, feldspathic siltstone with interbedded very fine-grained dolomite and shale. Lenses and layers of light gray, finely crystalline, sandy dolomite occur locally (Runkel and others, 2006). The

formation is 38 to 50 feet (12 to 15 meters) thick. The upper contact with the Jordan Sandstone is conformable and gradational. The gradational nature of the contact in well-cuttings and on natural gamma logs can make grading a precise contact between these formations from site to site problematic.


**Cc Tunnel City Group (Upper Cambrian)**—The Tunnel City Group, formerly named the Franconia Formation (Berg, 1954), varies from less than 135 to approximately 180 feet (41 to 55 meters) in thickness in the map area. It is divided into two formations: the upper Mazomanie Formation and the lower Lone Rock Formation (Mossler, 2008). Both the Mazomanie Formation and the members of the Lone Rock Formation were originally named and described by Berg (1954), who considered them to be members of the Franconia Formation. The uppermost non-glaucouitic formation, the Mazomanie Formation, is present everywhere in Anoka County beneath the St. Lawrence Formation. It is dominantly white to yellowish-gray, fine- to medium-grained, cross-stratified, generally friable, quartz sandstone. Glaucouitic grains typically are absent and never exceed 5 percent (Berg, 1954). Some beds contain brown, intergranular dolomite as cement. The Mazomanie Formation is about 60 to 80 feet (18 to 24 meters) thick in Anoka County. The Mazomanie Formation and the Jordan Sandstone may be mistaken for one another in Anoka County and neighboring areas because of their lithic similarities and similar thickness. The Lone Rock Formation underlies and intertongues with the Mazomanie Formation. The uppermost member of the Lone Rock Formation is the Reno Member, which is pale yellowish-green, very fine- to fine-grained, glauconitic, feldspathic sandstone with thin, greenish-gray shale partings. The sandstone is well sorted and contains thin zones with dolomitic intraclasts. The Reno Member is underlain by the Tomah Member, a thin unit composed of interbedded, grayish-yellow-green, feldspathic siltstone, very fine-grained sandstone, and pale green shale. The Tomah Member is sparsely glauconitic. The basal Birkmore Member is grayish-yellow-green, fine-grained sandstone that is cemented by dolomite. It is highly glauconitic and contains abundant dolomitic intraclasts. Most of the Lone Rock Formation consists of the Reno Member, which is 65 to 70 feet (20 to 21 meters) thick. The medial Tomah Member is less than 10 to 15 feet (3 to 5 meters) thick. The basal Birkmore Member is about 20 feet (6 meters) thick. The upper contact of the Tunnel City Group with the St. Lawrence Formation is conformable. In Anoka County the contact is fairly sharp and the contrast between the siltstone and shale of the St. Lawrence Formation and the underlying fine- to medium-grained, quartzose sandstone in the Mazomanie Formation of the Tunnel City Group is distinct.

**Cw Wenocese Sandstone (Upper Cambrian)**—This unit, formerly classified as two formations, the Ironton and Galeville Sandstones, is composed mostly of fine- to coarse-grained, moderately sorted to well sorted, light gray, quartz sandstone. The upper part is the coarsest-grained; the lower part is finer-grained, better sorted, and progressively finer-grained toward its base. The very fine-grained sandstone in the lower part is feldspathic. The sandstone contains abundant brachiopod valves locally along bedding planes. The thickness of the formation is 50 to 60 feet (15 to 18 meters). The Wenocese Sandstone is conformable with overlying and underlying formations; however, there is a subtle internal unconformity marked by a pebbly sandstone layer (Runkel and others, 1998).

**Ce Eau Claire Formation (Middle to Upper Cambrian)**—The formation is composed of yellowish-gray to pale olive-gray, very fine-grained, feldspathic sandstone, siltstone, and shale. The upper part is predominantly shale and siltstone, the lower part predominantly fine-grained sandstone and siltstone (Mossler, 1992). The formation ranges from 75 to 80 feet (23 to 24 meters) in thickness. The contact with the Mt. Simon Sandstone is conformable.

**Cm Mt. Simon Sandstone (Middle Cambrian)**—The Mt. Simon Sandstone is pale yellowish-brown to grayish-orange-pink to light gray, medium- to coarse-grained, quartz sandstone that contains interbedded siltstone and very fine-grained, feldspathic sandstone, particularly in its upper half to one third (Mossler, 1992). There is a thin conglomeratic zone at the base. It generally is from 125 to 200 feet (38 to 61 meters) thick in the map area, although its full thickness is not shown on cross sections due to lack of data. It unconformably overlies Mesoproterozoic rocks.

INDEX TO 7.5-MINUTE QUADRANGLES OF BEDROCK GEOLOGY IN THE TWIN CITIES METROPOLITAN AREA IN THE MINNESOTA GEOLOGICAL SURVEY MISCELLANEOUS MAP SERIES



The approximate trace of Hudson-Afton horst in the southeastern part of the Twin cities metropolitan area is shown (gray trace).

Every reasonable effort has been made to ensure the accuracy of the factual data on which this map interpretation is based; however, the Minnesota Geological Survey does not warrant or guarantee that there are no errors. Users may wish to verify critical information; sources include both the references listed here and information on file at the office of the Minnesota Geological Survey in St. Paul. In addition, effort has been made to ensure that the interpretation conforms to sound geologic and cartographic principles. No claim is made that the interpretation shown is rigorously correct; however, and it should not be used to guide engineering-scale decisions without site-specific verification.

The views and conclusions contained in this document are those of the author and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government. This map is submitted for publication with the understanding that the U.S. Government is authorized to reproduce and distribute reprints for governmental use. Supported by the U.S. Geological Survey, National Cooperative Geologic Mapping Program, under assistance Award No. G16AC00388.