

## **Sponsors**

---

### ***We thank the following sponsors:***

#### **Platinum**

Bayer Animal Health  
National Pork Board  
Pfizer Animal Health

#### **Silver**

Boehringer Ingelheim Vetmedica, Inc.

#### **Bronze**

Cargill  
Merck Animal Health  
Novartis Animal Health

#### **Copper**

AgStar Financial Services  
Elanco Animal Health  
IDEXX  
Newport Laboratories  
PIC USA  
PRRS CAP

#### **University of Minnesota Institutional Partners**

College of Veterinary Medicine  
University of Minnesota Extension  
College of Food, Agriculture and Natural Resources Sciences

#### **Formatting**

Tina Smith Graphics  
[www.tinasmithgraphics.com](http://www.tinasmithgraphics.com)

#### **CD-ROM**

David Brown  
[www.davidhbrown.us](http://www.davidhbrown.us)

#### **Logo Design**

Ruth Cronje, and Jan Swanson;  
based on the original design by Dr. Robert Dunlop

The University of Minnesota is committed to the policy that all persons shall have equal access to its programs, facilities, and employment without regard to race, color, creed, religion, national origin, sex, age, marital status, disability, public assistance status, or sexual orientation.

# Control and elimination of a porcine reproductive and respiratory syndrome virus field isolate from a continuous flow, single-site nursery-finisher unit using Ingelvac® PRRS MLV vaccination

Noel Garbes<sup>1</sup>, Brian Payne<sup>2</sup>, Laura Batista<sup>2</sup>

<sup>1</sup>Bethany Swine Health Services, Sycamore, IL; <sup>2</sup>Boehringer Ingelheim Vetmedica, St. Joseph, MO

## Introduction and Objectives

Porcine reproductive and respiratory syndrome (PRRS) Area Regional Control (ARC) projects have developed throughout the United States in the past year. The establishment of individual herd plans is a major component of these ARC projects. Herd plans for continuous flow growing pig sites are one of the more challenging to develop and implement.

A plan to control and subsequently eliminate an endemic PRRS virus isolate; (1-18-4 RFLP cut pattern), that infected a multi-age group growing pig herd was developed by the practicing veterinarian and then implemented by the farm employees. The objective of this study was first, to control the circulation and shedding of the endemic PRRS virus isolate and to subsequently eliminate it from the farm.

## Materials and Methods

The site houses three groups of 2800 pigs per group with nine-to-ten weeks of age difference between each group. The site is comprised of two nursery buildings with four and six nursery rooms in each building. There are two sets of finisher buildings with four buildings each dedicated to the grower and finisher age groups. Each set of finisher buildings has a dedicated hallway; however, exhaust air is shared between the buildings on the site. During the study period, pigs from the oldest group were aggressively marketed two weeks (day -14 to 0) prior to the first vaccination (day 0). On day 0, all pigs from the three groups housed in the farm (3-4, 12-13 and 21-22 weeks of age) were vaccinated intramuscularly with a full dose (2ml) of Ingelvac® PRRS MLV. The pigs to be marketed within the next 21 days (~200 head) were not vaccinated. Twenty-one days later (day 21), the same pigs were vaccinated with a second full dose of Ingelvac® PRRS MLV excluding pigs marketed (~200 head) or pigs to be marketed within 21 days (~1,000 head). Between day 42 and 52 a new group of PRRS virus negative weaned pigs from a PRRS

positive stable farm undergoing an elimination process; (Standard Herd Classification II-B)\*, was entered into the study farm. Each of the pigs in this group was vaccinated with a single full dose of Ingelvac® PRRS MLV. Between day 125 and 135 another PRRS virus negative wean group was entered into the nursery but was not vaccinated. Prior to and throughout the control and elimination process, internal biosecurity was increased. The protocol indicated the use of dedicated boots and clothes for each age group, no supply sharing between groups was allowed without previous disinfection and Synergize was used to disinfect the facility, equipment and supplies.

## Results

A random subset of pigs (n=10) from the group placed between day 125 to 135 (PRRS virus negative and not vaccinated) were bled on day 184 (9 weeks post placement) and sera was tested for the presence of PRRS virus antibodies by ELISA Herd Check PRRS 2XR (IDEXX Laboratories Westbrook, ME). All serum samples were negative.

## Discussion and Conclusions

The goal to control and eliminate the endemic PRRS virus isolate from the farm was achieved. Double-mass vaccination with Ingelvac® PRRS MLV of pigs already on the site, vaccination of the first incoming group after mass vaccination, and a short herd closure period between groups decreased shedding, and limited horizontal transmission. The negative PRRS virus pigs entered between day 125 to 135 remained negative to PRRSv antibodies, indicating no exposure to either field or modified live vaccine virus and therefore successful implementation of the protocol. The farm will continue monitoring PRRSv status and respecting internal and external biosecurity. Custom herd plans, such as this, are necessary in area regional control projects.

\*Holtkamp, D., Polson, D., et.al., *JSHAP*, 2011.