
Sponsors

University of Minnesota

College of Veterinary Medicine

College of Agricultural, Food and Environmental Sciences

Extension Service

Swine Center

Editors

W. Christopher Scruton

Stephen Claas

Layout

David Brown

Logo Design

Ruth Cronje, and Jan Swanson;

based on the original design by Dr. Robert Dunlop

Cover Design

Shawn Welch

The University of Minnesota is committed to the policy that all persons shall have equal access to its programs, facilities, and employment without regard to race, color, creed, religion, national origin, sex, age, marital status, disability, public assistance status, or sexual orientation.

An evaluation of the Thermo-Assisted Drying and Decontamination (TADD) system for the elimination of porcine reproductive and respiratory syndrome virus from contaminated livestock transport vehicles.

Scott Dee, Montserrat Torremorell, John Deen, Carlos Pijoan

Introduction

Livestock transport vehicles have long been considered a high-risk route for the introduction of swine pathogens to susceptible populations. Recent work using a scale model of a weaned pig transport vehicle indicated that exposing naïve pigs to porcine reproductive and respiratory syndrome (PRRSV)-contaminated interiors resulted in consistent reproduction of transmission and infection of PRRSV. This work further demonstrated the an overnight drying period after washing consistently eliminated PRRSV from the interior of the model and prevented infection of sentinels. Unfortunately, these prolonged periods of downtime for transport in commercial systems are cost-prohibitive. Therefore, means to reduce the time necessary for complete drying yet still insure virus elimination are needed.

Objective: The purpose of this experiment was to validate a new method for achieving a dry, PRRSV-free trailer in a reduced period of time entitled Thermo-Assisted Drying and Decontamination (TADD). The principle of TADD is to raise the interior of trailers to 71⁰C for 30 minutes, in order to promote drying and degradation of virus.

Hypothesis: Methods that enhance drying of trailers also promote elimination of PRRSV.

Materials and Methods

The study was conducted using the scale model trailers ¹. This model is constructed at a 1:150 scale of an actual weaned pig

transport vehicle. Animal density, materials and design use in the model were consistent with those found in actual trailers. To initiate the study, trailer interiors were artificially contaminated with a concentration of 5×10^5 TCID₅₀ of PRRSV strain MN 30-100 using a hand-held aerosol sprayer. After contamination, trailers were washed using 20⁰ C water, at 3000 psi for 72 seconds. After washing, trailers were treated with 1 of 4 treatments: a) TADD, b) forced air (no heat), c) overnight (8 hr) drying and d) no treatment post-wash. During the application of the TADD system, forced air and wash only treatments, swabs were collected from the trailer interiors at 0, 10, 20 and 30 minutes post-treatment. Swabs were collected for the overnight group after a drying period of 8 hours. Swabs were tested for the presence of PRRSV RNA by TaqMan PCR. A total of 10 replications were conducted per treatment. As a measure of the presence of infectious PRRSV post-treatment, sentinel pigs were housed in treated trailers for 2 hours post-treatment, removed and tested for evidence of PRRSV infection by PCR and ELISA at 7 and 14 days post-exposure. Finally, supernatants from swabs were pooled according to treatment and injected IM into naïve pigs (bioassay) and recipient pigs were tested as described for evidence of PRRSV infection.

Results

All trailers were PCR positive immediately after washing, prior to treatment (pt). At 10 minutes pt, 7/10 swabs were positive from the TADD trailers; however, all swabs collected at 20 and 30 minutes post

treatment were PCR-negative. In contrast, 9/19, 6/10 and 6/10 swabs collected at 10, 20, and 30 minutes from trailers treated with forced air (no heat) were positive. All swabs (10/10) collected from trailers treated with washing only were PCR positive and all swabs collected at 8 hours of drying were PCR negative. All tests for the presence of infectious PRRSV were negative for trailers treated with TADD and overnight drying. In contrast, infectious PRRSV was detected in sentinel pigs and bioassay pigs in the forced air treatment group and the wash only treatment group. Results are summarized in table 1. Air speeds recorded during the TADD treatment averaged 7.5-m/sec (15 mph) while velocities recorded from the forced air (no heat) treatment averaged 17.5-m/sec (36 mph). Finally, environmental temperature and relative humidity averaged 8°C and 79% throughout the study period.

Conclusions

Under the conditions of this study, the efficacy of the TADD system was equal to that of the overnight drying treatment, and it required a shorter period of time to complete its objective. Studies to evaluate the efficacy of TADD in full size trailers are underway and will be presented at the Leman conference. Further studies will also be conducted to evaluate the efficacy of the TADD system to decontaminate trailers from other swine pathogens and microorganisms of food safety concerns.

References

1. Dee SA, Deen J, Otake S, and Pijoan C. An assessment of transport vehicles as a source of porcine reproductive and respiratory syndrome virus transmission to susceptible pigs. *Can J Vet Res* 2004. 68:124-133.

Table 1. Summary of the PCR results following application of the different treatments and the assessment of viable PRRSV.

<u>Treatments</u>	<u>0 min</u>	<u>10 min</u>	<u>20 min</u>	<u>30 min</u>	<u>8 hrs</u>	<u>Bioassay</u>	<u>Sentinels</u>
TADD	10/10 pos	7/10 pos	0/10 pos	0/10 pos	NA	neg	0/3 pos
Forced air – no heat	10/10 pos	9/10 pos	6/10 pos	6/10 pos	NA	pos	2/3 pos
Wash only	10/10 pos	10/10 pos	10/10 pos	10/10 pos	NA	pos	3/3 pos
Overnight drying	10/10 pos	NA	NA	NA	0/10 pos	neg	0/3 pos