

Moving Coframes.

I. A Practical Algorithm

Mark Fels[†]
School of Mathematics
University of Minnesota
Minneapolis, MN 55455
fels@math.umn.edu

Peter J. Olver[‡]
School of Mathematics
University of Minnesota
Minneapolis, MN 55455
olver@ima.umn.edu
<http://www.math.umn.edu/~olver>

Abstract. This is the first in a series of papers devoted to the development and applications of a new general theory of moving frames. In this paper, we formulate a practical and easy to implement explicit method to compute moving frames, invariant differential forms, differential invariants and invariant differential operators, and solve general equivalence problems for both finite-dimensional Lie group actions and infinite Lie pseudo-groups. A wide variety of applications, ranging from differential equations to differential geometry to computer vision are presented. The theoretical justifications for the moving coframe algorithm will appear in the next paper in this series.

[†] *Supported in part by an NSERC Postdoctoral Fellowship.*

[‡] *Supported in part by NSF Grant DMS 95-00931.*

November 1, 1996

1. Introduction.

First introduced by Gaston Darboux, and then brought to maturity by Élie Cartan, [6], [8], the theory of moving frames (“repères mobiles”) is acknowledged to be a powerful tool for studying the geometrical properties of submanifolds under the action of a transformation group. While the basic ideas of moving frames for classical group actions are now ubiquitous in differential geometry, the theory and practice of the moving frame method for more general transformation group actions has remained relatively undeveloped and is as yet not well understood. The famous critical assessment by Weyl in his review, [45], of Cartan’s seminal book, [8], retains its perspicuity to this day:

“I did not quite understand how he [Cartan] does this in general, though in the examples he gives the procedure is clear.” “Nevertheless, I must admit I found the book, like most of Cartan’s papers, hard reading.”

Implementations of the method of moving frames for certain groups having direct geometrical significance — including the Euclidean, affine, and projective groups — can be found in both Cartan’s original treatise, [8], as well as many standard differential geometric texts; see, for example, the books of Guggenheimer, [17], which gives the method center stage, Sternberg, [42], and Willmore, [47]. The method continues to attract the attention of modern day researchers and has been successfully extended to some additional examples, including, for instance, the geometry of holomorphic curves in projective space and Grassmannians. The papers of M. Green, [15], Griffiths, [16], and the lecture notes of Jensen, [21], are particularly noteworthy, attempting to place Cartan’s intuitive constructions on a firm theoretical and differential geometric foundation. However, none of the proposed modern geometrical formulations of the theory incorporates the full scope or range of applicability of the method as originally envisioned by Cartan. To this day, both the formulation and construction of moving frames for general Lie group actions has remained obscure, particularly for anyone interested in new applications. Although they strive for generality, the range of examples treated remains rather limited, and Weyl’s pointed critique of Cartan’s original version still, in our opinion, applies to all of these later efforts.

There are two main goals of this series of papers devoted to a study of Cartan’s method of moving frames. The first, of utmost importance for applications and the subject of the present work, is to develop a practical and easily implementable algorithm for constructing moving frames, that can be systematically applied to concrete problems arising in a wide variety of applications. Our new algorithm, which we name the method of “moving coframes”, not only reproduces all of the classical moving frame constructions, often in a simpler and more direct fashion, but can be readily applied to a wide variety of new situations, including infinite-dimensional pseudo-groups, intransitive group actions, restricted reparametrization problems, joint group actions, and many more. Although one can see the germs of our ideas in the above mentioned references, our approach is different, and, we believe, significantly easier to implement in practical examples. Standard presentations of the method rely on an unusual hybrid of vector fields and differential forms. Our approach is inspired by the powerful Cartan equivalence method, [11], [14], [36], which has much of the flavor of moving frame-type computations, but relies solely on the use of differential forms, and the operation of exterior differentiation. The moving coframe method we de-

velop does have a complete analogy with the Cartan equivalence method; indeed, we shall see that the method includes not only all moving frame type equivalence problems, under both finite-dimensional Lie transformation groups and infinite Lie pseudo-groups, but also includes the standard Cartan equivalence problems in a very general framework.

Our second goal is to justify the moving coframe method by proposing a new theoretical foundation for the method of moving frames. The rigorous justification for our moving coframe algorithm will form the subject of the second paper in the series. The key new idea is to initially avoid the technically complicated normalization procedure during the implementation of the method, leading to a fully regularized version of the method that avoid branching and singularity complications inherent in the standard approach, and enabling one to treat both generic and singular submanifolds on the same general footing. Once the regularized solution to the problem has been properly implemented, the *a posteriori* justification for the usual normalization and reduction procedure can be readily provided. Details and further examples will appear in part II.

Beyond the traditional application to the differential geometry of curves and surfaces in certain homogeneous spaces, there are a host of applications of the method that lend great importance to its proper implementation. Foremost are the equivalence and symmetry theorems of Cartan, that characterize submanifolds up to a group transformation by the functional relationships among their fundamental differential invariants. The method provides an effective means of computing complete systems of differential invariants and associated invariant differential operators, which are used to generate all the higher order invariants. The fundamental differential invariants and their first order derived invariants serve to parametrize the “classifying manifold” associated with a given submanifold; the Cartan solution to the equivalence problem states that two submanifolds are (locally) congruent under a group transformation if and only if their classifying manifolds are identical. Moreover, the dimension of the classifying manifold completely determines the dimension of the symmetry subgroup of the submanifold in question. We note that the differential invariants also form the fundamental building blocks of basic physical theories, enabling to construct suitably invariant differential equations and variational principles, cf. [36]. Closely related to the determination of differential invariants are the rigidity theorems, [16], [21], that state that two submanifolds which are congruent by a group element are, in fact, identical, if and only if they have n^{th} order contact at a point for some well determined n . Typically the rigidity order n is one more than the order of the fundamental differential invariants, although there may exist exceptional submanifolds having a higher order of rigidity.

Additional motivation for pursuing this program comes from new applications of moving frames to computer vision promoted by Faugeras, [12], with applications to invariant curve and surface evolutions, and the use of the classifying (or “signature”) manifolds in the invariant characterization of object boundaries that forms the basis of a fully group-invariant object recognition visual processing system, [5]. Although differential invariants have evident direct applications to object recognition in images, the often high order of differentiation makes them difficult to compute in an accurate and stable manner. One alternative approach, [33], is to use joint differential invariants, or, as they are known in the computer vision literature, “semi-differential invariants”, which are based on several

points on the submanifold of interest. Although a few explicit examples of joint differential invariants are known, there is, as far as we know, no systematic classification of them in the literature. We show how the method of moving coframes can be readily used to compute complete systems of joint differential invariants, and illustrate with some examples of direct interest in image processing. The approximation of higher order differential invariants by joint differential invariants and, generally, ordinary joint invariants leads to fully invariant finite difference numerical schemes for their computation, first proposed in [5]. The moving coframe method should aid in the understanding and extension of such schemes to more complicated situations.

In this paper, we begin with a brief survey of the basic equivalence problems for submanifolds under transformation groups that form a primary motivation for the method of moving frames. Section 3 provides a brief introduction to one of the basic tools that the moving coframe method uses — the left-invariant Maurer–Cartan forms on a Lie group. Two practical means of computing the Maurer–Cartan forms, including a novel method based directly on the group transformation rules, are discussed. Section 4 begins the presentation of the moving coframe method for the simplest category of examples — finite-dimensional transitive group actions — and illustrates it with a non-conventional equivalence problem arising in the calculus of variations and in classical invariant theory. Section 5 extends the basic method to intransitive Lie group actions. The simplest example of an infinite-dimensional pseudo-group, namely the reparametrization pseudo-group for parametrized submanifolds, is discussed in Section 6 and illustrated with a simple geometrical example — the case of curves in the Euclidean plane. This is followed by a discussion of curves in affine and projective geometry, reproducing classical moving frame computations in a simple direct manner based on the moving coframe approach; in Section 7, the connections between the classical and moving coframe methods are explained in further detail. Section 8 employs the moving coframe method to completely analyze the joint differential invariants in two particular geometrical examples — two-point differential invariants for curves in the Euclidean and affine plane. Section 9 discusses how to analyze more general pseudo-group actions, illustrating the method with two examples arising in classical work of Lie, Vessiot and Medolaghi. In addition, we show how to solve the equivalence problem for second order ordinary differential equations under the pseudo-group of fiber-preserving transformations using the moving coframe method, thereby indicating how all Cartan equivalence problems can be treated by this method. Finally, we discuss some open problems that are under current investigation. In all cases, the paper is designed for a practically oriented reader, in that only the basic algorithmic steps are discussed in detail. In order not to cloud to exposition, all of the theoretical justifications for the algorithms proposed here will appear in the second paper in this series.

2. The Basic Equivalence Problems.

We begin our exposition with a discussion of the basic equivalence problems which can be handled by the method of moving frames; see also Jensen, [21; p. VI], for additional details. Suppose G is a transformation group acting smoothly on an m -dimensional manifold M . In classical applications, G is a finite-dimensional Lie group, but, as we shall see, the method can be extended to infinite-dimensional Lie pseudo-group actions, e.g.,

the group of conformal transformations on a Riemannian surface, the group of canonical transformations on a symplectic space, or the group of contact transformations on a jet space. In either situation, a basic equivalence problem is to determine whether two given submanifolds are congruent modulo a group transformation. We shall divide the basic problem into two different versions, depending on whether one allows reparametrizations of the submanifolds in question. Formally, these can be stated as follows.

The Fixed Parameter Equivalence Problem: Given two embeddings $\iota: X \rightarrow M$ and $\bar{\iota}: X \rightarrow M$ of an n -dimensional manifold X into M does there exist a group transformation $g \in G$ such that

$$\bar{\iota}(x) = g \cdot \iota(x) \quad \text{for all} \quad x \in X. \quad (2.1)$$

The Unparametrized Equivalence Problem: Given two submanifolds $N, \bar{N} \subset M$ of the same dimension n , determine whether there is exists a group transformation $g \in G$ such that

$$g \cdot N = \bar{N}. \quad (2.2)$$

In both problems we shall only consider the question in the small, meaning that (2.1) only needs to hold on an open subset of X , or that (2.2) holds in a suitable neighborhood of given points $z_0 \in N, \bar{z}_0 \in \bar{N}$. Global issues require global constructions that lie outside the scope of the Cartan approach to equivalence problems.

Note that the problem of determining the symmetries of a submanifold, meaning the set of all group elements that preserve the submanifold, forms a particular case of the equivalence problem. Indeed, a *symmetry* of a submanifold is merely a self-equivalence. For instance, the unparametrized symmetries of a given submanifold $N \subset M$ are those group elements that (locally) satisfy $g \cdot N = N$. Note that the symmetry group of a given submanifold forms a subgroup $H \subset G$ of the full transformation group.

Example 2.1. A classical example is inspired by the geometry of curves in the Euclidean plane. A curve $C \subset \mathbb{R}^2$ is parametrized by a smooth map $\mathbf{x}(t) = (x(t), y(t))$ defined on (a subinterval of) \mathbb{R} . The underlying group for Euclidean planar geometry is the Euclidean group $E(2) = O(2) \times \mathbb{R}^2$ consisting of translations, rotations and (in the non-oriented case) reflections.

In the fixed parametrization problem, we are given two parametrized curves $\mathbf{x}(t)$ and $\bar{\mathbf{x}}(t)$, and want to know when there exists a Euclidean motion such that $\bar{\mathbf{x}}(t) = R \cdot \mathbf{x}(t) + a$ for all t , where the rotation $R \in O(2)$ and translation $a \in \mathbb{R}^2$ are both independent of t . Physically, we are asking when two moving particles differ by a fixed Euclidean motion at all times, a problem that has significant applications to motion detection and recognition of moving objects.

In the unparametrized problem, we are interested in determining when two curves are congruent under a Euclidean motion, meaning $\bar{C} = R \cdot C + a$ for some fixed Euclidean transformation $(R, a) \in E(2)$. This occurs if and only if there exists a change of parameter $\bar{t} = \tau(t)$ such that $\bar{\mathbf{x}}(\bar{t}) = R \cdot \mathbf{x}(t) + a$ for some fixed Euclidean transformation (R, a) .

A Euclidean symmetry of a curve C is a Euclidean transformation (R, a) that preserves the curve: $R \cdot C + a = C$. For instance, the Euclidean symmetries of a circle consist of the

rotations around its center. In the fixed parameter version, the circle must be parametrized by a constant multiple of arc length for this to hold.

Example 2.2. Consider the action

$$A: (x, u) \mapsto \left(\frac{\alpha x + \beta}{\gamma x + \delta}, \frac{u}{\gamma x + \delta} \right), \quad A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \text{GL}(2), \quad (2.3)$$

of the general linear group $\text{GL}(2)$ on \mathbb{R}^2 . This forms a multiplier representation of $\text{GL}(2)$, cf. [13], [36], which lies at the heart of classical invariant theory. We restrict our attention to curves given by the graphs of functions $u = f(x)$, thereby avoiding issues of reparametrization. Two such curves are equivalent if and only if their defining functions f and \bar{f} are related by the formula

$$f(x) = (\gamma x + \delta) \bar{f} \left(\frac{\alpha x + \beta}{\gamma x + \delta} \right) = (\gamma x + \delta) \bar{f}(\bar{x}), \quad (2.4)$$

for some nonsingular matrix A . Equation (2.4) is the fundamental equivalence condition for first order Lagrangians that depend only on a derivative coordinate in the calculus of variations, cf. [34]. Moreover, if $f(x) = \sqrt[n]{P(x)}$, and $\bar{f}(\bar{x}) = \sqrt[n]{\bar{P}(\bar{x})}$, then (2.4) becomes[†]

$$P(x) = (\gamma x + \delta)^n \bar{P} \left(\frac{\alpha x + \beta}{\gamma x + \delta} \right) = (\gamma x + \delta)^n \bar{P}(\bar{x}), \quad (2.5)$$

which, when P and \bar{P} are polynomials of degree n , forms the fundamental equivalence problem of classical invariant theory.

In the general unparametrized equivalence problem, typically, the submanifolds N and \bar{N} are formulated via explicit parametrizations $\iota: X \rightarrow M$, with image $N = \iota(X)$ and $\bar{\iota}: X \rightarrow M$, with $\bar{N} = \bar{\iota}(X)$, where, for simplicity, the parameter spaces are taken to be the same. (Indeed, since our considerations are always local, we shall not lose any generality by assuming that $X \subset \mathbb{R}^n$ is an open subset of Euclidean space.) In such cases, we can easily reformulate the unparametrized equivalence problem in the following equivalent form.

The Reparametrization Equivalence Problem: Given two embeddings $\iota: X \rightarrow M$ and $\bar{\iota}: X \rightarrow M$ of an n -dimensional manifold X into M does there exist a local diffeomorphism $\Phi: X \rightarrow X$, i.e., a change of parameter, and a group transformation $g \in G$ such that

$$\bar{\iota}(\Phi(x)) = g \cdot \iota(x), \quad \text{for all } x \in X. \quad (2.6)$$

We shall see that by solving the fixed parametrization problem, first in the case of G being a finite dimensional Lie transformation group, then extending this to the case of G being an infinite Lie pseudo-group of transformations, that we will then be able to

[†] We are ignoring the branching of the n^{th} root here. See [34], [36] for a more precise version of this construction.

solve the reparametrization problem. For instance, we can reformulate the unparametrized equivalence problem for curves in the Euclidean plane as a fixed parametrization problem for curves in the extended space $E = \mathbb{R} \times \mathbb{R}^2$, which has coordinates $(t, \mathbf{x}) = (t, x, y)$. The extended curve is given as the graph $\{(t, \mathbf{x}(t))\}$ of the original parametrized curve, and the pseudo-group $\mathcal{G} = \mathcal{D}(1) \times \mathbb{E}(2)$ acting on E consists of a finite-dimensional group, the Euclidean group $\mathbb{E}(2)$ acting on \mathbb{R}^2 , together with the infinite-dimensional pseudo-group $\mathcal{D}(1)$ consisting of all smooth (local) diffeomorphisms $\bar{t} = \tau(t)$ of the parameter space \mathbb{R} .

The formulation of the reparametrization problem in the form (2.6) indicates an intermediate extension of the two cases, in which one only allows a subclass of all possible reparametrizations.

The Restricted Reparametrization Equivalence Problem: Given two embeddings $\iota: X \rightarrow M$ and $\bar{\iota}: X \rightarrow M$ of an n -dimensional manifold X into M and a Lie pseudo-group of transformations \mathcal{H} acting on X , determine whether there exists a group transformation $g \in G$ such that (2.6) holds for some reparametrization $\Phi \in \mathcal{H}$ in the prescribed pseudo-group.

For example, one might consider the problem of equivalence of surfaces in Euclidean space, in which one is only allowed conformal, or area preserving, or Euclidean reparametrizations. The general reparametrization equivalence problem is, of course, a special case when the pseudo-group $\mathcal{H} = \mathcal{D}(X)$ is the entire local diffeomorphism group.

In general, the solution to any equivalence problem is governed by a complete system of invariants. In the present context, the invariants are the fundamental differential invariants for the transformation group action in question. Thus, any solution method must, as a consequence, produce the differential invariants in question.

Example 2.3. In the case of curves in Euclidean geometry, the ordinary curvature[†] function $\kappa = |\mathbf{x}_t|^{-3} (\mathbf{x}_t \wedge \mathbf{x}_{tt})$ is the fundamental differential invariant. For the fixed parametrization problem, there is a second fundamental differential invariant — the speed $v = |\mathbf{x}_t|$. Furthermore, all higher order differential invariants are obtained by successively differentiating the curvature (and speed) with respect to arc length $ds = v dt = |\mathbf{x}_t| dt$, which is the fundamental Euclidean invariant one-form. (In the fixed problem, one can replace s derivatives by t derivatives since dt is also invariant if we disallow any changes in parameter.) A similar result holds for general transformation groups — one can obtain all higher order differential invariants from the fundamental ones by successively applying certain invariant differential operators to the basic differential invariants, cf. [36].

The functional relationships between the fundamental differential invariants will solve the equivalence problem. Roughly speaking, one uses the differential invariants to parametrize a “classifying” or “signature” manifold associated with the given submanifold, and the result is that, under suitable regularity hypotheses, two submanifolds will be congruent under a group transformation if and only if their classifying manifolds are *identical*. For example, in the unparametrized Euclidean curve problem, the classifying curve is parametrized by the two curvature invariants (κ, κ_s) , whereas in the fixed problem, one uses all

[†] Here $|\mathbf{a}|$ is the usual Euclidean norm and $\mathbf{a} \wedge \mathbf{b}$ is the scalar-valued cross product between vectors in the plane.

four invariants $(v, \kappa, v_s, \kappa_s)$ to parametrize the classifying curve. See [5] for applications of the classifying curve to the problem of object recognition in computer vision. Of course, this “solution” reduces one to another potentially difficult *identification problem* — when do two parametrized submanifolds coincide? One approach to the latter problem is to use the Implicit Function Theorem to realize the classifying submanifold as the graph of a function, which eliminates the reparametrization ambiguity. Alternatively, in an algebraic context, a solution can be provided by Gröbner basis techniques, cf. [4]. Neither approach can completely resolve the identification problem, but particular cases can often be handled effectively.

The differential invariants can also be used to effectively determine the structure of the symmetry group. In the case of an effectively acting Lie group G , the codimension of the symmetry subgroup H of the submanifold N , i.e., $\dim G - \dim H$, is the same as the number of functionally independent differential invariants on the submanifold. In particular, the maximally symmetric submanifolds occur when all differential invariants are constant; if G acts transitively, then these can be identified with the homogeneous submanifolds of M , i.e., the orbits of suitable closed subgroups of G . For instance, in the Euclidean case, the maximally symmetric curves are where the curvature is constant, which are the circles and straight lines, since these are the orbits of the one-parameter subgroups of $E(2)$. (Technically, these retain the infinite-dimensional reparametrization group $\mathcal{D}(1)$ as an additional symmetry group.) In the fixed parameter version, the circles and straight lines must be parametrized by a constant multiple of their arc length in order to retain their symmetry distinguished status.

For the Euclidean reparametrization equivalence problem, the associated rigidity theorem states that two curves which are congruent under a Euclidean transformation and have third order contact at a point are, in fact, identical. the rigidity order, namely 3, refers to the maximal order of the differential invariants parametrizing the classifying curve. Note that the symmetric curves have rigidity order 2, since they are uniquely determined, up to a Euclidean transformation, by their constant second order curvature invariant.

Finally, we remark that differential invariants can be used to construct the general invariant differential equations admitting the given transformation group. Specifically, if J_1, \dots, J_N form a complete system of functionally independent k^{th} order differential invariants, defined on an open subset $V^k \subset J^k$ of the jet space where the prolonged group action is regular, then, on V^k , any k^{th} order systems of differential equations admitting G as a symmetry group can be written in terms of the differential invariants: $H_\nu(J_1, \dots, J_N) = 0$. For example, the most general Euclidean-invariant third order differential equation has the form $d\kappa/ds = H(\kappa)$, equating the derivative of curvature with respect to arc length to a function of curvature. Similar comments apply to invariant variational problems, and we refer the reader to [35], [36], for details. These results form the foundations of modern physical field theories, in which one bases the differential equations or variational principle, on its invariance with respect to the theory’s underlying symmetry group. The groups in question range from basic Poincaré and conformal invariance, to the exceptional simple Lie groups lying at the foundations of string theory, as well as infinite-dimensional gauge groups and groups of Kac–Moody type. Remarkably, though, complete systems of differential invariants are known for only a small handful of physically relevant groups,

a collection that includes *none* of the above mentioned groups! Our moving coframe algorithm provides an direct and effective means for providing such classifications.

3. The Maurer–Cartan Forms.

In our approach to the theory and practical implementation of the method of moving frames, the left-invariant Maurer–Cartan forms on a finite-dimensional Lie group play an essential role. We therefore begin by reviewing the basic definition, and then present two computationally effective methods for finding the explicit formulae for the Maurer–Cartan forms. The theoretical justification for the second method will appear in part II.

Throughout this section, G will be an r -dimensional Lie group. We let $L_g: h \mapsto g \cdot h$ denote the standard left multiplication map.

Definition 3.1. A one-form μ on G is called a (left-invariant) *Maurer–Cartan form* if it satisfies

$$(L_g)^* \mu = \mu \quad \text{for all } g \in G. \quad (3.1)$$

The space of Maurer–Cartan forms on G is an r -dimensional vector space, which can naturally be identified with the dual to the Lie algebra \mathfrak{g} of left-invariant vector fields on G . If we choose a basis $\mathbf{v}_1, \dots, \mathbf{v}_r$ of the Lie algebra \mathfrak{g} , then there is a dual basis μ^1, \dots, μ^r , consisting of Maurer–Cartan forms, satisfying $\langle \mu^i; \mathbf{v}_j \rangle = \delta_j^i$, where δ_j^i is the usual Kronecker delta. The basis Maurer–Cartan forms satisfy the fundamental *structure equations*

$$d\mu^i = - \sum_{j < k} C_{jk}^i \mu^j \wedge \mu^k, \quad (3.2)$$

where the coefficients C_{ij}^k are the structure constants corresponding to our choice of basis of the Lie algebra \mathfrak{g} . The Maurer–Cartan forms are a *coframe* on the Lie group G , meaning that they form a pointwise basis for the cotangent space T^*G , or, equivalently, that we can write any one-form ω on G as a linear combination $\omega = \sum f_i \mu^i$ thereof, where the f_i are suitable smooth functions.

The most common method for explicitly determining the Maurer–Cartan forms on a given Lie group is to realize the group $G \subset \text{GL}(n)$ as a matrix Lie group. The independent entries of the $n \times n$ matrix of one-forms

$$\boldsymbol{\mu} = A^{-1} dA \quad (3.3)$$

form a basis for the left-invariant Maurer–Cartan forms on G . Here $A = A(g^1, \dots, g^r) \in G$ represents the general matrix in G , which we have parametrized by local coordinates (g^1, \dots, g^r) near the identity, and $dA = \sum (\partial A / \partial g^i) dg^i$ is its differential, which is an $n \times n$ matrix of one-forms.

For example, in the case $G = \text{GL}(2)$, the four independent Maurer–Cartan forms are the components of the matrix

$$\boldsymbol{\mu} = \begin{pmatrix} \mu_1 & \mu_2 \\ \mu_3 & \mu_4 \end{pmatrix} = A^{-1} dA = \frac{1}{\alpha\delta - \beta\gamma} \begin{pmatrix} \delta d\alpha - \beta d\gamma & \delta d\beta - \beta d\delta \\ \alpha d\gamma - \gamma d\alpha & \alpha d\delta - \gamma d\beta \end{pmatrix}. \quad (3.4)$$

Similarly, if $G = \mathbf{E}(2) = \mathbf{O}(2) \times \mathbb{R}^2$ is the Euclidean group in the plane, then we can identify $\mathbf{E}(2) \subset \mathbf{GL}(3)$ as a subgroup of $\mathbf{GL}(3)$ by identifying $(R, \mathbf{a}) \in \mathbf{E}(2)$ with the 3×3 matrix

$$\begin{pmatrix} R & \mathbf{a} \\ \mathbf{0} & 1 \end{pmatrix} = \begin{pmatrix} \cos \alpha & -\sin \alpha & a \\ \sin \alpha & \cos \alpha & b \\ 0 & 0 & 1 \end{pmatrix}.$$

Substituting into (3.3) leads to

$$\boldsymbol{\mu} = \begin{pmatrix} R^{-1} & -R^{-1}\mathbf{a} \\ \mathbf{0} & 1 \end{pmatrix} \begin{pmatrix} dR & d\mathbf{a} \\ \mathbf{0} & 0 \end{pmatrix} = \begin{pmatrix} 0 & -d\alpha & \cos \alpha da + \sin \alpha db \\ d\alpha & 0 & -\sin \alpha da + \cos \alpha db \\ 0 & 0 & 0 \end{pmatrix}.$$

Thus, the three independent Euclidean Maurer–Cartan forms are

$$\mu_1 = d\alpha, \quad \mu_2 = \cos \alpha da + \sin \alpha db, \quad \mu_3 = -\sin \alpha da + \cos \alpha db. \quad (3.5)$$

In cases when the group is explicitly realized as a local group of transformations on a manifold M , and not necessarily as a matrix Lie group, it is useful to have a direct method for determining the Maurer–Cartan forms. Given $g \in G$ and $z \in M$, we explicitly write the group transformation $\bar{z} = g \cdot z$ in coordinate form:

$$\bar{z}^i = H^i(z, g), \quad i = 1, \dots, n.$$

We then compute the differentials of the group transformations:

$$d\bar{z}^i = \sum_{k=1}^n \frac{\partial H^i}{\partial z^k} dz^k + \sum_{j=1}^r \frac{\partial H^i}{\partial g^j} dg^j, \quad i = 1, \dots, n,$$

of, more compactly,

$$d\bar{z} = H_z dz + H_g dg. \quad (3.6)$$

Next, set $d\bar{z} = 0$ in (3.6), and solve the resulting system of linear equations for the differentials dz^k . This leads to the formulae

$$-dz = F dg = (H_z^{-1} \cdot H_g) dg,$$

or, in full detail,

$$-dz^k = \sum_{j=1}^r F_j^k(z, g) dg^j, \quad k = 1, \dots, n. \quad (3.7)$$

Then, for each k and each fixed $z_0 \in M$, the one-form

$$\mu_0 = \sum_{j=1}^r F_j^k(z_0, g) dg^j, \quad (3.8)$$

is a left-invariant Maurer–Cartan form on the group G . Alternatively, if one expands the right hand side of (3.7) in a power series (or Fourier series, or ...) in z ,

$$\sum_{j=1}^r F_j^k(z, g) dg^j = \sum_{i=0}^{\infty} z^i \mu_i, \quad (3.9)$$

then each coefficient μ_i also forms a left-invariant Maurer–Cartan form on G . In particular, when G acts locally effectively, the resulting collection of one-forms spans the space of Maurer–Cartan forms.

Example 3.2. Consider the action of $\text{GL}(2)$ given by

$$\bar{x} = \frac{\alpha x + \beta}{\gamma x + \delta}, \quad \bar{u} = \frac{u}{\gamma x + \delta}, \quad (3.10)$$

as discussed above. Differentiating (3.10), we find, as in (3.6),

$$\begin{aligned} d\bar{x} &= \frac{(\gamma x + \delta)(\alpha dx + x d\alpha + d\beta) - (\alpha x + \beta)(\gamma dx + x d\gamma + d\delta)}{(\gamma x + \delta)^2} \\ &= \frac{(\alpha\delta - \beta\gamma) dx + (\gamma x + \delta)(x d\alpha + d\beta) - (\alpha x + \beta)(x d\gamma + d\delta)}{(\gamma x + \delta)^2}, \\ d\bar{u} &= \frac{(\gamma x + \delta) du + u(\gamma dx + x d\gamma + d\delta)}{(\gamma x + \delta)^2}. \end{aligned}$$

Setting $d\bar{x} = 0 = d\bar{u}$ and solving for dx and du , we obtain

$$\begin{aligned} -dx &= \frac{\delta d\beta - \beta d\delta}{\alpha\delta - \beta\gamma} + \left(\frac{\delta d\alpha + \gamma d\beta - \alpha d\delta - \beta d\gamma}{\alpha\delta - \beta\gamma} \right) x + \left(\frac{\gamma d\alpha - \alpha d\gamma}{\alpha\delta - \beta\gamma} \right) x^2, \\ -du &= \left(\frac{\alpha d\delta - \gamma d\beta}{\alpha\delta - \beta\gamma} \right) u + \left(\frac{\alpha d\gamma - \gamma d\alpha}{\alpha\delta - \beta\gamma} \right) xu. \end{aligned} \quad (3.11)$$

Note that the coefficients of 1, x and x^2 in the first formula, i.e.,

$$\hat{\mu}_1 = \frac{\delta d\beta - \beta d\delta}{\alpha\delta - \beta\gamma}, \quad \hat{\mu}_2 = \frac{\delta d\alpha + \gamma d\beta - \alpha d\delta - \beta d\gamma}{\alpha\delta - \beta\gamma}, \quad \hat{\mu}_3 = \frac{\gamma d\alpha - \alpha d\gamma}{\alpha\delta - \beta\gamma}, \quad (3.12)$$

recover three of the Maurer–Cartan forms in (3.4), while the coefficient of either u or xu provides the remaining one.

If G does not act effectively on M , then the forms computed by this method will form a basis for the annihilator

$$(\mathfrak{g}_M)^\perp = \{ \omega \in \mathfrak{g}^* \mid \langle \omega, \mathbf{v} \rangle = 0 \text{ for all } \mathbf{v} \in \mathfrak{g}_M \}$$

of the Lie algebra of the global isotropy subgroup

$$G_M = \{ g \in G \mid g \cdot z = z \text{ for all } z \in M \},$$

and thus can be identified with the Maurer–Cartan forms for the effectively acting quotient group $\tilde{G} = G/G_M$. For example, if we only treat the linear fractional transformations in x in (3.10), then the resulting three Maurer–Cartan forms (3.12) all annihilate the generator $\mathbf{v} = \alpha\partial_\alpha + \beta\partial_\beta + \gamma\partial_\gamma + \delta\partial_\delta$ of the isotropy subgroup $\{\lambda\mathbb{1}\} \subset \text{GL}(2)$ consisting of scalar multiples of the identity matrix. Hence, the three one-forms can be identified with a basis for the Maurer–Cartan forms of the effectively acting projective linear group $\text{PSL}(2) = \text{GL}(2)/\{\lambda\mathbb{1}\}$.

4. Compatible Lifts and Moving Coframes.

In this section, we start our development of the moving coframe method. We shall explain how to implement the computational steps in the procedure, which leads to a complete determination of the general moving frame, the associated invariant differential one-forms, which we call the moving coframe, and, consequently, the differential invariants, and the solution to the equivalence problem for submanifolds under the group action. We will begin with the simplest problems and gradually work our way up to more complicated situations.

Throughout this section, we assume that G is an r -dimensional Lie group which acts locally effectively and transitively on an m -dimensional manifold M . (As remarked above, we can always assume local effectiveness by quotienting by the global isotropy subgroup.) Let $z_0 \in M$ be a given point.

Definition 4.1. A smooth map $\gamma: M \rightarrow G$ is called a *compatible lift* with base point z_0 if it satisfies

$$\gamma(z) \cdot z_0 = z. \quad (4.1)$$

In order to compute the most general compatible lift, we solve the system of m equations (4.1) for m of the group parameters in terms of the coordinates z on M and the remaining $r - m = \dim G - \dim M$ group parameters, which we denote by h . This leads to a general formula $g = \gamma_0(z, h)$ for the solution to the compatibility equations (4.1). In other words, by solving the compatibility conditions (4.1), we have effectively “normalized” m of the original group parameters. Since our considerations are always local, in practice, we only need to solve the compatibility equations (4.1) near z_0 . We will call the general compatible lift $\gamma_0(z, h)$ the *moving frame of order zero* for the given transformation group. If $\iota: X \rightarrow M$ defines a parametrized submanifold $N = \iota(X)$, then one can view the composition $\gamma_0(\iota(x), h)$ as a restriction of the order zero moving frame to the submanifold N , where the unnormalized parameters h determine the degree of indeterminacy of the moving frame on N . In geometrical situations, such restrictions can be identified with the classical moving frames; see also Section 7 below.

Example 4.2. Consider the planar action (2.3) of the general linear group $\text{GL}(2)$:

$$A \cdot (x, u) = \left(\frac{\alpha x + \beta}{\gamma x + \delta}, \frac{u}{\gamma x + \delta} \right), \quad A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \text{GL}(2), \quad (4.2)$$

The action (4.2) is transitive on $M = \mathbb{R}^2 \setminus \{u = 0\}$. Choose the base point to be $z_0 = (0, 1)$. Since $A \cdot z_0 = (\beta/\delta, 1/\delta)$, any compatible lift $A = \gamma(x, u)$ must satisfy $\beta/\delta = x$, $1/\delta = u$, and hence the solution to (4.1) is

$$\beta = \frac{x}{u}, \quad \delta = \frac{1}{u}. \quad (4.3)$$

The most general compatible lift thus has the form

$$\gamma_0(x, u, \alpha, \gamma) = \begin{pmatrix} \alpha & x/u \\ \gamma & 1/u \end{pmatrix}, \quad (4.4)$$

where $\alpha = \alpha(x, u)$, $\gamma = \gamma(x, u)$ are arbitrary functions, subject only to the condition $\alpha \neq x\gamma$, so that the determinant of (4.4) does not vanish, and hence γ_0 does take its values in the group $\text{GL}(2)$.

Note that since G acts transitively, we can locally identify $M \simeq G/H$ with a homogeneous space, where $H = G_{z_0}$ is the isotropy group of the base point. Therefore, a compatible lift is merely a (local) section of the fiber bundle $G \rightarrow G/H$.

Proposition 4.3. *Two maps $\gamma, \hat{\gamma}: M \rightarrow G$ are compatible lifts with the same base point if and only if they satisfy*

$$\hat{\gamma}(z) = \gamma(z) \cdot \eta(z),$$

where $\eta: M \rightarrow H$ is an arbitrary map to the isotropy subgroup of the base point z_0 .

Thus, in the previous example, the isotropy subgroup H of the point $z_0 = (1, 0)$ consists of all invertible lower triangular matrices of the form $\begin{pmatrix} \alpha' & 0 \\ \gamma' & 1 \end{pmatrix}$. Indeed, we can rewrite (4.4) in the factored form

$$\gamma_0(x, u, \alpha, \gamma) = \begin{pmatrix} \alpha & x/u \\ \gamma & 1/u \end{pmatrix} = \begin{pmatrix} 1 & x/u \\ 0 & 1/u \end{pmatrix} \begin{pmatrix} \alpha' & 0 \\ \gamma' & 1 \end{pmatrix},$$

where $\alpha' = \alpha - x\gamma$, $\gamma' = u\gamma$, reconfirming Proposition 4.3 in this particular example.

Although the remaining unspecified group parameters can be identified with the isotropy subgroup coordinates, in any practical implementation of the moving coframe algorithm, it is not necessary to identify the isotropy subgroup explicitly, nor to adopt its particular coordinates to characterize the order zero moving frame. Thus, in the present example, the coordinates α, γ , are just as effective as the subgroup coordinates $\tilde{\alpha}, \tilde{\gamma}$. (The interested reader can follow through the ensuing calculations using the subgroup coordinates instead, reproducing the final result.)

The order zero moving frame $\gamma_0(z, h)$, which is the general solution to the compatible lift equations (4.1), defines a map from the zeroth order moving frame bundle $C_0 = M \times H \simeq G/H \times H$, coordinatized by (z, h) to the group G , which is, in fact, a local diffeomorphism $\gamma_0: C_0 \xrightarrow{\simeq} G$. There is an induced action of G on the moving frame bundle C_0 that makes γ_0 into a G -equivariant map: $\gamma_0(g \cdot (z, h)) = g \cdot \gamma_0(z, h)$. Thus, the action on the unnormalized group parameters h can be explicitly determined by multiplying the moving frame on the left by a group transformation. The action of G on C_0 projects to the original action of G on M , so that $g \cdot (z, h) = (g \cdot z, \eta(g, z, h))$ for $g \in G$.

In the present example, we can the induced action of $\text{GL}(2)$ on the unspecified parameters α, γ , by multiplying the moving frame (4.4) on the left by a group element $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}(2)$; we find

$$\begin{pmatrix} \bar{\alpha} & \bar{x}/\bar{u} \\ \bar{\gamma} & 1/\bar{u} \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \begin{pmatrix} \alpha & x/u \\ \gamma & 1/u \end{pmatrix} = \begin{pmatrix} a\alpha + b\gamma & (ax + b)/u \\ c\alpha + d\gamma & (cx + d)/u \end{pmatrix}, \quad (4.5)$$

Therefore, the induced action of $G = \text{GL}(2)$ on the moving frame bundle C_0 is given by

$$\bar{x} = \frac{ax + b}{cx + d}, \quad \bar{u} = \frac{u}{cx + d}, \quad \bar{\alpha} = a\alpha + b\gamma, \quad \bar{\gamma} = c\alpha + d\gamma. \quad (4.6)$$

Note that the (x, u) transformations coincide with the original action (2.3), as they should.

Remark: In practical implementations of the moving coframe algorithm, we do *not* have to explicitly compute this group action. We do this here so as to give the reader some justification for our claims.

Remark: The action of G on $C_0 = M \times H$ does *not* project to an action on the isotropy subgroup H , even if we use the associated subgroup coordinates. In the present example, we find (4.6) implies that the subgroup coordinates α' , γ' transform according to

$$\bar{\alpha}' = \frac{ad - bc}{cx + d} \alpha', \quad \bar{\gamma}' = \gamma' + \frac{cu}{cx + d} \alpha'.$$

The next step is to characterize the group transformations by a collection of differential forms. In the finite-dimensional situation we are currently considering, these will be obtained by pulling back the left-invariant Maurer–Cartan forms μ on G to the order zero moving frame bundle C_0 using the compatible lift. The resulting one-forms $\zeta_0 = \gamma_0^* \mu$ will provide a coframe on C_0 , which we name the *moving coframe* of order zero. The moving coframe forms ζ_0 clearly satisfy the same Maurer–Cartan structure equations (3.2).

Theorem 4.4. *The order zero moving coframe forms completely characterize the group transformations on the bundle C_0 . In other words, a map $\Psi: C_0 \rightarrow C_0$ satisfies $\Psi^* \zeta_0 = \zeta_0$ if and only if $\Psi(z, h) = g \cdot (z, h)$ coincides with the action of a group element $g \in G$ on C_0 .*

In the present example, we substitute the formulae (4.3) characterizing our compatible lift (4.4) into the Maurer–Cartan forms (3.4). The result is the order zero moving coframe

$$\begin{aligned} \zeta_1 &= \frac{d\alpha - x d\gamma}{\alpha - \gamma x}, & \zeta_2 &= \frac{dx}{u(\alpha - \gamma x)}, \\ \zeta_3 &= \frac{u(\alpha d\gamma - \gamma d\alpha)}{\alpha - \gamma x}, & \zeta_4 &= -\frac{\gamma dx}{\alpha - \gamma x} - \frac{du}{u}, \end{aligned} \tag{4.7}$$

which forms a basis for the space of one-forms on C_0 . The skeptical reader can explicitly check that these four one-forms really do completely characterize the group action (4.6), as described in Theorem 4.4.

Let us now consider a curve $N \subset M$. For simplicity, we shall assume that the curve coincides with the graph of a function $u = u(x)$. However, this restriction is not essential for the method to work, and later we show how general parametrized curves can also be readily handled by the general method. We restrict the moving coframe forms to the curve, which amounts to replacing the differential du by its “horizontal” component $u_x dx$. If we interpret the derivative u_x as a coordinate on the first jet space $J^1 = J^1(M) \simeq \mathbb{R}^3$ of curves in M , then the restriction of a differential form to the curve can be reinterpreted as the natural projection of the one-form du on J^1 to its horizontal component, using the canonical decomposition of differential forms on the jet space into horizontal and contact components. Indeed, the vertical component of the form du is the contact form $du - u_x dx$, which vanishes on all prolonged sections of the first jet bundle $J^1 M$. We refer the reader

to [36; Chapter 4] for a comprehensive review of the contact geometry of jet bundles. Therefore, the restricted (or horizontal) moving coframe forms are explicitly given by

$$\begin{aligned}\theta_1 &= \frac{d\alpha - x d\gamma}{\alpha - \gamma x}, & \theta_2 &= \frac{dx}{u(\alpha - \gamma x)}, \\ \theta_3 &= \frac{u(\alpha d\gamma - \gamma d\alpha)}{\alpha - \gamma x}, & \theta_4 &= \frac{\gamma(xu_x - u) - \alpha u_x}{u(\alpha - \gamma x)} dx,\end{aligned}\tag{4.8}$$

which now depend on first order derivatives.

The next step in the procedure is to look for invariant combinations of coordinates and group parameters. Each such invariant combination will either provide us with a basic differential invariant for the problem, or, in the case that it explicitly depends on the remaining group parameters, a “lifted invariant” which can be normalized and thereby eliminate one of the remaining group parameters, as discussed below. Specifically, in the present example, a function $J(\alpha, \gamma, x, u, u_x)$ will be a *lifted invariant* provided it is unaffected by the group action on its arguments, meaning that

$$J(\bar{\alpha}, \bar{\gamma}, \bar{x}, \bar{u}, \bar{u}_{\bar{x}}) = J(\alpha, \gamma, x, u, u_x),\tag{4.9}$$

wherever $\bar{\alpha}, \bar{\gamma}, \bar{x}, \bar{u}$, are related to α, γ, x, u , according to the induced action (4.6) of the group on the moving frame bundle, and $\bar{u}_{\bar{x}}$ is related to u_x according to the standard prolongation, [36], of the action of G on M to the first jet bundle J^1 . In the present case, if \bar{x}, \bar{u} are given by (4.6), then a straightforward chain rule computation provides the prolonged action of $GL(2)$ on the derivative coordinate:

$$\bar{u}_{\bar{x}} = \frac{(cx + d)u_x - cu}{ad - bc}.\tag{4.10}$$

In other words, we interpret $\alpha, \gamma, x, u, u_x$ as coordinates on a bundle $\tilde{C}_0 \rightarrow J^1$ over the first jet space, which is merely the pull-back $\tilde{C}_0 = (\pi_0^1)^* C_0$ of the zeroth order moving frame bundle via the standard projection $\pi_0^1: J^1 \rightarrow M$. There is an induced action of G on \tilde{C}_0 which projects to its prolonged action on J^1 . A (first order) lifted invariant, then, is just a function $J: \tilde{C}_0 \rightarrow \mathbb{R}$ which is invariant under the action of G on \tilde{C}_0 . If the lifted invariant $J = J(x, u, u_x)$ does not, in fact, depend on the group parameters α, γ , then it will be a (first order) differential invariant. (However, in the present example, there are no non-constant first order differential invariants, since $GL(2)$ acts transitively on J^1 .) Alternatively, if J actually depends on either α or γ then it can be used in the normalization procedure.

Fortunately, the lifted invariants can be determined without explicitly computing the prolonged group action, or solving any differential equations. They are systematically found by looking at the linear dependencies among the restricted (horizontal) moving coframe forms! Indeed, because the one-forms are invariant, each coefficient J_i in a linear relation $\theta_0 = J_1 \theta_1 + \cdots + J_k \theta_k$, in which the forms θ_i on the right hand side are linearly independent, is automatically under the action of the group. In our example, we note that,

among the restricted one-forms (4.8), there is one linear dependency, namely $\theta_4 = J\theta_2$, where

$$J = \gamma(xu_x - u) - \alpha u_x. \quad (4.11)$$

One can explicitly verify that J is indeed a lifted invariant, meaning that it satisfies (4.9) whenever $\bar{\alpha}, \bar{\gamma}, \bar{x}, \bar{u}, \bar{u}_x$, are related to $\alpha, \gamma, x, u, u_x$, according to (4.6), (4.10).

The ultimate goal of the moving frame method is to eliminate all the ambiguities, i.e., the undetermined group parameters, in the original moving frame, in a suitably invariant manner. Cartan's crucial observation is that, we can, without loss of generality, *normalize* any lifted invariant by setting it equal to any convenient constant value,

$$J(\alpha, \gamma, x, u, u_x) = c, \quad (4.12)$$

without affecting the equivalence problem. In (4.12), c can be any convenient constant, subject only to the requirement that the solutions to (4.12) remain in the group, e.g., that the determinant of any resulting matrix (4.4) remains nonzero. Typically, c is taken to be 0, 1, or -1 , although other values can be chosen to simplify the resulting formulae. Assuming that J does actually depend on the parameters α, γ , we can solve the normalization equation (4.12) for one of them; e.g., $\alpha = \alpha(\gamma, x, u, u_x)$. Because J is an invariant, such a normalization will not alter the solution to the equivalence problem, and hence we can use it to eliminate α from the original formulae for the moving frame and moving coframe. The result is a first order moving frame, depending on one fewer unnormalized group parameter. This produces to a corresponding first order moving coframe, to which one can apply the same procedure, leading to a chain of successive normalizations and reductions, eventually enabling one to completely eliminate all the undetermined parameters and specify a uniquely defined moving frame on some suitable jet bundle $J^n = J^n(M)$.

In accordance with the general procedure, then, we can normalize our particular lifted invariant (4.11) by setting it equal to zero, $J = 0$, which can be effected by setting

$$\alpha = \left(\frac{xu_x - u}{u_x} \right) \gamma. \quad (4.13)$$

Substituting (4.13) into (4.4) produces the *first order moving frame*

$$\gamma_1(x, u, u_x, \gamma) = \begin{pmatrix} (xu_x - u)\gamma/u_x & x/u \\ \gamma & 1/u \end{pmatrix}, \quad (4.14)$$

which now depends on first order derivatives of u , and just one unnormalized parameter. We can regard the coordinates (x, u, u_x, γ) as parametrizing a bundle $C_1 \rightarrow J^1$ sitting over the first jet space, which is realized as a G -invariant subbundle of \tilde{C}_0 , namely $C_1 = J^{-1}\{0\} \subset \tilde{C}_0$. As before, one can restrict the first order moving frame to a curve $u = u(x)$ by restricting the map γ_1 to the first prolongation or jet of the curve, i.e., we set $u = u(x)$, $u_x = u'(x)$, in (4.14), with γ indicating the remaining ambiguity. There is an induced action of $GL(2)$ on C_1 , which projects to the usual first prolonged action $G^{(1)}$ of the group on J^1 , cf. (4.6), (4.10), and makes the first order moving frame $\gamma_1: C_1 \xrightarrow{\sim} G$ into a local

G -equivariant diffeomorphism. In our case, the explicit transformation rules on C_1 are given by

$$\bar{x} = \frac{ax + b}{cx + d}, \quad \bar{u} = \frac{u}{cx + d}, \quad \bar{u}_x = \frac{(cx + d)u_x - cu}{ad - bc}, \quad \bar{\gamma} = \left(\frac{(cx + d)u_x - cu}{ad - bc} \right) \gamma, \quad (4.15)$$

which coincide with left multiplication of the first order moving frame (4.14) by the given group element. (Again, these are provided for illustration only, and are not essential for application of the method.) Furthermore, substituting (4.13) into (4.7), we find the first order moving coframe

$$\begin{aligned} \zeta_1 &= \frac{d\gamma}{\gamma} - \frac{du_x}{u_x} + \frac{du - u_x dx}{u}, & \zeta_2 &= -\frac{u_x dx}{\gamma u^2}, \\ \zeta_3 &= \frac{\gamma u du_x}{u_x} - \gamma(du - u_x dx), & \zeta_4 &= \frac{du - u_x dx}{u}. \end{aligned} \quad (4.16)$$

As in the order zero case, cf. Theorem 4.4, the first order moving coframe completely characterizes the group transformations on C_1 .

As before, we determine new lifted invariants by restricting the first order moving coframe one-forms to a curve $u = u(x)$. This amounts to replacing du and du_x by their horizontal components $u_x dx$ and $u_{xx} dx$ respectively, leading to the restricted forms

$$\theta_1 = \frac{d\gamma}{\gamma} - \frac{u_{xx} dx}{u_x}, \quad \theta_2 = -\frac{u_x dx}{\gamma u^2}, \quad \theta_3 = \frac{\gamma u u_{xx} dx}{u_x}, \quad \theta_4 = 0, \quad (4.17)$$

that now depend on second order derivatives. Alternatively, one could deduce these restricted forms by substituting the normalization (4.13) into the previous restricted forms (4.8). Note in particular that the fact that θ_4 vanishes is an automatic consequence of our normalization condition $\theta_4 = J\theta_2 = 0$; alternatively, we note that ζ_4 is an invariant contact form, which hence vanishes when restricted to any submanifold. Now there is an additional dependency, namely $\theta_3 = K\theta_2$, where

$$K = -\frac{\gamma^2 u^3 u_{xx}}{u_x^2},$$

is a new lifted invariant. Again, the reader can check that K is invariant under the prolonged action of $GL(2)$ on the bundle $\tilde{C}_1 = (\pi_1^2)^* C_1 \rightarrow J^2$, where $\pi_1^2: J^2 \rightarrow J^1$ is the natural projection, provided by (4.15) and the second order prolongation (chain rule) formula

$$\bar{u}_{\bar{x}\bar{x}} = \frac{(cx + d)^3 u_{xx}}{(ad - bc)^2}. \quad (4.18)$$

We can be normalize $K = -1$ by setting

$$\gamma = \frac{u_x}{\sqrt{u^3 u_{xx}}}. \quad (4.19)$$

Note that we *cannot* normalize $K = 0$ since this would require $\gamma = 0$, but then the lift (4.14) would have zero determinant, violating the group conditions. The final lift

$$\gamma_2(x, u, u_x, u_{xx}) = \begin{pmatrix} \frac{xu_x - u}{\sqrt{u^3 u_{xx}}} & \frac{x}{u} \\ \frac{u_x}{\sqrt{u^3 u_{xx}}} & \frac{1}{u} \end{pmatrix} \quad (4.20)$$

defines the second order moving frame on the jet space. Now the lift has been completely specified, and (4.20) contains a complete description of the geometry of curves (functions) under the transformation group (2.3). The second order moving frame provides an explicit G -equivariant identification $\gamma_2: V^2 \xrightarrow{\sim} G$ of the open subset $V^2 = \{uu_{xx} \neq 0\} \subset J^2$ of the second jet bundle with an open subset of the group G , identifying the prolonged action of $G^{(2)}$ on J^2 with the ordinary left multiplication on G ; thus

$$\gamma_2(g^{(2)} \cdot z^{(2)}) = g \cdot \gamma_2(z^{(2)}), \quad g \in \text{GL}(2), \quad z^{(2)} \in V^2.$$

Substituting (4.19) produces the final set of invariant one-forms

$$\begin{aligned} \zeta_1 &= -\frac{du_{xx}}{2u_{xx}} - \frac{du}{2u} - \frac{u_x dx}{u}, & \zeta_2 &= -\sqrt{\frac{u_{xx}}{u}} dx, \\ \zeta_3 &= \frac{du_x}{\sqrt{uu_{xx}}} - \frac{u_x(du - u_x dx)}{\sqrt{u^3 u_{xx}}}, & \zeta_4 &= \frac{du - u_x dx}{u}, \end{aligned} \quad (4.21)$$

which form the second order moving coframe. Note that the second order moving frame (4.20) provides an equivalence, $\gamma_2^* \mu_i = \zeta_i$, mapping the moving coframe forms on the second order jet space to the Maurer–Cartan forms (3.4) on the group. Consequently, the forms ζ_i uniquely characterize the second order prolonged action of $\text{GL}(2)$ on $V^2 \subset J^2$.

Finally, the restricted (horizontal) moving coframe forms become

$$\theta_1 = -\frac{uu_{xxx} + 3u_x u_{xx}}{2uu_{xx}} dx, \quad \theta_2 = \sqrt{\frac{u_{xx}}{u}} dx, \quad \theta_3 = -\theta_2, \quad \theta_4 = 0.$$

There is one final linear dependency, namely $\theta_1 = -I\theta_2$, where

$$I = \frac{uu_{xxx} + 3u_x u_{xx}}{2\sqrt{uu_{xx}^3}} \quad (4.22)$$

is the fundamental differential invariant of the transformation group, also known as the group invariant curvature. The remaining one-form $ds = \theta_2$ is the fundamental invariant one-form, or group-invariant arc length element. All higher order differential invariants can be found by differentiating the curvature invariant with respect to the invariant arc length; for instance, the fundamental fourth order differential invariant is

$$\begin{aligned} J &= \frac{\partial I}{\partial \theta^2} = \frac{dI}{ds} = \sqrt{\frac{u}{u_{xx}}} \frac{dI}{dx} \\ &= \frac{2u^2 u_{xx} u_{xxxx} - 3u^2 u_{xxx}^2 - 2uu_x u_{xx} u_{xxx} + 6uu_{xx}^3 - 3u_x^2 u_{xx}^2}{4uu_{xx}^3}. \end{aligned} \quad (4.23)$$

From the general theory, we conclude that every differential invariant for the group (2.3) is a function of the curvature and its successive derivatives with respect to the arc length. On the regular part V^2 of the jet space J^2 , all $GL(2)$ invariant ordinary differential equations can be written in terms of these invariants; for instance the most general invariant third order ordinary differential equation has the form

$$uu_{xxx} + 3u_x u_{xx} = k\sqrt{uu_{xx}^3}, \quad (4.24)$$

for some constant k .

Applications to the equivalence problem for curves (which include the equivalence problem for first order Lagrangians as well as that of classical invariant theory) follow directly from the general results. Given a function $u = u(x)$, we define its *classifying curve* \mathcal{C} to be the planar curve parametrized by the fundamental differential invariants $I(x), J(x)$. The general result states that two curves are mapped to each other by a group transformation (2.3), so $\bar{\mathcal{C}} = g \cdot \mathcal{C}$, if and only if their classifying curves are identical, $\bar{\mathcal{C}} = \mathcal{C}$. A curve \mathcal{C} is maximally symmetric if and only if its classifying reduces to a point; in this case the original curve is, in fact, is an orbit of a one-parameter group subgroup of $GL(2)$. Thus, we have, in a very simple and direct manner, recovered the results in [34] on the equivalence and symmetry of binary forms, which were found by a much less direct approach based on the standard Cartan equivalence problem for particle Lagrangians. A corresponding rigidity result also follows: two curves that are equivalent under a group transformation are identical if and only if they have fourth order contact at a point, four referring to the order of the fundamental differential invariant J .

There are a few technical points that should have been addressed during the preceding discussion. First, one needs to impose certain conditions on the function $u(x)$ in order to ensure that the computation is valid. For instance, the normalization (4.13) requires $u_x \neq 0$, i.e., the curve does not have a horizontal tangent. (We have already assumed that it does not have a vertical tangent by requiring that it be the graph of a smooth function.) If $u_x = 0$, then we can still normalize $J = 0$ as long as $u \neq xu_x$, in which case we normalize by solving for γ instead of α . Actually, both cases can be simultaneously handled by the normalization $\alpha = \lambda(xu_x - u)$, $\gamma = \lambda u_x$, where $\lambda \neq 0$ is a new parameter whose normalization will be specified at the next stage of the procedure. The reader can check that this alternative procedure leads to the same lift and differential invariants as before. In the second normalization, we have assumed[†] $u_{xx} > 0$ in order to take the square root. For $u_{xx} < 0$ we would need to normalize $K = +1$, and use $\sqrt{-u_{xx}}$ instead. Thus the problem actually separates into two branches, with the inflection points $u_{xx} = 0$ being interpreted as singular points for the group action. The straight lines, for which $u_{xx} \equiv 0$, form a special class and must be analyzed separately. Finally, the square root itself has a sign ambiguity (or, in the complex case, an ambiguity in its choice of branch). Both signs must, in fact, be allowed in the final expression for the lift and the differential invariants. Such branching and ambiguous sign phenomena will be familiar to practitioners of the Cartan equivalence method; see [36] for a detailed discussion of these issues.

[†] In the complex-valued problem, there is no sign restriction.

Let us finish this section by summarizing the basic method, in a form which will apply to much more general problems. The basic steps are:

- (a) Determine the general invariant lift, or moving frame of order zero, by solving (4.1) for the given group action.
- (b) Determine the invariant forms. In the finite-dimensional case, they are the Maurer–Cartan forms, which can be computed either by using the matrix approach, or by direct use of the transformation group formulae.
- (c) Use the invariant lift to pull-back the invariant forms, leading to the moving coframe of order zero.
- (d) Determine lifted invariants by finding linear dependencies among the restricted or horizontal components of the moving coframe forms.
- (e) Normalize any group-dependent invariants to convenient constant values by solving for some of the unspecified parameters.
- (f) Successively eliminate parameters by substituting the normalization formulae into the moving coframe and recomputing dependencies.
- (g) After the parameters have all been normalized, the differential invariants will appear through any remaining dependencies among the final moving coframe elements. The invariant differential operators are found as the dual differential operators to a basis for the invariant coframe forms.

Note that we do not need the explicit isotropy groups for the transformation group actions, nor do we need compute explicit formulae for the prolonged group action in order to successfully apply the method.

Remark: If one is solely interested in the final differential invariants and invariant horizontal one-forms (i.e., invariant forms on the submanifold itself), then one need only compute the effect of the normalizations on the horizontal components of the moving coframe forms during the computation. The moving coframe itself will also include invariant contact forms, which vanish upon restriction, but which, nevertheless, play an important role in other aspects of the geometry. See [36], [38], [18], for applications of invariant contact forms to the study of invariant evolution equations, with applications to image processing. Applications to the computation of the invariant cohomology of the variational bicomplex, cf. [2], are also of particular importance in the analysis of symmetries and conservation laws of variational problems.

Remark: The proposed method of moving coframes has the same basic structure as the Cartan equivalence method, [11], [14], [36], in that one deals with a system of differential forms depending on arbitrary parameters, and seeks to normalize all the parameters by a suitable collection of lifted invariants. One can, indeed, view the two methods as particular cases of a completely general equivalence procedure. However, it is worth pointing out a few of the differences between the two. First, the Cartan method only deals with lifted coframes, whose constituents are linearly independent differential forms, whereas the differential forms occurring in the moving coframe method are linearly dependent. The invariant combinations (lifted invariants) used to normalize the parameters are found via linear dependencies in the moving coframe method, whereas they arise as unabsorbed torsion coefficients in the differentials of the lifted coframe forms in the Cartan equivalence

method. In the moving coframe method, the differentials of the moving coframe one-forms satisfy the Maurer–Cartan structure equations, and hence do not provide any nonconstant invariants. Finally, and perhaps most significantly, the group parameters g only occur algebraically in the lifted coframe elements in the Cartan equivalence method, whereas in the moving frame problems their differentials dg occur as well, since they appear in the Maurer–Cartan forms. One can, of course, imagine solving hybrid equivalence problems, in which aspects of both problems occur during the normalization procedure, although we are not currently aware of any interesting examples where these occur naturally.

5. Intransitive Lie Group Actions.

Our next task is to extend the moving coframe method to the case of finite-dimensional Lie groups whose action is no longer transitive. In the intransitive case, we still assume that G is an r -dimensional Lie group acting effectively, and regularly, which implies that its orbits, which we take to have dimension s , form a foliation of M . We choose a local *cross-section* $K \subset M$ to this foliation, i.e., a submanifold of dimension $m - s$ intersecting the orbits transversally, and introduce a *compatible lift* $\gamma: M \rightarrow G$ by requiring that, for each z near K , the lift γ satisfies

$$z = \gamma(z) \cdot z_0, \quad \text{for some } z_0 \in K. \quad (5.1)$$

The general solution to the compatible lift equations (5.1) will be of the form $\gamma(z, h)$ depending on $r - s$ parameters h . Note that unless the isotropy subgroups at each point in the cross-section happen to be identical, we cannot identify the unspecified parameters as local coordinates on any subgroup $H \subset G$, leading us beyond any principal bundle-theoretic interpretation of the method. Nevertheless, the Implicit Function Theorem will allow us to locally write the general compatible lift in this form. In addition, the group admits (locally) $m - s$ functionally independent invariants, $I_1(z), \dots, I_{m-s}(z)$, whose level sets characterize the orbits. The zeroth order moving frame will then be the map

$$\gamma_0(z, h) = (\gamma(z, h), I(z)), \quad (5.2)$$

whose first components $g = \gamma(z, h)$ are those of the general compatible lift (for the given cross-section) and, in addition, has the invariants $w = I(z) = (I_1(z), \dots, I_{m-s}(z))$ as further components. Note that γ_0 is only locally defined, since z must lie near the cross-section K , and, moreover, the remaining parameters h are determined in accordance with the Implicit Function Theorem.

Note: We can view the range $G \times \mathbb{R}^{m-s}$ of γ_0 as having the structure of a Cartesian product Lie group, the additive group structure on the second factor formalizing the fact that we can add invariants.

The moving coframe forms in this case are constructed from the Maurer–Cartan forms μ on the group G , together with the coordinate one-forms $d\mathbf{w} = \{dw_1, \dots, dw_{m-s}\}$ on \mathbb{R}^{m-s} . The group transformations are then characterized by the conditions

$$\Phi^* \bar{\mathbf{w}} = \mathbf{w}, \quad \Phi^* d\bar{\mathbf{w}} = d\mathbf{w}, \quad \Phi^* \bar{\mu} = \mu. \quad (5.3)$$

Using the moving frame lift $g = \gamma(z, h)$, $w = I(z)$, to pull back these one-forms, we are led to the zeroth order moving coframe, consisting of the pulled-back Maurer–Cartan forms $\gamma_0^* \boldsymbol{\mu}$, along with the differentials $\gamma_0^* d\mathbf{w}_\kappa = dI_\kappa$ of the group invariants. At this stage, the set up of the intransitive problem is complete, and one proceeds to solve it, as in the transitive case, by looking for dependencies among the restricted coframe forms and then normalizing the resulting lifted invariants.

Example 5.1. The intransitive action

$$A : (x, u) \mapsto \left(x, \frac{\alpha u + \beta}{\gamma u + \delta} \right), \quad A = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \text{SL}(2), \quad (5.4)$$

of the special linear group $\text{SL}(2)$ on $M = \mathbb{R}^2$ arises in complex function theory, [19]. (We restrict to $\text{SL}(2)$ in order to maintain local effectiveness.) The group orbits are vertical lines and so the basic invariant is merely $I(x, u) = x$. We choose the cross-section $K = \{u = 0\}$. Solving the equation $A(x, u) \cdot z_0 = (x, u)$, where $z_0 = (y, 0)$, leads to the general compatible lift

$$A_0(x, u, \alpha, \delta) = \begin{pmatrix} \alpha & \delta u \\ \frac{\alpha\delta - 1}{\delta u} & \delta \end{pmatrix}, \quad (5.5)$$

which forms the group component of the zeroth order moving frame. The other component is just the invariant

$$w = I(x, u) = x. \quad (5.6)$$

Pulling back the Maurer–Cartan forms $\boldsymbol{\mu} = A^{-1} dA$ and dw via the lift (5.5), (5.6), leads to the order zero moving coframe

$$\begin{aligned} \zeta_1 &= (\alpha\delta - 1) \frac{du}{u} - \frac{d\delta}{\delta}, & \zeta_2 &= \delta^2 du, \\ \zeta_3 &= \frac{u d(\alpha\delta) + \alpha\delta(1 - \alpha\delta) du}{\delta^2 u^2}, & \zeta_4 &= dx. \end{aligned} \quad (5.7)$$

As before, we restrict to a curve $u = u(x)$ by replacing du by $u_x dx$. Letting θ_i denote the horizontal component of ζ_i , we find that there is one resulting linear dependency, namely

$$\theta_2 = \delta^2 u_x dx = J dx = J\theta_4.$$

This leads to the first normalization $\delta = 1/\sqrt{u_x}$ resulting from setting $J = 1$. Substituting this normalized value into (5.5), (5.6), provides the first order moving frame. Furthermore, substituting into (5.7) produces the second order moving coframe, with horizontal restrictions

$$\theta_1 = \left(\frac{2\alpha u_x^{3/2} + u u_{xx} - 2u_x^2}{2u u_x} \right) dx, \quad \theta_2 = \theta_4 = dx, \quad \theta_3 = \frac{\sqrt{u_x}}{u} (d\alpha - \alpha \theta_1). \quad (5.8)$$

Now we normalize $\theta_1 = 0$ by setting $\alpha = u_x^{-3/2}(u_x^2 - \frac{1}{2}uu_{xx})$. The final moving frame (of order 2) is

$$A_2 = \frac{1}{u_x^{3/2}} \begin{pmatrix} u_x^2 - \frac{1}{2}uu_{xx} & uu_x \\ -\frac{1}{2}u_{xx} & u_x \end{pmatrix}, \quad w = x. \quad (5.9)$$

The corresponding restricted moving coframe has reduced to

$$\theta_2 = \theta_4 = dx, \quad \theta_3 = -\frac{1}{4}S\theta_2, \quad \theta_1 = 0, \quad (5.10)$$

where

$$S = \frac{2u_x u_{xxx} - 3u_{xx}^2}{u_x^2} \quad (5.11)$$

is the classical Schwarzian derivative of the function $u(x)$, whose invariance under linear fractional transformations is of fundamental importance in complex function theory. Since the one-form dx is invariant, all the higher order differential invariants are found by differentiating S with respect to x .

Actually, the preceding computation can be slightly simplified by extending our general method to non-effective actions. We consider (5.4) as defining a non-effective (and intransitive) action of the general linear group $GL(2)$ on \mathbb{R}^2 . We may apply the second algorithm for computing the required Maurer–Cartan forms, leading to the three one-forms (3.12) that annihilate the global isotropy subalgebra. We substitute the compatible lift formulae $\beta = \delta u$ for the order zero moving coframe, which is now $A_0 = \begin{pmatrix} \alpha & \delta u \\ \gamma & \delta \end{pmatrix}$, into (3.12), leading to the restricted moving coframe forms

$$\begin{aligned} \widehat{\theta}_1 &= \frac{\delta u_x dx}{\alpha - u\gamma}, & \widehat{\theta}_2 &= \frac{d\alpha - u d\gamma + \gamma u_x dx}{\alpha - u\gamma} - \frac{d\delta}{\delta}, \\ \widehat{\theta}_3 &= \frac{\gamma d\alpha - \alpha d\gamma}{\delta(\alpha - u\gamma)}, & \widehat{\theta}_4 &= dx. \end{aligned} \quad (5.12)$$

The first dependency between $\widehat{\theta}_1$ and $\widehat{\theta}_4$ leads to the normalization $\delta = (\alpha - u\gamma)/u_x$. Substituting into $\widehat{\theta}_3$ leads to a second dependency, and the resulting normalization yields $\alpha = \gamma(u - 2u_x^2/u_{xx})$. At this stage, even though we have not normalized the final parameter γ , it no longer appears in the coframe, which coincides with our earlier one, (5.10). It does, of course, occur in the final moving frame lift, which is obtained by multiplying the matrix A_2 in (5.9) by γ . However, γ plays no other role in the problem, and merely reflects a final indeterminacy stemming from the ineffectiveness of the group action. The main point in this solution method is that one does not have to implement the effectivity restriction by hand, as was done in the original lift (5.5), in order to solve the problem. Indeed, in more complicated examples, it may be relatively straightforward to write down the compatible lift to an ineffective group action, whereas doing the same for the effectively acting quotient group G/G_M may be considerably more complicated.

Example 5.2. Consider the elementary similarity group $G = \mathbb{R}^+ \ltimes \mathbb{R}^2$ acting transitively on $M = \mathbb{R}^2$ via

$$A : (x, u) \longmapsto (\alpha x + a, \alpha u + b). \quad (5.13)$$

For the base point $z_0 = (0, 0)$, the associated moving frame of order 0 is the lift with $a = x$, $b = u$. The Maurer–Cartan forms $\{d\alpha/\alpha, da/\alpha, db/\alpha\}$ are pulled back to provide the zeroth order moving coframe, whose horizontal components are

$$\theta_1 = \frac{d\alpha}{\alpha}, \quad \theta_2 = \frac{dx}{\alpha}, \quad \theta_3 = \frac{u_x dx}{\alpha}. \quad (5.14)$$

There is a single linear dependency $\theta_3 = I\theta_2$, but the resulting invariant $I = u_x$ does *not* depend on the remaining group parameter, and hence *cannot* be used to normalize it. To proceed further in such cases, we work in analogy with the preceding intransitive case. Here the intransitivity is on the first order jet bundle, and is an indication of the fact that this particular group exhibits the pathology of “pseudo-stabilization” of its prolonged group orbits, cf. [36]. We therefore introduce an additional invariant one-form du_x , whose horizontal component is

$$\theta_4 = u_{xx} dx = K\theta_2.$$

The resulting dependency leads to the lifted invariant $K = \alpha u_{xx}$ which yields the desired normalization $\alpha = 1/u_{xx}$ and the second order moving frame. The associated invariant coframe is

$$\theta_2 = \theta_4 = u_{xx} dx, \quad \theta_1 = -J\theta_2, \quad \theta_3 = I\theta_2, \quad (5.15)$$

yielding two fundamental differential invariants

$$I = u_x, \quad J = u_{xx}^{-2} u_{xxx}. \quad (5.16)$$

The higher order invariants are found by differentiating J with respect to $\theta_4 = u_{xx} dx \simeq du_x$, so that a basic fourth order invariant is

$$K = \frac{dJ}{du_x} = \frac{1}{u_{xx}} \frac{dJ}{dx} = \frac{u_{xxxx}}{u_{xx}^2} - 2J^2.$$

Note that $dI/du_x = 1$, so that differentiating I produces nothing new. Thus, in this case, we find two fundamental differential invariants, and require three, namely (I, J, K) , to parametrize the classifying curve that solves the associated equivalence problem. We conclude that the phenomenon of pseudo-stabilization of group orbits is reflected in the moving coframe procedure by the premature appearance of differential invariants, whose differentials are required to finish the procedure. See [36], [37], for further discussion.

Remark: Interestingly, if the scaling acts differently on x and u , so the group is

$$A : (x, u) \longmapsto (\alpha x + a, \alpha^k u + b), \quad (5.17)$$

for $k \neq 1$, then pseudo-stabilization does not occur. Such cases can be readily handled via our basic method without any such intransitive normalizations.

6. Reparametrization Pseudo-Groups.

The classical differential geometric applications of moving frames to curves and surfaces in Euclidean, affine, and projective geometry, cf. [6], [8], [17], can all be readily implemented using the moving coframe algorithm. In each case, we consider the reparametrization equivalence problem for submanifolds, so that the underlying transformation group is the Cartesian product of an infinite Lie pseudo-group, namely the local diffeomorphism group $\mathcal{D}(X)$ of the parameter space, and a finite-dimensional Lie group acting on the manifold M . In this case, one can proceed by including the one-forms defining the diffeomorphism pseudo-group in addition to the Maurer–Cartan forms corresponding to the group action as in the original formulation. Once the problem has been formulated, one can proceed to reduce and normalize as before. For simplicity, we just deal with planar curves, although extensions to surfaces and curves in higher dimensional ambient spaces can also be handled.

Example 6.1. *Euclidean geometry of curves.* The most well-known classical example is the reparametrization equivalence problem for curves in the Euclidean plane, introduced in Example 2.1 above. In this case, we are dealing with a finite-dimensional group, the Euclidean group $E(2)$ on the plane, together with the pseudo-group $\mathcal{D}(1)$ consisting of all smooth (local) diffeomorphisms of the line representing the change of parameter. Thus, the entire pseudo-group $\mathcal{G} = \mathcal{D}(1) \times E(2)$ acts on the total space $M = \mathbb{R} \times \mathbb{R}^2$ with coordinates $(t, \mathbf{x}) = (t, x, y)$. For the Euclidean component, we use a compatible lift

$$A_0(x, y, \alpha) = \begin{pmatrix} R & \mathbf{x} \\ \mathbf{0} & 1 \end{pmatrix} = \begin{pmatrix} \cos \alpha & -\sin \alpha & x \\ \sin \alpha & \cos \alpha & y \\ 0 & 0 & 1 \end{pmatrix}, \quad (6.1)$$

and compute the pull-back of the associated Euclidean Maurer–Cartan forms:

$$\zeta = A_0^{-1} dA_0 = \begin{pmatrix} R^{-1} dR & R^{-1} d\mathbf{x} \\ \mathbf{0} & 0 \end{pmatrix} = \begin{pmatrix} 0 & -d\alpha & \cos \alpha dx + \sin \alpha dy \\ d\alpha & 0 & -\sin \alpha dx + \cos \alpha dy \\ 0 & 0 & 0 \end{pmatrix}. \quad (6.2)$$

On the other hand, the pseudo-group $\mathcal{D}(1)$ is characterized by the invariance of the canonical one-form σdt on the frame bundle $\mathcal{F}(\mathbb{R})$, cf. [25], and hence we include this additional one-form in our moving coframe formulation. Restricting these four one-forms to a parametrized curve $(x(t), y(t))$ leads to

$$\theta_1 = d\alpha, \quad \theta_2 = (x_t \cos \alpha + y_t \sin \alpha) dt, \quad \theta_3 = (-x_t \sin \alpha + y_t \cos \alpha) dt, \quad \theta_4 = \sigma dt. \quad (6.3)$$

Now $\theta_2 = J_1 \theta_4$ and $\theta_3 = J_2 \theta_4$ are the linear dependencies, with associated lifted invariants

$$J_1 = \frac{x_t \cos \alpha + y_t \sin \alpha}{\sigma}, \quad J_2 = \frac{-x_t \sin \alpha + y_t \cos \alpha}{\sigma}.$$

We normalize $J_1 = 1, J_2 = 0$ by setting

$$\alpha = \tan^{-1}(y_t/x_t), \quad \sigma = \sqrt{x_t^2 + y_t^2}. \quad (6.4)$$

This immediately produces the first order moving frame

$$R = \frac{1}{\sqrt{x_t^2 + y_t^2}} \begin{pmatrix} x_t & -y_t \\ y_t & x_t \end{pmatrix}, \quad \mathbf{a} = \begin{pmatrix} x \\ y \end{pmatrix}, \quad \sigma = \sqrt{x_t^2 + y_t^2}. \quad (6.5)$$

The canonical one-form σdt has been reduced to the fundamental arc length form $ds = \sqrt{x_t^2 + y_t^2} dt$ for the Euclidean group. Substituting into (6.3), we are left with a final set of one-forms on the parameter space

$$\theta_1 = \kappa \theta_4, \quad \theta_2 = \theta_4 = ds = \sqrt{x_t^2 + y_t^2} dt, \quad \theta_3 = 0. \quad (6.6)$$

Here

$$\kappa = \frac{x_t y_{tt} - x_{tt} y_t}{(x_t^2 + y_t^2)^{3/2}} = \frac{\mathbf{x}_t \wedge \mathbf{x}_{tt}}{|\mathbf{x}_t|^3} = \mathbf{x}_s \wedge \mathbf{x}_{ss} \quad (6.7)$$

is the fundamental differential invariant for the Euclidean group — the curvature of the plane curve. All higher order differential invariants are obtained by successively differentiating the curvature with respect to arc length.

The classical Frenet equations for curves in the Euclidean plane are reformulations of our final moving frame formulae. The rotational component in (6.5) is traditionally written as $R = (\mathbf{e}_1, \mathbf{e}_2)$, where \mathbf{e}_1 is the unit tangent and \mathbf{e}_2 the unit normal. (See Section 7 below for details on the connection with the classical theory.) The translation Maurer–Cartan forms $\theta_2 = ds$, $\theta_3 = 0$ are computed by the original formula as the entries of $R^{-1} d\mathbf{x} = \begin{pmatrix} ds \\ 0 \end{pmatrix}$, which reduces to the first Frenet equation $\frac{d\mathbf{x}}{ds} = \mathbf{e}_1$. Similarly, the Maurer–Cartan matrix

$$R^{-1} dR = \begin{pmatrix} 0 & -\kappa \\ \kappa & 0 \end{pmatrix} ds \quad \text{implies that} \quad \frac{dR}{ds} = R \begin{pmatrix} 0 & -\kappa \\ \kappa & 0 \end{pmatrix}.$$

The columns of the latter matrix differential equation complete the system of Frenet equations:

$$\frac{d\mathbf{x}}{ds} = \mathbf{e}_1, \quad \frac{d\mathbf{e}_1}{ds} = \kappa \mathbf{e}_2, \quad \frac{d\mathbf{e}_2}{ds} = -\kappa \mathbf{e}_1. \quad (6.8)$$

Finally, the Maurer–Cartan structure equations (3.2) for the Euclidean group reduce to the classical Frenet–Serret equations for curves. See [12; p. 23], [17; p. 20], for details.

Remark: The reader can check that, if we restrict to curves which are the graphs $y = u(x)$ of functions, then the moving frame and moving coframe immediately reduce to those obtained by the preceding method.

Remark: One can also compute, as in our original example, the full moving coframe forms on the jet bundle, leading to a corresponding set of fundamental Euclidean-invariant contact forms.

Remark: Actually, since we are dealing with the full pseudo-group $\mathcal{D}(1)$ consisting of *all* diffeomorphisms of \mathbb{R} , the final one-form $\theta_4 = \sigma dt$ in our moving coframe (6.3) is, in fact, irrelevant — one could perform the same normalization (6.4) of α based on the dependency between θ_2 and θ_3 , the lifted invariant now being J_2/J_1 which is normalized to zero by setting $J_2 = 0$, leading to the same final moving coframe and curvature invariant. Thus the calculations for parametrized curves and surfaces can, in fact, be done without invoking the diffeomorphism pseudo-group. Nevertheless, in all examples we have treated, the inclusion of the canonical one-form θ_4 on the parameter space leads to an immediate identification of the final invariant arc length element. Moreover, the restricted reparametrization equivalence problem does require the introduction of suitable one-forms that characterize the pseudo-group of reparametrizations.

Remark: The problem of Euclidean equivalence of curves with *fixed* parametrizations, as discussed in Example 2.1, can also be formulated and solve in the moving frame context. Now we are in the intransitive framework, where the parameter t provides a scalar invariant. Consequently, we retain the first three one-forms $\theta_1, \theta_2, \theta_3$ in (6.3), but replace θ_4 by dt to form the moving coframe. We normalize $J_1 = 0$ as before, but now $J_2 = v = \sqrt{x_t^2 + y_t^2}$ forms a first order differential invariant — the speed of the particle. The final moving frame has $\theta_1 = K dt$, $\theta_2 = 0$, $\theta_3 = v dt$, $\theta_4 = dt$, where $K = x_t y_{tt} - x_{tt} y_t = v^3 \kappa$ is a second order differential invariant. The higher order differential invariants are found by differentiating with respect to t . Note that the arc length $ds = v dt$ is also an invariant one-form, being an invariant multiple of dt , and hence one can, without loss of generality, apply the arc length derivative d/ds to produce the higher order differential invariants instead. Thus, in this case, a complete list of differential invariants is provided by v , κ , and their derivatives with respect to arc length.

Example 6.2. The equiaffine geometry of curves in the plane is governed by the special affine group $\text{SA}(2) = \text{SL}(2) \times \mathbb{R}^2$, acting on $M = \mathbb{R}^2$ according to

$$g : \mathbf{x} \mapsto A\mathbf{x} + \mathbf{a}, \quad \mathbf{x} \in M, \quad A \in \text{SL}(2), \quad \mathbf{a} \in \mathbb{R}^2. \quad (6.9)$$

We shall adopt a vector notation for the matrix $A = (\boldsymbol{\alpha}\boldsymbol{\beta}) \in \text{SL}(2)$, so that the column vectors are subject to the unimodularity constraint

$$\boldsymbol{\alpha} \wedge \boldsymbol{\beta} = 1. \quad (6.10)$$

It will be computationally convenient *not* to explicitly implement the unimodularity constraint (6.10) by solving for one of the parameters, but retain it as an additional constraint that is to be respected during the course of the calculation. This method, i.e., treating a subgroup of a larger Lie group via a collection of algebraic constraints, rather than parametrizing it directly, has general applicability, and can be readily implemented as is done in this particular case.

The Maurer–Cartan forms are computed directly as in Section 3, leading to

$$\mu_1 = \boldsymbol{\alpha} \wedge d\boldsymbol{\alpha}, \quad \mu_2 = \boldsymbol{\beta} \wedge d\boldsymbol{\alpha}, \quad \mu_3 = \boldsymbol{\beta} \wedge d\boldsymbol{\beta}, \quad \nu_1 = \boldsymbol{\alpha} \wedge d\mathbf{a}, \quad \nu_2 = \boldsymbol{\beta} \wedge d\mathbf{a}. \quad (6.11)$$

Note that the unimodularity constraint (6.10) implies that

$$\boldsymbol{\alpha} \wedge d\boldsymbol{\beta} = \boldsymbol{\beta} \wedge d\boldsymbol{\alpha}, \quad (6.12)$$

which means that the matrix of Maurer–Cartan forms $\boldsymbol{\mu} = A^{-1} dA$ must be trace free.

Choose the base point to be $\mathbf{x}_0 = 0$. Solving the compatible lift equations $\mathbf{x} = g \cdot \mathbf{x}_0 = \mathbf{a}$ yields the zeroth order moving frame, which sets $\mathbf{a} = \mathbf{x}$. Substituting into the Maurer–Cartan forms (6.11), we find that, for a parametrized curve $\mathbf{x}(t)$, the forms ν_1, ν_2 restrict to the following two horizontal forms:

$$\theta_1 = (\boldsymbol{\alpha} \wedge \mathbf{x}_t) dt, \quad \theta_2 = (\boldsymbol{\beta} \wedge \mathbf{x}_t) dt. \quad (6.13)$$

Their ratio produces the lifted invariant $(\boldsymbol{\alpha} \wedge \mathbf{x}_t)/(\boldsymbol{\beta} \wedge \mathbf{x}_t)$, which is normalized to 0 by setting

$$\boldsymbol{\alpha} = \lambda \mathbf{x}_t, \quad (6.14)$$

for some scalar parameter λ . Substituting (6.14) into the first Maurer–Cartan form $\mu_1 = \boldsymbol{\alpha} \wedge d\boldsymbol{\alpha}$, leads to the restricted form $\lambda^2(\mathbf{x}_t \wedge \mathbf{x}_{tt}) dt$. Assuming $\mathbf{x}_t \wedge \mathbf{x}_{tt} \neq 0$, the latter form can be normalized to equal $-\theta_2$ by setting

$$-\boldsymbol{\beta} \wedge \mathbf{x}_t = \lambda^2(\mathbf{x}_t \wedge \mathbf{x}_{tt}), \quad \text{or} \quad \boldsymbol{\beta} = \lambda^2 \mathbf{x}_{tt} + \mu \mathbf{x}_t, \quad (6.15)$$

for some scalar μ . However, applying the unimodularity constraint (6.10) to the normalizations (6.14), (6.15), we deduce that $\lambda^3(\mathbf{x}_t \wedge \mathbf{x}_{tt}) = 1$, and thus

$$\lambda = \frac{1}{\sqrt[3]{\mathbf{x}_t \wedge \mathbf{x}_{tt}}}. \quad (6.16)$$

Note that (6.15), (6.16) reduce the form θ_2 to be minus the equi-affine arc length form

$$ds = \sqrt[3]{\mathbf{x}_t \wedge \mathbf{x}_{tt}} dt. \quad (6.17)$$

Furthermore, substituting (6.15), (6.16) into the second Maurer–Cartan form, we find

$$\mu_2 = \boldsymbol{\beta} \wedge d\boldsymbol{\alpha} = J ds,$$

where the lifted invariant

$$J = \mu(\mathbf{x}_t \wedge \mathbf{x}_{tt})^{1/3} + \frac{\mathbf{x}_t \wedge \mathbf{x}_{ttt}}{3(\mathbf{x}_t \wedge \mathbf{x}_{tt})^{4/3}}$$

is normalized to zero in the obvious manner. Therefore, the final moving frame is given by

$$\boldsymbol{\alpha} = \frac{d\mathbf{x}}{ds} = \frac{\mathbf{x}_t}{\sqrt[3]{\mathbf{x}_t \wedge \mathbf{x}_{tt}}}, \quad \boldsymbol{\beta} = \frac{d^2\mathbf{x}}{ds^2} = \frac{\mathbf{x}_{tt}}{(\mathbf{x}_t \wedge \mathbf{x}_{tt})^{2/3}} - \frac{\mathbf{x}_t}{3(\mathbf{x}_t \wedge \mathbf{x}_{tt})^{5/3}}, \quad \mathbf{a} = \mathbf{x}. \quad (6.18)$$

The final Maurer–Cartan form becomes

$$\mu_3 = \beta \wedge d\beta = \kappa ds,$$

where

$$\kappa = \mathbf{x}_{ss} \wedge \mathbf{x}_{sss} = \frac{(\mathbf{x}_t \wedge \mathbf{x}_{ttt}) + 4(\mathbf{x}_{tt} \wedge \mathbf{x}_{ttt})}{3(\mathbf{x}_t \wedge \mathbf{x}_{tt})^{5/3}} - \frac{5(\mathbf{x}_t \wedge \mathbf{x}_{ttt})^2}{9(\mathbf{x}_t \wedge \mathbf{x}_{tt})^{8/3}} \quad (6.19)$$

defines the equi-affine curvature. As usual, all higher order differential invariants are obtained by differentiating κ with respect to the equi-affine arc length ds . This reproduces the basic invariants of the equi-affine geometry of curves, [17]; see also [5] for applications in computer vision.

As with the Euclidean case, we recover the classical Frenet equations as simple reformulations of the final moving frame formulae. We identify the linear part

$$A = (\mathbf{e}_1, \mathbf{e}_2) = (\mathbf{x}_s, \mathbf{x}_{ss})$$

of the final moving frame with the equi-affine frame at a point $\mathbf{x}(t)$ on the curve, so that $\mathbf{e}_1 = \mathbf{x}_s$ is the unit affine tangent vector, whereas $\mathbf{e}_2 = \mathbf{x}_{ss}$ is the unit equi-affine normal. Combining this with the Maurer–Cartan matrix $A^{-1} dA = \begin{pmatrix} 0 & 1 \\ \kappa & 0 \end{pmatrix} ds$ leads to the complete Frenet equations of planar equi-affine geometry, [12; p. 27]:

$$\frac{d\mathbf{x}}{ds} = \mathbf{e}_1, \quad \frac{d\mathbf{e}_1}{ds} = \mathbf{e}_2, \quad \frac{d\mathbf{e}_2}{ds} = \kappa \mathbf{e}_1. \quad (6.20)$$

See [17; § 7–3] for further details.

Example 6.3. The most complicated example treated in the literature, [7], is the projective geometry of curves in the plane. Here the group is $\mathrm{SL}(3)$, acting on $M = \mathbb{RP}^2$ according to

$$g: (x, u) \mapsto \left(\frac{\alpha x + \beta u + \gamma}{\rho x + \sigma u + \tau}, \frac{\lambda x + \mu u + \nu}{\rho x + \sigma u + \tau} \right), \quad \det A = \det \begin{vmatrix} \alpha & \beta & \gamma \\ \lambda & \mu & \nu \\ \rho & \sigma & \tau \end{vmatrix} = 1. \quad (6.21)$$

For simplicity, we deal with curves which can be expressed as the graphs of functions, $u = u(x)$, although the general case of parametrized curves can be handled via the same sequence of normalizations. Choose the base point to be $z_0 = (0, 0)$. Solving $g \cdot (0, 0) = (x, u)$ leads to the order zero moving frame in the form[†]

$$A = \begin{pmatrix} \alpha & \beta & \tau x \\ \lambda & \mu & \tau u \\ \rho & \sigma & \tau \end{pmatrix}, \quad \text{where} \quad \alpha = \frac{1 + \tau[\beta(\lambda - \rho u) + x(\mu\rho - \lambda\sigma)]}{\tau(\mu - \sigma x)}. \quad (6.22)$$

[†] In this example, we have chosen to implement the unimodularity constraint explicitly.

The one-forms in the first order moving coframe are the entries of the pull-back of the Maurer–Cartan matrix $A^{-1} dA$, which we label (in row order) as $\theta_1, \dots, \theta_8$, the final entry being $\theta_9 = -\theta_1 - \theta_5$, reflecting the unimodularity of A . For simplicity, we just indicate the salient features of the computation without dwelling on the details. (These computations were done with the aid of some MATHEMATICA routines written for this purpose.) The first normalization comes from the ratio θ_6/θ_3 , whose vanishing requires

$$\mu = \sigma(u - xu_x) - \beta u_x.$$

Plugging this normalization back into the moving coframe forms and recomputing, we find that we can normalize $\theta_3 = \theta_2$ by requiring

$$\beta = \sigma x - u_{xx}^{-1/3}.$$

In the next stage, we set θ_5 to zero by normalizing

$$\sigma = \frac{\tau(\rho u - \lambda)u_{xx}^{1/3}}{u_x} - \frac{u_x u_{xxx} - 3u_{xx}^2}{3u_x u_{xx}^{4/3}}.$$

At the next step, we can no longer just look at one-forms depending only on dx — these do not produce any further invariants. However, we discover that $\theta_8 = J\theta_2 + \theta_4$, and hence the rather complicated invariant J can be normalized to zero, leading to

$$\lambda = \rho u - \frac{u_x u_{xxx} - 3u_{xx}^2 + \sqrt{18\rho\tau u_{xx}^{8/3} - P_4}}{3\tau u_{xx}^{5/3}},$$

where

$$P_4 = 3u_{xx}u_{xxxx} - 5u_{xxx}^2.$$

Next, the normalization $\theta_2 = -\theta_7$ requires

$$\tau = \frac{\sqrt[3]{L_5}}{54u_{xx}^4}, \quad \text{where} \quad L_5 = 9u_{xx}^2 u_{xxxxx} - 45u_{xx}u_{xxx}u_{xxxx} + 40u_{xxx}^3.$$

The final normalization

$$\rho = \frac{M_6^2 + P_4 L_5^2}{3\sqrt[3]{4}u_{xx}^{4/3}L_5^{7/3}}, \quad \text{where} \quad M_6 = (u_{xx}D_x - 4u_{xxx})L_5$$

comes from setting θ_1 to zero. The final moving frame is explicitly given by

$$\begin{aligned} \alpha &= \frac{\lambda + \rho(xu_x - u)}{u_x} - \frac{3}{u_x} \sqrt[3]{\frac{2u_{xx}^5}{L_5}}, & \rho &= \frac{M_6^2 + P_4 L_5^2}{3\sqrt[3]{4}u_{xx}^4 L_5^7}, \\ \beta &= x\mu - u_{xx}^{-1/3}, & \sigma &= \frac{M_6}{3u_{xx}^{4/3}L_5}, \\ \lambda &= \frac{uM_6^2 + 6u_x u_{xx} L_5 M_6 + K_4 L_5^2}{3\sqrt[3]{4}u_{xx}^4 L_5^7}, & \tau &= \frac{L_5^{1/3}}{3\sqrt[3]{2}u_{xx}^4}, \\ \mu &= u\sigma - \frac{u_x}{u_{xx}^{1/3}}, \end{aligned} \tag{6.23}$$

The corresponding final coframe has

$$\theta_2 = \theta_6 = -\theta_7 = ds = \frac{\sqrt[3]{L_5}}{3\sqrt[3]{2}u_{xx}} dx = \sqrt[3]{\frac{9u_{xx}^2 u_{xxxxxx} - 45u_{xx}u_{xxx}u_{xxxx} + 40u_{xxx}^3}{54u_{xx}^3}} dx, \quad (6.24)$$

which determines the well-known projective arc length element, while $\theta_4 = \theta_8 = -\kappa\theta_3$ yields the projective curvature invariant

$$\kappa = \frac{6u_{xx}D_x M_6 - 7M_6^2 - 32u_{xxx}L_5 M_6 + P_4 L_5^2}{\sqrt[3]{2}L_5^{8/3}}. \quad (6.25)$$

Again, all higher order differential invariants are found by differentiating the projective curvature κ with respect to the projective arc length ds . This relatively straightforward computation reproduces the moving frame and the fundamental invariants for the projective geometry of curves. See Cartan, [7], for a variety of alternative methods to arrive at the same basic result, and [36] for a Lie-theoretic approach to the projective differential invariants.

In the classical moving frame method, one identifies the columns of the 3×3 moving frame matrix $A = (\mathbf{P}_2, \mathbf{P}_1, \mathbf{P})$ as homogeneous coordinates for three points in the projective plane $\mathbb{R}P^2$, the last column $\mathbf{P} = \tau(x, u, 1)$ representing the point on the curve. The Maurer–Cartan matrix

$$A^{-1} dA = \begin{pmatrix} 0 & 1 & 0 \\ -\kappa & 0 & 1 \\ -1 & -\kappa & 0 \end{pmatrix} ds$$

reduces to the full set of projective Frenet equations,

$$\frac{d\mathbf{P}}{ds} = \mathbf{P}_1, \quad \frac{d\mathbf{P}_1}{ds} = -\kappa\mathbf{P} + \mathbf{P}_2, \quad \frac{d\mathbf{P}_2}{ds} = -\mathbf{P} - \kappa\mathbf{P}_1. \quad (6.26)$$

See also [12; p. 33 ff.] for applications to projective curvature evolutions and computer vision.

7. Connections with the Classical Moving Frames Method.

Although our initial identification of a moving frame with a lift from the underlying space to the Lie group appears in several other places, e.g., [16], [21], it is not completely standard, so it is worth reviewing how it relates to the more usual geometric approaches, e.g., [17]. Typically, the moving frame is realized as a collection of vectors (or, in the projective case, points) in the underlying space. The reason that this works in the classical cases, including Euclidean, affine, and projective geometry of submanifolds, is that it is possible to identify the components of the group itself with objects in the underlying transformation space. For example, in the Euclidean case, one identifies a Euclidean group element $(R, \mathbf{a}) \in E(m) \simeq O(m) \times \mathbb{R}^m$ with a vector $\mathbf{a} \in \mathbb{R}^m$, together with an orthonormal frame determined by the columns of the orthogonal matrix R . The zeroth order moving frame, then, uses the lift $\mathbf{a} = \mathbf{x}$, where \mathbf{x} is a point on the submanifold $N \subset \mathbb{R}^m$, and

the orthogonal matrix is identified with an orthonormal frame in the ambient space based at the point. The remaining ambiguity in the frame is up to orthogonal transformations, which must then be resolved in an invariant manner. Similarly, in the equi-affine case, one identifies a group element $(A, \mathbf{a}) \in \text{SA}(m) \simeq \text{SL}(m) \times \mathbb{R}^m$ with a vector $\mathbf{a} \in \mathbb{R}^m$ together with a unimodular frame determined by the columns of the matrix A . Again, the zeroth order moving frame takes $\mathbf{a} = \mathbf{x}$ to be a point on the submanifold, and the unimodular frame becomes a set of vectors based at the point \mathbf{x} . In both cases, the moving coframe method introduces the Maurer–Cartan forms $\boldsymbol{\mu} = (\boldsymbol{\xi}, \boldsymbol{\eta})$, where $\boldsymbol{\xi} = A^{-1} dA$, $\boldsymbol{\eta} = A^{-1} d\mathbf{a}$, leading to the initial structure equations

$$d\mathbf{x} = A \cdot \boldsymbol{\eta}, \quad dA = A \cdot \boldsymbol{\xi}. \quad (7.1)$$

The moving coframe forms satisfy the usual Maurer–Cartan structure equations (3.2), which, in the classical cases, become the fundamental Cartan structure equations for Euclidean or affine geometry.

Usually, one bypasses the order zero moving frame entirely, and proceeds directly to the first order moving frame, in which the frame at the point $\mathbf{x} \in N$ is split into two parts, so that (using column vector notation)

$$A = (E, F) = (\mathbf{e}_1, \dots, \mathbf{e}_n, \mathbf{f}_1, \dots, \mathbf{f}_{m-n}), \quad (7.2)$$

where the first $n = \dim N$ frame vectors form a basis for the tangent space TN to the submanifold, while the remainder are left arbitrary, subject to the entire frame satisfying the proper orthonormality or unimodularity condition. Thus, in the Euclidean case, the vectors $\{\mathbf{f}_1, \dots, \mathbf{f}_{m-n}\}$ form an orthonormal basis for the normal space to N , whereas in the equi-affine case they are left arbitrary subject only to the condition that the determinant of the matrix (7.2) be unity. If we parametrize the submanifold by $\mathbf{x}(t^1, \dots, t^n)$, then the most general first order moving frame (7.2) will have the form

$$E = (\mathbf{e}_1, \dots, \mathbf{e}_n) = V \cdot B, \quad (7.3)$$

where

$$V = (\mathbf{v}_1, \dots, \mathbf{v}_n), \quad \mathbf{v}_i = \frac{\partial \mathbf{x}}{\partial t^i}, \quad (7.4)$$

is the $m \times n$ Jacobian matrix, whose columns span the tangent space to N , while B is an invertible $n \times n$ matrix. (In the Euclidean case, the matrix B is restricted so that the columns of E are orthonormal, leaving an $O(n)$ ambiguity.)

Let us show how this preliminary normalization to a first order moving frame is an immediate consequence of our general normalization procedure. Using the zeroth order moving frame lift, the pull-backs of the subset of Maurer–Cartan forms given by the entries of $\boldsymbol{\eta} = A^{-1} d\mathbf{a}$ can be written in matrix form as

$$\boldsymbol{\theta} = A^{-1} d\mathbf{x} = A^{-1} V dt.$$

Precisely n of the m one-forms θ are linearly independent, and hence we can normalize so that the last $m - n$ of these forms vanish. This requires that the matrix A satisfy the block matrix equation

$$A^{-1}V = \begin{pmatrix} D \\ 0 \end{pmatrix}, \quad (7.5)$$

where D is a nonsingular $n \times n$ matrix, while 0 denotes the zero matrix of size $(m - n) \times n$. Writing $A = (E, F)$ in block form (7.2), we see that (7.5) requires

$$V = E \cdot D, \quad \text{or} \quad E = V \cdot C, \quad \text{where} \quad C = D^{-1},$$

thereby recovering (7.3). Thus, we can see that in such cases, the first order frame recovered by an order zero normalization coincides with the traditional first order frame involving tangent and normal directions.

Similar considerations apply to the projective case. According to Cartan, [7], the zeroth order frame can be identified with a set of $n + 1$ linearly independent points in the projective space which are identified with the columns of the matrix $A \in \text{SL}(n + 1)$. The zeroth order lift, as in (6.22), amounts to identifying one of the columns with the point on the curve \mathbf{x} . More precisely, the column is a vector with $n + 1$ components, which are interpreted as the homogeneous coordinates of \mathbf{x} .

In more sophisticated versions, one realizes the moving frame on the submanifold $N \subset M$ as a section of the frame bundle $\mathcal{F}(M)$ over N , i.e., a section $\psi: N \rightarrow \mathcal{F}(M)$. One can continue in this manner, and try to realize cases that do not so readily fit into this simple framework by realizing them as sections of a suitable higher order frame bundle $\mathcal{F}^k(M)$ over N , cf. [25]. Although this is possible to realize for all (regular, transitive) transformation groups, the original geometrical realization has now been rather well obscured, and such a reinterpretation does not, we think, offer much insight or help in the explicit implementation of the method.

Consequently, the method of moving coframes includes all the classical constructions based on the indicated identification of group elements with geometric objects on the transformation space. However, once one steps beyond the traditional cases, such identifications become much less apparent, and, in our opinion, attempting to mimic the Euclidean, affine, and projective constructions directly on the transformation space has hindered the development of any significant extensions of the method. Furthermore, once one steps outside the realm of “classical” moving frame geometries, one can no longer use the identification of the first order frame with tangent and normal directions. Our non-traditional examples all illustrate this — the first order frames do not include the tangent spaces to the submanifolds in any obvious manner, because their naïve identification with subspaces of Euclidean space is not necessarily invariant with respect to the given transformation group. It is our view that, in order to attain their full range of applicability, the constructions must be viewed in the purely group- or, more generally, bundle-theoretic framework that we have presented here and develop in detail in part II.

8. Joint Differential Invariants.

New applications in computer vision, [33], have demonstrated the need for classification and computation of the joint differential invariants or, as they are known in computer

vision, *semi-differential invariants*, for a given transformation group. Specifically, one is given a Lie group (or pseudo-group) G acting on M and considers its diagonal action $g \cdot (z^1, \dots, z^k) = (g \cdot z^1, \dots, g \cdot z^k)$ on the k -fold Cartesian product $M^{\times k} = M \times \dots \times M$. The invariants $I(z^1, \dots, z^k)$ of such a Cartesian product action are known as the *k -point joint invariants* of the transformation group. Note that for $j < k$, any j -point invariant can be regarded as a k -point invariant, in several different ways. For example, the two-point invariant $I(z_1, z_2)$ produces three invariants on $M^{\times 3}$, namely $\hat{I}(z_1, z_2, z_3) = I(z_1, z_2)$, or $I(z_1, z_3)$ or $I(z_2, z_3)$. If I is not symmetric in its arguments, these in turn lead to 3 further invariants by interchanging the points. To avoid this trivial extension, we will reserve the term *k point invariant* for a joint invariant which cannot be written as as one depending on fewer than k arguments.

Similarly, the invariants of the prolonged diagonal action of $G^{(n)}$ on a k -fold Cartesian product of jet space $(J^n)^{\times k}$ are the *joint differential invariants* of k different submanifolds $N_1, \dots, N_k \subset M$, which we view as a single submanifold $N_1 \times \dots \times N_k$ of the Cartesian product space $M^{\times k}$. In applications, the submanifolds $N_j = N$ are identical, but the joint differential invariants are measured at k different points along the given submanifold.

The method of moving coframes readily adapts to this slightly more general situation, and immediately provides complete classifications of joint differential invariants for all of the standard geometric transformation groups. Further examples and applications will be discussed in a subsequent paper.

Example 8.1. *Euclidean joint differential invariants.* Consider the Euclidean group $E(2)$ acting on the plane $M = \mathbb{R}^2$. We consider two-point differential invariants, corresponding to the Cartesian product action

$$(\mathbf{x}, \mathbf{y}) \longmapsto (R \cdot \mathbf{x} + \mathbf{a}, R \cdot \mathbf{y} + \mathbf{a}), \quad \mathbf{x}, \mathbf{y} \in M, \quad (R, \mathbf{a}) \in E(2), \quad (8.1)$$

on $M^{\times 2} \simeq \mathbb{R}^4$. Note that the action is intransitive on $M^{\times 2}$, with the interpoint distance

$$r = |\mathbf{z}|, \quad \text{where} \quad \mathbf{z} = \mathbf{x} - \mathbf{y}, \quad (8.2)$$

is the fundamental joint Euclidean invariant. (See [46] for a proof that all Euclidean joint invariants can be written in terms of the elementary two-point invariants.) We can choose the cross section to the orbits given by $\mathbf{x}_0 = 0$, $\mathbf{y}_0 = (r, 0)$, which leads to the compatible lift with

$$\mathbf{a} = \mathbf{x}, \quad (r \cos \theta, r \sin \theta) = \mathbf{z} = \mathbf{x} - \mathbf{y}. \quad (8.3)$$

Therefore, all the group parameters are normalized by the initial compatible lift, and it only remains to substitute (8.3) into the Euclidean Maurer–Cartan forms (3.5). The net result is the following system of invariant forms:

$$\zeta_1 = \mathbf{z} \cdot d\mathbf{x}, \quad \zeta_2 = \mathbf{z} \cdot d\mathbf{y}, \quad \zeta_3 = r^2 d\theta = \mathbf{z} \wedge d\mathbf{z}, \quad (8.4)$$

where $\mathbf{a} \cdot \mathbf{b}$ and $\mathbf{a} \wedge \mathbf{b}$ are the standard dot and cross products between vectors in the plane. Note that the forms (8.4) include the differential of the joint invariant (8.2) since $r dr = \zeta_1 + \zeta_2$. Therefore, given two parametrized curves

$$\mathbf{x} = \mathbf{x}(t), \quad \mathbf{y} = \mathbf{y}(s), \quad (8.5)$$

the first two one-forms (8.4) restrict to define two invariant one-forms

$$\theta_1 = (\mathbf{z} \cdot \mathbf{x}_t) dt, \quad \theta_2 = (\mathbf{z} \cdot \mathbf{y}_s) ds, \quad (8.6)$$

while $\theta_3 = I_1 \theta_1 + I_2 \theta_2$, where

$$I_1 = \frac{\mathbf{z} \wedge \mathbf{x}_t}{\mathbf{z} \cdot \mathbf{x}_t}, \quad I_2 = \frac{\mathbf{z} \wedge \mathbf{y}_s}{\mathbf{z} \cdot \mathbf{y}_s}, \quad (8.7)$$

are the two fundamental first order differential invariants, which, along with the original joint invariant (8.2), form a complete system of first order joint differential invariants. The identity

$$(\mathbf{a} \cdot \mathbf{b})^2 + (\mathbf{a} \wedge \mathbf{b})^2 = |\mathbf{a}|^2 |\mathbf{b}|^2, \quad (8.8)$$

demonstrates that

$$J_1^2 = \frac{|\mathbf{x}_t|^2}{(\mathbf{z} \cdot \mathbf{x}_t)^2} = 1 + \frac{(\mathbf{z} \wedge \mathbf{x}_t)^2}{(\mathbf{z} \cdot \mathbf{x}_t)^2}$$

is also a joint differential invariant, and hence (in the orientation-preserving case) one can replace the one-forms (8.6) by the two Euclidean arc-length forms

$$\eta_1 = J_1 \theta_1 = |\mathbf{x}_t| dt, \quad \eta_2 = J_2 \theta_2 = |\mathbf{y}_s| ds. \quad (8.9)$$

Theorem 8.2. *Every two-point Euclidean joint differential invariant is a function of the interpoint distance $r = |\mathbf{x} - \mathbf{y}|$ and its derivatives with respect to the two arc length forms (8.9).*

For example, to recover the Euclidean curvature $\kappa_1 = |\mathbf{x}_t|^{-3} (\mathbf{x}_t \wedge \mathbf{x}_{tt})$ of the first curve, we differentiate

$$\begin{aligned} \frac{\partial I_1}{\partial \theta_1} &= \frac{\mathbf{z} \wedge \mathbf{x}_{tt}}{(\mathbf{z} \cdot \mathbf{x}_t)^2} - \frac{(\mathbf{z} \wedge \mathbf{x}_t)[(\mathbf{z} \cdot \mathbf{x}_{tt}) + |\mathbf{x}_t|^2]}{(\mathbf{z} \cdot \mathbf{x}_t)^3} = \frac{(\mathbf{z} \wedge \mathbf{x}_{tt})(\mathbf{z} \cdot \mathbf{x}_t) - (\mathbf{z} \wedge \mathbf{x}_t)(\mathbf{z} \cdot \mathbf{x}_{tt})}{(\mathbf{z} \cdot \mathbf{x}_t)^3} - I_1 J_1^2 \\ &= \frac{(\mathbf{x}_t \wedge \mathbf{x}_{tt}) |\mathbf{z}|^2}{(\mathbf{z} \cdot \mathbf{x}_t)^3} - I_1 J_1^2 = \kappa_1 r^2 - I_1 J_1^2, \end{aligned}$$

where we have used the first of the following equivalent identities

$$\begin{aligned} (\mathbf{a} \wedge \mathbf{b})(\mathbf{c} \cdot \mathbf{d}) + (\mathbf{b} \wedge \mathbf{c})(\mathbf{a} \cdot \mathbf{d}) - (\mathbf{a} \wedge \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) &= 0, \\ (\mathbf{a} \wedge \mathbf{b})(\mathbf{c} \wedge \mathbf{d}) + (\mathbf{b} \wedge \mathbf{c})(\mathbf{a} \wedge \mathbf{d}) - (\mathbf{a} \wedge \mathbf{c})(\mathbf{b} \wedge \mathbf{d}) &= 0. \end{aligned} \quad (8.10)$$

Example 8.3. A more substantial example is provided by the two-point differential invariants for the special affine group $\text{SA}(2) = \text{SL}(2) \ltimes \mathbb{R}^2$, acting on $M = \mathbb{R}^2$. The Cartesian product action

$$(\mathbf{x}, \mathbf{y}) \longmapsto (A\mathbf{x} + \mathbf{a}, A\mathbf{y} + \mathbf{a}), \quad \mathbf{x}, \mathbf{y} \in M, \quad A \in \text{SL}(2), \quad \mathbf{a} \in \mathbb{R}^2, \quad (8.11)$$

is transitive on $M^{\times 2}$. As in Example 6.2, we use the vector notation $A = (\alpha\beta) \in \text{SL}(2)$, where $\alpha \wedge \beta = 1$.

In view of (8.11), we can choose the base point $\mathbf{x}_0 = 0$, $\mathbf{y}_0 = (1, 0)$, noting that the diagonal $\Delta = \{\mathbf{x} = \mathbf{y}\} \subset M^{\times 2}$ is a singular two-dimensional orbit. This leads to the compatible lift with

$$\mathbf{a} = \mathbf{x}, \quad \alpha = \mathbf{z} = \mathbf{x} - \mathbf{y}. \quad (8.12)$$

Substituting into the Maurer–Cartan forms (6.11), we find that, for a pair of parametrized curves as in (8.5), the following horizontal forms:

$$(\mathbf{z} \wedge \mathbf{x}_t) dt, \quad (\beta \wedge \mathbf{x}_t) dt, \quad (\mathbf{z} \wedge \mathbf{y}_s) ds, \quad (\beta \wedge \mathbf{y}_s) ds,$$

the first two being the pull-backs of ν_1, ν_2 , and the latter being that of $\nu_1 - \mu_1, \nu_2 - \mu_2$. Generically (i.e., provided $\mathbf{x} - \mathbf{y}$ is not parallel to the tangent \mathbf{x}_t) we can normalize the second form to zero, leading, in view of (8.12) and the unimodularity constraint, to

$$\beta = \frac{\mathbf{x}_t}{\mathbf{z} \wedge \mathbf{x}_t}, \quad (8.13)$$

which, combined with (8.12) provides the complete moving frame. The remaining one-forms are

$$\theta_1 = \mathbf{z} \wedge d\mathbf{x} = (\mathbf{z} \wedge \mathbf{x}_t) dt, \quad \theta_2 = \mathbf{z} \wedge d\mathbf{y} = (\mathbf{z} \wedge \mathbf{y}_s) ds, \quad (8.14)$$

which provide the fundamental invariant one-forms, and

$$\theta_3 = -\beta \wedge d\mathbf{y} = \left[\frac{\mathbf{x}_t \wedge \mathbf{y}_s}{\mathbf{z} \wedge \mathbf{x}_t} \right] ds, \quad \theta_4 = -\beta \wedge d\beta = \left[\frac{\mathbf{x}_t \wedge \mathbf{x}_{tt}}{(\mathbf{z} \wedge \mathbf{x}_t)^2} \right] dt.$$

The resulting linear dependencies provide the two basic differential invariants, consisting of a single first order invariant

$$I = \frac{\mathbf{x}_t \wedge \mathbf{y}_s}{(\mathbf{z} \wedge \mathbf{x}_t)(\mathbf{z} \wedge \mathbf{y}_s)}, \quad (8.15)$$

and the first of the two second order invariants

$$J_1 = \frac{\mathbf{x}_t \wedge \mathbf{x}_{tt}}{(\mathbf{z} \wedge \mathbf{x}_t)^3}, \quad J_2 = \frac{\mathbf{y}_s \wedge \mathbf{y}_{ss}}{(\mathbf{z} \wedge \mathbf{y}_s)^3}. \quad (8.16)$$

Clearly J_2 can be obtained from J_1 by the interchange symmetry $\mathbf{x} \leftrightarrow \mathbf{y}$. Alternatively, we use (8.10) to compute

$$\frac{\partial I}{\partial \theta_1} - I^2 = -\frac{(\mathbf{x}_{tt} \wedge \mathbf{y}_s)(\mathbf{z} \wedge \mathbf{x}_t) - (\mathbf{x}_t \wedge \mathbf{y}_s)(\mathbf{z} \wedge \mathbf{x}_{tt})}{(\mathbf{z} \wedge \mathbf{x}_t)^3(\mathbf{z} \wedge \mathbf{y}_s)} = \frac{(\mathbf{x}_{tt} \wedge \mathbf{x}_t)(\mathbf{z} \wedge \mathbf{y}_s)}{(\mathbf{z} \wedge \mathbf{x}_t)^3(\mathbf{z} \wedge \mathbf{y}_s)} = J_1,$$

so that (8.16) are equivalent to the invariant first order derivatives of the single basic joint invariant I .

Theorem 8.4. *Every two-point equi-affine joint differential invariant is a function of the fundamental first order invariant (8.15) and its derivatives with respect to the two “joint arc length” forms (8.14).*

The reader is invited to try to express the ordinary affine curvature in terms of the derivatives of I . the same general method readily extends to multi-point invariants of more general groups, including the projective group, as well as joint invariants for surfaces and higher dimensional submanifolds. Additional examples and applications will appear elsewhere.

9. Pseudo-Group Actions.

The next case is that of infinite Lie pseudo-groups, cf. [27], [29], [10], [40], [41]. See also [28], [43], for classical results on differential invariants of Lie pseudo-groups. These are readily fit into the same general framework as follows. Assume, initially, that the pseudo-group \mathcal{G} acts transitively on the space M . By definition, a Lie pseudo-group consists of an infinite-dimensional family of invertible (local) transformations that form the general solution to an involutive system of partial differential equations. We can always characterize the transformations $\psi: M \rightarrow M$ in \mathcal{G} as the projections of bundle maps $\Psi: B \rightarrow B$, defined on a principal fiber bundle $B \rightarrow M$, that preserve a system of one-forms $\zeta = \{\zeta_1, \dots, \zeta_k\}$ defined on B :

$$\Psi^* \zeta = \zeta. \quad (9.1)$$

The forms ζ will play the role of the moving coframe forms for the pseudo-group, and the fiber coordinates of the bundle B will play the role of the undetermined group parameters. Of course, in this case ζ does not form a full coframe on B . (It cannot, because the symmetry group of a coframe is necessarily a finite-dimensional Lie group, [36].) A compatible lift, or moving frame of order zero, is just an arbitrary section $\gamma_0: M \rightarrow B$. Such a section defines a corresponding moving frame $\gamma = \gamma_0 \circ \iota: X \rightarrow B$ on any parametrized submanifold $\iota: X \rightarrow M$. With these provisos, the normalization and reduction procedure proceeds as in the finite-dimensional situation.

Example 9.1. Consider the pseudo-group \mathcal{G} consisting of (local) diffeomorphisms on $M = \mathbb{R}^2$ of the form

$$\bar{x} = f(x), \quad \bar{u} = \frac{u}{f'(x)}. \quad (9.2)$$

The Lie algebra of \mathcal{G} is generated by vector fields of the form

$$\mathbf{v}_h = h(x)\partial_x - uh'(x)\partial_u.$$

This pseudo-group was first introduced by Lie, [27; p. 353], [31], in his classification of infinite-dimensional pseudo-groups acting on the plane. We are interested in the action of \mathcal{G} on curves.

The first step is to construct a bundle B and one-forms on the bundle whose invariance characterizes the pseudo-group transformations. In this case, away the axis $u = 0$, the group transformations (9.2) form the general solution to the defining system of partial differential equations

$$z_u = 0, \quad z_x = \frac{u}{w}, \quad w_u = \frac{w}{u}, \quad (9.3)$$

for $\bar{x} = z(x, u)$, $\bar{u} = w(x, u)$, cf. [44; p. 325]. The system (9.3) defines a submanifold $\Phi: \mathcal{R} \hookrightarrow J^1(\mathbb{R}^2, \mathbb{R}^2)$ of the first jet space, parametrized by the coordinates (x, u, z, w, w_x) . The pull-backs of the basic contact forms on $J^1(\mathbb{R}^2, \mathbb{R}^2)$ to the equation submanifold \mathcal{R} are given by

$$\begin{aligned} \theta_z &= \Phi^*(dz - z_x dx - z_u du) = dz - \frac{u}{w} dx, \\ \theta_w &= \Phi^*(dw - w_x dx - w_u du) = dw - w_x dx - \frac{w}{u} du. \end{aligned} \quad (9.4)$$

The Pfaffian system

$$\theta_z = 0, \quad \theta_w = 0,$$

with independence condition $dx \wedge du \neq 0$ is involutive on \mathcal{R} , cf. [9], [3], [36]. Indeed, the first Cartan character is $s_1 = 1$, as it should be. Following a general procedure[†] presented by Kamran, [23], we set $dz = dw = 0$, which amounts to pulling back to a level set of \mathcal{R} where $u = u_0$ and $w = w_0$ are constant. Choosing $w_0 = 1$ we find that the contact forms (9.4) reduce to the invariant one-forms

$$\eta_1 = -u dx, \quad \eta_2 = -w_x dx - \frac{du}{u}.$$

Therefore, the desired bundle $B \simeq M \times \mathbb{R}$ will be coordinatized by x, u , and the remaining jet coordinate, which we rewrite as $\alpha = w_x$ for clarity. In other words, the zeroth order moving coframe forms for the pseudo-group (9.2) will be

$$\zeta_1 = u dx, \quad \zeta_2 = \alpha dx + \frac{du}{u}. \quad (9.5)$$

Restricting to a curve $u = u(x)$, and letting θ_i denote the horizontal component of ζ_i , we have the relation

$$\theta_2 = (u\alpha + u_x) dx = (u\alpha + u_x)\theta_1,$$

and so we normalize $\alpha = -u_x/u$. Thus the final invariant moving coframe is

$$\zeta_1 = u dx, \quad \zeta_2 = \frac{du - u_x dx}{u}, \quad (9.6)$$

the first providing a pseudo-group invariant arc length form, and the latter an invariant contact form. Note that there are no dependencies among these one-forms, and hence there are *no* differential invariants in this example. Indeed, it is not hard to see that the prolonged actions of \mathcal{G} are transitive on every jet space $J^n M$, justifying the preceding statement.

Example 9.2. We now extend the pseudo-group discussed in the previous example to an intransitive action obtained by augmenting the transformation rules (9.2) by an additional invariant coordinate y , so that the pseudo-group now has the form

$$\bar{x} = f(x), \quad \bar{y} = y, \quad \bar{u} = \frac{u}{f'(x)}. \quad (9.7)$$

This pseudo-group was introduced by Lie, [30; p. 373], in his study of second order partial differential equations integrable by the method of Darboux. In his paper on group splitting and automorphic systems, Vessiot, [44], used (9.7) as one of two principal examples illustrating his method. More recently, Kumpera, [26] again employed this pseudo-group

[†] Interestingly, this method is similar to our construction of the Maurer–Cartan forms in the finite-dimensional case.

to illustrate his formalization of the Lie theory of differential invariants. Now we are interested in the equivalence problem and differential invariants for surfaces $u = u(x, y)$ under the pseudo-group (9.7). The Maurer–Cartan forms are given by supplementing (9.5) by an additional coframe element $\zeta_0 = dy$. The linear dependency

$$\theta_2 = -(u\alpha + u_x)\theta_1 - \frac{u_y}{u} dy$$

again produces the normalization $\alpha = -u_x/u$, along with the basic first order differential invariant

$$I = \frac{u_y}{u}.$$

The final invariant moving coframe is

$$\zeta_0 = dy, \quad \zeta_1 = u dx, \quad \zeta_2 = \frac{du - u_x dx}{u}. \quad (9.8)$$

The invariant total differential operators associated with the first two forms are

$$\frac{\partial}{\partial \zeta_0} = D_y, \quad \frac{\partial}{\partial \zeta_1} = \frac{1}{u} D_x. \quad (9.9)$$

Applying them to the fundamental invariant I produce the second order differential invariants

$$J_1 = \frac{uu_{yy} - u_y^2}{u^2}, \quad J_2 = \frac{uu_{xy} - u_y u_x}{u^3},$$

agreeing with the classical formulae. All higher order differential invariants are obtained by successively applying the invariant total derivative operators (9.9) to the invariant I . Similarly, the classifying surface associated with a generic surface $u(x, y)$ is parametrized by the three invariants (I, J_1, J_2) ; two surfaces are congruent under a pseudo-group transformation if and only if their classifying surfaces are identical. Surfaces with higher order symmetry occur when J_1, J_2 are functionally dependent upon I , in which case the classifying surface, reduces to a curve, or when I is constant, so that $u(x, y) = f(x)e^{ky}$. Finally, the most general second order partial differential equation admitting (9.7) (supplemented by $\bar{y} = y$) can be written in the form

$$H \left(y, \frac{u_y}{u}, \frac{uu_{yy} - u_y^2}{u^2}, \frac{uu_{xy} - u_y u_x}{u^3} \right) = 0. \quad (9.10)$$

These are the class of equations considered by Lie, [30; p. 374].

In his classification of planar second order partial differential equations which admit symmetry pseudo-groups, Medolaghi, [32], treats the same example, but rewritten in a slightly different coordinate system. The group transformations take the form

$$\bar{x} = f(x), \quad \bar{y} = y + f'(x), \quad \bar{u} = u. \quad (9.11)$$

Applying the same method (or merely changing variables) leads to the invariant moving coframe

$$\zeta_1 = e^{-y} dx, \quad \zeta_2 = \frac{u_y}{u_x} dx + dy, \quad \zeta_3 = du.$$

The basic differential invariants are

$$u, \quad I = u_y, \quad J_1 = u_{yy}, \quad J_2 = e^y(u_y u_{xy} - u_x u_{yy}),$$

the latter two being obtained by applying the invariant differential operators

$$D_y, \quad e^y(D_x - (u_x/u_y)D_y),$$

to I . This recovers Medolaghi's form, [32; p. 249],

$$H(u, u_y, u_{yy}, e^y(u_y u_{xy} - u_x u_{yy})) = 0, \quad (9.12)$$

of Lie's equation (9.10). The pseudo-group (9.11) is the second of nine different pseudo-groups acting on a three-dimensional space that are isomorphic to the diffeomorphism pseudo-group $\mathcal{D}(1)$, as classified by Medolaghi, [32; p. 242]. The other eight pseudo-groups can be handled by the same method, reproducing the differential invariants and invariant differential equations catalogued there.

Example 9.3. Consider the infinite Lie pseudo-group

$$\bar{x} = f(x), \quad \bar{y} = yf'(x) + g(x), \quad \bar{u} = u + \frac{f''(x)y + g'(x)}{f'(x)}. \quad (9.13)$$

acting on the space $M \simeq \mathbb{R}^3$ with coordinates (x, y, u) . Here $f(x)$ and $g(x)$ are arbitrary smooth functions of a single variable x . The case $g \equiv 0$ corresponds to the third of Medolaghi's pseudo-groups, [32]; the present generalization was introduced by J. Pohjanpelto. The pseudo-group transformations can be characterized in terms of an involutive system of invariant one-forms on a five-dimensional bundle $B \rightarrow M$, with coordinates $(x, y, u, \alpha, \beta, \gamma, \delta, \varepsilon)$. These can be found by a similar method to that used in Example 9.1:

$$\begin{aligned} \zeta_1 &= -\alpha dx, & \zeta_4 &= \frac{d\alpha}{\alpha} - \frac{\gamma}{\alpha} dx, \\ \zeta_2 &= -\alpha dy + u\alpha dx, & \zeta_5 &= \frac{d\beta}{\alpha} + \frac{u}{\alpha} d\gamma - \frac{\delta - u\varepsilon}{\alpha} dx - \frac{\varepsilon}{\alpha} dy, \\ \zeta_3 &= -du - \beta dx - \gamma dy, & \zeta_6 &= \frac{d\gamma}{\alpha} - \frac{\varepsilon}{\alpha} dx. \end{aligned} \quad (9.14)$$

It is easy to checked that a local diffeomorphism $\Psi: B \rightarrow B$ satisfies $\Psi^*\zeta_i = \zeta_i$, $i = 1, \dots, 6$, if and only if it is a bundle map whose projection $\psi: M \rightarrow M$ has the form (9.13).

We now consider the equivalence problem for surfaces $u = u(x, y)$ under the pseudo-group (9.13). In order to invariantly normalize the bundle parameters, we replace du by its horizontal component $u_x dx + u_y dy$, which leads to the linear relation

$$\theta_3 = J_1 \theta_1 + J_2 \theta_2,$$

between the horizontal components θ_i of ζ_i . The lifted invariants are

$$J_1 = \frac{u_x + \beta + u(u_y + \gamma)}{\alpha}, \quad J_2 = \frac{u_y + \gamma}{\alpha}.$$

Both J_1 and J_2 can be normalized to zero by choosing $\beta = -u_x$ and $\gamma = -u_y$, which defines the first order moving frame. Substituting these values in the last two moving coframe forms yields

$$\begin{aligned} \theta_5 &= -\frac{u_{xx} dx + u_{xy} dy}{\alpha} - \frac{u(u_{xy} dx + u_{yy} dy)}{\alpha} - \frac{\delta - u\varepsilon}{\alpha} dx - \frac{\varepsilon}{\alpha} dy \\ &= \frac{u_{xx} + 2uu_{xy} + u^2u_{yy} + \delta}{\alpha^2} \theta_1 - \frac{u_{xy} + uu_{yy} + \varepsilon}{\alpha^2} \theta_2, \\ \theta_6 &= -\frac{(u_{xy} + \varepsilon) dx + u_{yy} dy}{\alpha} = -\frac{u_{xy} + uu_{yy} + \varepsilon}{\alpha^2} \theta_1 + \frac{u_{yy}}{\alpha^2} \theta_2. \end{aligned}$$

We can normalize the coefficients of θ_1, θ_2 in both formulae by choosing

$$\alpha = \sqrt{u_{yy}}, \quad \varepsilon = -u_{xy} - uu_{yy}, \quad \delta = -u_{xx} - 2uu_{xy} - u^2u_{yy},$$

which produces the second order moving frame, given by

$$\alpha = \sqrt{u_{yy}}, \quad \beta = -u_x, \quad \gamma = -u_y, \quad \delta = -u_{xx} - 2uu_{xy} - u^2u_{yy}, \quad \varepsilon = -u_{xy} - uu_{yy}.$$

Finally, substituting into the last moving coframe form leads to $\theta_4 = -I_1\theta_1 - I_2\theta_2$, where

$$I_1 = \frac{uu_{yyy} + u_{xyy} + 2u_y u_{yy}}{2u_{yy}^{3/2}}, \quad I_2 = \frac{u_{yyy}}{2u_{yy}^{3/2}} \quad (9.15)$$

are the principal differential invariants of the pseudo-group. The fundamental invariant one-forms are

$$\theta_1 = -\sqrt{u_{yy}} dx, \quad \theta_2 = -\sqrt{u_{yy}} (dy - u dx),$$

so that the invariant total differential operators are

$$\mathcal{D}_1 = \frac{1}{\sqrt{u_{yy}}} (D_x + uD_y), \quad \mathcal{D}_2 = \frac{1}{\sqrt{u_{yy}}} D_y.$$

As above, these can be applied to the basic differential invariants (9.15) to generate all higher order differential invariants.

Example 9.4. In this example, we show how the well-known equivalence problem of characterizing second order ordinary differential equations under the pseudo-group of fiber-preserving transformations, cf. [20], [36], can be recast into the moving frame formulation, and thereby solved by our moving coframe techniques. This example indicates a general procedure for reformulating all Cartan-type equivalence problems, as in [11], [14], [36], as moving frame equivalence problems under a suitable infinite-dimensional Lie pseudo-group. The main idea is to cast the original pseudo-group of equivalences into the involutive bundle formulation presented above, and then view the original equivalence as a suitable equivalence of submanifolds under the pseudo-group.

We consider the trivial bundle $M \simeq \mathbb{R} \times \mathbb{R}$, with coordinates x, u . Let \mathcal{G} denote the pseudo-group of fiber-preserving transformations, i.e., bundle maps

$$\bar{x} = \phi(x), \quad \bar{u} = \psi(x, u). \quad (9.16)$$

We let $\mathcal{G}^{(2)}$ denote the associated second prolongation acting on J^2 , cf. [36]. A (regular) second-order differential equation

$$\Delta(x, u, u_x, u_{xx}) = 0 \quad (9.17)$$

can be identified with a hypersurface $\mathcal{S}_\Delta \subset J^2$. Two such second order ordinary differential equations are equivalent if and only if their associated surfaces are mapped to each other,

$$g^{(2)}(\mathcal{S}_{\bar{\Delta}}) = \mathcal{S}_\Delta, \quad (9.18)$$

by a prolonged fiber-preserving transformation $g^{(2)} \in \mathcal{G}^{(2)}$.

In order to use the method of moving frames we need the structure equations of the pseudo-group. These can be found by the Cartan prolongation algorithm, [11], [14], [36], leading to the structure equations

$$\begin{aligned} d\zeta_1 &= \eta_1 \wedge \zeta_1, \\ d\zeta_2 &= \eta_2 \wedge \zeta_2 - \zeta_3 \wedge \zeta_1, \\ d\zeta_3 &= (\eta_2 - \eta_1) \wedge \zeta_3 + \eta_3 \wedge \zeta_2 - \zeta_4 \wedge \zeta_1, \\ d\zeta_4 &= (\eta_2 - 2\eta_1) \wedge \zeta_4 + \eta_4 \wedge \zeta_1 + \eta_5 \wedge \zeta_2 + \eta_6 \wedge \zeta_3, \\ d\eta_1 &= (\eta_6 - 2\eta_3) \wedge \zeta_1, \\ d\eta_2 &= -\pi_2 \wedge \zeta_2 - \eta_3 \wedge \zeta_1, \\ d\eta_3 &= -\pi_1 \wedge \zeta_2 - \pi_2 \wedge \zeta_3 + \eta_3 \wedge \eta_1 - \eta_5 \wedge \zeta_1, \\ d\eta_4 &= -\pi_3 \wedge \zeta_1 - \pi_4 \wedge \zeta_3 - \pi_5 \wedge \zeta_2 - 3\eta_1 \wedge \eta_4 - \eta_4 \wedge \eta_2 + 3(\eta_3 - \eta_6) \wedge \zeta_4, \\ d\eta_5 &= -2\pi_1 \wedge \zeta_3 - \pi_2 \wedge \zeta_4 - \pi_5 \wedge \zeta_1 - \pi_6 \wedge \zeta_2 + 2\eta_5 \wedge \eta_1 - \eta_3 \wedge \eta_6, \\ d\eta_6 &= -2\pi_1 \wedge \zeta_2 - 2\pi_2 \wedge \zeta_3 - \pi_4 \wedge \zeta_1 - \eta_1 \wedge \eta_6 + \eta_5 \wedge \zeta_1. \end{aligned}$$

The Cartan characters are $s_1 = 5$ and $s_2 = 1$, the kernel dimension is 7, hence this system is involutive.

The parametric values of the one-forms ζ_i, η_j , are determined by introducing the group transformation matrix

$$S = \begin{pmatrix} \alpha_1 & 0 & 0 & 0 \\ 0 & \alpha_2 & 0 & 0 \\ 0 & \alpha_3 \alpha_2 \alpha_1^{-1} & \alpha_2 \alpha_1^{-1} & 0 \\ \alpha_4 \alpha_2 \alpha_1^{-2} & \alpha_5 \alpha_2 \alpha_1^{-2} & \alpha_6 \alpha_2 \alpha_1^{-2} & \alpha_2 \alpha_1^{-2} \end{pmatrix}, \quad (9.19)$$

where α_i, β_i are the fiber coordinates on the prolonged bundle. Equation (9.19) parametrizes the structure group corresponding to the action of the fiber-preserving pseudo-group

on J^2 ; see, for instance, [36; p. 398] for the corresponding group on J^1 . The first set of lifted forms are

$$\begin{pmatrix} \zeta_1 \\ \zeta_2 \\ \zeta_3 \\ \zeta_4 \end{pmatrix} = \begin{pmatrix} \alpha_1 & 0 & 0 & 0 \\ 0 & \alpha_2 & 0 & 0 \\ 0 & \alpha_3 \alpha_2 \alpha_1^{-1} & \alpha_2 \alpha_1^{-1} & 0 \\ \alpha_4 \alpha_2 \alpha_1^{-2} & \alpha_5 \alpha_2 \alpha_1^{-2} & \alpha_6 \alpha_2 \alpha_1^{-2} & \alpha_2 \alpha_1^{-2} \end{pmatrix} \begin{pmatrix} dx \\ du - u_x dx \\ du_x - u_{xx} dx \\ du_{xx} \end{pmatrix}.$$

Furthermore,

$$\begin{pmatrix} \eta_1 & 0 & 0 & 0 \\ 0 & \eta_2 & 0 & 0 \\ 0 & \eta_3 & \eta_2 - \eta_1 & 0 \\ \eta_6 & \eta_5 & \eta_4 & \eta_2 - 2\eta_1 \end{pmatrix} = S^{-1} dS + \Omega,$$

where Ω represents the absorbed torsion terms. The explicit formulas are

$$\begin{aligned} \eta_1 &= \frac{d\alpha_1}{\alpha_1} + \frac{\alpha_6 - 2\alpha_3}{\alpha_2} \zeta_1, \\ \eta_2 &= \frac{d\alpha_2}{\alpha_2} + \frac{\alpha_3}{\alpha_1} \zeta_1 - \beta_2 \zeta_2, \\ \eta_3 &= \frac{d\alpha_3}{\alpha_2} + \frac{\alpha_3 \alpha_6 - \alpha_5 - \alpha_3^2}{\alpha_1^2} \zeta_1 - \beta_1 \zeta_2 - \beta_2 \zeta_3, \\ \eta_4 &= \frac{\alpha_2}{\alpha_1^3} d\alpha_4 - \beta_3 \zeta_1 + \frac{\alpha_6 \alpha_5 + \alpha_3 \alpha_5 - \alpha_3 \alpha_6^2 - \alpha_2^2 \beta_5}{\alpha_2^2} \zeta_2 + \frac{\alpha_6^2 - \alpha_5 - \beta_4 \alpha_1^2}{\alpha_1^2} \zeta_3 + 3 \frac{\alpha_3 - \alpha_6}{\alpha_1} \zeta_4, \\ \eta_5 &= \frac{d\alpha_5}{\alpha_1^2} - \frac{\alpha_3}{\alpha_1^2} d\alpha_4 + \beta_6 \zeta_2 - 2\beta_1 \zeta_3 - \beta_2 \zeta_4 - \beta_5 \eta_1, \\ \eta_6 &= \frac{d\alpha_6}{\alpha_1} - \beta_4 \zeta_1 - 2\beta_1 \zeta_2 - 2\beta_2 \zeta_3. \end{aligned}$$

We now assume, for simplicity, that the second order ordinary differential equation (9.17) is given by the graph of a section $\sigma: J^1 \rightarrow J^2$; this is equivalent to assuming that the equation is normal, and solved

$$u_{xx} = Q(x, u, u_x), \tag{9.20}$$

for its highest order derivative. (However, the moving frame method could be applied without this assumption; doing the corresponding problem for non-normal equations using the Cartan equivalence approach would be harder.) Pulling back the Maurer-Cartan forms under the map σ amounts to substituting for u_{xx} according to (9.20) where-ever it occurs. We denote the pull-back of ζ_i by θ_i and of η_i by ϖ_i . To apply the moving frame method, we look for dependencies among the resulting one-forms. The first of these is

$$\theta_4 = J_1 \theta_1 + J_2 \theta_2 + J_3 \theta_3,$$

where

$$\begin{aligned} J_1 &= \frac{\alpha_2}{\alpha_1^3} \left(\alpha_4 + \frac{dQ}{dx} \right), \\ J_2 &= \frac{1}{\alpha_1^2} \left(\alpha_5 - \alpha_6 \alpha_3 + \frac{\partial Q}{\partial u} - \alpha_3 \frac{\partial Q}{\partial u_x} \right), \\ J_3 &= \frac{1}{\alpha_1} \left(\alpha_6 + \frac{\partial Q}{\partial u_x} \right). \end{aligned} \tag{9.21}$$

Here

$$\frac{dQ}{dx} = \frac{\partial Q}{\partial x} + u_x \frac{\partial Q}{\partial u} + Q \frac{\partial Q}{\partial u_x}$$

denotes the total derivative of Q , restricted to the equation manifold (9.20). The lifted invariants (9.21) can all be translated to zero by choosing

$$\alpha_4 = -\frac{dQ}{dx}, \quad \alpha_5 = -\frac{\partial Q}{\partial u}, \quad \alpha_6 = -\frac{\partial Q}{\partial u_x}.$$

We then pull-back the forms η_5, η_6 , leading to

$$\begin{aligned} \varpi_5 &\equiv -\left(2\beta_1 + \frac{1}{\alpha_2 \alpha_1} Q_{uu_x} - \frac{\alpha_3}{\alpha_2^2} Q_{u_x u_x} \right) \theta_3, \\ \varpi_6 &\equiv -\left(2\beta_2 + \frac{Q_{u_x u_x}}{\alpha_2} \right) \theta_3, \end{aligned} \quad \text{mod}\{\theta_1, \theta_2\}$$

Translating the coefficients of θ_3 to zero in ϖ_5 and ϖ_6 gives

$$\beta_1 = -\frac{1}{2\alpha_1 \alpha_2} Q_{uu_x} + \frac{\alpha_3}{2\alpha_2^2} Q_{u_x u_x}, \quad \beta_2 = -\frac{1}{2\alpha_2} Q_{u_x u_x},$$

which then leads to the pulled-back forms

$$\begin{aligned} \varpi_1 &= \frac{d\alpha_1}{\alpha_1} - \left(\frac{Q_{u_x} + 2\alpha_3}{\alpha_1} \right) \theta_1, \\ \varpi_2 &= \frac{d\alpha_2}{\alpha_2} - \frac{\alpha_3}{\alpha_1} \theta_1 + \left(\frac{1}{2\alpha_2} Q_{u_x u_x} \right) \theta_2, \\ \varpi_3 &= \frac{d\alpha_3}{\alpha_1} + \left(\frac{Q_u - \alpha_3^2 - \alpha_3 Q_{u_x}}{\alpha_1^2} \right) \theta_1 + \frac{1}{2\alpha_1 \alpha_2} (Q_{uu_x} - \alpha_3 Q_{u_x u_x}) \theta_2 + \left(\frac{1}{2\alpha_2} Q_{u_x u_x} \right) \theta_3. \end{aligned}$$

At this stage, we have reproduced the one-forms produced by the equivalence method in [20; p. 394], to which we refer the reader for a further discussion of this example.

10. Conclusions.

In this paper we have described, albeit without theoretical justification, a systematic procedure for determining moving frames and invariant differential forms for very general Lie group and Lie pseudo-group actions. The moving frame and moving coframe can be used to directly determine a complete system of fundamental differential invariants and invariant differential operators for the given transformation group. These, in turn, have immediate applications, including the solution to equivalence problems, classification of symmetry groups, rigidity theorems, construction of invariant equations and variational principles, and so on. As we have demonstrated, the method not only readily reproduces all of the standard examples of moving frames known in the literature, but is also in a form that can immediately be applied to a host of new and interesting group actions, including intransitive and ineffective actions, infinite-dimensional Lie pseudo-groups, joint actions, and so on. The theoretical foundations of our method will be presented in the second paper in this series. Additional applications, to differential invariants, to the theory of Lie pseudo-groups, to automorphic systems, and to computer vision, will be the subject of subsequent papers in this series. Some additional extensions that we intend to investigate include:

- (1) The moving coframe method, as described in this paper, parallels the explicit “parametric” approach to the solution of Cartan equivalence problems. Gardner, [14], showed how, in such situations, one could perform an “intrinsic” computation, based on the infinitesimal group action on the torsion coefficients, and thereby determine the general structure of the solution. An interesting question is whether one can implement an intrinsic version of the moving coframe algorithm.
- (2) In [24] a inductive approach to complicated equivalence problems, based on the solution to a simpler problem based on a subgroup of the full structure group, was proposed; see also [36]. This method not only simplifies the computations, but also provides direct correspondences between the invariants of the two problems. Is there a similar inductive version of the moving coframe method? For example, does the computation of the moving frame for curves in the plane under, say, the equi-affine group help simplify the corresponding projective computation, thereby expressing the projective arc length and curvature directly in terms of its equiaffine counterparts?
- (3) In [5], a new scheme for generating invariant numerical approximations to differential invariants based on the use of joint invariants was proposed, and illustrated in the planar Euclidean and equi-affine cases. The computation of joint differential invariants using the moving coframe method strongly indicates that it could be applied to the general problem of invariant numerical formulae for more complicated transformation groups. In particular, determining how joint invariants converge to differential invariants as the points coalesce would be of great importance.
- (4) An immediate and important application of the moving method would be to the classification of the differential invariants associated many of the fundamental groups arising in physics. As remarked above, to date such classifications have not been completed, even for some of the most fundamental groups of physical import.

Acknowledgments: We particularly thank Ian Anderson for inspiration, enlightening discussions, and provocative comments. One of us (P.J.O.) would like to thank Allen Tannenbaum and Olivier Faugeras for stimulating remarks on moving frames and differential invariants in computer vision.

References

- [1] Ackerman, M., and Hermann, R., *Sophus Lie's 1884 Differential Invariant Paper*, Math Sci Press, Brookline, Mass., 1976.
- [2] Anderson, I.M., Introduction to the variational bicomplex, *Contemp. Math.* **132** (1992), 51–73.
- [3] Bryant, R.L., Chern, S.-S., Gardner, R.B., Goldschmidt, H.L., and Griffiths, P.A., *Exterior Differential Systems*, Math. Sci. Res. Inst. Publ., Vol. 18, Springer–Verlag, New York, 1991.
- [4] Buchberger, B., Applications of Gröbner bases in non-linear computational geometry, in: *Mathematical Aspects of Scientific Software*, J.R. Rice, ed., IMA Volumes in Mathematics and its Applications, Vol. 14, Springer–Verlag, New York, 1988, pp. 59–87.
- [5] Calabi, E., Olver, P.J., Shakiban, C., Tannenbaum, A., and Haker, S., Differential and numerically invariant signature curves applied to object recognition, preprint, University of Minnesota, 1996.
- [6] Cartan, É., *La Méthode du Repère Mobile, la Théorie des Groupes Continus, et les Espaces Généralisés*, Exposés de Géométrie No. 5, Hermann, Paris, 1935.
- [7] Cartan, É., *Leçons sur la Théorie des Espaces à Connexion Projective*, Cahiers Scientifiques, Vol. 17, Gauthier–Villars, Paris, 1937.
- [8] Cartan, É., *La Théorie des Groupes Finis et Continus et Lie algebra Géométrie Différentielle traitées par la Méthode du Repère Mobile*, Cahiers Scientifiques, Vol. 18, Gauthier–Villars, Paris, 1937.
- [9] Cartan, É., *Les Systèmes Différentiels Extérieurs et Leurs Applications Géométriques*, Exposés de Géométrie No. 14, Hermann, Paris, 1945.
- [10] Cartan, É., Sur la structure des groupes infinis de transformations, in: *Oeuvres Complètes*, Part. II, Vol. 2, Gauthier–Villars, Paris, 1953, pp. 571–624.
- [11] Cartan, É., Les problèmes d'équivalence, in: *Oeuvres Complètes*, Part. II, Vol. 2, Gauthier–Villars, Paris, 1953, pp. 1311–1334.
- [12] Faugeras, O., Cartan's moving frame method and its application to the geometry and evolution of curves in the euclidean, affine and projective planes, in: *Applications of Invariance in Computer Vision*, J.L. Mundy, A. Zisserman, D. Forsyth (eds.), Springer–Verlag Lecture Notes in Computer Science, Vol. 825, 1994, pp. 11–46.
- [13] Fels, M., and Olver, P.J., On relative invariants, *Math. Ann.*, to appear.

- [14] Gardner, R.B., *The Method of Equivalence and Its Applications*, SIAM, Philadelphia, 1989.
- [15] Green, M.L., The moving frame, differential invariants and rigidity theorems for curves in homogeneous spaces, *Duke Math. J.* **45** (1978), 735–779.
- [16] Griffiths, P.A., On Cartan’s method of Lie groups and moving frames as applied to uniqueness and existence questions in differential geometry, *Duke Math. J.* **41** (1974), 775–814.
- [17] Guggenheimer, H.W., *Differential Geometry*, McGraw–Hill, New York, 1963.
- [18] Heredero, R.H., and Olver, P.J., Classification of invariant wave equations, *J. Math. Phys.*, to appear.
- [19] Hille, E., *Ordinary Differential Equations in the Complex Domain*, John Wiley & Sons, New York, 1976.
- [20] Hsu, L., and Kamran, N., Classification of second-order ordinary differential equations admitting Lie groups of fiber-preserving symmetries, *Proc. London Math. Soc.* **58** (1989), 387–416.
- [21] Jensen, G.R., *Higher order contact of submanifolds of homogeneous spaces*, Lecture Notes in Math., No. 610, New York, Springer–Verlag, 1977.
- [22] Johnson, H.H., Classical differential invariants and applications to partial differential equations, *Math. Ann.* **148** (1962), 308–329.
- [23] Kamran, N., Contributions to the study of the equivalence problem of Elie Cartan and its applications to partial and ordinary differential equations, *Mém. Cl. Sci. Acad. Roy. Belg.* **45** (1989), Fac. 7.
- [24] Kamran, N., and Olver, P.J., Equivalence problems for first order Lagrangians on the line, *J. Diff. Eq.* **80** (1989), 32–78.
- [25] Kobayashi, S., Canonical forms on frame bundles of higher order contact, *Proc. Sympos. Pure Math.* **3** (1961), 186–19.
- [26] Kumpera, A., Invariants différentiels d’un pseudogroupe de Lie, *J. Diff. Geom.* **10** (1975), 289–416.
- [27] Lie, S., Über unendlichen kontinuierliche Gruppen, *Christ. Forh. Aar.* **8** (1883), 1–47; also *Gesammelte Abhandlungen*, Vol. 5, B.G. Teubner, Leipzig, 1924, pp. 314–360.
- [28] Lie, S., Über Differentialinvarianten, *Math. Ann.* **24** (1884), 537–578; also *Gesammelte Abhandlungen*, Vol. 6, B.G. Teubner, Leipzig, 1927, pp. 95–138; see [1] for an English translation.
- [29] Lie, S., Die Grundlagen für die Theorie der unendlichen kontinuierlichen Transformationsgruppen, *Leipzig. Ber.* **43** (1891), 316–393; also *Gesammelte Abhandlungen*, Vol. 6, B.G. Teubner, Leipzig, 1927, pp. 300–364.
- [30] Lie, S., Zur allgemeinen Theorie der partiellen Differentialgleichungen beliebiger Ordnung, *Leipz. Berich.* **47** (1895), 53–128; also *Gesammelte Abhandlungen*, Vol. 4, B.G. Teubner, Leipzig, 1929, 320–384.
- [31] Lie, S., Gruppenregister, in: *Gesammelte Abhandlungen*, Vol. 5, B.G. Teubner, Leipzig, 1924, pp. 767–773.

- [32] Medolaghi, P., Classificazione delle equazioni alle derivate parziali del secondo ordine, che ammettono un gruppo infinito di trasformazioni puntuali, *Ann. Mat. Pura Appl.* **1** (3) (1898), 229–263.
- [33] Moons, T., Pauwels, E., Van Gool, L., and Oosterlinck, A., Foundations of semi-differential invariants, *Int. J. Comput. Vision* **14** (1995), 25–48.
- [34] Olver, P.J., Classical invariant theory and the equivalence problem for particle Lagrangians. I. Binary Forms, *Adv. in Math.* **80** (1990), 39–77.
- [35] Olver, P.J., *Applications of Lie Groups to Differential Equations*, Second Edition, Graduate Texts in Mathematics, vol. 107, Springer–Verlag, New York, 1993.
- [36] Olver, P.J., *Equivalence, Invariants, and Symmetry*, Cambridge University Press, Cambridge, 1995.
- [37] Olver, P.J., Pseudo-stabilization of prolonged group actions. I. The order zero case, preprint, University of Minnesota, 1996.
- [38] Olver, P.J., Sapiro, G., and Tannenbaum, A., Invariant geometric evolutions of surfaces and volumetric smoothing, *SIAM J. Appl. Math.*, to appear.
- [39] Ovsiannikov, L.V., *Group Analysis of Differential Equations*, Academic Press, New York, 1982.
- [40] Robart, T., and Kamran, N., Sur Lie algebra théorie locale des pseudogroupes de transformations continus infinis I, preprint, McGill University, 1996.
- [41] Singer, I.M., and Sternberg, S., The infinite groups of Lie and Cartan. Part I (the transitive groups), *J. Analyse Math.* **15** (1965), 1–114.
- [42] Sternberg, S., *Lectures on Differential Geometry*, Prentice-Hall, Englewood Cliffs, N.J., 1964.
- [43] Tresse, A., Sur les invariants différentiels des groupes continus de transformations, *Acta Math.* **18** (1894), 1–88.
- [44] Vessiot, E., Sur l’intégration des systèmes différentiels qui admettent des groupes continus de transformations, *Acta. Math.* **28** (1904), 307–349.
- [45] Weyl, H., Cartan on groups and differential geometry, *Bull. Amer. Math. Soc.* **44** (1938), 598–601.
- [46] Weyl, H., *Classical Groups*, Princeton Univ. Press, Princeton, N.J., 1946.
- [47] Willmore, T.J., *Riemannian Geometry*, Oxford University Press, Oxford, 1993.