

THE HP-LOCAL DISCONTINUOUS GALERKIN METHOD FOR LOW-FREQUENCY TIME-HARMONIC MAXWELL'S EQUATIONS

ILARIA PERUGIA * AND DOMINIK SCHÖTZAU †

Version of June 11, 2001

Abstract. The local discontinuous Galerkin method for the numerical approximation of the time-harmonic Maxwell equations in low-frequency regime is introduced and analyzed. We consider topologically non-trivial domains and heterogeneous media, containing both conducting and insulating materials. The presented method involves discontinuous Galerkin discretizations of the curl-curl and grad-div operators, based on a mixed formulation of the problem and on the introduction of the so-called numerical fluxes. An hp -analysis is carried out and error estimates that are optimal in the meshsize h and slightly suboptimal in the approximation degree p are obtained.

Key words. hp -finite elements, discontinuous Galerkin methods, low-frequency time-harmonic Maxwell's equations, heterogeneous media

AMS subject classifications. 65N30

1. Introduction. In this paper, we propose and analyze an hp -local discontinuous Galerkin (LDG) method for the low-frequency time-harmonic Maxwell equations in heterogeneous media, containing both conducting and insulating materials. The method is based on the following formulation of the problem: find the complex field \mathbf{E} that satisfies

$$\nabla \times (\mu^{-1} \nabla \times \mathbf{E}) + i\omega\sigma\mathbf{E} = -i\omega\mathbf{J}_s =: \mathbf{J} \quad \text{in } \Omega \subset \mathbb{R}^3 \quad (1.1)$$

$$\nabla \cdot \mathbf{E} = 0 \quad \text{in } \Omega_0 \subset \Omega, \quad (1.2)$$

together with suitable boundary conditions (see Alonso and Valli [1]). The complex field \mathbf{E} is actually related to the electric field \mathcal{E} by the identity $\mathcal{E}(\mathbf{x}, t) = \text{Re}(\mathbf{E}(\mathbf{x})e^{i\omega t})$, where $\omega \neq 0$ is a given frequency. The parameter $\mu = \mu(\mathbf{x})$ is the magnetic permeability, \mathbf{J}_s is the phasor associated with a given current density and $\sigma = \sigma(\mathbf{x})$ is the electric conductivity, which is zero in the subdomain Ω_0 occupied by insulating materials. We remark that the electric field-based formulation in (1.1)–(1.2) is only one of several field and potential-based formulations proposed in the literature for the solution of eddy current problems (see, e.g., Bryan, Emson, Fernandes and Trowbridge [12], Bossavit [11], Hiptmair [33] and the references therein).

There are two main reasons for using discontinuous Galerkin (DG) methods for the numerical approximation of the above problem.

The first is that these methods, being based on discontinuous finite element spaces, can easily handle meshes with hanging nodes, elements of general shape and local spaces of different types. They are thus ideally suited for hp -adaptivity and multi-physics or multi-material problems. This flexibility in the mesh-design is not shared in

*Dipartimento di Matematica, Università di Pavia, Via Ferrata 1, 27100 Pavia, Italy, email: perugia@dimat.unipv.it. Supported in part by the National Science Foundation (Grant DMS-9807491) and by the Supercomputing Institute of the University of Minnesota. This work was carried out when the author was visiting the School of Mathematics, University of Minnesota.

†School of Mathematics, University of Minnesota, Vincent Hall, Minneapolis, MN 55455, email: schoetza@math.umn.edu.

a straightforward way by standard edge or face elements commonly used in computational electromagnetics. Indeed, these elements are designed to enforce the continuity of certain momenta in tangential or normal direction across interelement boundaries (see, e.g., Nédélec [37, 38], Bossavit [9, 10], and Monk [36]). This makes the handling of non-matching grids and high-order approximations rather inconvenient from an implementational point of view. Nevertheless, efficient hp -adaptive edge element methods have been developed recently by Demkowicz and Vardapetyan in [29, 45].

The second reason, of no less importance, is related to the difficulties in obtaining numerical approximations in the subregion Ω_0 containing the insulators. In fact, when conforming elements are considered, the continuity of both the tangential and normal components of the discrete field has to be imposed and, therefore, continuous elements have to be used. It is well known that for such approximations convergence to the exact solution might not be achieved, for instance, if Ω_0 is a non-convex polyhedron or in the presence of singularities of the material properties. These situations are in fact excluded in Alonso and Valli [1], where an iteration-by-subdomain procedure is studied, using edge elements in the conducting region $\Omega \setminus \Omega_0$ and continuous elements in Ω_0 . Possible remedies can be found, in the context of the full Maxwell problem, in Bonnet-BenDhia, Hazard and Lohrengel [8], where a singular field method is introduced; in [27], where the original variational forms are modified near solution singularities; or in Chen, Du and Zou [20], where the divergence constraint is taken into account by a mixed method. These problems are overcome in a natural way by DG methods, since they are based on completely discontinuous finite element spaces and the continuity requirements are enforced in a weak sense only.

The LDG method has been introduced by Cockburn and Shu [26] for convection-diffusion systems, and has been further developed and analyzed in Cockburn and Dawson [22], Castillo, Cockburn, Schötzau and Schwab [19], Castillo, Cockburn, Perugia and Schötzau [18], Cockburn, Kanschat, Perugia and Schötzau [23]; see also the review by Cockburn and Shu [25]. It is one of several DG methods that have been proposed in the literature for diffusion problems. We only mention here the DG methods of Baumann and Oden [7, 39], and the interior penalty (IP) methods and their variants which have been recently studied, e.g., in Rivière, Wheeler and Girault [43], Rivière and Wheeler [42] and Houston, Schwab and Süli [34]. A comparison of DG methods from a computational point of view can be found in Castillo [17]. Recent works have aimed at unifying the presentation and the analysis of all these methods for elliptic problems. In Prudhomme, Pascal, Oden and Romkes [41], an hp -analysis of different DG methods has been given, including the Baumann-Oden method and interior penalty methods. Furthermore, in Arnold, Brezzi, Cockburn and Marini [5], a framework has been presented within which virtually all the DG methods found in the literature can be analyzed; it is based on a mixed formulation of the second-order problem and on the so-called numerical fluxes.

The LDG method for the discretization of (1.1)–(1.2) is designed by adapting to the curl-curl and grad-div operators the definition of the numerical fluxes considered in [26, 18] for the Laplacian. This is done in a consistent way and such that the auxiliary variables needed to define the LDG formulation can be eliminated from the equations in an element-by-element manner. For discontinuity stabilization parameters of the order p^2/h , we prove error estimates that are optimal in the mesh-size h and slightly suboptimal in the polynomial degree p (half a power of p is lost). This analysis is the first hp -error analysis for the LDG method in several space dimensions and in this sense extends previous work in [19, 18]. For elliptic problems on unstructured

two- or three-dimensional domains no better p -bounds can be found in the literature; see, e.g., Rivière, Wheeler and Girault [43], Prudhomme, Pascal, Oden and Romkes [41] and Houston, Schwab and Süli [34], where the same rates of convergence as in our case are obtained by different analysis techniques. We mention, however, that optimal hp -bounds have been proved by Castillo, Cockburn, Schötzau and Schwab [19] for one-dimensional convection-diffusion problems, and recently by Georgoulis and Süli [32] for two-dimensional reaction-diffusion problems on structured quadrilateral grids.

The outline of the paper is as follows. In section 2, we present the low-frequency time-harmonic Maxwell equations in heterogeneous media, under quite general and realistic assumptions on the domain and the data. We need to extend to our case the existence and uniqueness results established in [1] for a more particular situation. The proof of these extensions is developed in detail in the appendix, and relies on the existence of a continuous lifting of tangential traces, which is divergence free in Ω_0 and satisfies certain homogeneous flux conditions through the cavities of Ω_0 . In section 3, we derive the LDG method and show that it defines a unique approximate solution. An hp -error analysis is carried out in section 4. Possible extensions of our work and concluding remarks are presented in section 5.

2. The model problem in heterogeneous media. In this section, we specify our assumptions on the domain and the data, and present the complete model problem in heterogeneous media. The proof of the well-posedness of the continuous problem is postponed to the appendix.

2.1. Preliminaries. We start by making precise the assumptions on the domain and on the data, and by introducing the functional spaces used throughout the paper.

Assumptions on the domain. Let Ω be a connected, bounded, open Lipschitz polyhedron in \mathbb{R}^3 , whose boundary may contain several connected, not necessarily simply connected components. Throughout the paper, whenever referring to a non-simply connected domain, we assume that there exists an “admissible set of cuts” in the sense of [3], whose removal reduces the domain to a simply connected one (see also [31] for further comments). Let Ω_0 be the subdomain of Ω occupied by insulating materials. We define $\Omega_\sigma = \Omega \setminus \Omega_0$, and denote by Γ the interface $\partial\Omega_0 \cap \partial\Omega_\sigma$. We assume Ω_0 and Ω_σ to be open Lipschitz polyhedra such that the closure of Γ is a collection of closed faces of $\partial\Omega_0$ and $\partial\Omega_\sigma$. For the sake of simplicity, we assume Ω_0 to be connected. The extension to the general case where Ω_0 is not connected can be done easily by dealing with each of the connected components of Ω_0 as done with Ω_0 in this paper. Let $\Gamma_{0,j}$, $j = 0, \dots, J$, be the connected, not necessarily simply connected components of $\partial\Omega_0$. We denote by $\Gamma_{0,0}$ the “external” connected component of $\partial\Omega_0$, defined as the boundary of the only unbounded component of $\mathbb{R}^3 \setminus \overline{\Omega_0}$, and by $\Gamma_{0,j}$, $j = 1, \dots, J$, the possible “cavities” of Ω_0 , which are boundaries of connected, bounded Lipschitz polyhedra in $\mathbb{R}^3 \setminus \overline{\Omega_0}$.

Assumptions on the data. The magnetic permeability and reluctivity, μ and μ^{-1} , are symmetric, uniformly positive definite tensors with bounded coefficients. The electric conductivity σ is a symmetric tensor with bounded coefficients, uniformly positive definite in the conducting region Ω_σ and zero in Ω_0 . These tensors are smooth within any subdomain occupied by a single material, and might be discontinuous across the interfaces between different materials. Finally, the current density \mathbf{J} satisfies $\mathbf{J} = \mathbf{0}$ in Ω_σ , $\nabla \cdot \mathbf{J} = 0$ in Ω_0 and $\mathbf{J} \cdot \mathbf{n}_0 = 0$ on $\partial\Omega_0$, where \mathbf{n}_0 is the outward normal unit vector to $\partial\Omega_0$.

Functional spaces. Given a domain D in \mathbb{R}^2 or \mathbb{R}^3 , we denote, as usual, by $H^s(D)^d$, $d = 1, 2, 3$, the Sobolev space of real or complex functions with integer or fractional regularity exponent $s \geq 0$, endowed with the norm $\|\cdot\|_{s,D}$; see, e.g., [35]. For $D \subset \mathbb{R}^3$, $H(\text{curl}; D)$ and $H(\text{div}; D)$ are the spaces of real or complex vector functions $\mathbf{u} \in L^2(D)^3$ with $\nabla \times \mathbf{u} \in L^2(D)^3$ and $\nabla \cdot \mathbf{u} \in L^2(D)$, respectively, endowed with the graph norms. We denote by $H_0^1(D)$, $H_0(\text{curl}; D)$ and $H_0(\text{div}; D)$ the subspaces of $H^1(D)$, $H(\text{curl}; D)$ and $H(\text{div}; D)$ of functions with zero trace, tangential trace and normal trace, respectively, and by $H(\text{curl}^0; D)$ and $H(\text{div}^0; D)$ the subspaces of $H(\text{curl}; D)$ and $H(\text{div}; D)$ of curl-free and divergence-free functions, respectively. We also define $H_0(\text{curl}, \text{div}; D) = H_0(\text{curl}; D) \cap H_0(\text{div}; D)$ and $H_0(\text{curl}^0, \text{div}^0; D) = H_0(\text{curl}^0; D) \cap H_0(\text{div}^0; D)$. Finally, we denote by $\mathcal{H}(\partial D)$ the space of tangential traces of $H(\text{curl}; D)$ functions endowed with the norm $\|\cdot\|_{\mathcal{H}(\partial D)}$, and refer to [13] for its complete characterization in non-simply connected domains.

2.2. Low-frequency time-harmonic Maxwell's equations in heterogeneous media. The physical problem we are interested in is the low-frequency time-harmonic Maxwell system (1.1)–(1.2), completed with Dirichlet boundary conditions on $\partial\Omega$ and flux conditions through the cavities of Ω_0 ; see [1]. Renaming the unknown field, the complete problem reads as follows: find $\mathbf{u} \in H(\text{curl}; \Omega) \cap H(\text{div}; \Omega_0)$ such that

$$\nabla \times (\mu^{-1} \nabla \times \mathbf{u}) + i\omega \sigma \mathbf{u} = \mathbf{J} \quad \text{in } \Omega \quad (2.1)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } \Omega_0 \quad (2.2)$$

$$\mathbf{n} \times \mathbf{u} = \mathbf{g} \quad \text{on } \partial\Omega \quad (2.3)$$

$$\langle (\mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = 0 \quad \forall j = 1, \dots, J, \quad (2.4)$$

where \mathbf{n} is the outward normal unit vector to $\partial\Omega$, \mathbf{g} is the tangential trace in $\mathcal{H}(\partial\Omega)$ of a function in $H(\text{curl}; \Omega)$, $\mathbf{n}_{0,j}$ is the normal unit vector to $\Gamma_{0,j}$ pointing outside Ω_0 , and $\langle \cdot, \cdot \rangle_{\Gamma_{0,j}}$ denotes the duality product between $H^{-\frac{1}{2}}(\Gamma_{0,j})$ and $H^{\frac{1}{2}}(\Gamma_{0,j})$, with $L^2(\Gamma_{0,j})$ as pivot space.

Although \mathbf{J} is divergence-free in Ω_0 , possible errors in the experimental recovering and/or numerical representation of \mathbf{J} may give rise to source terms components that are not divergence-free. In order to address this issue, we consider the following variational formulation of (2.1)–(2.4): find $\mathbf{u} \in H(\text{curl}; \Omega) \cap H(\text{div}; \Omega_0)$ such that $\mathbf{n} \times \mathbf{u} = \mathbf{g}$ on $\partial\Omega$, $\langle (\mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = 0$ for $j = 1, \dots, J$, and

$$\int_{\Omega} \mu^{-1} \nabla \times \mathbf{u} \cdot \nabla \times \bar{\mathbf{v}} \, d\mathbf{x} + i\omega \int_{\Omega} \sigma \mathbf{u} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\Omega_0} \nu \nabla \cdot \mathbf{u} \nabla \cdot \bar{\mathbf{v}} \, d\mathbf{x} = \int_{\Omega} \mathbf{J} \cdot \bar{\mathbf{v}} \, d\mathbf{x}, \quad (2.5)$$

for all $\mathbf{v} \in H_0(\text{curl}; \Omega) \cap H(\text{div}; \Omega_0)$ with $\langle (\mathbf{v}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = 0$, $j = 1, \dots, J$. Here, $\nu = \nu(\mathbf{x})$ is any positive bounded scalar function, bounded away from zero, with the same physical dimension as μ^{-1} , that should be chosen in such a way that the magnitudes of the different terms at the left-hand side are balanced. The LDG method we are going to introduce is actually based on a discrete counterpart of (2.5). Now, we can account for perturbations of the data by considering the strong problem corresponding to (2.5) with \mathbf{J} replaced by a generic $\mathbf{F} \in L^2(\Omega)^3$. This is done in section 2.3.

REMARK 2.1. *A conforming finite element discretization of the variational formulation in (2.5) requires elements in Ω_0 that are both curl- and div-conforming, and therefore also H^1 -conforming. Whenever Ω_0 is a non-convex polyhedron, it is*

well-known that these conforming approximations may fail to converge to the exact solution. In [1], for instance, where the above problem is discretized by a domain decomposition method with conforming elements, this situation is forbidden. See also [8] for a discussion of this issue in the context of the full time-harmonic Maxwell problem in a closed cavity. The use of discontinuous Galerkin methods is a way to overcome these difficulties since these methods are based on completely discontinuous finite element spaces and the continuity requirements are enforced in a weak sense only.

REMARK 2.2. Conditions (2.2) and (2.4) are actually motivated in [1], in the case $\mathbf{J} = \mathbf{0}$ in Ω , by means of a physical perturbation argument. Let us adapt this argument to our case. To this aim, we introduce the perturbed tensor σ_ϑ defined as $\sigma_\vartheta = \sigma$ in Ω_σ and $\sigma_\vartheta = \vartheta I$ in Ω_0 , where I is the identity and ϑ is a positive constant that we will let tend to zero. By Lemma 3.5 in [3], we can represent the current density \mathbf{J} in Ω_0 as $\mathbf{J} = \nabla \times \psi$, with a field $\psi \in H^1(\Omega_0)^3$. Consequently, from equation (2.1) with σ replaced by σ_ϑ , we have $i\omega\vartheta\mathbf{u} = -\nabla \times (i\omega\psi + \mu^{-1}\nabla \times \mathbf{u})$ in Ω_0 . Therefore, $\mathbf{u}|_{\Omega_0}$ belongs to $\nabla \times (H(\text{curl}; \Omega_0))$. From the second part of Lemma 3.5 in [3], which holds true also for $H(\text{curl}; \Omega_0)$ functions, we have that \mathbf{u} satisfies (2.2) and (2.4) for any $\vartheta > 0$. Then, as $\vartheta \rightarrow 0$, we obtain problem (2.1)–(2.4).

2.3. The model problem. In this section, we introduce a slightly more general model problem than the one in (2.1)–(2.4), in order to allow for general source terms $\mathbf{F} \in L^2(\Omega)^3$ in the corresponding variational formulation. To this end, we use the fact that any function $\mathbf{F} \in L^2(\Omega)^3$ admits the orthogonal decomposition

$$\mathbf{F} = \mathbf{F}' + \mathbf{F}'', \quad (2.6)$$

where $\mathbf{F}' \in L^2(\Omega)^3$ is such that $\nabla \cdot \mathbf{F}'|_{\Omega_0} = 0$ and $\langle (\mathbf{F}'|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = 0$ for $j = 1, \dots, J$, while \mathbf{F}'' satisfies $\mathbf{F}''|_{\Omega_\sigma} = \mathbf{0}$ and $\mathbf{F}''|_{\Omega_0} = \nabla f$, with $f \in H^1(\Omega_0)$, $f = 0$ on $\Gamma_{0,0}$ and f constant, say $f = f_j$, on each $\Gamma_{0,j}$ for $j = 1, \dots, J$. This is a consequence of the decomposition in [3, Section 3.5, point (iii)].

We consider the following problem: find $\mathbf{u} \in H(\text{curl}; \Omega) \cap H(\text{div}; \Omega_0)$ such that

$$\nabla \times (\mu^{-1}\nabla \times \mathbf{u}) + i\omega\sigma\mathbf{u} = \mathbf{F}' \quad \text{in } \Omega \quad (2.7)$$

$$\nu\nabla \cdot \mathbf{u} = -f \quad \text{in } \Omega_0 \quad (2.8)$$

$$\mathbf{n} \times \mathbf{u} = \mathbf{g} \quad \text{on } \partial\Omega \quad (2.9)$$

$$\lambda \langle (\mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = f_j \quad \forall j = 1, \dots, J, \quad (2.10)$$

where λ is any positive constant, whose purpose, again, is to achieve the correct physical dimensions. Notice that for $\mathbf{F} = \mathbf{J}$, we have $\mathbf{F}' = \mathbf{J}$ and $f = 0$ in the decomposition (2.6), and problem (2.7)–(2.10) reduces to (2.1)–(2.4).

We point out that in the LDG discretization of problem (2.7)–(2.10) that will be introduced in the following section, we need to compute neither the elements \mathbf{F}' and f in the decomposition of \mathbf{F} , nor the constants f_j . The only data that enter explicitly the formulation of the method are \mathbf{F} and \mathbf{g} . This is due to the variational character of the method and the particular choice of the inhomogeneous flux conditions in (2.10); see Remark 3.3 below.

We define the space $\mathbf{V} = H(\text{curl}; \Omega) \cap H(\text{div}; \Omega_0)$, endowed with the norm

$$\|\mathbf{u}\|_{\mathbf{V}}^2 = |\omega| \|\sigma_\vartheta^{\frac{1}{2}} \mathbf{u}\|_{0,\Omega}^2 + \|\mu^{-\frac{1}{2}} \nabla \times \mathbf{u}\|_{0,\Omega}^2 + \|\nu^{\frac{1}{2}} \nabla \cdot \mathbf{u}\|_{0,\Omega_0}^2 + \lambda \sum_{j=1}^J |\langle (\mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}}|^2,$$

with $\sigma_\vartheta = \sigma$ in Ω_σ and $\sigma_\vartheta = \vartheta I$ in Ω_0 , where I is the identity and ϑ is a fixed positive dimensional constant.

The variational formulation corresponding to (2.7)–(2.10) is as follows: find $\mathbf{u} \in \mathbf{V}$ such that $\mathbf{n} \times \mathbf{u} = \mathbf{g}$ on $\partial\Omega$, $\lambda \langle (\mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = f_j$ for $j = 1, \dots, J$, and

$$\int_{\Omega} \mu^{-1} \nabla \times \mathbf{u} \cdot \nabla \times \bar{\mathbf{v}} \, d\mathbf{x} + i\omega \int_{\Omega} \sigma \mathbf{u} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\Omega_0} \nu \nabla \cdot \mathbf{u} \nabla \cdot \bar{\mathbf{v}} \, d\mathbf{x} = \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x}, \quad (2.11)$$

for all $\mathbf{v} \in \mathbf{V}$, with $\mathbf{n} \times \mathbf{v} = \mathbf{0}$ on $\partial\Omega$ and $\langle (\mathbf{v}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = 0$, $j = 1, \dots, J$.

Well-posedness of the above formulation is established in the following theorem.

THEOREM 2.3. *For any $\mathbf{F} \in L^2(\Omega)^3$ and $\mathbf{g} \in \mathcal{H}(\partial\Omega)$, the variational formulation (2.11) admits a unique solution and there exists a positive constant C such that*

$$\|\mathbf{u}\|_{\mathbf{V}} \leq C (\|\mathbf{F}\|_{0,\Omega} + \|\mathbf{g}\|_{\mathcal{H}(\partial\Omega)}).$$

Moreover, \mathbf{u} is solution to problem (2.7)–(2.10) if and only if \mathbf{u} is solution to (2.11).

In the case where the domain is such that $H(\text{curl}; \Omega_0) \cap H_0(\text{div}; \Omega_0) \hookrightarrow H^1(\Omega_0)^3$ and the problem is driven by boundary conditions only, this result has been proved in [1]. The extension to our more general case is rather technical and will be given in all details in the appendix. One of the key ingredients necessary to prove Theorem 2.3 is to construct, under our assumptions on the domain, a continuous lifting of tangential traces with zero divergence in Ω_0 and zero flux conditions through $\Gamma_{0,j}$, $j = 1 \dots, J$. We do this in Proposition A.1, by using trace theorems recently proved in [14] and [15], and extended in [13] to domains with non-simply connected boundaries.

3. The local discontinuous Galerkin method. In this section, we formulate the LDG method for the discretization of problem (2.7)–(2.10). We assume from now on that

$$\mathbf{g} \in L^2(\partial\Omega)^3. \quad (3.1)$$

3.1. Traces and discontinuous finite element spaces. We start by introducing certain trace operators and finite element spaces used in the definition of the method. Let \mathcal{T}_h be a shape regular triangulation of the domain Ω into tetrahedra and/or parallelepipeds, with possible hanging nodes and aligned with the interfaces between different materials, so that μ , μ^{-1} and σ are smooth within each element of \mathcal{T}_h . We set $\mathcal{T}_h^0 := \mathcal{T}_h|_{\Omega_0}$ and have $\bar{\Omega} = \cup_{K \in \mathcal{T}_h} \bar{K}$ and $\bar{\Omega}_0 = \cup_{K \in \mathcal{T}_h^0} \bar{K}$. We will denote by h_K the diameter of the element $K \in \mathcal{T}_h$.

Faces. We define and characterize the *faces* of the triangulation \mathcal{T}_h . An *interior face* of \mathcal{T}_h is defined as the (non-empty) two-dimensional interior of $\partial K^+ \cap \partial K^-$, where K^+ and K^- are two adjacent elements of \mathcal{T}_h , not necessarily matching. A *boundary face* of \mathcal{T}_h is defined as the (non-empty) two-dimensional interior of $\partial K \cap \partial\Omega$, where K is a boundary element of \mathcal{T}_h . We denote by $\mathcal{E}_{\mathcal{I}}$ the union of all interior faces of \mathcal{T}_h , by $\mathcal{E}_{\mathcal{D}}$ the union of all the boundary faces of \mathcal{T}_h , and by $\mathcal{E} = \mathcal{E}_{\mathcal{I}} \cup \mathcal{E}_{\mathcal{D}}$ the union of all faces of \mathcal{T}_h . Similarly, we denote by \mathcal{E}^0 the union of all faces of \mathcal{T}_h^0 , and we set $\mathcal{E}_{\mathcal{I}}^0$ and $\mathcal{E}_{\mathcal{D}}^0$ for the interior and boundary faces of \mathcal{T}_h^0 .

Traces. Let $H^s(\mathcal{T}_h) := \{v : v|_K \in H^s(K), K \in \mathcal{T}_h\}$ for $s > \frac{1}{2}$, endowed with the norm $\|v\|_{s,\mathcal{T}_h}^2 = \sum_{K \in \mathcal{T}_h} \|v\|_{s,K}^2$. Then, the elementwise traces of functions in $H^s(\mathcal{T}_h)$ belong to $\text{TR}(\mathcal{E}) := \prod_{K \in \mathcal{T}_h} L^2(\partial K)$; they are double-valued on $\mathcal{E}_{\mathcal{I}}$ and single-valued on $\mathcal{E}_{\mathcal{D}}$. The space $L^2(\mathcal{E})$ can be identified with the functions in $\text{TR}(\mathcal{E})$ for which the two trace values coincide. We define similarly $H^s(\mathcal{T}_h^0)$, $\text{TR}(\mathcal{E}^0)$ and $L^2(\mathcal{E}^0)$.

Trace operators. Let us introduce the following trace operators for piecewise smooth functions. First, let $\mathbf{v} \in \text{TR}(\mathcal{E})^3$ and $e \in \mathcal{E}$. If e is an interior face in $\mathcal{E}_{\mathcal{I}}$, we denote by K_1 and K_2 the elements sharing e , by \mathbf{n}_i the normal unit vector pointing exterior to K_i , and we set $\mathbf{v}_i = \mathbf{v}|_{\partial K_i}$, $i = 1, 2$. We define the *average* and *tangential jump* of \mathbf{v} at $\mathbf{x} \in e$ as

$$\{\!\!\{ \mathbf{v} \}\!\!\} = \begin{cases} \frac{1}{2}(\mathbf{v}_1 + \mathbf{v}_2) & \text{if } e \in \mathcal{E}_{\mathcal{I}} \\ \mathbf{v} & \text{if } e \in \mathcal{E}_{\mathcal{D}} \end{cases} \quad \llbracket \mathbf{v} \rrbracket_T = \begin{cases} \mathbf{n}_1 \times \mathbf{v}_1 + \mathbf{n}_2 \times \mathbf{v}_2 & \text{if } e \in \mathcal{E}_{\mathcal{I}} \\ \mathbf{n} \times \mathbf{v} & \text{if } e \in \mathcal{E}_{\mathcal{D}}, \end{cases}$$

and, if $e \in \mathcal{E}_{\mathcal{I}}^0$, the *normal jump* of \mathbf{v} at $\mathbf{x} \in e$ as

$$\llbracket \mathbf{v} \rrbracket_N = \mathbf{v}_1 \cdot \mathbf{n}_1 + \mathbf{v}_2 \cdot \mathbf{n}_2 \quad \text{if } e \in \mathcal{E}_{\mathcal{I}}^0.$$

The normal jump of \mathbf{v} will not be used on faces outside \mathcal{T}_h^0 , and thus is left undefined. Similarly, we define for $\psi \in \text{TR}(\mathcal{E}^0)$ the *average* and *jump* at $\mathbf{x} \in e$ as

$$\{\!\!\{ \psi \}\!\!\} = \begin{cases} \frac{1}{2}(\psi_1 + \psi_2) & \text{if } e \in \mathcal{E}_{\mathcal{I}}^0 \\ \psi & \text{if } e \in \mathcal{E}_{\partial}^0 \end{cases} \quad \llbracket \psi \rrbracket = \begin{cases} \psi_1 \mathbf{n}_1 + \psi_2 \mathbf{n}_2 & \text{if } e \in \mathcal{E}_{\mathcal{I}}^0 \\ \psi \mathbf{n}_0 & \text{if } e \in \mathcal{E}_{\partial}^0, \end{cases}$$

where we recall that \mathbf{n}_0 denotes the outward normal unit to $\partial\Omega_0$. Note that the averages and jumps above defined are single-valued functions.

If $\mathbf{v} \in H(\text{curl}; \Omega)$, then, for all $e \in \mathcal{E}_{\mathcal{I}}$, the jump condition $\mathbf{n}_1 \times \mathbf{v}_1 + \mathbf{n}_2 \times \mathbf{v}_2 = \mathbf{0}$ holds true in $H_{00}^{-\frac{1}{2}}(e)^3$, and thus also in $L^2(e)^3$. For the definition of $H_{00}^{-\frac{1}{2}}(e)$, see, e.g., [35]. Therefore $\llbracket \mathbf{v} \rrbracket_T$ is well-defined and equal to zero on $\mathcal{E}_{\mathcal{I}}$. Similarly, for $\mathbf{v} \in H(\text{div}; \Omega_0)$, we have that $\llbracket \mathbf{v} \rrbracket_N$ is well-defined and equal to zero on $\mathcal{E}_{\mathcal{I}}^0$. Furthermore, for the exact solution $\mathbf{u} \in \mathbf{V}$, owing to assumption (3.1), we have for a boundary face $e \in \mathcal{E}_{\mathcal{D}}$ that $\llbracket \mathbf{u} \rrbracket_T = \mathbf{g}$ in $L^2(e)^3$, in addition to $\llbracket \mathbf{u} \rrbracket_T = \mathbf{0}$ on $\mathcal{E}_{\mathcal{I}}$ and $\llbracket \mathbf{u} \rrbracket_N = 0$ on $\mathcal{E}_{\mathcal{I}}^0$.

Finite element spaces. Let $\underline{p} = \{p_K\}_{K \in \mathcal{T}_h}$ be a degree vector that assigns to each element $K \in \mathcal{T}_h$ a polynomial approximation order $p_K \geq 1$. The generic hp -finite element space of piecewise polynomials is then given by

$$S^{\underline{p}, 0}(\mathcal{T}_h) := \{u \in L^2(\Omega) : u|_K \in S^{p_K}(K), \forall K \in \mathcal{T}_h\},$$

where $S^{p_K}(K)$ is the space $\mathcal{P}^{p_K}(K)$ of complex polynomials of degree at most p_K in K , if K is a tetrahedron, and the space $\mathcal{Q}^{p_K}(K)$ of complex polynomials of degree at most p_K in each variable in K , if K is a parallelepiped. The superscript 0 indicates that $S^{\underline{p}, 0}(\mathcal{T}_h) \subset L^2(\Omega) = H^0(\Omega)$. We define $S^{\underline{p}, 0}(\mathcal{T}_h^0)$ similarly.

3.2. Derivation of the LDG method. We introduce the auxiliary variables

$$\mathbf{s} = \mu^{-1} \mathbf{w} \quad \mathbf{w} = \nabla \times \mathbf{u} \quad \text{in } \Omega \quad (3.2)$$

$$\varphi = \nu \rho \quad \rho = \nabla \cdot \mathbf{u} \quad \text{in } \Omega_0. \quad (3.3)$$

Notice that $\mathbf{s} \in H(\text{curl}; \Omega)$, $\mathbf{w} \in L^2(\Omega)^3$, $\varphi \in H^1(\Omega_0)$ and $\rho \in L^2(\Omega_0)$. By subtracting the gradient of equation (2.8) from equation (2.7), taking into account the above identities and that $\mathbf{F} = \mathbf{F}'$ in Ω_σ and $\mathbf{F} = \mathbf{F}' + \nabla f$ in Ω_0 , we obtain

$$\nabla \times \mathbf{s} + i\omega\sigma\mathbf{u} - \widetilde{\nabla}\varphi = \mathbf{F} \quad \text{in } \Omega, \quad (3.4)$$

where $\widetilde{\nabla}\varphi = \nabla\varphi$ in Ω_0 and $\widetilde{\nabla}\varphi = \mathbf{0}$ in Ω_σ .

The LDG method is obtained by discretizing the first order equations in (3.2)–(3.4) in a discontinuous way. Notice that \mathbf{s} is actually related to the magnetic field phasor given by $i\omega^{-1}\mu^{-1}\nabla \times \mathbf{u}$. In this context, however, \mathbf{s} , \mathbf{w} , φ and ρ are auxiliary variables introduced in order to derive the method and will be eliminated from the equations *locally* in an element by element manner. This local solvability gives the name to the LDG method. We refer to [16] and [17] for a discussion of this elimination process from the computational point of view.

Since the LDG method is defined elementwise, we fix $K \in \mathcal{T}_h$ and, in order to account for the fact that the term $\widehat{\nabla}\varphi$ is different from zero in the elements contained in Ω_0 only, we set $K_0 = K_0(K) = K$, if $K \subset \Omega_0$, and $K_0 = K_0(K) = \emptyset$, if $K \subset \Omega_\sigma$. We proceed formally by multiplying in K the first identities in (3.2) and (3.3) by test functions \mathbf{z} and τ , the second identities in (3.2) and (3.3) by test functions \mathbf{t} and ψ , and equation (3.4) by a test function \mathbf{v} . By integration by parts and varying $K \in \mathcal{T}_h$, we obtain the following weak formulation:

$$\begin{aligned}
\int_K \mu^{-1} \mathbf{w} \cdot \bar{\mathbf{z}} \, d\mathbf{x} &= \int_K \mathbf{s} \cdot \bar{\mathbf{z}} \, d\mathbf{x} \\
\int_{K_0} \nu \rho \bar{\tau} \, d\mathbf{x} &= \int_{K_0} \varphi \bar{\tau} \, d\mathbf{x} \\
\int_K \mathbf{w} \cdot \bar{\mathbf{t}} \, d\mathbf{x} &= \int_K \mathbf{u} \cdot \nabla \times \bar{\mathbf{t}} \, d\mathbf{x} - \int_{\partial K} \mathbf{u} \cdot \mathbf{n}_K \times \bar{\mathbf{t}} \, ds \\
\int_{K_0} \rho \bar{\psi} \, d\mathbf{x} &= - \int_{K_0} \mathbf{u} \cdot \nabla \bar{\psi} \, d\mathbf{x} + \int_{\partial K} \mathbf{u} \cdot (\bar{\psi} \mathbf{n}_K) \, ds \\
\int_K \mathbf{s} \cdot \nabla \times \bar{\mathbf{v}} \, d\mathbf{x} &- \int_{\partial K} \mathbf{s} \cdot \mathbf{n}_K \times \bar{\mathbf{v}} \, ds + i\omega \int_K \sigma \mathbf{u} \cdot \bar{\mathbf{v}} \, d\mathbf{x} \\
&+ \int_{K_0} \varphi \nabla \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\partial K_0} \varphi \bar{\mathbf{v}} \cdot \mathbf{n}_K \, ds = \int_K \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x},
\end{aligned} \tag{3.5}$$

for any $K \in \mathcal{T}_h$, where \mathbf{n}_K is the outward normal unit vector to ∂K . The boundary integrals in (3.5) have to be understood as duality pairings.

We approximate $(\mathbf{w}, \rho, \mathbf{s}, \varphi, \mathbf{u})$ in (3.5) by functions $(\mathbf{w}_h, \rho_h, \mathbf{s}_h, \varphi_h, \mathbf{u}_h)$ in the hp -finite element space $\mathbf{W}_h \times M_h \times \Sigma_h \times Q_h \times \mathbf{V}_h$ chosen as

$$\mathbf{W}_h = \Sigma_h = \mathbf{V}_h = S^{\underline{p},0}(\mathcal{T}_h)^3 \quad M_h = Q_h = S^{\underline{p},0}(\mathcal{T}_h^0), \tag{3.6}$$

for a given degree distribution \underline{p} . Note that this choice actually implies that, for all $\mathbf{v} \in \mathbf{V}_h$, $\nabla_h \times \mathbf{v} \in \Sigma_h = \mathbf{W}_h$ and $\nabla_h \cdot \mathbf{v}|_{\Omega_0} \in Q_h = M_h$, where $\nabla_h \times$ and $\nabla_h \cdot$ denote the elementwise curl and divergence operators, respectively.

The discrete version of (3.5) then reads as follows: find $(\mathbf{w}_h, \rho_h, \mathbf{s}_h, \varphi_h, \mathbf{u}_h) \in \mathbf{W}_h \times M_h \times \Sigma_h \times Q_h \times \mathbf{V}_h$ such that, for any $K \in \mathcal{T}_h$ and for any choice of test

functions $(\mathbf{z}, \tau, \mathbf{t}, \psi, \mathbf{v}) \in \mathbf{W}_h \times M_h \times \Sigma_h \times Q_h \times \mathbf{V}_h$, we have

$$\begin{aligned}
\int_K \mu^{-1} \mathbf{w}_h \cdot \bar{\mathbf{z}} \, d\mathbf{x} &= \int_K \mathbf{s}_h \cdot \bar{\mathbf{z}} \, d\mathbf{x} \\
\int_{K_0} \nu \rho_h \bar{\tau} \, d\mathbf{x} &= \int_{K_0} \varphi_h \bar{\tau} \, d\mathbf{x} \\
\int_K \mathbf{w}_h \cdot \bar{\mathbf{t}} \, d\mathbf{x} &= \int_K \mathbf{u}_h \cdot \nabla \times \bar{\mathbf{t}} \, d\mathbf{x} - \int_{\partial K} \hat{\mathbf{u}}_h \cdot \mathbf{n}_K \times \bar{\mathbf{t}} \, ds \\
\int_{K_0} \rho_h \bar{\psi} \, d\mathbf{x} &= - \int_{K_0} \mathbf{u}_h \cdot \nabla \bar{\psi} \, d\mathbf{x} + \int_{\partial K_0} \hat{\hat{\mathbf{u}}}_h \cdot (\bar{\psi} \mathbf{n}_K) \, ds \\
\int_K \mathbf{s}_h \cdot \nabla \times \bar{\mathbf{v}} \, d\mathbf{x} &- \int_{\partial K} \hat{\mathbf{s}}_h \cdot \mathbf{n}_K \times \bar{\mathbf{v}} \, ds + i\omega \int_K \sigma \mathbf{u}_h \cdot \bar{\mathbf{v}} \, d\mathbf{x} \\
&+ \int_{K_0} \varphi_h \nabla \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\partial K_0} \hat{\hat{\varphi}}_h \bar{\mathbf{v}} \cdot \mathbf{n}_K \, ds = \int_K \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x}.
\end{aligned} \tag{3.7}$$

Here, $\hat{\mathbf{u}}_h$, $\hat{\mathbf{s}}_h$, $\hat{\hat{\mathbf{u}}}_h$ and $\hat{\hat{\varphi}}_h$ denote the so-called *numerical fluxes* which are approximations to the traces of \mathbf{u} , \mathbf{s} and φ on ∂K . They are crucial for the stability as well as for the accuracy of the method and will be defined in the next section. The fluxes $\hat{\mathbf{u}}_h$ and $\hat{\mathbf{s}}_h$ are related to the curl-curl operator, whereas the fluxes $\hat{\hat{\mathbf{u}}}_h$ and $\hat{\hat{\varphi}}_h$ are associated with the grad-div operator in Ω_0 .

REMARK 3.1. *If μ and ν are piecewise constant, the auxiliary variables \mathbf{w} and ρ are not needed, and the method can be defined by introducing directly $\mathbf{s} = \mu^{-1} \nabla \times \mathbf{u}$ and $\varphi = \nu \nabla \cdot \mathbf{u}$.*

3.3. The LDG numerical fluxes. Similar to [5], we understand the numerical fluxes as follows. Given \mathbf{u} and \mathbf{s} in $H^s(\mathcal{T}_h)^3$ for $s > \frac{1}{2}$, the fluxes $\hat{\mathbf{u}} = \hat{\mathbf{u}}(\mathbf{u})$ and $\hat{\mathbf{s}} = \hat{\mathbf{s}}(\mathbf{s}, \mathbf{u})$ belong to $L^2(\mathcal{E})^3$. Similarly, for $\mathbf{u}|_{\Omega_0} \in H^s(\mathcal{T}_h^0)^3$ and $\varphi \in H^s(\mathcal{T}_h^0)$, $\hat{\hat{\mathbf{u}}} = \hat{\hat{\mathbf{u}}}(\mathbf{u}|_{\Omega_0})$ belongs to $L^2(\mathcal{E}^0)^3$ and $\hat{\hat{\varphi}} = \hat{\hat{\varphi}}(\varphi, \mathbf{u}|_{\Omega_0})$ to $L^2(\mathcal{E}^0)$. The fluxes are thus single-valued on the union of faces. Furthermore, the fluxes in \mathbf{u} are assumed to be independent of the auxiliary variables in order to be able to eliminate them from the system of equations.

We define the fluxes $\hat{\mathbf{s}}$ and $\hat{\mathbf{u}}$ face by face by adapting to the curl-curl operator the numerical fluxes considered in [18] and [23] for the Laplacian:

$$\begin{aligned}
\hat{\mathbf{s}} &= \begin{cases} \{\{\mathbf{s}\}\} - \mathbf{a}[\mathbf{u}]_T + \mathbf{b}[\mathbf{s}]_T & \text{if } e \in \mathcal{E}_I \\ \mathbf{s} - \mathbf{a}(\mathbf{n} \times \mathbf{u} - \mathbf{g}) & \text{if } e \in \mathcal{E}_D \end{cases} \\
\hat{\mathbf{u}} &= \begin{cases} \{\{\mathbf{u}\}\} + \mathbf{b}[\mathbf{u}]_T & \text{if } e \in \mathcal{E}_I \\ \mathbf{g} & \text{if } e \in \mathcal{E}_D. \end{cases}
\end{aligned}$$

We use a similar recipe for the grad-div fluxes and set

$$\begin{aligned}
\hat{\hat{\varphi}} &= \begin{cases} \{\{\varphi\}\} - \mathbf{c}[\mathbf{u}]_N + \mathbf{d} \cdot \llbracket \varphi \rrbracket & \text{if } e \in \mathcal{E}_I^0 \\ -\lambda \langle \mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} & \text{if } e \in \Gamma_{0,j} \quad j = 1, \dots, J \\ 0 & \text{if } e \in \Gamma_{0,0} \end{cases} \\
\hat{\hat{\mathbf{u}}} &= \begin{cases} \{\{\mathbf{u}\}\} - \mathbf{d}[\mathbf{u}]_N & \text{if } e \in \mathcal{E}_I^0 \\ \mathbf{u} & \text{if } e \in \mathcal{E}_D^0. \end{cases}
\end{aligned}$$

Here, $\mathbf{a} \in L^\infty(\mathcal{E})$, $\mathbf{b} \in L^\infty(\mathcal{E}_I)$, $\mathbf{c} \in L^\infty(\mathcal{E}_I^0)$ and $\mathbf{d} \in L^\infty(\mathcal{E}_I^0)^3$ are real valued functions still at our disposal. This completes the definition of the LDG method.

Let us make some comments about these fluxes.

- The fluxes introduced above are *conservative* in the sense of [5], and give rise to a consistent formulation (see Theorem 3.2 below).

- The parameters \mathbf{a} and \mathbf{c} are referred to as *discontinuity stabilization parameters*. They have to be positive and will be chosen depending on the local meshsize, polynomial degree, and on the coefficients μ and ν . The parameters \mathbf{b} and \mathbf{d} , on the other hand, are chosen of order one; their purpose is to enhance the accuracy in the approximation of the auxiliary variables \mathbf{s} and φ that might be computed in a post-processing step. Indeed, in [23] it has been shown for the Laplacian that a parameter like \mathbf{b} and \mathbf{d} can be selected in such a way that the auxiliary variable superconverges on Cartesian grids.

- The numerical flux $\widehat{\mathbf{u}}$ enforces the boundary condition (2.3) in a weak sense. Namely, for any $\mathbf{u} \in H^s(\mathcal{T}_h)^3$, we have that

$$\mathbf{n} \times \widehat{\mathbf{u}} = \mathbf{g} \quad \text{on } \mathcal{E}_D. \quad (3.8)$$

The flux $\widehat{\varphi}$ imposes the condition $\varphi = 0$ on $\Gamma_{0,0}$ and $\varphi = -\lambda(\mathbf{u}_h|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1)_{\Gamma_{0,j}}$ on $\Gamma_{0,j}$, $j = 1, \dots, J$. Since for the exact solution $\lambda(\mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1)_{\Gamma_{0,j}} = f_j$ on $\Gamma_{0,j}$, the flux $\widehat{\varphi}$ approximates the boundary condition $\varphi = -f_j$ on $\Gamma_{0,j}$. This is the reason why the constants f_j do not appear explicitly in the formulation; see also Remark 3.3 below.

- Since the trace on $\Gamma_{0,j}$ of a function $\mathbf{v} \in H^s(\mathcal{T}_h^0)^3$ with $s > \frac{1}{2}$ actually belongs to $L^2(\Gamma_{0,j})^3$, we have that $\langle \mathbf{v}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} = \int_{\Gamma_{0,j}} \mathbf{v}|_{\Omega_0} \cdot \mathbf{n}_{0,j} ds$, $j = 1, \dots, J$.

3.4. The mixed formulation of the LDG method. In this section, we cast the LDG method in a mixed form, as in [18], and prove existence and uniqueness of discrete solutions. To do this, we sum the equations in (3.7) over all elements. Using integration by parts, the identities

$$\begin{aligned} \sum_{K \in \mathcal{T}_h} \int_{\partial K} \mathbf{v} \cdot \mathbf{n}_K \times \bar{\mathbf{t}} ds &= - \sum_{K \in \mathcal{T}_h} \int_{\partial K} \bar{\mathbf{t}} \cdot \mathbf{n}_K \times \mathbf{v} ds \\ &= - \int_{\mathcal{E}} [[\mathbf{v}]]_T \cdot \{\{\bar{\mathbf{t}}\}\} ds + \int_{\mathcal{E}_I} \{\{\mathbf{v}\}\} \cdot [[\bar{\mathbf{t}}]]_T ds \\ \sum_{K \in \mathcal{T}_h^0} \int_{\partial K} \mathbf{w} \cdot (\bar{\psi} \mathbf{n}_K) ds &= \int_{\mathcal{E}_I^0} (\{\{\mathbf{w}\}\} \cdot [[\bar{\psi}]] + [[\mathbf{w}]]_N \{\{\bar{\psi}\}\}) ds + \int_{\mathcal{E}_\theta^0} \mathbf{w} \cdot (\bar{\psi} \mathbf{n}_\theta) ds \end{aligned} \quad (3.9)$$

that hold true for all $\mathbf{v}, \mathbf{t} \in \text{TR}(\mathcal{E})^3$, $\mathbf{w} \in \text{TR}(\mathcal{E}^0)^3$ and $\psi \in \text{TR}(\mathcal{E}^0)$, as well as the form of the numerical fluxes, we obtain the following formulation.

Mixed formulation. Find $(\mathbf{w}_h, \rho_h, \mathbf{s}_h, \varphi_h, \mathbf{u}_h) \in \mathbf{W}_h \times M_h \times \Sigma_h \times Q_h \times \mathbf{V}_h$ such that

$$\begin{aligned}
\int_{\Omega} \mu^{-1} \mathbf{w}_h \cdot \bar{\mathbf{z}} \, d\mathbf{x} &= \int_{\Omega} \mathbf{s}_h \cdot \bar{\mathbf{z}} \, d\mathbf{x} \\
\int_{\Omega_0} \nu \rho_h \bar{\tau} \, d\mathbf{x} &= \int_{\Omega_0} \varphi_h \bar{\tau} \, d\mathbf{x} \\
\int_{\Omega} \mathbf{w}_h \cdot \bar{\mathbf{t}} \, d\mathbf{x} &= \int_{\Omega} \nabla_h \times \mathbf{u}_h \cdot \bar{\mathbf{t}} \, d\mathbf{x} - \int_{\mathcal{E}_T} \mathbf{b}[\mathbf{u}_h]_T \cdot [\bar{\mathbf{t}}]_T \, ds \\
&\quad - \int_{\mathcal{E}} [\mathbf{u}_h]_T \cdot \{\{\bar{\mathbf{t}}\}\} \, ds + \int_{\mathcal{E}_D} \mathbf{g} \cdot \bar{\mathbf{t}} \, ds \\
\int_{\Omega_0} \rho_h \bar{\psi} \, d\mathbf{x} &= \int_{\Omega_0} \nabla_h \cdot \mathbf{u}_h \bar{\psi} \, d\mathbf{x} - \int_{\mathcal{E}_T^0} \mathbf{d}[\mathbf{u}_h]_N \cdot [\bar{\psi}] \, ds - \int_{\mathcal{E}_T^0} [\mathbf{u}_h]_N \{\{\bar{\psi}\}\} \, ds \\
\int_{\Omega} \mathbf{s}_h \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} &- \int_{\mathcal{E}} \{\{\mathbf{s}_h\}\} \cdot [\bar{\mathbf{v}}]_T \, ds - \int_{\mathcal{E}_T} \mathbf{b}[\mathbf{s}_h]_T \cdot [\bar{\mathbf{v}}]_T \, ds + \int_{\mathcal{E}} \mathbf{a}[\mathbf{u}_h]_T \cdot [\bar{\mathbf{v}}]_T \, ds \\
&+ i\omega \int_{\Omega} \sigma \mathbf{u}_h \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\Omega_0} \varphi_h \nabla_h \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\mathcal{E}_T^0} \{\{\varphi_h\}\} [\bar{\mathbf{v}}]_N \, ds - \int_{\mathcal{E}_T^0} \mathbf{d} \cdot [\varphi_h] [\bar{\mathbf{v}}]_N \, ds \\
&+ \int_{\mathcal{E}_T^0} \mathbf{c}[\mathbf{u}_h]_N [\bar{\mathbf{v}}]_N \, ds + \lambda \sum_{j=1}^J \langle \mathbf{u}_h|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} \langle \bar{\mathbf{v}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} \\
&= \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\mathcal{E}_D} \mathbf{a} \mathbf{g} \cdot (\mathbf{n} \times \bar{\mathbf{v}}) \, ds,
\end{aligned} \tag{3.10}$$

for all $(\mathbf{z}, \tau, \mathbf{t}, \psi, \mathbf{v}) \in \mathbf{W}_h \times M_h \times \Sigma_h \times Q_h \times \mathbf{V}_h$.

We prove existence and uniqueness of solutions and consistency of (3.10) in the following theorem. Notice that, in order to have consistency, we do not need any smoothness assumption on the exact solution in addition to (3.1).

THEOREM 3.2. *For strictly positive discontinuity stabilization parameters \mathbf{a} and \mathbf{c} , the LDG method defines a unique approximate solution $(\mathbf{w}_h, \rho_h, \mathbf{s}_h, \varphi_h, \mathbf{u}_h)$ in the space $\mathbf{W}_h \times M_h \times \Sigma_h \times Q_h \times \mathbf{V}_h$. Furthermore, the LDG formulation (3.10) is consistent, i.e., the exact solution $(\mathbf{w}, \rho, \mathbf{s}, \varphi, \mathbf{u})$ satisfies (3.10), for all test functions $(\mathbf{z}, \tau, \mathbf{t}, \psi, \mathbf{v}) \in \mathbf{W}_h \times M_h \times \Sigma_h \times Q_h \times \mathbf{V}_h$.*

Proof. Since problem (3.10) is linear and finite dimensional, in order to prove existence and uniqueness of solutions, it is sufficient to prove that if $\mathbf{F} = \mathbf{0}$ and $\mathbf{g} = \mathbf{0}$, then $\mathbf{w}_h = \mathbf{s}_h = \mathbf{u}_h = \mathbf{0}$ and $\rho_h = \varphi_h = 0$. Taking $(\mathbf{z}, \tau, \mathbf{t}, \psi, \mathbf{v}) = (\mathbf{w}_h, \rho_h, \mathbf{s}_h, \varphi_h, \mathbf{u}_h)$ in (3.10), subtracting the first and the second equations from the third and the fourth ones, respectively, and then subtracting the results from the fifth equation, we obtain

$$\begin{aligned}
\int_{\Omega} \mu^{-1} \mathbf{w}_h^2 \, d\mathbf{x} &+ \int_{\mathcal{E}} \mathbf{a} [\mathbf{u}_h]_T^2 \, ds + i\omega \int_{\Omega} \sigma \mathbf{u}_h^2 \, d\mathbf{x} + \int_{\Omega_0} \nu \rho_h^2 \, d\mathbf{x} \\
&+ \int_{\mathcal{E}_T^0} \mathbf{c} [\mathbf{u}_h]_N^2 \, ds + \lambda \sum_{j=1}^J \langle \mathbf{u}_h|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}}^2 = 0.
\end{aligned}$$

Taking into account that μ^{-1} is positive definite in Ω and ν is positive in Ω_0 , we have $\mathbf{w}_h = \mathbf{0}$ in Ω and $\rho_h = 0$ in Ω_0 ; since σ is positive definite in Ω_σ , then $\mathbf{u}_h = \mathbf{0}$ in Ω_σ , and since $\mathbf{a} > 0$, $\mathbf{c} > 0$ and $\lambda > 0$, then $[\mathbf{u}_h]_T = \mathbf{0}$ on \mathcal{E} , $[\mathbf{u}_h]_N = 0$ on \mathcal{E}_T^0 and

$\langle \mathbf{u}_h|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} = 0$, $j = 1, \dots, J$. Now, since $\Sigma_h = \mathbf{W}_h$ and $Q_h = M_h$, taking \mathbf{s}_h and φ_h as test functions in the first and second equations of (3.10), respectively, from $\mathbf{w}_h = \mathbf{0}$ and $\rho_h = 0$, we have $\mathbf{s}_h = \mathbf{0}$ in Ω and $\varphi_h = 0$ in Ω_σ . Then, the third equation reduces to $\int_\Omega \nabla_h \times \mathbf{u}_h \cdot \bar{\mathbf{t}} \, d\mathbf{x} = 0$, for all $\mathbf{t} \in \Sigma_h$. Since $\nabla_h \times \mathbf{V}_h \subseteq \Sigma_h$, we have $\nabla_h \times \mathbf{u}_h = \mathbf{0}$ in Ω . Similarly, the fourth equation becomes $\int_{\Omega_0} \nabla_h \cdot \mathbf{u}_h \bar{\psi} \, d\mathbf{x} = 0$, for all $\psi \in Q_h$. From $\nabla_h \cdot \mathbf{V}_h|_{\Omega_0} \subseteq Q_h$, we have $\nabla_h \cdot \mathbf{u}_h = 0$ in Ω_0 . From $\mathbf{u}_h = \mathbf{0}$ in Ω_σ and $[\mathbf{u}_h]_T = \mathbf{0}$ on \mathcal{E} , we get $\mathbf{n}_0 \times \mathbf{u}_h = \mathbf{0}$ on $\partial\Omega_0$. We can summarize the above conditions on \mathbf{u}_h in Ω_0 as $\mathbf{u}_h|_{\Omega_0} \in H_0(\text{curl}^0, \text{div}^0; \Omega_0)$ and $\langle \mathbf{u}_h|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} = 0$, $j = 1, \dots, J$. This implies that $\mathbf{u}_h = \mathbf{0}$ also in Ω_0 (see [30], formula (4.14) with $\Gamma_\tau = \partial\Omega_0$, $\Gamma_\nu = \emptyset$ and weight $\omega = 1$). This concludes the proof of the first part of the theorem.

As far as the consistency of the formulation (3.10) is concerned, let $(\mathbf{w}, \rho, \mathbf{s}, \varphi, \mathbf{u})$ be the exact solution. From $\mathbf{s} = \mu^{-1} \mathbf{w}$ and $\varphi = \nu \rho$, it is obvious that the first two equations are fulfilled, for any $\mathbf{z} \in \mathbf{W}_h$ and $\tau \in M_h$. Since $[\mathbf{u}]_T = \mathbf{0}$ on \mathcal{E}_I and $[\mathbf{u}]_T = \mathbf{g} \in L^2(\mathcal{E}_D)^3$ on \mathcal{E}_D , due to (3.1), taking into account that $\mathbf{w} = \nabla \times \mathbf{u}$, we have that the third equation is satisfied by \mathbf{w} and \mathbf{u} , for all $\mathbf{t} \in \Sigma_h$. Similarly, since $[\mathbf{u}]_N = 0$ on \mathcal{E}_I^0 , taking into account that $\varphi = \nabla \cdot \mathbf{u}$, we have that the fourth equation is satisfied by φ and \mathbf{u} , for all $\psi \in Q_h$. Finally, consider the fifth equation. Express the term $\int_{\mathcal{E}} \{\{\mathbf{s}_h\}\} \cdot [\bar{\mathbf{v}}]_T \, ds$ by $-\sum_{K \in \mathcal{T}_h} \int_{\partial K} \bar{\mathbf{v}} \cdot \mathbf{n}_K \times \mathbf{s}_h \, ds + \int_{\mathcal{E}_I} \{\{\bar{\mathbf{v}}\}\} \cdot [\mathbf{s}_h]_T \, ds$, according to (3.9), and substitute the exact solution into the obtained equation. Understanding the boundary integral as duality pairings, integrating by parts and observing that $\mathbf{u} \in \mathbf{V}$, $\mathbf{s} \in H(\text{curl}; \Omega)$ and $\varphi \in H^1(\Omega_0)$, together with the definition of $\{\{\varphi\}\}$, we get

$$\begin{aligned} & \int_\Omega \nabla \times \mathbf{s} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + i\omega \int_\Omega \sigma \mathbf{u} \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\Omega_0} \nabla \varphi \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\mathcal{E}_D^0} \varphi \bar{\mathbf{v}} \cdot \mathbf{n} \, ds \\ & + \lambda \sum_{j=1}^J \langle \mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} \langle \bar{\mathbf{v}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} = \int_\Omega \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x}. \end{aligned}$$

From (3.4) and the flux conditions (2.10), we obtain

$$\int_{\mathcal{E}_D^0} \varphi \bar{\mathbf{v}} \cdot \mathbf{n} \, ds + \sum_{j=1}^J f_j \langle \bar{\mathbf{v}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} = 0,$$

which is satisfied because $\varphi|_{\partial\Omega_0} = (\nu \nabla \cdot \mathbf{u})|_{\partial\Omega_0} = -f|_{\partial\Omega_0}$, and f is zero on $\Gamma_{0,0}$ and constant f_j on $\Gamma_{0,j}$, $j = 1, \dots, J$. This completes the proof of the theorem. \square

REMARK 3.3. *The constants f_j do not appear explicitly in the LDG formulation (3.10). As can be inferred from the proof of Theorem 3.2, this is due to the particular choice of the flux conditions in (2.10), whose purpose is, in fact, to cancel the terms containing the constants f_j , since they are not easily computable from the datum \mathbf{F} . If we consider problem (2.7)–(2.10) with more general flux conditions*

$$\lambda \langle (\mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = \alpha_j,$$

for given constants α_j , $j = 1, \dots, J$, the numerical flux $\widehat{\varphi}$ on the faces belonging to $\Gamma_{0,j}$, $j = 1, \dots, J$, must be adjusted accordingly by

$$\widehat{\varphi} = (\alpha_j - f_j) - \lambda \langle \mathbf{u}_h|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}}.$$

Consequently, the right-hand side in the last equation of (3.10) becomes

$$\int_\Omega \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\mathcal{E}_D} \mathbf{a} \mathbf{g} \cdot (\mathbf{n} \times \bar{\mathbf{v}}) \, ds + \sum_{j=1}^J (\alpha_j - f_j) \langle \bar{\mathbf{v}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}}.$$

3.5. The primal formulation of the LDG method. In this subsection, we eliminate the auxiliary variables \mathbf{w} , \mathbf{s} , ρ and φ from the mixed system in (3.10) and derive the primal formulation of the LDG method. This is possible since the fluxes $\widehat{\mathbf{u}}$ and $\widehat{\mathbf{u}}$ are chosen independently of \mathbf{s} and φ .

Let us start by introducing the lifting operators $\mathcal{L}_1 : L^2(\mathcal{E}_T)^3 \rightarrow \Sigma_h$, $\mathcal{L}_2 : L^2(\mathcal{E})^3 \rightarrow \Sigma_h$, $\mathcal{M}_1 : L^2(\mathcal{E}_T^0)^3 \rightarrow Q_h$ and $\mathcal{M}_2 : L^2(\mathcal{E}_T^0) \rightarrow Q_h$ defined by

$$\begin{aligned} \int_{\Omega} \mathcal{L}_1(\mathbf{v}) \cdot \bar{\mathbf{t}} \, d\mathbf{x} &= \int_{\mathcal{E}_T} \mathbf{v} \cdot [\bar{\mathbf{t}}]_T \, ds & \int_{\Omega} \mathcal{L}_2(\mathbf{v}) \cdot \bar{\mathbf{t}} \, d\mathbf{x} &= \int_{\mathcal{E}} \mathbf{v} \cdot \{\{\bar{\mathbf{t}}\}\} \, ds & \forall \mathbf{t} \in \Sigma_h, \\ \int_{\Omega_0} \mathcal{M}_1(\mathbf{v}) \bar{\psi} \, d\mathbf{x} &= \int_{\mathcal{E}_T^0} \mathbf{v} \cdot [\bar{\psi}] \, ds & \int_{\Omega_0} \mathcal{M}_2(v) \bar{\psi} \, d\mathbf{x} &= \int_{\mathcal{E}_T^0} v \{\{\bar{\psi}\}\} \, ds & \forall \psi \in Q_h, \end{aligned}$$

as well as the lifting $\mathbf{G}_D \in \Sigma_h$ of the boundary datum given by

$$\int_{\Omega} \mathbf{G}_D \cdot \bar{\mathbf{t}} \, d\mathbf{x} = \int_{\mathcal{E}_D} \mathbf{g} \cdot \bar{\mathbf{t}} \, d\mathbf{x} \quad \forall \mathbf{t} \in \Sigma_h.$$

Denoting by Π_{Σ_h} and Π_{Q_h} the L^2 -projections onto $\mathbf{W}_h = \Sigma_h$ and $M_h = Q_h$, the first and second equation in (3.10) can then be written as $\mathbf{s}_h = \Pi_{\Sigma_h}(\mu^{-1}\mathbf{w}_h)$ and $\varphi_h = \Pi_{Q_h}(\nu\rho_h)$. Then, from the third and fourth equations in (3.10), we obtain

$$\mathbf{s}_h = \Pi_{\Sigma_h} [\mu^{-1}(\nabla_h \times \mathbf{u}_h - \mathcal{L}([\mathbf{u}_h]_T) + \mathbf{G}_D)], \quad (3.11)$$

$$\varphi_h = \Pi_{Q_h} [\nu(\nabla_h \cdot \mathbf{u}_h - \mathcal{M}([\mathbf{u}_h]_N))], \quad (3.12)$$

where we used the compact notation $\mathcal{L}([\mathbf{u}_h]_T) := \mathcal{L}_1(\mathbf{b}[\mathbf{u}_h]_T) + \mathcal{L}_2([\mathbf{u}_h]_T)$, with $\mathbf{b}[\mathbf{u}_h]_T$ understood as restricted to \mathcal{E}_T , and $\mathcal{M}([\mathbf{u}_h]_N) := \mathcal{M}_1(\mathbf{d}[\mathbf{u}_h]_N) + \mathcal{M}_2([\mathbf{u}_h]_N)$. Since $\nabla_h \times \mathbf{V}_h \subseteq \Sigma_h$ and $\nabla_h \cdot \mathbf{V}_h|_{\Omega_0} \subseteq Q_h$, identities (3.11) and (3.12) can be used in the fifth equation of (3.10), giving rise to the so-called primal formulation of the LDG discretization of (2.7)–(2.10), in the variable \mathbf{u} only.

Primal formulation. Find $\mathbf{u}_h \in \mathbf{V}_h$ such that, for all $\mathbf{v} \in \mathbf{V}_h$,

$$\mathcal{B}_h(\mathbf{u}_h, \mathbf{v}) := \mathcal{A}_h(\mathbf{u}_h, \mathbf{v}) + \mathcal{I}_h(\mathbf{u}_h, \mathbf{v}) + i\omega \int_{\Omega} \sigma \mathbf{u}_h \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \mathcal{J}(\mathbf{u}_h, \mathbf{v}) = \mathcal{F}_h(\mathbf{v}), \quad (3.13)$$

where the forms \mathcal{A}_h , \mathcal{I}_h (*interior penalty* form) and \mathcal{J} are defined by

$$\begin{aligned} \mathcal{A}_h(\mathbf{u}, \mathbf{v}) &= \int_{\Omega} \mu^{-1}(\nabla_h \times \mathbf{u} - \mathcal{L}([\mathbf{u}]_T)) \cdot (\nabla_h \times \bar{\mathbf{v}} - \mathcal{L}([\bar{\mathbf{v}}]_T)) \, d\mathbf{x} \\ &\quad + \int_{\Omega_0} \nu(\nabla_h \cdot \mathbf{u} - \mathcal{M}([\mathbf{u}]_N)) (\nabla_h \cdot \bar{\mathbf{v}} - \mathcal{M}([\bar{\mathbf{v}}]_N)) \, d\mathbf{x} \\ \mathcal{I}_h(\mathbf{u}, \mathbf{v}) &= \int_{\mathcal{E}} \mathbf{a}[\mathbf{u}]_T \cdot [\bar{\mathbf{v}}]_T \, ds + \int_{\mathcal{E}_T^0} \mathbf{c}[\mathbf{u}]_N \cdot [\bar{\mathbf{v}}]_N \, ds \\ \mathcal{J}(\mathbf{u}, \mathbf{v}) &= \lambda \sum_{j=1}^J \langle \mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, \mathbf{1} \rangle_{\Gamma_{0,j}} \langle \bar{\mathbf{v}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, \mathbf{1} \rangle_{\Gamma_{0,j}}, \end{aligned}$$

and the linear form \mathcal{F}_h by

$$\mathcal{F}_h(\mathbf{v}) = \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\Omega} \mu^{-1} \mathbf{G}_D \cdot (\nabla_h \times \bar{\mathbf{v}} - \mathcal{L}([\bar{\mathbf{v}}]_T)) \, d\mathbf{x} + \int_{\mathcal{E}_D} \mathbf{a} \mathbf{g} \cdot (\mathbf{n} \times \bar{\mathbf{v}}) \, ds.$$

For *discrete* test and trial functions, the primal form (3.13) of the LDG method is equivalent to the mixed system (3.10). However, unlike (3.10), the formulation (3.13) is no longer consistent, due to the discrete nature of the lifting operators. Nevertheless, the form $\mathcal{B}_h(\cdot, \cdot)$ has the continuity and coercivity properties that allow us to carry out an error analysis in a straightforward way by using Strang's lemma. Regarding this point, our approach differs from the analysis in [5].

REMARK 3.4. *Other DG methods can be obtained by modifying the definitions of $\mathcal{A}_h(\mathbf{u}, \mathbf{v})$, $\mathcal{I}_h(\mathbf{u}, \mathbf{v})$ and $\mathcal{F}_h(\mathbf{v})$ in (3.13). The interior penalty (IP) method and its nonsymmetric variant (NIP), for instance, can be obtained by taking in (3.13) the same $\mathcal{I}_h(\mathbf{u}, \mathbf{v})$ as in the LDG method, and instead of $\mathcal{A}_h(\mathbf{u}, \mathbf{v})$ and $\mathcal{F}_h(\mathbf{v})$,*

$$\begin{aligned} \mathcal{A}_h^{\text{IP}}(\mathbf{u}, \mathbf{v}) &= \int_{\Omega} \mu^{-1} \nabla_h \times \mathbf{u} \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\mathcal{E}} \llbracket \mathbf{u} \rrbracket_T \cdot \{\mu^{-1} \nabla_h \times \bar{\mathbf{v}}\} \, ds \\ &\quad - \int_{\mathcal{E}} \llbracket \bar{\mathbf{v}} \rrbracket_T \cdot \{\mu^{-1} \nabla_h \times \mathbf{u}\} \, ds + \int_{\Omega_0} \nu \nabla_h \cdot \mathbf{u} \nabla_h \cdot \bar{\mathbf{v}} \, d\mathbf{x} \\ &\quad - \int_{\mathcal{E}_T^0} \llbracket \mathbf{u} \rrbracket_N \{\nu \nabla_h \cdot \bar{\mathbf{v}}\} \, ds - \int_{\mathcal{E}_T^0} \llbracket \bar{\mathbf{v}} \rrbracket_N \{\nu \nabla_h \cdot \mathbf{u}\} \, ds \\ \mathcal{F}_h^{\text{IP}}(\mathbf{v}) &= \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\Omega} \mu^{-1} (\mathbf{n} \times \mathbf{g}) \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\mathcal{E}_D} \mathbf{a} \mathbf{g} \cdot (\mathbf{n} \times \bar{\mathbf{v}}) \, ds, \end{aligned}$$

and

$$\begin{aligned} \mathcal{A}_h^{\text{NIP}}(\mathbf{u}, \mathbf{v}) &= \int_{\Omega} \mu^{-1} \nabla_h \times \mathbf{u} \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\mathcal{E}} \llbracket \mathbf{u} \rrbracket_T \cdot \{\mu^{-1} \nabla_h \times \bar{\mathbf{v}}\} \, ds \\ &\quad - \int_{\mathcal{E}} \llbracket \bar{\mathbf{v}} \rrbracket_T \cdot \{\mu^{-1} \nabla_h \times \mathbf{u}\} \, ds + \int_{\Omega_0} \nu \nabla_h \cdot \mathbf{u} \nabla_h \cdot \bar{\mathbf{v}} \, d\mathbf{x} \\ &\quad + \int_{\mathcal{E}_T^0} \llbracket \mathbf{u} \rrbracket_N \{\nu \nabla_h \cdot \bar{\mathbf{v}}\} \, ds - \int_{\mathcal{E}_T^0} \llbracket \bar{\mathbf{v}} \rrbracket_N \{\nu \nabla_h \cdot \mathbf{u}\} \, ds \\ \mathcal{F}_h^{\text{NIP}}(\mathbf{v}) &= \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\Omega} \mu^{-1} (\mathbf{n} \times \mathbf{g}) \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\mathcal{E}_D} \mathbf{a} \mathbf{g} \cdot (\mathbf{n} \times \bar{\mathbf{v}}) \, ds, \end{aligned}$$

or, in terms of the lifting operators \mathcal{L}_2 and \mathcal{M}_2 ,

$$\begin{aligned} \mathcal{A}_h^{\text{IP}}(\mathbf{u}, \mathbf{v}) &= \int_{\Omega} \mu^{-1} (\nabla_h \times \mathbf{u} \cdot \nabla_h \times \bar{\mathbf{v}} - \mathcal{L}_2(\llbracket \mathbf{u} \rrbracket_T) \cdot \nabla_h \times \bar{\mathbf{v}} - \mathcal{L}_2(\llbracket \bar{\mathbf{v}} \rrbracket_T) \cdot \nabla_h \times \mathbf{u}) \, d\mathbf{x} \\ &\quad + \int_{\Omega_0} \nu (\nabla_h \cdot \mathbf{u} \nabla_h \cdot \bar{\mathbf{v}} - \mathcal{M}_2(\llbracket \mathbf{u} \rrbracket_N) \nabla_h \cdot \bar{\mathbf{v}} - \mathcal{M}_2(\llbracket \bar{\mathbf{v}} \rrbracket_N) \nabla_h \cdot \mathbf{u}) \, d\mathbf{x} \\ \mathcal{F}_h^{\text{IP}}(\mathbf{v}) &= \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\Omega} \mu^{-1} \mathbf{G}_D \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\mathcal{E}_D} \mathbf{a} \mathbf{g} \cdot (\mathbf{n} \times \bar{\mathbf{v}}) \, ds, \end{aligned}$$

and

$$\begin{aligned} \mathcal{A}_h^{\text{NIP}}(\mathbf{u}, \mathbf{v}) &= \int_{\Omega} \mu^{-1} (\nabla_h \times \mathbf{u} \cdot \nabla_h \times \bar{\mathbf{v}} + \mathcal{L}_2(\llbracket \mathbf{u} \rrbracket_T) \cdot \nabla_h \times \bar{\mathbf{v}} - \mathcal{L}_2(\llbracket \bar{\mathbf{v}} \rrbracket_T) \cdot \nabla_h \times \mathbf{u}) \, d\mathbf{x} \\ &\quad + \int_{\Omega_0} \nu (\nabla_h \cdot \mathbf{u} \nabla_h \cdot \bar{\mathbf{v}} + \mathcal{M}_2(\llbracket \mathbf{u} \rrbracket_N) \nabla_h \cdot \bar{\mathbf{v}} - \mathcal{M}_2(\llbracket \bar{\mathbf{v}} \rrbracket_N) \nabla_h \cdot \mathbf{u}) \, d\mathbf{x} \\ \mathcal{F}_h^{\text{NIP}}(\mathbf{v}) &= \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\Omega} \mu^{-1} \mathbf{G}_D \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\mathcal{E}_D} \mathbf{a} \mathbf{g} \cdot (\mathbf{n} \times \bar{\mathbf{v}}) \, ds. \end{aligned}$$

These formulations can also be derived by using the same mixed formulation as for the LDG method, and defining appropriately the numerical fluxes, see [5]. The analysis of

the IP and NIP methods is almost identical to the one of the LDG method presented in the next section and the obtained error estimates hold true verbatim also for these methods. The only difference consists in a restriction on the choice of the stabilization parameters in the IP method, as pointed out in Remark 4.6 below. We refer to [5] and [17] for an extensive discussion and comparison of different DG methods, from a theoretical and a computational point of view.

4. Error analysis. The aim of this section is to present an hp -error analysis of the LDG method introduced in section 3, based on its primal formulation (3.13). Although we use the same setting of [5], our analysis differs from the one presented there since we directly work on the discrete form (3.13), taking into account non-consistency terms by Strang's lemma. This approach in the analysis of DG methods seems to be new and might be of interest on its own.

The main result (see Theorem 4.12 below) consists in error estimates, in a suitable energy-norm, of the form

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}_h\|_h^2 &\leq C \sum_{K \in \mathcal{T}_h} \frac{h_K^{2 \min(p_K, s_K)}}{p_K^{2s_K-1}} (\|\mathbf{u}\|_{s_K+1, K}^2 + \|\mu^{-1} \nabla \times \mathbf{u}\|_{s_K, K}^2) \\ &\quad + C \sum_{K \in \mathcal{T}_h^0} \frac{h_K^{2 \min(p_K, s_K)}}{p_K^{2s_K-1}} \|\nu \nabla \cdot \mathbf{u}\|_{s_K, K}^2, \end{aligned}$$

for exact solutions \mathbf{u} that satisfy $\mathbf{u} \in H^{s_K+1}(K)^3$, $\mu^{-1} \nabla \times \mathbf{u} \in H^{s_K}(K)^3$, for all $K \in \mathcal{T}_h$, and $\nu \nabla \cdot \mathbf{u} \in H^{s_K}(K)$, for all $K \in \mathcal{T}_h^0$, with local regularity exponents $s_K \geq 1$. These estimates are optimal in the local meshsizes h_K and slightly suboptimal in the local approximation degree p_K . Furthermore, in Theorem 4.12, we also make explicit the dependence on the local material properties.

The outline of this section is as follows. In section 4.1, we define the discontinuity stabilization parameters \mathbf{a} and \mathbf{c} in terms of the local meshsize, approximation degree and magnetic permeability. Section 4.2 is devoted to establish hp -stability estimates for the lifting operators \mathcal{L} and \mathcal{M} in the definition of the primal formulation. These estimates will be crucial in section 4.3 where we prove continuity and coercivity properties of the bilinear form $\mathcal{B}_h(\cdot, \cdot)$. Based on Strang's lemma, we derive hp -error estimates in section 4.4, and in section 4.5 we recover error estimates for the auxiliary variables \mathbf{s} and φ used in the derivation of the LDG method. Recall that the variable \mathbf{s} is related to the magnetic field, and therefore its computation might be of interest. We conclude in section 4.6 by investigating the stability of the discrete problem with respect to the data.

4.1. The discontinuity stabilization parameters. In this section, we define the discontinuity stabilization parameters \mathbf{a} and \mathbf{c} in terms of the "local meshsize", "local polynomial degree" and "local magnetic permeability". This allows us to obtain continuity and coercivity constants independent of global bounds for these quantities.

Let us start by introducing the functions \mathbf{h} and \mathbf{p} in $L^\infty(\mathcal{E})$, related to the local meshsize and polynomial degree, defined as

$$\mathbf{h} = \mathbf{h}(\mathbf{x}) := \begin{cases} \min\{h_K, h_{K'}\} & \text{if } \mathbf{x} \text{ in the interior of } \partial K \cap \partial K' \\ h_K & \text{if } \mathbf{x} \text{ in the interior of } \partial K \cap \partial \Omega \end{cases}$$

$$\mathbf{p} = \mathbf{p}(\mathbf{x}) := \begin{cases} \max\{p_K, p_{K'}\} & \text{if } \mathbf{x} \text{ in the interior of } \partial K \cap \partial K' \\ p_K & \text{if } \mathbf{x} \text{ in the interior of } \partial K \cap \partial \Omega. \end{cases}$$

Regarding the magnetic permeability, we assume μ to be Lipschitz continuous in K , for any $K \in \mathcal{T}_h$. This implies that $\mu|_K$ can be extended up to ∂K , and we denote this extension by μ_K . Therefore, for any $K \in \mathcal{T}_h$, there are positive constants m_K and M_K such that

$$m_K \leq \lambda_i(\mu_K(\mathbf{x})) \leq M_K \quad \forall \mathbf{x} \in \bar{K}, \quad (4.1)$$

where $\lambda_i(\mu_K(\mathbf{x}))$, $i = 1, 2, 3$, are the eigenvalues of $\mu_K(\mathbf{x})$. For any $K \in \mathcal{T}_h$, the constants m_K and M_K satisfy $0 < m \leq m_K$ and $M_K \leq M < +\infty$, where m is the uniform ellipticity constant of μ and M is the reciprocal of the uniform ellipticity constant of μ^{-1} .

We choose the scalar function ν in the formulation of the problem as $\nu(\mathbf{x}) = 1/|\mu(\mathbf{x})|$, for all $\mathbf{x} \in \Omega_0$, where $|\mu(\mathbf{x})|$ is the spectral norm of the tensor $\mu(\mathbf{x})$ ($|\mu(\mathbf{x})|$ simply reduces to $\mu(\mathbf{x})$ whenever μ is a scalar function). Then we also have that ν satisfies

$$\frac{1}{M_K} \leq \nu_K(\mathbf{x}) \leq \frac{1}{m_K} \quad \forall \mathbf{x} \in \bar{K},$$

for any $K \in \mathcal{T}_h^0$, where we have defined ν_K in the same way as μ_K .

We make the additional assumption that there exists $\kappa > 0$ such that

$$\frac{M_K}{m_K} \leq \kappa \quad \forall K \in \mathcal{T}_h, \forall \mathcal{T}_h. \quad (4.2)$$

Whenever μ is a piecewise constant scalar function, (4.2) holds true with $\kappa = 1$. For μ piecewise constant tensor, κ in (4.2) expresses the maximum anisotropy among the different materials. We set

$$\mathbf{m} = \mathbf{m}(\mathbf{x}) := \begin{cases} \min\{|\mu_K(\mathbf{x})|, |\mu_{K'}(\mathbf{x})|\} & \text{if } \mathbf{x} \text{ is in the interior of } \partial K \cap \partial K' \\ |\mu_K(\mathbf{x})| & \text{if } \mathbf{x} \text{ is in the interior of } \partial K \cap \partial \Omega. \end{cases}$$

We are now ready to define the discontinuity stabilization parameters \mathbf{a} and \mathbf{c} in terms of \mathbf{h} , \mathbf{p} and \mathbf{m} . They are chosen as

$$\mathbf{a} = \alpha \mathbf{h}^{-1} \mathbf{p}^2 \mathbf{m}^{-1} \quad \text{in } L^\infty(\mathcal{E}) \quad \mathbf{c} = \alpha \mathbf{h}^{-1} \mathbf{p}^2 \mathbf{m}^{-1} \quad \text{in } L^\infty(\mathcal{E}_T^0), \quad (4.3)$$

with $\alpha > 0$ independent of the meshsize, approximation order and the magnetic permeability. The parameters \mathbf{b} and \mathbf{d} are taken to be of order one, i.e.,

$$\|\mathbf{b}\|_{\infty, \mathcal{E}_T} \leq \delta \quad \|\mathbf{d}\|_{\infty, \mathcal{E}_T^0} \leq \delta, \quad (4.4)$$

with $\delta > 0$ independent of h , p and μ .

REMARK 4.1. *The choice of the stabilization parameters of order p^2/h is the hp -extension of the choice in [5] for h -version DG methods for the Laplacian. This choice balances the interior penalty terms in $\mathcal{I}_h(\cdot, \cdot)$ with the stability estimates in the following Proposition 4.2 for the lifting operators \mathcal{L} and \mathcal{M} , or, equivalently, with the inverse estimate (4.5) below.*

Stabilization parameters of order p^2/h can also be found in the hp -literature on DG methods for diffusion problems, see, e.g., [34], [41] and [43], where different error analyzes are developed. The choice p/h is investigated in [34] for the NIP method, still leading to a suboptimal error bound in p . Furthermore, an optimal hp -bound has been recently obtained in [32] for two-dimensional reaction-diffusion problems on structured quadrilateral grids for solutions belonging to certain “augmented” Sobolev spaces. The same result can be established in our case, leading to hp -optimal bounds on structured grids, provided that the corresponding arguments can be extended to three space-dimensions.

4.2. The lifting operators. In this section, we derive hp -stability estimates for the lifting operators introduced in section 3.5. To do this, we define the space

$$\mathbf{V}(h) := \{ \mathbf{v} = \mathbf{w}_h + \mathbf{w} \mid \mathbf{w}_h \in \mathbf{V}_h, \mathbf{w} \in \mathbf{V} \text{ with } \mathbf{n} \times \mathbf{w} \in L^2(\partial\Omega)^3 \}.$$

Owing to (3.1), the exact solution \mathbf{u} belongs to $\mathbf{V}(h)$.

PROPOSITION 4.2. *Let \mathcal{L} and \mathcal{M} be the lifting operators defined in section 3.5. Under the above assumptions on μ and ν , and assumption (4.4) on the parameters \mathbf{b} and \mathbf{d} , we have that, for all $\mathbf{v} \in \mathbf{V}(h)$,*

$$\begin{aligned} \|\mu^{-\frac{1}{2}} \mathcal{L}(\llbracket \mathbf{v} \rrbracket_T)\|_{0,\Omega} &\leq C_{\text{lift}} \kappa (\delta + 1) \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} \llbracket \mathbf{v} \rrbracket_T\|_{0,\mathcal{E}} \\ \|\nu^{\frac{1}{2}} \mathcal{M}(\llbracket \mathbf{v} \rrbracket_N)\|_{0,\Omega_0} &\leq C_{\text{lift}} \kappa (\delta + 1) \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} \llbracket \mathbf{v} \rrbracket_N\|_{0,\mathcal{E}_T^0}, \end{aligned}$$

with a constant $C_{\text{lift}} > 0$ only depending on the shape regularity of the mesh. Moreover, for $\mathbf{G}_{\mathcal{D}}$ defined in section 3.5, we have

$$\|\mu^{-\frac{1}{2}} \mathbf{G}_{\mathcal{D}}\|_{0,\Omega} \leq C_{\text{lift}} \kappa \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} \mathbf{g}\|_{0,\mathcal{E}_{\mathcal{D}}}.$$

Proof. Let us first recall the following inverse inequality:

$$\|q\|_{0,\partial K}^2 \leq C_{\text{inv}} \frac{p_K^2}{h_K} \|q\|_{0,K}^2 \quad \forall q \in \mathcal{S}^{p_K}(K), \quad (4.5)$$

with a constant $C_{\text{inv}} > 0$ only depending on the shape regularity of the mesh. For two-dimensional elements, the proof of (4.5) can be found in [44, formula (4.6.4) of Theorem 4.76]; for three-space dimensions, the proof is analogous, see also [34].

From the definition of \mathcal{L} and \mathcal{M} in terms of \mathcal{L}_i and \mathcal{M}_i , $i = 1, 2$ (see section 3.5), the bounds for \mathcal{L} and \mathcal{M} can be proved by combining estimates for \mathcal{L}_i and \mathcal{M}_i , $i = 1, 2$. We develop in detail the proof of the following estimate for \mathcal{L}_1 :

$$\|\mu^{-\frac{1}{2}} \mathcal{L}_1(\mathbf{b} \llbracket \mathbf{v} \rrbracket_T)\|_{0,\Omega} \leq C_{\text{lift}} \kappa \delta \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} \llbracket \mathbf{v} \rrbracket_T\|_{0,\mathcal{E}_T} \quad (4.6)$$

Recall that, for $\mathbf{v} = \mathbf{w}_h + \mathbf{w} \in \mathbf{V}(h)$, we have $\llbracket \mathbf{v} \rrbracket_T = \llbracket \mathbf{w}_h \rrbracket_T$ on \mathcal{E}_T . Denoting by $\mathbf{\Pi}_{\Sigma_h}$ the L^2 -projection onto Σ_h , by the definition of the operator \mathcal{L}_1 and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \|\mu^{-\frac{1}{2}} \mathcal{L}_1(\mathbf{b} \llbracket \mathbf{v} \rrbracket_T)\|_{0,\Omega} &= \sup_{\mathbf{z} \in L^2(\Omega)^3} \frac{\int_{\Omega} \mathcal{L}_1(\mathbf{b} \llbracket \mathbf{v} \rrbracket_T) \cdot \mu^{-\frac{1}{2}} \bar{\mathbf{z}} \, d\mathbf{x}}{\|\mathbf{z}\|_{0,\Omega}} \\ &= \sup_{\mathbf{z} \in L^2(\Omega)^3} \frac{\int_{\Omega} \mathcal{L}_1(\mathbf{b} \llbracket \mathbf{v} \rrbracket_T) \cdot \mathbf{\Pi}_{\Sigma_h}(\mu^{-\frac{1}{2}} \bar{\mathbf{z}}) \, d\mathbf{x}}{\|\mathbf{z}\|_{0,\Omega}} \\ &= \sup_{\mathbf{z} \in L^2(\Omega)^3} \frac{\int_{\mathcal{E}_T} \mathbf{b} \llbracket \mathbf{v} \rrbracket_T \cdot \llbracket \mathbf{\Pi}_{\Sigma_h}(\mu^{-\frac{1}{2}} \bar{\mathbf{z}}) \rrbracket_T \, ds}{\|\mathbf{z}\|_{0,\Omega}} \\ &\leq \delta \sup_{\mathbf{z} \in L^2(\Omega)^3} \frac{\|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} \llbracket \mathbf{v} \rrbracket_T\|_{0,\mathcal{E}_T} \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} \llbracket \mathbf{\Pi}_{\Sigma_h}(\mu^{-\frac{1}{2}} \bar{\mathbf{z}}) \rrbracket_T\|_{0,\mathcal{E}_T}}{\|\mathbf{z}\|_{0,\Omega}}. \end{aligned}$$

Then, by using conditions (4.1) on μ , the definitions of $\llbracket \cdot \rrbracket_T$, \mathbf{m} , \mathbf{h} and \mathbf{p} , the inverse

inequality (4.5), and properties of the L^2 -projection, we obtain

$$\begin{aligned} \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} [\mathbf{\Pi}_{\Sigma_h}(\mu^{-\frac{1}{2}} \mathbf{z})]_T\|_{0, \mathcal{E}_x}^2 &\leq 2 \sum_{K \in \mathcal{T}_h} \frac{h_K M_K}{p_K^2} \|\mathbf{n}_K \times \mathbf{\Pi}_{\Sigma_h}(\mu^{-\frac{1}{2}} \mathbf{z})\|_{0, \partial K}^2 \\ &\leq 2 C_{\text{inv}} \sum_{K \in \mathcal{T}_h} M_K \|\mathbf{\Pi}_{\Sigma_h}(\mu^{-\frac{1}{2}} \mathbf{z})\|_{0, K}^2 \leq 2 C_{\text{inv}} \sum_{K \in \mathcal{T}_h} M_K \|\mu^{-\frac{1}{2}} \mathbf{z}\|_{0, K}^2 \\ &\leq 2 C_{\text{inv}} \sum_{K \in \mathcal{T}_h} \frac{M_K}{m_K} \|\mathbf{z}\|_{0, K}^2 \leq 2 C_{\text{inv}} \kappa \|\mathbf{z}\|_{0, \Omega}^2, \end{aligned}$$

where in the last step we used (4.2). This proves the desired estimate for \mathcal{L}_1 in (4.6). Analogous estimates can be obtained for \mathcal{M}_1 and \mathcal{M}_2 , as well as for \mathcal{L}_2 , recalling that $[\mathbf{v}]_T \in L^2(\mathcal{E}_D)^3$ on \mathcal{E}_D . Then the bounds for \mathcal{L} and \mathcal{M} immediately follow. Since $\mathbf{G}_D = \mathcal{L}([\mathbf{u}]_T)$, the same arguments give the bound for \mathbf{G}_D . \square

4.3. Continuity, coercivity and error bound. In this section, we establish continuity and coercivity properties of the form B_h . To do this, we introduce the seminorm $|\cdot|_h$ given by

$$\begin{aligned} |\mathbf{v}|_h^2 &= \|\mu^{-\frac{1}{2}} \nabla_h \times \mathbf{v}\|_{0, \Omega}^2 + \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} [\mathbf{v}]_T\|_{0, \mathcal{E}}^2 \\ &\quad + \|\nu^{\frac{1}{2}} \nabla_h \cdot \mathbf{v}\|_{0, \Omega_0}^2 + \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} [\mathbf{v}]_N\|_{0, \mathcal{E}_D^0}^2, \end{aligned} \quad (4.7)$$

as well as the norm $\|\cdot\|_h$

$$\|\mathbf{v}\|_h^2 = |\mathbf{v}|_h^2 + |\omega| \|\sigma^{\frac{1}{2}} \mathbf{v}\|_{0, \Omega}^2 + \lambda \sum_{j=1}^J |\langle \mathbf{v}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}}|^2. \quad (4.8)$$

That (4.8) is actually a norm in $\mathbf{V}(h)$ is proved in the following proposition.

PROPOSITION 4.3. *The quantity defined in (4.8) is a norm in $\mathbf{V}(h)$.*

Proof. From $\|\mathbf{v}\|_h = 0$, we immediately have $\mathbf{v} = \mathbf{0}$ in Ω_σ , $[\mathbf{v}]_T = 0$ on \mathcal{E} and $[\mathbf{v}]_N = 0$ on \mathcal{E}_T^0 , i.e., $\mathbf{v} \in H_0(\text{curl}; \Omega) \cap H(\text{div}; \Omega_0)$. Now, from $\mathbf{v} = \mathbf{0}$ in Ω_σ and $\mathbf{v} \in H_0(\text{curl}; \Omega)$, it follows that $\mathbf{n}_0 \times \mathbf{v}|_{\Omega_\sigma} = \mathbf{0}$ on the interface $\Gamma = \partial\Omega_\sigma \cap \partial\Omega_0$, and therefore $\mathbf{n}_0 \times \mathbf{v}|_{\Omega_0} = \mathbf{0}$ on $\partial\Omega_0$. From $\nabla \times \mathbf{v} = \mathbf{0}$ in Ω , $\mathbf{n}_0 \times \mathbf{v} = \mathbf{0}$ on $\partial\Omega_0$, $\nabla \cdot \mathbf{v} = 0$ in Ω_0 and $\langle (\mathbf{v}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = 0$, $j = 1, \dots, J$, we get $\mathbf{v} = \mathbf{0}$ also in Ω_0 (see, e.g., formula (4.14) with $\Gamma_\tau = \partial\Omega_0$ and $\Gamma_\nu = \emptyset$ in [30]), which concludes the proof. \square

Let us first prove continuity and coercivity properties for the LDG forms in (3.13).

LEMMA 4.4. *Assume the above hypotheses on μ , ν and on the coefficients in the definition of the numerical fluxes. Then the following continuity property holds true:*

$$|\mathcal{A}_h(\mathbf{w}, \mathbf{v}) + \mathcal{I}_h(\mathbf{w}, \mathbf{v})| \leq C |\mathbf{w}|_h |\mathbf{v}|_h \quad \forall \mathbf{w}, \mathbf{v} \in \mathbf{V}(h),$$

with a constant C only depending on α , δ , κ , and C_{lift} .

Proof. For $\mathbf{w}, \mathbf{v} \in \mathbf{V}(h)$, we have

$$\begin{aligned} |\mathcal{A}_h(\mathbf{w}, \mathbf{v}) + \mathcal{I}_h(\mathbf{w}, \mathbf{v})| &\leq \|\mu^{-\frac{1}{2}} [\nabla_h \times \mathbf{w} - \mathcal{L}([\mathbf{w}]_T)]\|_{0, \Omega} \|\mu^{-\frac{1}{2}} [\nabla_h \times \mathbf{v} - \mathcal{L}([\mathbf{v}]_T)]\|_{0, \Omega} \\ &\quad + \|\nu^{\frac{1}{2}} [\nabla_h \cdot \mathbf{w} - \mathcal{M}([\mathbf{w}]_N)]\|_{0, \Omega_0} \|\nu^{\frac{1}{2}} [\nabla_h \cdot \mathbf{v} - \mathcal{M}([\mathbf{v}]_N)]\|_{0, \Omega_0} \\ &\quad + \alpha \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} [\mathbf{w}]_T\|_{0, \mathcal{E}} \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} [\mathbf{v}]_T\|_{0, \mathcal{E}} \\ &\quad + \alpha \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} [\mathbf{w}]_N\|_{0, \mathcal{E}_T^0} \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} [\mathbf{v}]_N\|_{0, \mathcal{E}_T^0}. \end{aligned}$$

From Proposition 4.2, we have $\|\mathcal{L}([\mathbf{z}]_T)\|_{0,\Omega} \leq C|\mathbf{z}|_h$ and $\|\mathcal{M}([\mathbf{z}]_N)\|_{0,\Omega_0} \leq C|\mathbf{z}|_h$, for $\mathbf{z} = \mathbf{w}$ and $\mathbf{z} = \mathbf{v}$, and the result immediately follows. \square

LEMMA 4.5. *Assume the above hypotheses on μ , ν and on the coefficients in the definition of the numerical fluxes. The coercivity property*

$$\mathcal{A}_h(\mathbf{v}, \mathbf{v}) + \mathcal{I}_h(\mathbf{v}, \mathbf{v}) \geq C|\mathbf{v}|_h^2 \quad \forall \mathbf{v} \in \mathbf{V}_h,$$

holds true for any choice of $\alpha > 0$. The constant C depends on α , δ , κ and C_{lift} .

Proof. We have

$$\begin{aligned} \mathcal{A}_h(\mathbf{v}, \mathbf{v}) + \mathcal{I}_h(\mathbf{v}, \mathbf{v}) &= \int_{\Omega} \{\mu^{-1}[\nabla_h \times \mathbf{v} - \mathcal{L}([\mathbf{v}]_T)]\}^2 dx \\ &\quad + \int_{\Omega_0} \{\nu[\nabla_h \cdot \mathbf{v} - \mathcal{M}([\mathbf{v}]_N)]\}^2 dx \\ &\quad + \alpha \|h^{-\frac{1}{2}} \text{pm}^{-\frac{1}{2}} [\mathbf{v}]_T\|_{0,\varepsilon}^2 + \alpha \|h^{-\frac{1}{2}} \text{pm}^{-\frac{1}{2}} [\mathbf{v}]_N\|_{0,\varepsilon_x^0}^2. \end{aligned}$$

The first term at right-hand side can be bounded by

$$\begin{aligned} &\|\mu^{-\frac{1}{2}} \nabla_h \times \mathbf{v}\|_{0,\Omega}^2 - 2 \int_{\Omega} \mu^{-1} \nabla_h \times \mathbf{v} \cdot [\mathcal{L}([\mathbf{v}]_T)] dx + \|\mu^{-\frac{1}{2}} [\mathcal{L}([\mathbf{v}]_T)]\|_{0,\Omega}^2 \\ &\geq (1 - \varepsilon) \|\mu^{-\frac{1}{2}} \nabla_h \times \mathbf{v}\|_{0,\Omega}^2 + \left(1 - \frac{1}{\varepsilon}\right) \|\mu^{-\frac{1}{2}} [\mathcal{L}([\mathbf{v}]_T)]\|_{0,\Omega}^2, \end{aligned}$$

with $\varepsilon > 0$ still at our disposal. Similarly, the second term at right-hand side can be bounded by

$$(1 - \varepsilon) \|\nu^{\frac{1}{2}} \nabla_h \cdot \mathbf{v}\|_{0,\Omega_0}^2 + \left(1 - \frac{1}{\varepsilon}\right) \|\nu^{\frac{1}{2}} [\mathcal{M}([\mathbf{v}]_N)]\|_{0,\Omega_0}^2.$$

Therefore, using the estimates of Proposition 4.2, and taking ε that satisfies the inequalities

$$\frac{C_{\text{lift}}^2 \kappa^2 (\delta + 1)^2}{C_{\text{lift}}^2 \kappa^2 (\delta + 1)^2 + \alpha} < \varepsilon < 1,$$

we obtain the result. \square

REMARK 4.6. *The continuity result of Lemma 4.4 holds true verbatim for the IP and NIP methods discussed in Remark 3.4. The coercivity property of Lemma 4.5, instead, is trivially satisfied for any choice of $\alpha > 0$ for the NIP method, but only under the condition $\alpha > C_{\text{lift}}^2 \kappa^2$ for the IP method.*

As already pointed out, the primal formulation (3.13) our analysis is based on is not consistent, due to the discrete nature of the lifting operators. However, we can obtain error bounds by using Strang's lemma.

PROPOSITION 4.7. *Assume the above hypotheses on μ , ν and on the coefficients in the definition of the numerical fluxes. The following continuity and coercivity properties hold true:*

$$\begin{aligned} |\mathcal{B}_h(\mathbf{w}, \mathbf{v})| &\leq C_{\text{cont}} \|\mathbf{w}\|_h \|\mathbf{v}\|_h & \forall \mathbf{w}, \mathbf{v} \in \mathbf{V}(h) \\ \mathcal{B}_h(\mathbf{v}, \mathbf{v}) &\geq C_{\text{coer}} \|\mathbf{v}\|_h^2 & \forall \mathbf{v} \in \mathbf{V}_h, \end{aligned}$$

with C_{cont} and C_{coer} only depending on α , δ , κ and C_{lift} .

Proof. Since $|\mathcal{B}_h(\mathbf{v}, \mathbf{v})| = [(\mathcal{A}_h(\mathbf{v}, \mathbf{v}) + \mathcal{I}_h(\mathbf{v}, \mathbf{v}) + \mathcal{J}(\mathbf{v}, \mathbf{v}))^2 + \omega^2 \|\sigma^{\frac{1}{2}} \mathbf{v}\|_{0,\Omega}^4]^{\frac{1}{2}}$, the continuity and coercivity properties follow from Lemma 4.4, Lemma 4.5 and the definition of the norm $\|\cdot\|_h$. \square

From Proposition 4.7 and from Strang's lemma (see, e.g., [21, Theorem 4.2.2]), we immediately have the following error bound.

THEOREM 4.8. *Assume the above hypotheses on μ , ν and on the coefficients in the definition of the numerical fluxes. Then we have*

$$\|\mathbf{u} - \mathbf{u}_h\|_h \leq \left(1 + \frac{C_{\text{cont}}}{C_{\text{coer}}}\right) \inf_{\mathbf{v} \in \mathbf{V}_h} \|\mathbf{u} - \mathbf{v}\|_h + \frac{1}{C_{\text{coer}}} \sup_{\mathbf{w} \in \mathbf{V}_h} \frac{|\mathcal{B}_h(\mathbf{u}, \mathbf{w}) - \mathcal{F}_h(\mathbf{w})|}{\|\mathbf{w}\|_h}.$$

REMARK 4.9. *In order to analyze stability properties of discrete solutions with respect to the data, the continuity of the functional $\mathcal{F}_h(\cdot)$ with respect to the norm $\|\cdot\|_h$ has to be investigated. This is not straightforward since $\|\cdot\|_h$ does not contain the L^2 -norm over Ω_0 . In section 4.6, we prove a discrete Poincaré inequality that allows us to address this issue in the particular case where μ is the identity. In the general case of discontinuous permeabilities, we obtain the same stability estimates provided that the datum \mathbf{F} satisfies certain restrictions. This point is also addressed in section 4.6.*

4.4. hp-error estimates. In this section, we estimate the terms at the right-hand side in the error bound established in Theorem 4.8 and derive a-priori error estimates for piecewise smooth solutions. In order to do that, we need the following hp-approximation result.

PROPOSITION 4.10. *Let $K \in \mathcal{T}_h$ and suppose that $u \in H^{t_K}(K)$, $t_K \geq 0$. Then there exists a sequence of polynomials $\pi_{p_K}^{h_K} u$ in $\mathcal{S}^{p_K}(K)$, $p_K = 1, 2, \dots$, satisfying*

$$\|u - \pi_{p_K}^{h_K} u\|_{q,K} \leq C \frac{h_K^{\min(p_K+1, t_K) - q}}{p_K^{t_K - q}} \|u\|_{t_K, K} \quad \forall 0 \leq q \leq t_K. \quad (4.9)$$

Furthermore, if $t_K \geq 1$,

$$\|u - \pi_{p_K}^{h_K} u\|_{0, \partial K} \leq C \frac{h_K^{\min(p_K+1, t_K) - \frac{1}{2}}}{p_K^{t_K - \frac{1}{2}}} \|u\|_{t_K, K}. \quad (4.10)$$

The constant C is independent of u , h_K and p_K , but depends on the shape regularity of the mesh and on $t = \max_{K \in \mathcal{T}_h} t_K$.

Proof. The assertion (4.9) has been proved in [6, Lemma 4.5] for two-dimensional domains. For three-dimensional domains, the proof is analogous, see also [34]. In order to prove (4.10), we use the multiplicative trace inequality (see, e.g., [41, Lemma A.3])

$$\|\eta\|_{0, \partial K}^2 \leq C (\|\eta\|_{0, K} \|\nabla \eta\|_{0, K} + h_K^{-1} \|\eta\|_{0, K}^2) \quad (4.11)$$

that holds true for any $\eta \in H^1(K)$ with a constant $C > 0$ only depending on the shape regularity of the mesh. The second assertion now follows by applying in (4.11) the approximation result (4.9) for $q = 0, 1$. \square

We will denote by Π_p^h the operator defined by $\Pi_p^h(u)|_K = \pi_{p_K}^{h_K}(u|_K)$, for any $K \in \mathcal{T}_h$, with $\pi_{p_K}^{h_K}(u|_K)$ as in Proposition 4.10, and by $\mathbf{\Pi}_p^h$ the operator that maps $\mathbf{u} = (u_1, u_2, u_3)$ into $(\Pi_p^h(u_1), \Pi_p^h(u_2), \Pi_p^h(u_3))$.

Next, we give an estimate of the residual $\mathcal{R}_h(\mathbf{u}, \mathbf{w}) := \mathcal{B}_h(\mathbf{u}, \mathbf{w}) - \mathcal{F}_h(\mathbf{w})$.

LEMMA 4.11. *Let \mathbf{u} be the exact solution. Assume $(\mu^{-1}\nabla \times \mathbf{u})|_K \in H^{s_K}(K)^3$, for all $K \in \mathcal{T}_h$, and $(\nu\nabla \cdot \mathbf{u})|_K \in H^{s_K}(K)$, for all $K \in \mathcal{T}_h^0$, with local regularity exponents $s_K \geq 1$. Then, for any $\mathbf{w} \in \mathbf{V}_h$, the following estimate holds true:*

$$\begin{aligned} |\mathcal{R}_h(\mathbf{u}, \mathbf{w})| &\leq C \left(\sum_{K \in \mathcal{T}_h} \frac{h_K^{2\min(p_K+1, s_K)} M_K}{p_K^{2s_K}} \|\mu^{-1}\nabla \times \mathbf{u}\|_{s_K, K}^2 \right)^{\frac{1}{2}} \|\mathbf{w}\|_h \\ &\quad + C \left(\sum_{K \in \mathcal{T}_h^0} \frac{h_K^{2\min(p_K+1, s_K)} M_K}{p_K^{2s_K}} \|\nu\nabla \cdot \mathbf{u}\|_{s_K, K}^2 \right)^{\frac{1}{2}} \|\mathbf{w}\|_h, \end{aligned}$$

where M_K are the constants in (4.1).

Proof. By straightforward calculations involving integration by parts, taking into account that $\mathbf{u} \in \mathbf{V}$, $\mathcal{L}([\mathbf{u}]_T) = \mathbf{G}_{\mathcal{D}}$, along with boundary and flux conditions and the characterization of the data, we have that, for any $\mathbf{w} \in \mathbf{V}_h$,

$$\begin{aligned} \mathcal{R}_h(\mathbf{u}, \mathbf{w}) &= \int_{\mathcal{E}} \{ \mu^{-1}\nabla \times \mathbf{u} \} \cdot [\bar{\mathbf{w}}]_T ds - \int_{\Omega} \mu^{-1}\nabla \times \mathbf{u} \cdot \mathcal{L}([\bar{\mathbf{w}}]_T) dx \\ &\quad + \int_{\mathcal{E}_T^0} \{ \nu\nabla \cdot \mathbf{u} \} [\bar{\mathbf{w}}]_N ds - \int_{\Omega_0} \nu\nabla \cdot \mathbf{u} \mathcal{M}([\bar{\mathbf{w}}]_N) dx. \end{aligned}$$

Since

$$\begin{aligned} \int_{\Omega} \mu^{-1}\nabla \times \mathbf{u} \cdot \mathcal{L}([\bar{\mathbf{w}}]_T) dx &= \int_{\Omega} \Pi_{\Sigma_h}(\mu^{-1}\nabla \times \mathbf{u}) \cdot \mathcal{L}([\bar{\mathbf{w}}]_T) dx, \\ \int_{\Omega_0} \nu\nabla \cdot \mathbf{u} \mathcal{M}([\bar{\mathbf{w}}]_N) dx &= \int_{\Omega_0} \Pi_{Q_h}(\nu\nabla \cdot \mathbf{u}) \mathcal{M}([\bar{\mathbf{w}}]_N) dx, \end{aligned} \tag{4.12}$$

for the L^2 -projections Π_{Σ_h} and Π_{Q_h} onto Σ_h and Q_h , respectively, we can write

$$\mathcal{R}_h(\mathbf{u}, \mathbf{w}) =: T_1 + T_2 + T_3 + T_4,$$

where

$$\begin{aligned} T_1 &= \int_{\mathcal{E}} \{ \mu^{-1}\nabla \times \mathbf{u} - \Pi_{\Sigma_h}(\mu^{-1}\nabla \times \mathbf{u}) \} \cdot [\bar{\mathbf{w}}]_T ds \\ T_2 &= \int_{\mathcal{E}_T} [\mu^{-1}\nabla \times \mathbf{u} - \Pi_{\Sigma_h}(\mu^{-1}\nabla \times \mathbf{u})]_T \cdot \mathbf{b}[\bar{\mathbf{w}}]_T ds \\ T_3 &= \int_{\mathcal{E}_T^0} \{ \nu\nabla \cdot \mathbf{u} - \Pi_{Q_h}(\nu\nabla \cdot \mathbf{u}) \} [\bar{\mathbf{w}}]_N ds \\ T_4 &= \int_{\mathcal{E}_T^0} [\nu\nabla \cdot \mathbf{u} - \Pi_{Q_h}(\nu\nabla \cdot \mathbf{u})] \cdot \mathbf{d}[\bar{\mathbf{w}}]_N ds. \end{aligned}$$

Let us bound the term T_1 ; the other terms are bounded similarly, observing our assumptions on \mathbf{b} and \mathbf{d} . By the Cauchy-Schwarz and triangle inequalities, and the definition of $\|\cdot\|_h$, we obtain the following bound:

$$\begin{aligned} T_1 &\leq \|\mathbf{w}\|_h (\|h^{\frac{1}{2}} p^{-1} m^{\frac{1}{2}} \{ \mu^{-1}\nabla \times \mathbf{u} - \Pi_p^h(\mu^{-1}\nabla \times \mathbf{u}) \} \|_{0, \mathcal{E}} \\ &\quad + \|h^{\frac{1}{2}} p^{-1} m^{\frac{1}{2}} \{ \Pi_p^h(\mu^{-1}\nabla \times \mathbf{u}) - \Pi_{\Sigma_h}(\mu^{-1}\nabla \times \mathbf{u}) \} \|_{0, \mathcal{E}}). \end{aligned}$$

From the definitions of \mathbf{h} , \mathbf{p} and \mathbf{m} , and (4.10) with $t_K = s_K$, we conclude that

$$\begin{aligned} & \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} \{\{\mu^{-1} \nabla \times \mathbf{u} - \mathbf{\Pi}_p^h(\mu^{-1} \nabla \times \mathbf{u})\}\}\|_{0,\mathcal{E}}^2 \\ & \leq C \sum_{K \in \mathcal{T}_h} \frac{h_K M_K}{p_K^2} \|\mu^{-1} \nabla \times \mathbf{u} - \mathbf{\Pi}_p^h(\mu^{-1} \nabla \times \mathbf{u})\|_{0,\partial K}^2 \\ & \leq C \sum_{K \in \mathcal{T}_h} \frac{h_K^{2 \min(p_K+1, s_K)} M_K}{p_K^{2s_K+1}} \|\mu^{-1} \nabla \times \mathbf{u}\|_{s_K, K}^2, \end{aligned}$$

and, similarly,

$$\begin{aligned} & \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} \{\{\mathbf{\Pi}_p^h(\mu^{-1} \nabla \times \mathbf{u}) - \mathbf{\Pi}_{\Sigma_h}(\mu^{-1} \nabla \times \mathbf{u})\}\}\|_{0,\mathcal{E}}^2 \\ & \leq C \sum_{K \in \mathcal{T}_h} \frac{h_K M_K}{p_K^2} \|\mathbf{\Pi}_p^h(\mu^{-1} \nabla \times \mathbf{u}) - \mathbf{\Pi}_{\Sigma_h}(\mu^{-1} \nabla \times \mathbf{u})\|_{0,\partial K}^2 \\ & \leq C \sum_{K \in \mathcal{T}_h} M_K \|\mathbf{\Pi}_p^h(\mu^{-1} \nabla \times \mathbf{u}) - \mathbf{\Pi}_{\Sigma_h}(\mu^{-1} \nabla \times \mathbf{u})\|_{0,K}^2 \\ & \leq C \sum_{K \in \mathcal{T}_h} M_K \|\mathbf{\Pi}_p^h(\mu^{-1} \nabla \times \mathbf{u}) - \mu^{-1} \nabla \times \mathbf{u}\|_{0,K}^2 \\ & \leq C \sum_{K \in \mathcal{T}_h} \frac{h_K^{2 \min(p_K+1, s_K)} M_K}{p_K^{2s_K}} \|\mu^{-1} \nabla \times \mathbf{u}\|_{s_K, K}^2, \end{aligned}$$

where we have used the inverse estimate (4.5), the fact that $\mathbf{\Pi}_p^h(\mu^{-1} \nabla \times \mathbf{u}) = \mathbf{\Pi}_{\Sigma_h} \mathbf{\Pi}_p^h(\mu^{-1} \nabla \times \mathbf{u})$, the stability of the L^2 -projection $\mathbf{\Pi}_{\Sigma_h}$ and (4.9) with $t_K = s_K$ and $q = 0$. Therefore, we obtain

$$T_1^2 \leq C \|w\|_h^2 \sum_{K \in \mathcal{T}_h} \frac{h_K^{2 \min(p_K+1, s_K)} M_K}{p_K^{2s_K}} \|\mu^{-1} \nabla \times \mathbf{u}\|_{s_K, K}^2.$$

This, together with similar estimates for the terms T_2 , T_3 and T_4 in the above expression for $\mathcal{R}_h(\mathbf{u}, \mathbf{w})$, proves the result. \square

In order to estimate the infimum at the right-hand side of the bound in Theorem 4.8, we make the assumption that the local meshsizes and approximation degrees have bounded variation, i.e., that there exist a constant $\ell > 0$ such that

$$\ell^{-1} h_K \leq h_{K'} \leq \ell h_K, \quad \ell^{-1} p_K \leq p_{K'} \leq \ell p_K \quad (4.13)$$

for all K and K' sharing a two-dimensional face. In particular, this assumption forbids the situation where the mesh is indefinitely refined in only one of two adjacent subdomains. Nevertheless, the above hypothesis is not restrictive in practice, and allows, for instance, for geometric refinement and linearly increasing approximation orders. For any element K , we denote by $m_{\delta K}$ the quantity defined by

$$m_{\delta K} = \min\{m_{K'} : K \text{ and } K' \text{ share at least one face}\}. \quad (4.14)$$

We are now able to prove the main approximation result.

THEOREM 4.12. *Assume the above hypotheses on μ , ν and on the coefficients in the definition of the numerical fluxes. Consider shape regular meshes and polynomial degree distributions obeying (4.13). Furthermore, denote by \mathbf{u}_h the discrete solution*

of the LDG method defined in section 3 and let the exact solution \mathbf{u} satisfy $\mathbf{u}|_K \in H^{s_K+1}(K)^3$, $(\mu^{-1}\nabla \times \mathbf{u})|_K \in H^{s_K}(K)^3$, for all $K \in \mathcal{T}_h$, and $(\nu\nabla \cdot \mathbf{u})|_K \in H^{s_K}(K)$, for all $K \in \mathcal{T}_h^0$, with local regularity exponents $s_K \geq 1$. Then we have the a-priori error estimate

$$\begin{aligned} \|\mathbf{u} - \mathbf{u}_h\|_h^2 &\leq C \sum_{K \in \mathcal{T}_h} \frac{h_K^{2\min(p_K, s_K)}}{p_K^{2s_K-1}} \left(N_K \|\mathbf{u}\|_{s_K+1, K}^2 + M_K \|\mu^{-1}\nabla \times \mathbf{u}\|_{s_K, K}^2 \right) \\ &\quad + C \sum_{K \in \mathcal{T}_h^0} \frac{h_K^{2\min(p_K, s_K)}}{p_K^{2s_K-1}} M_K \|\nu\nabla \cdot \mathbf{u}\|_{s_K, K}^2, \end{aligned}$$

where M_K are the constants in (4.1), and $N_K = \frac{1}{m_{\delta K}} + |\omega| \sup_{\mathbf{x} \in K} |\sigma(\mathbf{x})| + \lambda$, with $m_{\delta K}$ defined by (4.14), and $|\sigma(\mathbf{x})|$ denoting the spectral norm of the tensor $\sigma(\mathbf{x})$. The constant C depends on Ω , Ω_0 , $\{s_K\}$, κ , ℓ , α , δ , and on the shape regularity of the mesh, but is independent of the local meshsizes h_K and the polynomial degrees p_K .

Proof. We start by estimating $\|\mathbf{u} - \mathbf{\Pi}_p^h(\mathbf{u})\|_h$, where $\mathbf{\Pi}_p^h$ is the operator defined after Proposition 4.10. From the definition of $\|\cdot\|_h$, the assumptions on μ , ν and on the coefficients in the definition of the numerical fluxes, and hypothesis (4.13), we have

$$\begin{aligned} \|\mathbf{u} - \mathbf{\Pi}_p^h \mathbf{u}\|_h^2 &\leq C \left(\sum_{K \in \mathcal{T}_h} \frac{1}{m_{\delta K}} \|\mathbf{u} - \mathbf{\Pi}_p^h \mathbf{u}\|_{1, K}^2 + \sum_{K \in \mathcal{T}_h} \frac{p_K^2}{h_K m_{\delta K}} \|\mathbf{u} - \mathbf{\Pi}_p^h \mathbf{u}\|_{0, \partial K}^2 \right. \\ &\quad \left. + |\omega| \sum_{K \in \mathcal{T}_h} \sup_{\mathbf{x} \in K} |\sigma(\mathbf{x})| \|\mathbf{u} - \mathbf{\Pi}_p^h \mathbf{u}\|_{0, K}^2 + \lambda \sum_{K \in \mathcal{T}_h^0: \partial K \cap \partial \Omega_0 \neq \emptyset} \|\mathbf{u} - \mathbf{\Pi}_p^h \mathbf{u}\|_{0, \partial K}^2 \right) \end{aligned}$$

with $m_{\delta K}$ defined by (4.14) (C depends on Ω , Ω_0 , κ , ℓ , α , δ and on the shape regularity of the mesh). The hp -approximation results with $t_K = s_K + 1$ in Proposition 4.10 yield

$$\|\mathbf{u} - \mathbf{\Pi}_p^h \mathbf{u}\|_h^2 \leq C \sum_{K \in \mathcal{T}_h} \frac{h_K^{2\min(p_K, s_K)}}{p_K^{2s_K-1}} N_K \|\mathbf{u}\|_{s_K+1, K}^2.$$

By inserting this and the estimate of Lemma 4.11 in the inequality of Theorem 4.8, we obtain the result. \square

Notice that for solutions $\mathbf{u} \in H^{s+1}(\mathcal{T}_h)^3$, with $\mu^{-1}\nabla \times \mathbf{u} \in H^s(\mathcal{T}_h)^3$ and $\nu\nabla \cdot \mathbf{u} \in H^s(\mathcal{T}_h^0)$, $s \geq 1$, assuming constant approximation orders $p_K = p$ for all $K \in \mathcal{T}_h$, setting $h = \max_{K \in \mathcal{T}_h} h_K$, and incorporating bounds related to μ , ν , σ and λ in the constant C , the estimate in Theorem 4.12 simplifies to

$$\|\mathbf{u} - \mathbf{u}_h\|_h \leq C \frac{h^{\min(s, p)}}{p^{s-\frac{1}{2}}} \left(\|\mathbf{u}\|_{s+1, \mathcal{T}_h} + \|\mu^{-1}\nabla \times \mathbf{u}\|_{s, \mathcal{T}_h} + \|\nu\nabla \cdot \mathbf{u}\|_{s, \mathcal{T}_h^0} \right).$$

This estimate is optimal in the meshsize h , and slightly suboptimal in p (half a power of p is lost). In the case of elliptic diffusion problems in two- or three-dimensional domains, no better p -bound can be found in the literature for general unstructured grids (see, e.g., the hp -version analyzes in [34, 41, 43]). Optimal p -bounds have been obtained in [19] for one-dimensional convection-diffusion problems, and recently in [32] for two-dimensional reaction-diffusion problems on structured quadrilateral grids.

REMARK 4.13. *For solutions that are elementwise analytic, we have in fact exponential convergence as $p \rightarrow \infty$. This can be seen from the error bound in Theorem 4.8 and standard approximation properties for analytic functions; see, e.g., [44].*

Note also that the restriction $s_K \geq 1$ has been made for convenience only, and it is possible to prove error estimates for $s_K > \frac{1}{2}$ as well. This minimal regularity assumption is still unrealistic when strong edge and corner singularities are present in the solutions (see [28]). On the other hand, the use of appropriate hp -mesh design principles might resolve these singularities at exponential convergence; see, e.g., [44]. The extension of our analysis to such low-regularity cases remains to be done.

REMARK 4.14. *The result of Theorem 4.12 holds true verbatim for the IP and NIP method. However, in the IP case, the constant α has to be sufficiently large to guarantee the existence of discrete solutions; see Remark 4.6.*

4.5. Error estimates for the auxiliary variables. By invoking the expressions in (3.11) and (3.12), we are able to derive error estimates for the auxiliary variables \mathbf{s} and φ . This is important, in particular, because the variable \mathbf{s} is related to the magnetic field that might be of interest. These estimates are a straightforward consequence of the following result.

PROPOSITION 4.15. *Under the same assumptions as in Theorem 4.12, we have*

$$\begin{aligned} \|\mu^{-1}\nabla \times \mathbf{u} - \mathbf{s}_h\|_{0,\Omega} &\leq C(\|\mu^{-1}\nabla \times \mathbf{u} - \Pi_p^h(\mu^{-1}\nabla \times \mathbf{u})\|_{0,\Omega} + \|\mathbf{u} - \mathbf{u}_h\|_h) \\ \|\nu\nabla \cdot \mathbf{u} - \varphi_h\|_{0,\Omega_0} &\leq C(\|\nu\nabla \cdot \mathbf{u} - \Pi_p^h(\nu\nabla \cdot \mathbf{u})\|_{0,\Omega_0} + \|\mathbf{u} - \mathbf{u}_h\|_h), \end{aligned}$$

with C depending on α , δ , κ and the shape regularity of the mesh.

Proof. Let us denote again by Π_{Σ_h} the L^2 -projection onto Σ_h . Taking into account the identity (3.11), the triangle inequality and that, for the exact solution \mathbf{u} , $\mathcal{L}(\llbracket \mathbf{u} \rrbracket_T) = \mathbf{G}_D$, we obtain

$$\|\mu^{-1}\nabla \times \mathbf{u} - \mathbf{s}_h\|_{0,\Omega} \leq T_1 + T_2,$$

where

$$\begin{aligned} T_1 &= \|\mu^{-1}\nabla \times \mathbf{u} - \Pi_{\Sigma_h}(\mu^{-1}\nabla \times \mathbf{u})\|_{0,\Omega} \\ T_2 &= \|\Pi_{\Sigma_h}(\mu^{-1}\mathcal{L}(\llbracket \mathbf{u} - \mathbf{u}_h \rrbracket_T))\|_{0,\Omega}. \end{aligned}$$

Using the stability of Π_{Σ_h} and the estimates in Proposition 4.2, we obtain $T_2 \leq C\|\mathbf{u} - \mathbf{u}_h\|_h$. We estimate T_1 as follows:

$$\begin{aligned} T_1 &\leq \|\mu^{-1}\nabla \times \mathbf{u} - \Pi_{\Sigma_h}(\mu^{-1}\nabla \times \mathbf{u})\|_{0,\Omega} + \|\Pi_{\Sigma_h}(\mu^{-1}\nabla \times \mathbf{u} - \mu^{-1}\nabla \times \mathbf{u}_h)\|_{0,\Omega} \\ &\leq C\|\mu^{-1}\nabla \times \mathbf{u} - \Pi_p^h(\mu^{-1}\nabla \times \mathbf{u})\|_{0,\Omega} + \|\mathbf{u} - \mathbf{u}_h\|_h, \end{aligned}$$

where we used the triangle inequality, the fact that $\Pi_p^h(\mu^{-1}\nabla \times \mathbf{u}) = \Pi_{\Sigma_h}\Pi_p^h(\mu^{-1}\nabla \times \mathbf{u})$, the stability of Π_{Σ_h} and the definition of $\|\cdot\|_h$. This completes the proof of the first estimate. The second one can be obtained in a similar way. \square

Proposition 4.15, together with Theorem 4.12 and Proposition 4.10, yields immediately hp -bounds for the error in \mathbf{s} and φ . For instance, for solutions $\mathbf{u} \in H^{s+1}(\mathcal{T}_h)^3$, with $\mu^{-1}\nabla \times \mathbf{u} \in H^s(\mathcal{T}_h)^3$ and $\nu\nabla \cdot \mathbf{u} \in H^s(\mathcal{T}_h^0)$, $s \geq 1$, and for constant approximation orders $p_K = p$, for all $K \in \mathcal{T}_h$, we get

$$\begin{aligned} \|\mu^{-1}\nabla \times \mathbf{u} - \mathbf{s}_h\|_{0,\Omega} &\leq C \frac{h^{\min(s,p)}}{p^{s-\frac{1}{2}}} (\|\mathbf{u}\|_{s+1,\mathcal{T}_h} + \|\mu^{-1}\nabla \times \mathbf{u}\|_{s,\mathcal{T}_h} + \|\nu\nabla \cdot \mathbf{u}\|_{s,\mathcal{T}_h^0}) \\ \|\nu\nabla \cdot \mathbf{u} - \varphi_h\|_{0,\Omega_0} &\leq C \frac{h^{\min(s,p)}}{p^{s-\frac{1}{2}}} (\|\mathbf{u}\|_{s+1,\mathcal{T}_h^0} + \|\mu^{-1}\nabla \times \mathbf{u}\|_{s,\mathcal{T}_h} + \|\nu\nabla \cdot \mathbf{u}\|_{s,\mathcal{T}_h^0}). \end{aligned}$$

We remark that for the second estimate the global error $\|\mathbf{u} - \mathbf{u}_h\|_h$ has been invoked for convenience. This bound can also be expressed in terms of local quantities related to Ω_0 only.

From the approximation point of view, the above error bounds for the auxiliary variables are suboptimal in h and in p . In [23], a special numerical flux has been identified for the Laplacian that gives better rates of convergence for Q^p elements on Cartesian grids. A similar result remains to be established in the context of the present paper.

We also remark that it is possible to use polynomials of elemental degree $p_K - 1$ in the approximation of \mathbf{s} and φ , resulting in *optimal* error estimates. However, numerical results in [24] and [40], for the Stokes problem and the Laplacian, show that this alternative does not lead to a gain in terms of computational cost and produces slightly less accurate approximations.

4.6. Stability of discrete solutions. In this section, we investigate the stability of discrete solutions with respect to the data. First, we do this by proving a discrete Poincaré inequality, based on a duality argument similar to the one in [4]. We restrict ourselves to the case of $\mu = I$, with I being the identity, since the elliptic regularity result needed for this argument does not hold true if μ is piecewise smooth (see [28]). Then, in the general case of discontinuous coefficients, the analogous stability result can be obtained, provided that the source term \mathbf{F} satisfies certain restrictions.

PROPOSITION 4.16. *Assume that $\mu = I$, and take $\nu = 1$, $\lambda = 1$. Moreover, assume the above hypotheses (4.3) and (4.4) on the coefficients in the definition of the numerical fluxes. Then we have that*

$$\|\mathbf{v}\|_{0,\Omega} \leq C \|\mathbf{v}\|_h$$

for any $\mathbf{v} \in H^s(\mathcal{T}_h)^3$, $s > \frac{1}{2}$. The constant C is independent of the meshsizes and the approximation degrees.

Proof. Fix $\mathbf{v} \in H^s(\mathcal{T}_h)^3$. Since $\mathbf{v} \in L^2(\Omega)^3$, we can decompose \mathbf{v} according to (2.6) into $\mathbf{v} = \mathbf{F}' + \mathbf{F}''$, with $\mathbf{F}'' = \nabla f$ in Ω_0 and $f = f_j$ on $\Gamma_{0,j}$, $j = 1, \dots, J$. We consider the following dual problem: find $\mathbf{z} \in H(\text{curl}; \Omega) \cap H(\text{div}; \Omega_0)$ such that

$$\nabla \times \nabla \times \mathbf{z} + i\omega\sigma\mathbf{z} = \mathbf{F}' \quad \text{in } \Omega \quad (4.15)$$

$$\nabla \cdot \mathbf{z} = -f \quad \text{in } \Omega_0 \quad (4.16)$$

$$\mathbf{n} \times \mathbf{z} = \mathbf{0} \quad \text{on } \partial\Omega \quad (4.17)$$

$$\langle (\mathbf{z}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = f_j \quad \forall j = 1, \dots, J. \quad (4.18)$$

First, we claim that

$$\nabla \times \mathbf{z} \in H^{s_0}(\Omega)^3, \quad \|\nabla \times \mathbf{z}\|_{s_0,\Omega} \leq C \|\mathbf{v}\|_{0,\Omega}, \quad \text{for } s_0 > \frac{1}{2}. \quad (4.19)$$

To prove this, set $\mathbf{w} = \nabla \times \mathbf{z}$. From equation (4.15), we have $\nabla \times \mathbf{w} = -i\omega\sigma\mathbf{z} + \mathbf{F}' \in L^2(\Omega)^3$. Furthermore, $\nabla \cdot \mathbf{w} = 0$ and $\mathbf{w} \cdot \mathbf{n} = \nabla \times \mathbf{z} \cdot \mathbf{n} = 0$ on $\partial\Omega$. Hence, from [3, Proposition 3.7], it follows that $\mathbf{w} \in H^{s_0}(\Omega)^3$ for a regularity exponent $s_0 > \frac{1}{2}$, as well as $\|\mathbf{w}\|_{s_0,\Omega} \leq C \|\mathbf{w}\|_{H(\text{curl}; \Omega)}$. However, $\|\mathbf{w}\|_{H(\text{curl}; \Omega)} \leq C \|\mathbf{z}\|_{H(\text{curl}; \Omega)} + \|\mathbf{F}'\|_{0,\Omega}$. Then, from the stability estimate in Theorem 2.3 and the L^2 -orthogonality of the decomposition of \mathbf{v} , we also have $\|\mathbf{w}\|_{s_0,\Omega} \leq C \|\mathbf{v}\|_{0,\Omega}$, which completes the proof of (4.19).

Subtracting the gradient of equation (4.16) from (4.15), multiplying the result by $\bar{\mathbf{v}}$, and integrating over Ω and Ω_0 , owing to the decomposition of \mathbf{v} , we obtain that

$$\|\mathbf{v}\|_{0,\Omega}^2 = \int_{\Omega} (\nabla \times \nabla \mathbf{z} + i\omega\sigma\mathbf{z}) \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\Omega_0} (\nabla \nabla \cdot \mathbf{z}) \cdot \bar{\mathbf{v}} \, d\mathbf{x}.$$

Let us first consider the integral containing the curl–curl term. Integration by parts, together with the first rule in (3.9), as well as the Cauchy-Schwarz inequality and the definition of $\|\cdot\|_h$, gives

$$\begin{aligned} \left| \int_{\Omega} \nabla \times \nabla \mathbf{z} \cdot \bar{\mathbf{v}} \, d\mathbf{x} \right| &\leq \left| \int_{\Omega} \nabla \times \mathbf{z} \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} \right| + \left| \int_{\mathcal{E}} \{\{\nabla \times \mathbf{z}\}\} \cdot \llbracket \bar{\mathbf{v}} \rrbracket_T \, ds \right| \\ &\leq \|\mathbf{z}\|_{H(\text{curl};\Omega)} \|\mathbf{v}\|_h + C \left(\sum_{K \in \mathcal{T}_h} \frac{h_K}{p_K^2} \|\nabla \times \mathbf{z}\|_{0,\partial K}^2 \right)^{\frac{1}{2}} \|\mathbf{v}\|_h. \end{aligned}$$

Using similar scaling arguments as the ones in [2, Lemma 5.2], together with the result of [2, Lemma 5.5], we can see that

$$\|\nabla \times \mathbf{z}\|_{0,\partial K}^2 \leq Ch_K^{-1} \|\nabla \times \mathbf{z}\|_{s_0,K}^2,$$

with a constant C only depending on the shape regularity of the meshes. This, together with $p_K^{-1} \leq 1$, yields $\sum_{K \in \mathcal{T}_h} \frac{h_K}{p_K^2} \|\nabla \times \mathbf{z}\|_{0,\partial K}^2 \leq C \|\nabla \times \mathbf{z}\|_{s_0,\Omega}^2 \leq C \|\mathbf{v}\|_{0,\Omega}^2$. Therefore, since also $\|\mathbf{z}\|_{H(\text{curl};\Omega)} \leq C \|\mathbf{v}\|_{0,\Omega}$ (see Theorem 2.3), we get

$$\left| \int_{\Omega} \nabla \times \nabla \mathbf{z} \cdot \bar{\mathbf{v}} \, d\mathbf{x} \right| \leq C \|\mathbf{v}\|_{0,\Omega} \|\mathbf{v}\|_h.$$

Similarly, by integration by parts, using the second identity in (3.9), we obtain

$$\left| \int_{\Omega_0} (\nabla \nabla \cdot \mathbf{z}) \cdot \bar{\mathbf{v}} \, d\mathbf{x} \right| \leq \|\mathbf{z}\|_{H(\text{div};\Omega_0)} \|\mathbf{v}\|_h + \left| \int_{\mathcal{E}_T^0} \{\{\nabla \cdot \mathbf{z}\}\} \llbracket \bar{\mathbf{v}} \rrbracket_N \, ds \right| + \sum_{j=1}^J |f_j| \int_{\Gamma_{0,j}} |\bar{\mathbf{v}} \cdot \mathbf{n}_{0,j}| \, ds,$$

where we also used the fact that $\nabla \cdot \mathbf{z}$ is zero on $\Gamma_{0,0}$ and equal to $-f_j$ on $\Gamma_{0,j}$, for $j = 1, \dots, J$. Since $\|\nabla \cdot \mathbf{z}\|_{0,\partial K} \leq Ch_K^{-\frac{1}{2}} \|\nabla \cdot \mathbf{z}\|_{1,K}$, for all $K \in \mathcal{T}_h^0$, we get

$$\left| \int_{\mathcal{E}_T^0} \{\{\nabla \cdot \mathbf{z}\}\} \llbracket \bar{\mathbf{v}} \rrbracket_N \, ds \right| \leq C \|\nabla \cdot \mathbf{z}\|_{1,\Omega_0} \|\mathbf{v}\|_h.$$

The Cauchy-Schwarz inequality, the trace theorem and the standard Poincaré inequality then yield

$$\sum_{j=1}^J |f_j| \int_{\Gamma_{0,j}} |\bar{\mathbf{v}} \cdot \mathbf{n}_{0,j}| \, ds \leq C \|\nabla f\|_{0,\Omega_0} \|\mathbf{v}\|_h.$$

Combining the above estimates, together with $\|\mathbf{z}\|_{H(\text{div};\Omega_0)} + \|\nabla \cdot \mathbf{z}\|_{1,\Omega_0} \leq C \|\mathbf{v}\|_{0,\Omega}$ and a similar argument for the term $i\omega \int_{\Omega} \sigma \mathbf{z} \bar{\mathbf{v}} \, d\mathbf{x}$, shows that

$$\|\mathbf{v}\|_{0,\Omega}^2 \leq C \|\mathbf{v}\|_{0,\Omega} \|\mathbf{v}\|_h.$$

This completes the proof. \square

From Proposition 4.16, we are able to establish the continuity of the functional $\mathcal{F}_h(\cdot)$ in (3.13) with respect to the norm $\|\cdot\|_h$, at least in the particular case where $\mu = I$.

COROLLARY 4.17. *Under the same assumptions as in Proposition 4.16, we have for all $\mathbf{v} \in \mathbf{V}_h$,*

$$|\mathcal{F}_h(\mathbf{v})| \leq C(\|\mathbf{F}\|_{0,\Omega}^2 + \|\mathbf{h}^{-\frac{1}{2}}\mathbf{p}\mathbf{g}\|_{0,\partial\Omega}^2)^{\frac{1}{2}}\|\mathbf{v}\|_h,$$

with a constant C independent of the meshsizes and the approximation degrees.

Proof. From the definition of \mathcal{F}_h , Proposition 4.16 and the Cauchy-Schwarz inequality, we obtain

$$|\mathcal{F}_h(\mathbf{v})| \leq C(\|\mathbf{F}\|_{0,\Omega}^2 + \|\mathbf{G}_{\mathcal{D}}\|_{0,\Omega}^2 + \|\mathbf{h}^{-\frac{1}{2}}\mathbf{p}\mathbf{g}\|_{0,\partial\Omega}^2)^{\frac{1}{2}}\|\mathbf{v}\|_h,$$

for any $\mathbf{v} \in \mathbf{V}_h$. The result follows from the estimate for $\mathbf{G}_{\mathcal{D}}$ in Proposition 4.2. \square

Consider now the case where μ satisfies our more general assumptions. The part of the functional \mathcal{F}_h involving the boundary term can be dealt with as in Corollary 4.17, therefore, we will focus on the term $\int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x}$. We can prove the following result.

PROPOSITION 4.18. *Assume our general hypotheses on μ , ν and the coefficients in the definition of the numerical fluxes. Whenever Ω_0 is not simply-connected, let $\{\Sigma_\ell\}_{\ell=1,\dots,L}$ be an admissible set of cuts for Ω_0 in the sense of [3], and denote by \mathbf{n}_ℓ the normal unit vector to Σ_ℓ , pointing in one of the two possible directions. Given $\mathbf{F} \in L^2(\Omega)^3$, let $\mathbf{F} = \mathbf{F}' + \mathbf{F}''$ be its orthogonal decomposition, according to (2.6). If*

$$\begin{aligned} \mathbf{F}' \cdot \mathbf{n}_0 &= 0 && \text{on } \partial\Omega_0 \\ \langle \mathbf{F}' \cdot \mathbf{n}_\ell, 1 \rangle_{\Sigma_\ell} &= 0 && \forall \ell = 1, \dots, L, \end{aligned} \quad (4.20)$$

then, for all $\mathbf{v} \in \mathbf{V}_h$,

$$\int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} \leq C\|\mathbf{F}\|_{0,\Omega}\|\mathbf{v}\|_h. \quad (4.21)$$

with a constant C independent of the meshsizes and the approximation degrees.

We first remark that, without the above restrictions, the continuity property (4.21) still holds true, but with a constant C that depends weakly on the meshsizes and the approximation degrees.

Second, we notice that, if Ω_0 is simply-connected, the second condition in (4.20) is empty. In this case, if \mathbf{F} coincides with the physical source field \mathbf{J} , then (4.20) is satisfied (and $\mathbf{F}'' = \mathbf{0}$). If Ω_0 is not simply-connected, the case $\mathbf{F} = \mathbf{J}$ is covered, provided that $\langle \mathbf{J}' \cdot \mathbf{n}_\ell, 1 \rangle_{\Sigma_\ell} = 0$, $\ell = 1, \dots, L$.

Finally, we point out that in the case where $\Omega = \Omega_0$ and $\Omega_\sigma = \emptyset$ no restriction on \mathbf{F} needs to be assumed.

Proof of Proposition 4.18. Recall that in the orthogonal decomposition (2.6) of $\mathbf{F} = \mathbf{F}' + \mathbf{F}''$, the function \mathbf{F}'' is such that $\mathbf{F}''|_{\Omega_\sigma} = \mathbf{0}$ and $\mathbf{F}''|_{\Omega_0} = \nabla f$, with $f \in H^1(\Omega_0)$, $f = 0$ on $\Gamma_{0,0}$ and f constant, say $f = f_j$, on each $\Gamma_{0,j}$ for $j = 1, \dots, J$. The standard Poincaré inequality implies that $\|f\|_{0,\Omega_0}$ and $\|f\|_{1,\Omega_0}$ are equivalent to $\|\nabla f\|_{0,\Omega_0} = \|\mathbf{F}''\|_{0,\Omega_0}$. From [3, Theorem 3.17], the assumptions (4.20) on \mathbf{F}' , together with $\nabla \cdot \mathbf{F}'|_{\Omega_0} = 0$, imply that there exists $\mathbf{w} \in H_0(\text{curl}, \text{div}; \Omega_0)$ with $\nabla \cdot \mathbf{w} = 0$ in Ω_0 and $\langle \mathbf{w} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}} = 0$, $j = 1, \dots, J$, such that $\mathbf{F}' = \nabla \times \mathbf{w}$. Corollary 3.19 in [3] implies that $\|\mathbf{w}\|_{0,\Omega_0}$ and $\|\mathbf{w}\|_{H(\text{curl}, \Omega_0)}$ are equivalent to $\|\nabla \times \mathbf{w}\|_{0,\Omega_0} = \|\mathbf{F}'\|_{0,\Omega_0}$.

Moreover, from the continuous imbedding of $H_0(\text{curl}, \text{div}; \Omega_0)$ in $H^s(\Omega_0)^3$, for some $s > \frac{1}{2}$, we also have that $\mathbf{w} \in H^s(\Omega_0)^3$ and $\|\mathbf{w}\|_{s, \Omega_0}$ can be controlled by $\|\mathbf{w}\|_{H(\text{curl}; \Omega_0)}$. After these preliminaries, we can proceed by estimating $\int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x}$. We can write

$$\begin{aligned} \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} &= \int_{\Omega_{\sigma}} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\Omega_0} (\nabla \times \mathbf{w} + \nabla f) \cdot \bar{\mathbf{v}} \, d\mathbf{x} \\ &= \int_{\Omega_{\sigma}} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \int_{\Omega_0} \mathbf{w} \cdot \nabla_h \times \bar{\mathbf{v}} \, d\mathbf{x} - \sum_{K \in \mathcal{T}_h^0} \int_{\partial K} \mathbf{w} \cdot \mathbf{n}_K \times \bar{\mathbf{v}} \, ds \\ &\quad - \int_{\Omega_0} f \nabla_h \cdot \bar{\mathbf{v}} \, d\mathbf{x} + \sum_{K \in \mathcal{T}_h^0} \int_{\partial K} f \bar{\mathbf{v}} \cdot \mathbf{n}_K \, ds. \end{aligned}$$

Therefore,

$$\begin{aligned} \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} &\leq |\omega|^{-1} \|\sigma^{-\frac{1}{2}} \mathbf{F}\|_{0, \Omega_{\sigma}} |\omega| \|\sigma^{\frac{1}{2}} \mathbf{v}\|_{0, \Omega_{\sigma}} + \|\mu^{\frac{1}{2}} \mathbf{w}\|_{0, \Omega_0} \|\mu^{-\frac{1}{2}} \nabla_h \times \mathbf{v}\|_{0, \Omega_0} \\ &\quad + \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} \mathbf{w}\|_{0, \mathcal{E}_x^0} \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} \llbracket \mathbf{v} \rrbracket_T\|_{0, \mathcal{E}_x^0} + \|\nu^{-\frac{1}{2}} f\|_{0, \Omega_0} \|\nu^{\frac{1}{2}} \nabla_h \cdot \mathbf{v}\|_{0, \Omega_0} \\ &\quad + \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} f\|_{0, \mathcal{E}_x^0} \|\mathbf{h}^{-\frac{1}{2}} \mathbf{p} \mathbf{m}^{-\frac{1}{2}} \llbracket \mathbf{v} \rrbracket_N\|_{0, \mathcal{E}_x^0} \\ &\quad + \left(\lambda^{-1} \sum_{j=1}^J f_j^2 \right)^{\frac{1}{2}} \left(\lambda \sum_{j=1}^J |\langle \mathbf{v} \cdot \mathbf{n}, \mathbf{1} \rangle_{\Gamma_{0,j}}|^2 \right)^{\frac{1}{2}}, \end{aligned}$$

where we have used the fact that $\mathbf{n} \times \mathbf{w} = \mathbf{0}$ on $\partial\Omega_0$. Consequently,

$$\begin{aligned} \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} &\leq C \|\mathbf{v}\|_h \left(|\omega|^{-1} \|\sigma^{-\frac{1}{2}} \mathbf{F}\|_{0, \Omega_{\sigma}}^2 + \|\mu^{\frac{1}{2}} \mathbf{w}\|_{0, \Omega_0}^2 + \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} \mathbf{w}\|_{0, \mathcal{E}^0}^2 \right. \\ &\quad \left. + \|\nu^{-\frac{1}{2}} f\|_{0, \Omega_0}^2 + \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} f\|_{0, \mathcal{E}_x^0}^2 + \lambda^{-1} \sum_{j=1}^J f_j^2 \right). \end{aligned}$$

For the volume terms at right-hand side, we have

$$\begin{aligned} |\omega|^{-1} \|\sigma^{-\frac{1}{2}} \mathbf{F}\|_{0, \Omega_{\sigma}}^2 &\leq C \|\mathbf{F}\|_{0, \Omega_{\sigma}}^2 \\ \|\mu^{\frac{1}{2}} \mathbf{w}\|_{0, \Omega_0}^2 &\leq C \|\mathbf{w}\|_{0, \Omega_0}^2 \leq C \|\mathbf{F}\|_{0, \Omega_0}^2 \\ \|\nu^{-\frac{1}{2}} f\|_{0, \Omega_0}^2 &\leq C \|f\|_{0, \Omega_0}^2 \leq C \|\mathbf{F}\|_{0, \Omega_0}^2, \end{aligned}$$

with C independent of the meshsizes and the approximation degrees. Using standard scaling arguments, we obtain

$$\begin{aligned} \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} \mathbf{w}\|_{0, \mathcal{E}^0}^2 &\leq \sum_{K \in \mathcal{T}_h^0} \frac{h_K M_K}{p_K} \|\mathbf{w}\|_{0, \partial K}^2 \leq \sum_{K \in \mathcal{T}_h^0} \frac{h_K M_K}{p_K} h_K^{-1} \|\mathbf{w}\|_{s, K}^2 \\ &\leq C \|\mathbf{w}\|_{s, \Omega_0}^2 \leq C \|\mathbf{w}\|_{H(\text{curl}; \Omega_0)}^2 \leq C \|\mathbf{F}\|_{0, \Omega_0}^2, \end{aligned}$$

again with C independent of the meshsizes and the approximation degrees. Similarly,

$$\begin{aligned} \|\mathbf{h}^{\frac{1}{2}} \mathbf{p}^{-1} \mathbf{m}^{\frac{1}{2}} f\|_{0, \mathcal{E}_x^0}^2 &\leq \sum_{K \in \mathcal{T}_h^0} \frac{h_K M_K}{p_K} \|f\|_{0, \partial K}^2 \leq \sum_{K \in \mathcal{T}_h^0} \frac{h_K M_K}{p_K} h_K^{-1} \|f\|_{1, K}^2 \\ &\leq C \|f\|_{1, \Omega_0}^2 \leq C \|F\|_{0, \Omega_0}^2, \end{aligned}$$

and

$$\lambda^{-1} \sum_{j=1}^J f_j^2 \leq C \lambda^{-1} \|f\|_{1, \Omega_0} \leq C \|\mathbf{F}\|_{0, \Omega_0}.$$

This completes the proof of the proposition. \square

5. Conclusions and extensions. In this paper, we propose a local discontinuous Galerkin method for the discretization of the time-harmonic Maxwell equations in low-frequency regime and carry out its hp -error analysis. We consider heterogeneous materials and topologically non-trivial domains; assumptions that are realistic in practice. In particular, since our method enforces interelement continuity only in weak sense, we are able to allow for domains where standard conforming approximations may fail to converge to the exact solutions.

Although we use the setting of [5] to cast the LDG method in its primal form, we propose a new technique to actually derive error estimates which is based on Strang's lemma and which might be of independent interest in the analysis of DG methods. For triangulations containing hanging nodes, we derive hp -error estimates that are optimal in the meshsize and suboptimal in the approximation order. This analysis is the first hp -error analysis for the LDG method in several space dimensions and in this sense also extends our previous work in [18].

Let us also indicate some related issues that are still open:

- For the Laplacian, the parameters \mathbf{b} and \mathbf{d} can be chosen in such a way that the LDG method superconverges on Cartesian grids and for tensor product polynomials, see [23]. Whether or not a similar phenomenon actually takes place in the context of the p -version of the method is an open question and has to be addressed in future work.

- The study of hp -refinement towards edge and corner singularities, in order to resolve them at exponential convergence, remains to be done.

- One of the drawbacks of discontinuous Galerkin methods is the relatively high number of degrees of freedom due to the discontinuous nature of the finite element spaces. This problem can be overcome by coupling discontinuous and conforming elements, following [40]. The approach there combines the ease with which the LDG method handles hanging nodes with the lower computational cost of standard finite elements.

- The extension of the LDG method to mixed boundary conditions can be done in a straightforward way (see, e.g, [18]). However, in order to avoid further complications in the analysis of the corresponding continuous problems, this point is omitted in this paper.

We conclude by pointing out that the analysis of discontinuous Galerkin methods for time-harmonic Maxwell's equation in the high-frequency case is the subject of a forthcoming paper.

Appendix. The proof of Theorem 2.3.

The appendix is devoted to the proof of Theorem 2.3. We start by establishing, in section A.1, the existence of a continuous and divergence-free lifting of tangential traces. This result is a key tool in the proof of existence and uniqueness of solutions to our model problem which is developed in section A.2.

A.1. Divergence-free continuous lifting of tangential traces. Recall that we denote by $\mathcal{H}(\partial\Omega)$ the space of tangential traces of $H(\text{curl}; \Omega)$ functions. We refer to

[14] for its complete characterization and for the definition of its norm. The existence of a continuous lifting of the tangential traces of $H(\text{curl}; \Omega)$ functions is guaranteed by Theorem 3.1 in [13] in the case where Ω is a bounded, open Lipschitz polyhedron in \mathbb{R}^3 , not necessarily simply connected. In this section, we show that we can actually require this lifting to be divergence-free in a Lipschitz polyhedral subdomain Ω_0 of Ω , and to satisfy homogeneous flux conditions through surfaces of the possible cavities of Ω_0 . This result is stated in the following proposition. We define, for convenience, the space

$$H_{\text{flux}}(\text{div}; \Omega_0) = \{\mathbf{v} \in H(\text{div}; \Omega_0) : \langle (\mathbf{v}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = 0, j = 1, \dots, J\}, \quad (\text{A.1})$$

along with its subspace $H_{\text{flux}}(\text{div}^0; \Omega_0)$ consisting of divergence-free functions.

PROPOSITION A.1. *Let Ω be a connected, bounded, open Lipschitz polyhedron in \mathbb{R}^3 , $\Omega_0 \subseteq \Omega$ a Lipschitz polyhedral subdomain. Then, given $\tilde{\mathbf{g}} \in H(\text{curl}; \Omega)$, there is $\tilde{\mathbf{u}} \in H(\text{curl}; \Omega) \cap H_{\text{flux}}(\text{div}^0; \Omega_0)$ such that $\mathbf{n} \times \tilde{\mathbf{u}} = \mathbf{n} \times \tilde{\mathbf{g}} =: \mathbf{g}$ on $\partial\Omega$ and $\|\tilde{\mathbf{u}}\|_{H(\text{curl}; \Omega)} \leq C\|\mathbf{g}\|_{\mathcal{H}(\partial\Omega)}$.*

In order to prove Proposition A.1, we need the following two lemmas.

LEMMA A.2. *Let Ω be a connected, bounded, open Lipschitz polyhedron in \mathbb{R}^3 . Then, given $\tilde{\mathbf{g}} \in H(\text{curl}; \Omega)$, there is $\tilde{\mathbf{u}}_{\text{bc}} \in H(\text{curl}; \Omega) \cap H(\text{div}^0; \Omega)$ such that $\mathbf{n} \times \tilde{\mathbf{u}}_{\text{bc}} = \mathbf{n} \times \tilde{\mathbf{g}} =: \mathbf{g}$ on $\partial\Omega$ and $\|\tilde{\mathbf{u}}_{\text{bc}}\|_{H(\text{curl}; \Omega)} \leq C\|\mathbf{g}\|_{\mathcal{H}(\partial\Omega)}$.*

Proof. Let $\tilde{\mathbf{g}} \in H(\text{curl}; \Omega)$. Since $\nabla \times H(\text{curl}; \Omega) = \nabla \times H^1(\Omega)^3$ (see [30]), there exists $\mathbf{g}_1 \in H^1(\Omega)^3$ such that $\nabla \times \mathbf{g}_1 = \nabla \times \tilde{\mathbf{g}}$. Then, we can write $\tilde{\mathbf{g}} = \mathbf{g}_1 + \mathbf{g}_2$, with $\mathbf{g}_2 \in H(\text{curl}; \Omega)$ and $\nabla \times \mathbf{g}_2 = 0$. Consequently, \mathbf{g}_2 admits the orthogonal decomposition $\mathbf{g}_2 = \nabla\varphi + \mathbf{w}$, with $\varphi \in H^1(\Omega)$, $\mathbf{w} \in H_0(\text{div}^0; \Omega) \cap H(\text{curl}^0; \Omega)$, (see [30], formula (4.14) with $\Gamma_\nu = \partial\Omega$ and $\Gamma_\tau = \emptyset$). Let $\tilde{\varphi} \in H^1(\Omega)^3$ be the unique solution to $\Delta\tilde{\varphi} = -\nabla \cdot \mathbf{g}_1$ in Ω , $\tilde{\varphi} = \varphi|_{\partial\Omega}$ on $\partial\Omega$, where $\varphi|_{\partial\Omega}$ is the trace in $H^{\frac{1}{2}}(\partial\Omega)$ of $\varphi \in H^1(\Omega)$. Define $\tilde{\mathbf{u}}_{\text{bc}} = \mathbf{g}_1 + \nabla\tilde{\varphi} + \mathbf{w}$. By construction, $\mathbf{n} \times \tilde{\mathbf{u}}_{\text{bc}} = \mathbf{g}$ on $\partial\Omega$, since $\mathbf{n} \times \nabla\tilde{\varphi} = \mathbf{n} \times \nabla\varphi$ on $\partial\Omega$, and $\nabla \cdot \tilde{\mathbf{u}}_{\text{bc}} = 0$ in Ω immediately follows from $\Delta\tilde{\varphi} = -\nabla \cdot \mathbf{g}_1$ and $\nabla \cdot \mathbf{w} = 0$. Therefore, the tangential trace operator is linear continuous and surjective from the divergence-free subspace of $H(\text{curl}; \Omega) \cap H(\text{div}; \Omega)$ onto $\mathcal{H}(\partial\Omega)$, and then, up to its kernel, its inverse is continuous, owing to the open mapping theorem. \square

LEMMA A.3. *Let Ω and Ω_0 be as in Proposition A.1, $\tilde{\mathbf{u}}_{\text{bc}}$ be as in Lemma A.2, and let $\alpha \in \mathbb{R}^p$ be the vector with components $\alpha_j = \langle (\tilde{\mathbf{u}}_{\text{bc}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}}$, $j = 1, \dots, J$. Then there is a function $\tilde{\mathbf{u}}_{\text{flux}} \in H_0(\text{curl}^0; \Omega) \cap H(\text{div}^0; \Omega_0)$ such that $\langle (\tilde{\mathbf{u}}_{\text{flux}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = \alpha_j$, $j = 1, \dots, J$, and*

$$\|\tilde{\mathbf{u}}_{\text{flux}}\|_{0,\Omega} \leq C\|\tilde{\mathbf{u}}_{\text{bc}}\|_{0,\Omega}. \quad (\text{A.2})$$

Proof. Conditions $\langle (\mathbf{u}_{\text{flux}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = \alpha_j$, $j = 1, \dots, J$, define a unique $\mathbf{u}_{\text{flux}} \in H_0(\text{curl}^0; \text{div}^0; \Omega_0)$ satisfying $\|\mathbf{u}_{\text{flux}}\|_{0,\Omega_0} \leq C|\alpha|$ (see, e.g., [31], Lemma 3.2, with $\Gamma_\nu = \partial\Omega_0$ and $\Gamma_\tau = \emptyset$). Actually, $\|\mathbf{u}_{\text{flux}}\|_{0,\Omega_0} \leq C\|\tilde{\mathbf{u}}_{\text{bc}}\|_{0,\Omega}$. This can be seen in the following way. By definition of dual norms, by the fact that $\Gamma_{0,j}$ is a surface without boundary, and by continuity of the normal trace operator from $H(\text{div}; \Omega_0)$

onto $H^{-\frac{1}{2}}(\partial\Omega_0)$, we have

$$\begin{aligned} \frac{|(\langle \tilde{\mathbf{u}}_{\text{bc}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}})|}{\|1\|_{\frac{1}{2}, \Gamma_{0,j}}} &\leq \sup_{\varphi \in H^{\frac{1}{2}}(\Gamma_{0,j})} \frac{|(\langle \tilde{\mathbf{u}}_{\text{bc}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, \varphi \rangle_{\Gamma_{0,j}})|}{\|\varphi\|_{\frac{1}{2}, \Gamma_{0,j}}} \\ &= \|\tilde{\mathbf{u}}_{\text{bc}}|_{\Omega_0} \cdot \mathbf{n}_{0,j}\|_{H^{-\frac{1}{2}}(\Gamma_{0,j})} \leq \|\tilde{\mathbf{u}}_{\text{bc}}|_{\Omega_0} \cdot \mathbf{n}\|_{H^{-\frac{1}{2}}(\partial\Omega_0)} \\ &\leq C \|\tilde{\mathbf{u}}_{\text{bc}}\|_{H(\text{div}; \Omega_0)} = C \|\tilde{\mathbf{u}}_{\text{bc}}\|_{0, \Omega}. \end{aligned}$$

Therefore, $|\alpha| \leq C \|\tilde{\mathbf{u}}_{\text{bc}}\|_{0, \Omega}$, from where $\|\mathbf{u}_{\text{flux}}\|_{0, \Omega_0} \leq C \|\tilde{\mathbf{u}}_{\text{bc}}\|_{0, \Omega}$ follows. Taking the trivial extension $\tilde{\mathbf{u}}_{\text{flux}}$ of \mathbf{u}_{flux} to Ω completes the proof. \square

Proof of Proposition A.1. It is enough to define $\tilde{\mathbf{u}} = \tilde{\mathbf{u}}_{\text{bc}} - \tilde{\mathbf{u}}_{\text{flux}}$, where $\tilde{\mathbf{u}}_{\text{bc}}$ and $\tilde{\mathbf{u}}_{\text{flux}}$ are as in Lemma A.2 and Lemma A.3, respectively. \square

A.2. Proof of Theorem 2.3. Recalling that $\mathbf{V} = H(\text{curl}; \Omega) \cap H(\text{div}; \Omega_0)$, endowed with the norm

$$\|\mathbf{u}\|_{\mathbf{V}}^2 = |\omega| \|\sigma_{\vartheta}^{\frac{1}{2}} \mathbf{u}\|_{0, \Omega}^2 + \|\mu^{-\frac{1}{2}} \nabla \times \mathbf{u}\|_{0, \Omega}^2 + \|\nu^{\frac{1}{2}} \nabla \cdot \mathbf{u}\|_{0, \Omega_0}^2 + \lambda \sum_{j=1}^J |(\langle \mathbf{u}|_{\Omega_0} \cdot \mathbf{n}_{0,j}, 1 \rangle_{\Gamma_{0,j}})|^2, \quad (\text{A.3})$$

we set $\mathbf{V}_0 = H_0(\text{curl}; \Omega) \cap H_{\text{flux}}(\text{div}; \Omega_0)$, with $H_{\text{flux}}(\text{div}; \Omega_0)$ defined in (A.1).

Consider the bilinear form in (2.11)

$$a(\mathbf{u}, \mathbf{v}) = \int_{\Omega} \mu^{-1} \nabla \times \mathbf{u} \cdot \nabla \times \bar{\mathbf{v}} \, dx + i\omega \int_{\Omega} \sigma \mathbf{u} \cdot \bar{\mathbf{v}} \, dx + \int_{\Omega_0} \nu \nabla \cdot \mathbf{u} \nabla \cdot \bar{\mathbf{v}} \, dx.$$

In the following proposition, we prove \mathbf{V} -ellipticity of $a(\cdot, \cdot) : \mathbf{V}_0 \times \mathbf{V}_0 \rightarrow \mathbb{R}$. The proof essentially follows the lines of [1, Theorem 3.1], but uses the result of Proposition A.1.

PROPOSITION A.4. *There is $C > 0$ such that $a(\mathbf{u}, \mathbf{u}) \geq C \|\mathbf{u}\|_{\mathbf{V}}^2$ for all $\mathbf{u} \in \mathbf{V}_0$.*

Proof. Since, for $\mathbf{u} \in \mathbf{V}_0$, $a(\mathbf{u}, \mathbf{u})$ differs from $\|\mathbf{u}\|_{\mathbf{V}}^2$ only in that it does not contain $|\omega| \|\vartheta^{\frac{1}{2}} \mathbf{u}\|_{0, \Omega_0}^2$, it is enough to prove that, for all $\mathbf{u} \in \mathbf{V}_0$,

$$\|\mathbf{u}\|_{0, \Omega_0}^2 \leq C a(\mathbf{u}, \mathbf{u}). \quad (\text{A.4})$$

In order to do this, we need to establish the existence of a continuous lifting of tangential traces on $\Gamma = \partial\Omega_{\sigma} \cap \partial\Omega_0$ with zero divergence and flux conditions. To this end, we introduce the space $H_{0, \Gamma_-}(\text{curl}; \Omega_0) = \{\mathbf{v} \in H(\text{curl}; \Omega_0) : \mathbf{n}_0 \times \mathbf{v}|_{\Gamma_-} = \mathbf{0} \text{ in } H_{00}^{-\frac{1}{2}}(\Gamma_-)\}$, where $\Gamma_- = \partial\Omega_0 \setminus \bar{\Gamma}$, i.e., Γ_- is the part of $\partial\Omega_0$ contained in $\partial\Omega$. For the definition of $H_{00}^{-\frac{1}{2}}(\Gamma_-)$, see, e.g., [35]. Let γ^- be the restriction of the tangential trace operator to $\bar{\Gamma}$ and $\mathcal{H}(\Gamma)$ its image. A complete characterization of this space, as well as the precise definition of its norm can be found in [14]. Fix $\mathbf{v} \in H_{0, \Gamma_-}(\text{curl}; \Omega_0)$. According to Proposition A.1, with $\Omega = \Omega_0$, let $\mathbf{v}_{\text{bc}} \in H(\text{curl}; \Omega_0) \cap H_{\text{flux}}(\text{div}^0; \Omega_0)$ be the lifting of $\mathbf{n}_0 \times \mathbf{v}$ on $\partial\Omega_0$, i.e., $\mathbf{n}_0 \times \mathbf{v} = \mathbf{n}_0 \times \mathbf{v}_{\text{bc}}$ on $\partial\Omega_0$. Then, we also have $\mathbf{v}_{\text{bc}} \in H_{0, \Gamma_-}(\text{curl}; \Omega_0)$ and $\gamma^- \mathbf{v}_{\text{bc}} = \gamma^- \mathbf{v}$. This shows that γ^- is linear continuous (see [15] and [13]) and surjective from $H_{0, \Gamma_-}(\text{curl}; \Omega_0) \cap H_{\text{flux}}(\text{div}^0; \Omega_0)$ onto $\mathcal{H}(\Gamma)$, and then, up to its kernel, its inverse is continuous, owing to the open mapping theorem.

Now, let $\mathbf{u} \in \mathbf{V}_0$, and denote by \mathbf{u}_0 the restriction of \mathbf{u} to Ω_0 . Then, due to the previous considerations, \mathbf{u}_0 can be decomposition as $\mathbf{u}_0 = \mathbf{w} + \mathbf{u}_{\text{bc}}$, where $\mathbf{w} \in H_0(\text{curl}; \Omega_0) \cap H_{\text{flux}}(\text{div}; \Omega_0)$ and \mathbf{u}_{bc} is such that $\mathbf{u}_{\text{bc}} \in H_{0, \Gamma_-}(\text{curl}; \Omega_0) \cap H_{\text{flux}}(\text{div}^0; \Omega_0)$, $\mathbf{n}_0 \times \mathbf{u}_{\text{bc}} = \mathbf{n}_0 \times \mathbf{u}_0$ on Γ and $\|\mathbf{u}_{\text{bc}}\|_{H(\text{curl}; \Omega_0)} \leq C \|\gamma^- \mathbf{u}\|_{\mathcal{H}(\Gamma)}$. From [3, Corollary 3.19], we have

$$\|\mathbf{w}\|_{0, \Omega_0} \leq C (\|\nabla \times \mathbf{w}\|_{0, \Omega_0} + \|\nabla \cdot \mathbf{w}\|_{0, \Omega_0}) \leq C a(\mathbf{u}, \mathbf{u})^{\frac{1}{2}}. \quad (\text{A.5})$$

As far as \mathbf{u}_{bc} is concerned, we can obtain a bound in terms of $a(\mathbf{u}, \mathbf{u})$ as follows. Since $\mathbf{u} \in H(\text{curl}; \Omega)$, we have that $\mathbf{n}_0 \times \mathbf{u}_0 = \mathbf{n}_0 \times \mathbf{u}|_{\Omega_\sigma}$ on Γ . This implies that $\|\gamma^- \mathbf{u}_0\|_{\mathcal{H}(\Gamma)} = \|\tilde{\gamma}^- \mathbf{u}\|_{\mathcal{H}(\Gamma)}$, where $\tilde{\gamma}^-$ is the restriction to Γ of the tangential trace operator taken from $\Omega_\sigma = \Omega \setminus \Omega_0$. By the continuity of $\tilde{\gamma}^-$, we conclude that

$$\|\mathbf{u}_{bc}\|_{H(\text{curl}; \Omega_0)} \leq C \|\mathbf{u}\|_{H(\text{curl}; \Omega_\sigma)} \leq C a(\mathbf{u}, \mathbf{u})^{\frac{1}{2}}.$$

This, together with (A.5), implies (A.4). The statement of the proposition then follows. \square

Well-posedness of formulation (2.11) can now be proved in a standard way as follows. Let \mathbf{u}_f be the (unique) function in $H_0(\text{curl}^0, \text{div}^0; \Omega_0)$ such that $\langle (\mathbf{u}_f|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = \lambda^{-1} f_j$, $j = 1, \dots, J$ (see, e.g., [31, Lemma 3.2]), and define $\tilde{\mathbf{u}}_f$ as the trivial extension of \mathbf{u}_f to Ω . By the orthogonal decomposition $\mathbf{F} = \mathbf{F}' + \mathbf{F}''$, as in (2.6), with $\mathbf{F}'' = \nabla f$ in Ω_0 , and the trace theorem in $H^1(\Omega_0)$, we have $\|\tilde{\mathbf{u}}_f\|_{\mathbf{V}} \leq C \|\mathbf{F}\|_{0, \Omega}$.

Furthermore, let $\tilde{\mathbf{u}} \in \mathbf{V}$ be such that $\nabla \cdot \tilde{\mathbf{u}} = 0$ in Ω_0 , $\mathbf{n} \times \tilde{\mathbf{u}} = \mathbf{g}$ on $\partial\Omega$ and $\|\tilde{\mathbf{u}}\|_{H(\text{curl}; \Omega)} \leq C \|\mathbf{g}\|_{\mathcal{H}(\partial\Omega)}$, according to Proposition A.1. By defining $\mathbf{u}_0 = \mathbf{u} - \tilde{\mathbf{u}}_f - \tilde{\mathbf{u}} \in \mathbf{V}_0$, the variational problem (2.11) can be written as: find $\mathbf{u}_0 \in \mathbf{V}_0$ such that, for any $\mathbf{v} \in \mathbf{V}_0$,

$$a(\mathbf{u}_0, \mathbf{v}) = L(\mathbf{v}),$$

where

$$L(\mathbf{v}) = \int_{\Omega} \mathbf{F} \cdot \bar{\mathbf{v}} \, d\mathbf{x} - \int_{\Omega} \mu^{-1} \nabla \times \tilde{\mathbf{u}} \cdot \nabla \times \bar{\mathbf{v}} \, d\mathbf{x} - i\omega \int_{\Omega} \sigma \tilde{\mathbf{u}} \cdot \bar{\mathbf{v}} \, d\mathbf{x}.$$

From Proposition A.4, the bilinear form $a(\cdot, \cdot) : \mathbf{V}_0 \times \mathbf{V}_0 \rightarrow \mathbb{R}$ is continuous and \mathbf{V} -elliptic. Moreover, the linear functional $L(\cdot) : \mathbf{V}_0 \rightarrow \mathbb{R}$ is continuous in the \mathbf{V} -norm. Existence and uniqueness of the solution in $\mathbf{u}_0 \in \mathbf{V}_0$, as well continuous dependence in the \mathbf{V} -norm on the data \mathbf{F} and $\tilde{\mathbf{u}}$ then follow from Lax–Milgram’s lemma. We conclude the existence and uniqueness of the solution $\mathbf{u} \in \mathbf{V}$ and the stability estimate

$$\|\mathbf{u}\|_{\mathbf{V}} \leq C \|\mathbf{F}\|_{0, \Omega} + C \|\tilde{\mathbf{u}}\|_{\mathbf{V}} + \|\tilde{\mathbf{u}}_f\|_{\mathbf{V}} \leq C \|\mathbf{F}\|_{0, \Omega} + C \|\mathbf{g}\|_{\mathcal{H}(\partial\Omega)},$$

where we used Proposition A.1 and the properties of $\tilde{\mathbf{u}}_f$.

To obtain the equivalence of strong and variational form of our model problem we proceed as follows. Integrating by parts and taking into account the properties of f , it is obvious that, if \mathbf{u} solves (2.7)–(2.10), then it also solves (2.11), for any $\mathbf{v} \in \mathbf{V}_0$. In order to prove the converse, define

$$\begin{aligned} \mathbf{V}_1 &= \{\mathbf{v} \in H_0(\text{curl}; \Omega) : \mathbf{v}|_{\Omega_0} \in \nabla H_0^1(\Omega_0), \mathbf{v}|_{\Omega_\sigma} = \mathbf{0}\} \\ \mathbf{V}_2 &= \{\mathbf{v} \in H_0(\text{curl}; \Omega) : \mathbf{v}|_{\Omega_0} \in H_0(\text{curl}^0, \text{div}^0; \Omega_0), \mathbf{v}|_{\Omega_\sigma} = \mathbf{0}\} \\ \mathbf{V}_3 &= \{\mathbf{v} \in H_0(\text{curl}; \Omega) : \mathbf{v}|_{\Omega_0} \in H_{\text{flux}}(\text{div}^0; \Omega_0)\}. \end{aligned}$$

Then

$$H_0(\text{curl}; \Omega) = \mathbf{V}_1 \oplus \mathbf{V}_2 \oplus \mathbf{V}_3 \tag{A.6}$$

$$\mathbf{V}_0 = (\mathbf{V}_1 \oplus \mathbf{V}_2) \cap H_{\text{flux}}(\text{div}; \Omega_0) \oplus \mathbf{V}_3. \tag{A.7}$$

For test functions $\mathbf{v} \in \mathbf{V}_3$, the variational problem (2.11) becomes

$$\int_{\Omega} \mu^{-1} \nabla \times \mathbf{u} \cdot \nabla \times \bar{\mathbf{v}} \, d\mathbf{x} + i\omega \int_{\Omega_\sigma} \sigma \mathbf{u} \cdot \bar{\mathbf{v}} \, d\mathbf{x} = \int_{\Omega} \mathbf{F}' \cdot \bar{\mathbf{v}} \, d\mathbf{x}, \tag{A.8}$$

recalling that $\mathbf{F} = \mathbf{F}' + \nabla f$. Since for any $\mathbf{v} \in \mathbf{V}_1 \oplus \mathbf{V}_2$, $\nabla \times \mathbf{v} = \mathbf{0}$ in Ω , $\mathbf{v} = \mathbf{0}$ in Ω_σ and \mathbf{v} is L^2 -orthogonal to \mathbf{F}' , owing to decomposition (A.6) we can take any $\mathbf{v} \in H_0(\text{curl}; \Omega)$ in (A.8) without altering the problem, and obtain that (2.7) is satisfied in the sense of the distributions. Since $\mathbf{F}' \in L^2(\Omega)^3$ and $\sigma \mathbf{u} \in L^2(\Omega)^3$, we have that $\nabla \times (\mu^{-1} \nabla \times \mathbf{u}) \in L^2(\Omega)^3$ and (2.7) holds true almost everywhere.

For test functions $\mathbf{v} \in (\mathbf{V}_1 \oplus \mathbf{V}_2) \cap H_{\text{flux}}(\text{div}; \Omega_0)$, (2.11) becomes

$$\int_{\Omega_0} \nu \nabla \cdot \mathbf{u} \nabla \cdot \bar{\mathbf{v}} \, d\mathbf{x} = \int_{\Omega_0} \nabla f \cdot \bar{\mathbf{v}} \, d\mathbf{x}. \quad (\text{A.9})$$

From (2.6), $(\mathbf{F}_1 + \mathbf{F}_2)|_{\Omega_0} = \nabla f$, with $f \in H^1(\Omega_0)$, $f = 0$ on $\Gamma_{0,0}$ and f constant on each $\Gamma_{0,j}$, $j = 1, \dots, J$. By integrating by parts the right-hand side, taking into account the properties of f and the flux conditions on \mathbf{v} , we have

$$\int_{\Omega_0} \nu \nabla \cdot \mathbf{u} \nabla \cdot \bar{\mathbf{v}} \, d\mathbf{x} = - \int_{\Omega_0} f \nabla \cdot \bar{\mathbf{v}} \, d\mathbf{x}, \quad (\text{A.10})$$

for any $\mathbf{v} \in (\mathbf{V}_1 \oplus \mathbf{V}_2) \cap H_{\text{flux}}(\text{div}; \Omega_0)$. We can take in (A.10) test functions $\mathbf{v} \in (\mathbf{V}_1 \oplus \mathbf{V}_2) \cap H(\text{div}; \Omega_0)$ without altering the problem. In fact, given $\mathbf{v} \in (\mathbf{V}_1 \oplus \mathbf{V}_2) \cap H(\text{div}; \Omega_0)$, there is a (unique) $\mathbf{v}_\beta \in H_0(\text{curl}^0, \text{div}^0; \Omega_0)$ such that $\langle (\mathbf{v}_\beta|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}} = \langle (\mathbf{v}|_{\Omega_0} \cdot \mathbf{n}_{0,j}), 1 \rangle_{\Gamma_{0,j}}$, $j = 1, \dots, J$ (see, e.g., [31], Lemma 3.2, with $\Gamma_\nu = \partial\Omega_0$ and $\Gamma_\tau = \emptyset$). Naming again \mathbf{v}_β the trivial extension of \mathbf{v}_β to Ω_σ , we have that $\mathbf{v}_0 = \mathbf{v} - \mathbf{v}_\beta$ belongs to $(\mathbf{V}_1 \oplus \mathbf{V}_2) \cap H_{\text{flux}}(\text{div}; \Omega_0)$, and $\nabla \cdot \mathbf{v} = \nabla \cdot \mathbf{v}_0$ in Ω_0 . Consequently, equation (A.10) holds true also for $\mathbf{v} \in (\mathbf{V}_1 \oplus \mathbf{V}_2) \cap H(\text{div}; \Omega_0)$.

Now, for any $\varphi \in L^2(\Omega_0)$, let ψ be the (unique) solution in $H_0^1(\Omega_0)$ of the problem $\Delta\psi = \varphi$ in Ω_0 , $\psi = 0$ on $\partial\Omega_0$. Denoting again by ψ the trivial extension of ψ to Ω_σ , we have that $\nabla\psi$ belong to $H_0(\text{curl}; \Omega) \cap (\nabla H_0^1(\Omega_0) \cap H(\text{div}; \Omega_0))$, which is contained in $(\mathbf{V}_1 \oplus \mathbf{V}_2) \cap H(\text{div}; \Omega_0)$. Then, we can take $\mathbf{v} = \nabla\psi$ as test function in (A.10) and obtain that, for any $\varphi \in L^2(\Omega_0)$, $\int_{\Omega_0} \nu \nabla \cdot \mathbf{u} \bar{\varphi} \, d\mathbf{x} = - \int_{\Omega_0} f \bar{\varphi} \, d\mathbf{x}$, from which (2.8) follows, along with the regularity property $\nu \nabla \cdot \mathbf{u}|_{\Omega_0} \in H^1(\Omega_0)$.

This completes the proof of Theorem 2.3.

REFERENCES

- [1] A. Alonso and A. Valli, *A domain decomposition approach for heterogeneous time-harmonic Maxwell equations*, Comput. Meth. Appl. Mech. Engrg. **143** (1997), 97–112.
- [2] ———, *An optimal domain decomposition preconditioner for low-frequency time-harmonic Maxwell equations*, Math. Comp. **68** (1999), 607–631.
- [3] C. Amrouche, C. Bernardi, M. Dauge, and V. Girault, *Vector potentials in three-dimensional non-smooth domains*, Math. Models Appl. Sci. **21** (1998), 823–864.
- [4] D.N. Arnold, *An interior penalty finite element method with discontinuous elements*, SIAM J. Numer. Anal. **19** (1982), 742–760.
- [5] D.N. Arnold, F. Brezzi, B. Cockburn, and L.D. Marini, *Unified analysis of discontinuous Galerkin methods for elliptic problems*, SIAM J. Numer. Anal. (2001), to appear.
- [6] I. Babuška and M. Suri, *The hp -version of the finite element method with quasiuniform meshes*, Modél. Math. Anal. Numér. **21** (1987), 199–238.
- [7] C.E. Baumann and J.T. Oden, *A discontinuous hp -finite element method for convection-diffusion problems*, Comput. Methods Appl. Mech. Engrg. **175** (1999), 311–341, Special issue on Spectral, Spectral Element, and hp Methods in CFD, edited by G.E. Karniadakis, M. Ainsworth and C. Bernardi.
- [8] A.S. Bonnet-BenDhia, C. Hazard, and S. Lohrengel, *A singular field method for the solution of Maxwell's equations in polyhedral domains*, SIAM J. Appl. Math. **59** (1999), 2028–2044.
- [9] A. Bossavit, *A rationale for edge-elements in 3D fields computation*, IEEE Trans. on Magnetics **24** (1988), 74–79.
- [10] ———, *Computational electromagnetism*, Academic Press, San Diego, 1998.

- [11] A. Bossavit, “Hybrid” electric–magnetic methods in eddy–current problems, *Comput. Methods Appl. Mech. Engrg.* **178** (1999), 383–391.
- [12] C.F. Bryant, C.R.I. Emson, P. Fernandes, and C.W. Trowbridge, *Lorentz gauge eddy current formulations for multiply connected piecewise homogeneous conductors*, *COMPEL* **17** (1998), 732–740.
- [13] A. Buffa, *Hodge decomposition on the boundary of a polyhedron: the non–simply connected case*, *Math. Models Methods Appl. Sci.* (2000), to appear.
- [14] A. Buffa and Jr. P. Ciarlet, *On traces for functional spaces related to Maxwell’s equations. Part I: An integration by parts formula in Lipschitz polyhedra*, *Math. Models Appl. Sci.* **24** (2001), to appear.
- [15] ———, *On traces for functional spaces related to Maxwell’s equations. Part II: Hodge decomposition on the boundary of Lipschitz polyhedra and applications polyhedra*, *Math. Models Appl. Sci.* **24** (2001), to appear.
- [16] P. Castillo, *Local discontinuous Galerkin methods for convection–diffusion and elliptic problems*, Ph.D. thesis, University of Minnesota, Minneapolis, MN, 2001.
- [17] P. Castillo, *Performance of discontinuous Galerkin methods for elliptic partial differential equations*, Tech. Report 1764, IMA, University of Minnesota, 2001.
- [18] P. Castillo, B. Cockburn, I. Perugia, and D. Schötzau, *An a priori error analysis of the local discontinuous Galerkin method for elliptic problems*, *SIAM J. Numer. Anal.* **38** (2000), 1676–1706.
- [19] P. Castillo, B. Cockburn, D. Schötzau, and C. Schwab, *Optimal a priori error estimates for the hp–version of the local discontinuous Galerkin method for convection–diffusion problems*, *Math. Comp.* (2001), to appear.
- [20] Z. Chen, Q. Du, and J. Zou, *Finite element methods with matching and nonmatching meshes for Maxwell equations with discontinuous coefficients*, *SIAM J. Numer. Anal.* **37** (2000), 1542–1570.
- [21] P. Ciarlet, *The finite element method for elliptic problems*, North Holland, 1978.
- [22] B. Cockburn and C. Dawson, *Some extensions of the local discontinuous Galerkin method for convection–diffusion equations in multidimensions*, The Proceedings of the 10th Conference on the Mathematics of Finite Elements and Applications: MAFELAP X (J. Whitemann, ed.), Elsevier, 2000, pp. 225–238.
- [23] B. Cockburn, G. Kanschat, I. Perugia, and D. Schötzau, *Superconvergence of the local discontinuous Galerkin method for elliptic problems on Cartesian grids*, *SIAM J. Numer. Anal.*, to appear.
- [24] B. Cockburn, G. Kanschat, D. Schötzau, and C. Schwab, *Local discontinuous Galerkin methods for the Stokes system*, Tech. Report 214, Minnesota Supercomputing Institute, University of Minnesota, 2000.
- [25] B. Cockburn and C.-W. Shu, *Runge–Kutta discontinuous Galerkin methods for convection–dominated problems*, *J. Sci. Comput.*, to appear.
- [26] ———, *The local discontinuous Galerkin method for time–dependent convection–diffusion systems*, *SIAM J. Numer. Anal.* **35** (1998), 2440–2463.
- [27] M. Costabel and M. Dauge, *Weighted regularization of Maxwell equations in polyhedral domains*, Tech. Report 26, IRMAR, Université de Rennes, 2001.
- [28] M. Costabel, M. Dauge, and S. Nicaise, *Singularities of Maxwell interface problems*, *Modél. Math. Anal. Numér.* **33** (1999), 627–649.
- [29] L. Demkowicz and L. Vardapetyan, *Modeling of electromagnetic absorption/scattering problems using hp–adaptive finite elements*, *Comput. Methods Appl. Mech. Engrg.* **152** (1998), 103–124.
- [30] P. Fernandes and G. Gilardi, *Magnetostatic and electrostatic problems in inhomogeneous anisotropic media with irregular boundary and mixed boundary conditions*, *Math. Models Methods Appl. Sci.* **7** (1997), 957–991.
- [31] P. Fernandes and I. Perugia, *Vector potential formulation for magnetostatics and modeling of permanent magnets*, *IMA J. Appl. Math.* (2001), to appear.
- [32] E.H. Georgoulis and E. Süli, *hp–DGFEM on shape–irregular meshes: reaction–diffusion*, Tech. Report NA 01–09, Oxford University Computing Laboratory, 2001.
- [33] R. Hiptmair, *Symmetric coupling for eddy–current problems*, Tech. Report 148, SFB382, Universität Tübingen, 2000, (submitted).
- [34] P. Houston, C. Schwab, and E. Süli, *Discontinuous hp finite element methods for advection–diffusion problems*, Tech. Report NA 00–15, Oxford University Computing Laboratory, 2000, (submitted).
- [35] J. L. Lions and E. Magenes, *Problèmes aux limites non–homogènes et applications*, Dunod, Paris, 1968.

- [36] P. Monk, *A finite element method for approximating the time-harmonic Maxwell equations*, Numer. Math. **63** (1992), 243–261.
- [37] J. C. Nédélec, *Mixed finite elements in \mathbb{R}^3* , Numer. Math. **35** (1980), 315–341.
- [38] ———, *Éléments finis mixtes incompressibles pour l'équation de Stokes dans \mathbb{R}^3* , Numer. Math. **39** (1982), 97–112.
- [39] J.T. Oden, Ivo Babuška, and C.E. Baumann, *A discontinuous hp finite element method for diffusion problems*, J. Comput. Phys. **146** (1998), 491–519.
- [40] I. Perugia and D. Schötzau, *On the coupling of local discontinuous Galerkin and conforming finite element methods*, J. Sci. Comp. (2001), to appear.
- [41] S. Prudhomme, F. Pascal, J.T. Oden, and A. Romkes, *Review of a priori error estimation for discontinuous Galerkin methods*, Tech. Report 2000-27, TICAM, University of Texas at Austin, 2000.
- [42] B. Rivière and M.F. Wheeler, *A discontinuous Galerkin method applied to nonlinear parabolic equations*, Discontinuous Galerkin Methods: Theory, Computation and Applications (B. Cockburn, G.E. Karniadakis, and C.-W. Shu, eds.), Lect. Notes Comput. Sci. Eng., vol. 11, Springer-Verlag, 2000, pp. 231–244.
- [43] B. Rivière, M.F. Wheeler, and V. Girault, *Part I: Improved energy estimates for interior penalty, constrained and discontinuous Galerkin methods for elliptic problems*, Computational Geosciences **3** (4) (1999), 337–360.
- [44] C. Schwab, *p - and hp -FEM – Theory and Application to Solid and Fluid Mechanics*, Oxford University Press, Oxford, 1998.
- [45] L. Vardapetyan and L. Demkowicz, *hp -adaptive finite elements in electromagnetics*, Comput. Methods Appl. Mech. Engrg. **152** (1998), 103–124.