

Models for the 3D singular isotropic oscillator quadratic algebra

E. G. Kalnins,¹ W. Miller, Jr.,² and S. Post²

¹*Department of Mathematics, University of Waikato, Hamilton, New Zealand.*

²*School of Mathematics, University of Minnesota, Minneapolis, Minnesota, U.S.A.*

We give the first explicit construction of the quadratic algebra for a 3D quantum superintegrable system with nondegenerate (4-parameter) potential together with realizations of irreducible representations of the quadratic algebra in terms of differential-differential or differential-difference and difference-difference operators in two variables. The example is the singular isotropic oscillator. We point out that the quantum models arise naturally from models of the Poisson algebras for the corresponding classical superintegrable system. These techniques extend to quadratic algebras for superintegrable systems in n dimensions and are closely related to Hecke algebras and multivariable orthogonal polynomials.

PACS numbers: 02.00.00, 02.20.Qs, 02.30.Ik, 03.65.Fd

I. INTRODUCTION

The distinct classical and quantum second order superintegrable systems on real or complex 3D flat space and with nondegenerate (4-parameter) potentials have been classified [1]. (Recall that a second order superintegrable system in n dimensions is one that admits $2n - 1$ functionally independent constants of the motion quadratic in the momentum variables, the maximum possible, [2–4].) Indeed, the classification for nondegenerate potentials on 3D conformally flat spaces is virtually complete, [5–9]. Characteristic of these systems in all dimensions is that the second order constants of the motion generate a finite dimensional algebra, polynomially closed under commutation, the quadratic algebra. In several recent papers [10, 11] for the 2D cases, the authors have launched a study of the irreducible representations of these algebras and their applications via models of the representations, in terms of differential and difference operators. (Some earlier work on this subject can be found in [13–18].) For the 3D case where $2n-1=5$, we have shown that in fact there are always 6 linearly independent second order symmetries and that these generate a quadratic

algebra closing at order 6 in the momenta. The second order generators are always functionally dependent via a polynomial relation at order 8. In this paper we present, for the first time, the details for a nontrivial quadratic algebra in 3D: the singular isotropic oscillator. Two-variable models for irreducible representations of the quantum system follow directly from models for the classical system. There are three possible models expressed in differential or difference operators, corresponding to separation of the eigenvalue equation for the Schrödinger operator in Cartesian, cylindrical or spherical coordinates. The models have important connections with the theory of dual Hahn and Wilson polynomials. This is a prototype for the general study of the representations of 3D quadratic algebras.

II. THE QUANTUM SUPERINTEGRABLE SYSTEM

The Hamiltonian operator is

$$H = \partial_1^2 + \partial_2^2 + \partial_3^2 + a^2(x_1^2 + x_2^2 + x_3^2) + \frac{b_1}{x_1^2} + \frac{b_2}{x_2^2} + \frac{b_3}{x_3^2} \quad \partial_i \equiv \partial_{x_i}. \quad (1)$$

A basis for the second order constants of the motion is (with $H = M_1 + M_2 + M_3$.)

$$M_\ell = \partial_\ell^2 + a^2 x_\ell^2 + \frac{b_\ell}{x_\ell^2}, \quad \ell = 1, 2, 3, \quad L_i = (x_j \partial_k - x_k \partial_j)^2 + \frac{b_j x_k^2}{x_j^2} + \frac{b_k x_j^2}{x_k^2}, \quad (2)$$

where i, j, k are pairwise distinct and run from 1 to 3. There are 4 linearly independent commutators of the second order symmetries:

$$S_1 = [L_1, M_2] = [M_3, L_1], \quad S_2 = -[M_3, L_2] = [M_1, L_2], \quad (3)$$

$$S_3 = -[M_1, L_3] = [M_2, L_3], \quad R = [L_1, L_2] = [L_2, L_3] = [L_3, L_1],$$

$$[M_i, M_j] = [M_i, L_i] = 0, \quad 1 \leq i, j \leq 3.$$

Here we define the commutator of linear operators F, G by $[F, G] = FG - GF$. (Thus a second order constant of the motion is a second order partial differential operator K in the variables x_j such that $[K, H] = 0$, where 0 is the zero operator.)

The fourth order structure equations are $[M_i, S_i] = 0$, $1 = 1, 2, 3$, and

$$\epsilon_{ijk}[M_i, S_j] = 8M_i M_k - 16a^2 L_j + 8a^2, \quad \epsilon_{ijk}[M_i, R] = 8(M_j L_j - M_k L_k) + 4(M_k - M_j), \quad (4)$$

$$\epsilon_{ijk}[S_i, L_j] = 8M_i L_i - 8M_k L_k + 4(M_k - M_i),$$

$$\epsilon_{ijk}[L_i, S_i] = 4\{L_i, M_k - M_j\} + 16b_j M_k - 16b_k M_j + 8(M_k - M_j),$$

$$\epsilon_{ijk}[L_i, R] = 4\{L_i, L_k - L_j\} - 16b_j L_j + 16b_k L_k + 8(L_k - L_j + b_j - b_k).$$

Here, $\{F, G\} = FG + GF$ and ϵ_{ijk} is the completely antisymmetric tensor.

The fifth order structure equations are obtainable directly from the fourth order equations and the Jacobi identity. The sixth order structure equations are

$$S_i^2 - \frac{8}{3}\{L_j, M_j, M_k\} + 16a^2 L_i^2 + (16b_k + 12)M_j^2 + (16b_j + 12)M_k^2 - \frac{104}{3}M_j M_k - \frac{176}{3}a^2 L_i - \frac{16}{3}a^2(2 + 9b_j + 9b_k + 12b_j b_k) = 0, \quad (5)$$

$$\frac{1}{2}\{S_i, S_j\} + \frac{4}{3}\{L_i, M_k, M_i\} + \frac{4}{3}\{L_j, M_k, M_j\} - 8L_k M_k^2 - 8a^2\{L_i, L_j\} - (16b_k + 12)M_i M_j$$

$$+ 4M_k^2 - 4M_k(M_i + M_j) + a^2(32b_k + 24)L_k + 8a^2(L_i + L_j) - 16a^2(b_k + 1) = 0,$$

$$\frac{1}{2}\{S_i, R\} - 8L_i^2 M_i + \frac{4}{3}\{L_k, L_i, M_k\} + \frac{4}{3}\{L_i, L_j, M_j\} - (8b_k + 6)\{L_k, M_j\}$$

$$- (8b_j + 6)\{L_j, M_k\} - 2\{L_i, M_k + M_j\} + \frac{88}{3}L_i M_i + \frac{52}{3}(L_k M_k + L_j M_j)$$

$$+ (32b_k b_j + 24b_k + 24b_j + \frac{16}{3})M_i + (8b_j - \frac{8}{3})M_k + (8b_k - \frac{8}{3})M_j = 0,$$

$$R^2 - \frac{8}{3}\{L_1, L_2, L_3\} + (16b_1 + 12)L_1^2 + (16b_2 + 12)L_2^2 + (16b_3 + 12)L_3^2 - \frac{52}{3}\{L_1, L_2\}$$

$$- \frac{52}{3}\{L_1, L_3\} - \frac{52}{3}\{L_2, L_3\} - \frac{16}{3}(11b_1 + 1)L_1 - \frac{16}{3}(11b_2 + 1)L_2$$

$$- \frac{16}{3}(11b_3 + 1)L_3 - \frac{32}{3}\left(6b_1 b_2 b_3 + \frac{9}{2}(b_1 b_2 + b_1 b_3 + b_2 b_3) + b_1 + b_2 + b_3\right).$$

Here, $\{A, B, C\} = ABC + ACB + BAC + BCA + CAB + CBA$ and i, j, k are pairwise distinct.

The eighth order functional relation is

$$L_1^2 M_1^2 + L_2^2 M_2^2 + L_3^2 M_3^2 - \frac{1}{12}\{L_1, L_2, M_1, M_2\} - \frac{1}{12}\{L_1, L_3, M_1, M_3\} \quad (6)$$

$$- \frac{1}{12}\{L_2, L_3, M_2, M_3\} - \frac{7}{3}L_1 M_1^2 - \frac{7}{3}L_2 M_2^2 - \frac{7}{3}L_3 M_3^2 + \frac{2}{3}a\{L_1, L_2, L_3\}$$

$$- \frac{1}{18}\{L_1, M_1, M_2\} - \frac{1}{18}\{L_1, M_1, M_3\} - \frac{1}{18}\{L_2, M_1, M_2\} - \frac{1}{18}\{L_2, M_2, M_3\}$$

$$- \frac{1}{18}\{L_3, M_1, M_3\} - \frac{1}{18}\{L_3, M_2, M_3\} + \frac{1}{6}(4b_1 + 3)\{L_1, M_2, M_3\}$$

$$+ \frac{1}{6}(4b_2 + 3)\{L_2, M_1, M_3\} + \frac{1}{6}(4b_3 + 3)\{L_3, M_1, M_2\} - a^2(4b_1 + 3)L_1^2 - a^2(4b_2 + 3)L_2^2 - a^2(4b_3 + 3)L_3^2$$

$$\begin{aligned}
& +\frac{a^2}{3}(\{L_1, L_2\} + \{L_1, L_3\} + \{L_2, L_3\}) - (4b_2b_3 + 3b_2 + 3b_3 + \frac{4}{3})M_1^2 \\
& - (4b_1b_3 + 3b_1 + 3b_3 + \frac{4}{3})M_2^2 - (4b_1b_2 + 3b_1 + 3b_2 + \frac{4}{3})M_3^2 + \frac{2}{3}(b_3 + 2)M_1M_2 \\
& + \frac{2}{3}(b_2 + 2)M_1M_3 + \frac{2}{3}(b_1 + 2)M_2M_3 + \frac{4}{3}a^2(7b_1 + 4)L_1 + \frac{4}{3}a^2(7b_2 + 4)L_2 \\
& + \frac{4}{3}a^2(7b_3 + 4)L_3 + \frac{4}{3}a^2(12b_1b_2b_3 + 9b_1b_2 + 9b_1b_3 + 9b_2b_3 + 4b_1 + 4b_2 + 4b_3) = 0.
\end{aligned}$$

Here, $\{A, B, C, D\}$ is the 24 term symmetrizer of 4 operators.

A. Cartesian case: A quantum model with M_1, M_2 diagonal

For the model we choose variables u, v in which the eigenfunctions are polynomials, and write the parameters as $b_j = 1/4 - k_j^2$. Then we have

$$M_1 = 2ia(2u\partial_u + k_1 + 1), \quad M_2 = 2ia(2v\partial_v + k_2 + 1), \quad M_1 + M_2 + M_3 = E \quad (7)$$

$$\begin{aligned}
L_1 = & 4v \left(u^2 \partial_u^2 + 2uv \partial_u \partial_v + (v^2 + 1) \partial_v^2 - \left(\frac{E}{2ia} - k_1 - k_2 - 4 \right) u \partial_u - \left(\frac{E}{2ia} - k_1 - k_2 - 4 \right) \partial_v \right) \\
& + 4(1 + k_2) \partial_v + v \left(\left(-\frac{E}{2ia} + k_1 + k_2 + 3 \right)^2 - k_3^2 \right) + \frac{1}{2a^2} M_2 M_3 + \frac{1}{2}
\end{aligned} \quad (8)$$

$$\begin{aligned}
L_2 = & 4u \left(v^2 \partial_v^2 + 2uv \partial_u \partial_v + (u^2 + 1) \partial_u^2 - \left(\frac{E}{2ia} - k_1 - k_2 - 4 \right) v \partial_v - \left(\frac{E}{2ia} - k_1 - k_2 - 4 \right) u \partial_u \right) \\
& + 4(1 + k_1) \partial_u + u \left(\left(-\frac{E}{2ia} + k_1 + k_2 + 3 \right)^2 - k_3^2 \right) + \frac{1}{2a^2} M_1 M_3 + \frac{1}{2}
\end{aligned} \quad (9)$$

$$L_3 = 4(uv\partial_u^2 + uv\partial_v^2 + (k_1 + 1)v\partial_u + (k_2 + 1)u\partial_v) + \frac{1}{2a^2}M_1M_2 + \frac{1}{2}. \quad (10)$$

In the model, the monomials $f_{N,j} = u^j v^{N-j}$ are simultaneous eigenfunctions of the operators M_j :

$$M_1 f_{N,j} = 2ia(2j + k_1 + 1) f_{N,j}, \quad M_2 f_{N,j} = 2ia(2N - 2j + k_2 + 1) f_{N,j} \quad (11)$$

Further, we have the expansion formulas

$$\begin{aligned}
L_1 f_{N,j} = & \left((2N + 3 - \frac{E}{2ia} + k_1 + k_2)^2 - k_3^2 \right) f_{N+1,j} + 4(N - j)(N - j + k_2) f_{N-1,j} \\
& + \left(2\left(\frac{E}{2ia} - 2N - 2k_1 - 2k_2 - 2 \right) (2N - 2j + k_2 + 1) + \frac{1}{2} \right) f_{N,j}
\end{aligned} \quad (12)$$

$$L_2 f_{N,j} = \left((2N + 3 - \frac{E}{2ia} + k_1 + k_2)^2 - k_3^2 \right) f_{N+1,j+1} + 4j(j + k_1) f_{N-1,j-1} + \left(2\left(\frac{E}{2ia} - 2N - 2k_1 - 2k_2 - 2\right)(2j + k_1 + 1) + \frac{1}{2} \right) f_{N,j} \quad (13)$$

$$L_3 f_{N,j} = 4(N - j)(N - j + k_2) f_{N,j+1} + 4j(j + k_2) f_{N,j-1} + \left(2(2j + k_1 + 1)(2N - 2j + k_2 + 1) + \frac{1}{2} \right) f_{N,j} \quad (14)$$

From the model we can find a family of finite dimensional irreducible representations, labeled by the nonnegative integer M . A basis of eigenfunctions is given by $\{f_{N,j} = u^j v^{N-j}\}$, such that the N and j are nonnegative integers satisfying $0 \leq j \leq N \leq M$. The energy satisfies

$$E = 2ia(2M + k_1 + k_2 + k_3 + 3). \quad (15)$$

The dimension of the representation is $(M + 2)(M + 1)/2$. Now we introduce an inner product such that the operators M_j, L_j are formally self-adjoint for $j = 1, 2, 3$. This forces a to be pure imaginary.

Normalization coefficients: Let $\hat{f}_{N,j} = K_{N,j} u^j v^{N-j}$ such that $\|\hat{f}_{N,j}\| = 1$. If we assume $K_{0,0} = 1$ then the coefficients become,

$$K_{N,j} = \frac{(-M)_N (-M - k_3)_N}{(N - j)! j! (k_2 + 1)_{N-j} (k_1 + 1)_j}$$

B. Recurrence relations for Wilson polynomials

The spherical case is intimately bound up with recurrence relations for Wilson polynomials and the cylindrical case with the dual Hahn polynomials. To see this we modify some of the results of [19]. The unnormalized Wilson polynomials are

$$w_n(y^2) \equiv w_n(y^2, \alpha, \beta, \gamma, \delta) = (\alpha + \beta)_n (\alpha + \gamma)_n (\alpha + \delta)_n \times {}_4F_3 \left(\begin{matrix} -n, & \alpha + \beta + \gamma + \delta - n - 1, & \alpha - y, & \alpha + y \\ \alpha + \beta, & \alpha + \gamma, & \alpha + \delta & \end{matrix} ; 1 \right) = (\alpha + \beta)_n (\alpha + \gamma)_n (\alpha + \delta)_n \Phi_n^{(\alpha, \beta, \gamma, \delta)}(y^2), \quad (16)$$

where $(a)_n$ is the Pochhammer symbol and ${}_4F_3(1)$ is a generalized hypergeometric function of unit argument.

For fixed $\alpha, \beta, \gamma, \delta > 0$ the Wilson polynomials are orthogonal with respect to the inner product

$$\begin{aligned} \langle w_n, w_{n'} \rangle &= \frac{1}{2\pi} \int_0^\infty w_n(-y^2) w_{n'}(-y^2) \left| \frac{\Gamma(\alpha + iy)\Gamma(\beta + iy)\Gamma(\gamma + iy)\Gamma(\delta + iy)}{\Gamma(2iy)} \right|^2 dy \quad (17) \\ &= \delta_{nn'} n! (\alpha + \beta + \gamma + \delta + n - 1)_n \times \\ &\quad \frac{\Gamma(\alpha + \beta + n)\Gamma(\alpha + \gamma + n)\Gamma(\alpha + \delta + n)\Gamma(\beta + \gamma + n)\Gamma(\beta + \delta + n)\Gamma(\gamma + \delta + n)}{\Gamma(\alpha + \beta + \gamma + \delta + 2n)}. \end{aligned}$$

The Wilson polynomials $\Phi_n(y^2) \equiv \Phi_n^{(\alpha, \beta, \gamma, \delta)}(y^2)$, satisfy the three term recurrence formula

$$y^2 \Phi_n(y^2) = K(n+1, n) \Phi_{n+1}(y^2) + K(n, n) \Phi_n(y^2) + K(n-1, n) \Phi_{n-1}(y^2) \quad (18)$$

where

$$K(n+1, n) = \frac{\alpha + \beta + \gamma + \delta + n - 1}{(\alpha + \beta + \gamma + \delta + 2n - 1)(\alpha + \beta + \gamma + \delta + 2n)}, \quad (19)$$

$$\begin{aligned} K(n-1, n) &= \frac{n(\alpha + \beta + n - 1)(\alpha + \gamma + n - 1)(\alpha + \delta + n - 1)}{(\alpha + \beta + \gamma + \delta + 2n - 2)(\alpha + \beta + \gamma + \delta + 2n - 1)} \times \\ &\quad (\beta + \gamma + n - 1)(\beta + \delta + n - 1)(\gamma + \delta + n - 1), \end{aligned} \quad (20)$$

$$\begin{aligned} K(n, n) &= \alpha^2 - K(n+1, n)(\alpha + \beta + n)(\alpha + \gamma + n)(\alpha + \delta + n) \\ &\quad - \frac{K(n-1, n)}{(\alpha + \beta + n - 1)(\alpha + \gamma + n - 1)(\alpha + \delta + n - 1)}. \end{aligned} \quad (21)$$

Moreover, they satisfy the following parameter-changing recurrence relations when acting on the basis polynomials $\Phi_n \equiv \Phi_n^{(\alpha, \beta, \gamma, \delta)}$. Here $T^\tau f(y) = f(y + \tau)$.

1.

$$R = \frac{1}{2y} [T^{1/2} - T^{-1/2}], \quad R\Phi_n = \frac{n(n + \alpha + \beta + \gamma + \delta - 1)}{(\alpha + \beta)(\alpha + \gamma)(\alpha + \delta)} \Phi_{n-1}^{(\alpha+1/2, \beta+1/2, \gamma+1/2, \delta+1/2)}.$$

2.

$$\begin{aligned} L &= \frac{1}{2y} \left[(\alpha - \frac{1}{2} + y)(\beta - \frac{1}{2} + y)(\gamma - \frac{1}{2} + y)(\delta - \frac{1}{2} + y) T^{1/2} \right. \\ &\quad \left. - (\alpha - \frac{1}{2} - y)(\beta - \frac{1}{2} - y)(\gamma - \frac{1}{2} - y)(\delta - \frac{1}{2} - y) T^{-1/2} \right]. \end{aligned}$$

$$L\Phi_n = (\alpha + \beta - 1)(\alpha + \gamma - 1)(\alpha + \delta - 1) \Phi_{n+1}^{(\alpha-1/2, \beta-1/2, \gamma-1/2, \delta-1/2)}.$$

3.

$$L_{\alpha\beta} = \frac{1}{2y} \left[-(\alpha - \frac{1}{2} + y)(\beta - \frac{1}{2} + y) T^{1/2} + (\alpha - \frac{1}{2} - y)(\beta - \frac{1}{2} - y) T^{-1/2} \right].$$

$$L_{\alpha\beta}\Phi_n = -(\alpha + \beta - 1) \Phi_n^{(\alpha-1/2, \beta-1/2, \gamma+1/2, \delta+1/2)}.$$

4.

$$R^{\alpha\beta} = \frac{1}{2y} \left[-\left(\gamma - \frac{1}{2} + y\right)\left(\delta - \frac{1}{2} + y\right)T^{1/2} + \left(\gamma - \frac{1}{2} - y\right)\left(\delta - \frac{1}{2} - y\right)T^{-1/2} \right].$$

$$R^{\alpha\beta}\Phi_n = -\frac{(n + \gamma + \delta - 1)(n + \alpha + \beta)}{\alpha + \beta} \Phi_n^{(\alpha+1/2, \beta+1/2, \gamma-1/2, \delta-1/2)}.$$

The operators $L_{\alpha\gamma}, L_{\alpha\delta}, R_{\alpha\gamma}, R_{\alpha\delta}$ are obtained by obvious substitutions.

C. Recurrence relations for dual Hahn polynomials

The dual Hahn polynomials are given by the formula

$$\begin{aligned} h_n(y^2) \equiv h_n(y^2, \alpha, \beta, \gamma) &= (\alpha + \beta)_n (\alpha + \gamma)_n {}_3F_2 \left(\begin{matrix} -n, & \alpha - y, & \alpha + y \\ & \alpha + \beta, & \alpha + \gamma \end{matrix}; 1 \right) \\ &= (\alpha + \beta)_n (\alpha + \gamma)_n \Omega_n^{(\alpha, \beta, \gamma)}(y^2), \end{aligned} \quad (22)$$

Note that the discrete dual Hahn polynomials [20] are $p_j(y^2) = \Omega^{(-Q/2, (A+1)/2, (B+1)/2)}(y^2)$ in this notation. The dual Hahn polynomials are obtained from the Wilson polynomials by letting $\delta \rightarrow \infty$. Indeed

$$h_n(y^2, \alpha, \beta, \gamma) = \lim_{\delta \rightarrow \infty} \frac{w_n(y^2, \alpha, \beta, \gamma, \delta)}{(\alpha + \delta)_n}, \quad \Omega_n^{(\alpha, \beta, \gamma)}(y^2) = \lim_{\delta \rightarrow \infty} \Phi_n^{(\alpha, \beta, \gamma, \delta)}(y^2). \quad (23)$$

It follows immediately that $h_n(y^2, \alpha, \beta, \gamma)$ is symmetric in α, β, γ .

The recurrence relations for Wilson polynomials presented in the previous section go in the limit to parameter changing recurrences for dual Hahn polynomials. The three term recurrence relation for the dual Hahn polynomials is

$$(-\alpha^2 + y^2)\Omega_n(y^2) = K(n+1, n)\Omega_{n+1}(y^2) + K(n, n)\Omega_n(y^2) + K(n-1, n)\Omega_{n-1}(y^2) \quad (24)$$

where $K(n+1, n) = (n + \alpha + \beta)(n + \alpha + \gamma)$, $K(n-1, n) = n(n + \beta + \gamma - 1)$, $K(n, n) = -K(n+1, n) - K(n-1, n)$. There are 8 basic raising and lowering operators for the dual Hahn polynomials. We list them here and describe their actions on the basis polynomials $\Omega_n \equiv \Omega_n^{(\alpha, \beta, \gamma)}$.

1.

$$R = \frac{1}{2y} [T^{1/2} - T^{-1/2}], \quad R\Omega_n = \frac{n}{(\alpha + \beta)(\alpha + \gamma)} \Omega_{n-1}^{(\alpha+1/2, \beta+1/2, \gamma+1/2)}.$$

2.

$$L = \frac{1}{2y} \left[P\left(y - \frac{1}{2}\right)T^{1/2} - P\left(-y - \frac{1}{2}\right)T^{-1/2} \right].$$

$$P(y) = (\alpha + y)(\beta + y)(\gamma + y), \quad L\Omega_n = (\alpha + \beta - 1)(\alpha + \gamma - 1)\Omega_{n+1}^{(\alpha-1/2, \beta-1/2, \gamma-1/2)}.$$

3.

$$R^\alpha = \frac{1}{2y} \left[-(\beta + y - \frac{1}{2})(\gamma + y - \frac{1}{2})T^{1/2} + (\beta - y - \frac{1}{2})(\gamma - y - \frac{1}{2})T^{-1/2} \right]$$

$$R^\alpha \Omega_n = -(\beta + \gamma + n - 1)\Omega_n^{(\alpha+1/2, \beta-1/2, \gamma-1/2)},$$

4.

$$R^\beta = \frac{1}{2y} \left[-(\alpha + y - \frac{1}{2})(\gamma + y - \frac{1}{2})T^{1/2} + (\alpha - y - \frac{1}{2})(\gamma - y - \frac{1}{2})T^{-1/2} \right]$$

$$R^\beta \Omega_n = -(\alpha + \gamma - 1)\Omega_n^{(\alpha-1/2, \beta+1/2, \gamma-1/2)},$$

5.

$$R^\gamma = \frac{1}{2y} \left[-(\alpha + y - \frac{1}{2})(\beta + y - \frac{1}{2})T^{1/2} + (\alpha - y - \frac{1}{2})(\beta - y - \frac{1}{2})T^{-1/2} \right].$$

$$R^\gamma \Omega_n = -(\alpha + \beta - 1)\Omega_n^{(\alpha-1/2, \beta-1/2, \gamma+1/2)},$$

6.

$$L_\alpha = \frac{1}{2y} \left[-(\alpha - \frac{1}{2} + y)T^{1/2} + (\alpha - \frac{1}{2} - y)T^{-1/2} \right], \quad L_\alpha \Omega_n = -\Omega_n^{(\alpha-1/2, \beta+1/2, \gamma+1/2)}$$

7.

$$L_\beta = \frac{1}{2y} \left[-(\beta - \frac{1}{2} + y)T^{1/2} + (\beta - \frac{1}{2} - y)T^{-1/2} \right],$$

$$L_\beta \Omega_n = -\left(\frac{n + \alpha + \gamma}{\alpha + \gamma} \right) \Omega_n^{(\alpha+1/2, \beta-1/2, \gamma+1/2)}$$

8.

$$L_\gamma = \frac{1}{2y} \left[-(\gamma - \frac{1}{2} + y)T^{1/2} + (\gamma - \frac{1}{2} - y)T^{-1/2} \right],$$

$$L_\gamma \Omega_n = -\left(\frac{n + \alpha + \beta}{\alpha + \beta} \right) \Omega_n^{(\alpha+1/2, \beta+1/2, \gamma-1/2)}$$

D. Cylindrical case: A quantum model with L_3, M_3 diagonal

For the model of the finite dimensional irreducible representation (15) the basis eigenfunctions of M_1 and M_2 have the form $\psi_{N,j} = d_{N,j} t^N \delta(y - \lambda_j)$, $\lambda_j = j + \frac{k_1+k_2+1}{2}$. We use the differential operator ∂_t and the difference operator $T^\tau(f(y, t)) = f(y + \tau, t)$ to construct our model operators. Setting,

$$\alpha = -N - \frac{k_1 + k_2 + 1}{2} \quad \beta = \frac{k_1 + k_2 + 1}{2} \quad \gamma = \frac{k_1 - k_2 + 1}{2},$$

we have the model

$$M_1 = 2ia(2\mathbf{L}R - 2t\partial_t + \beta - \gamma) \quad (25)$$

$$M_3 = 2ia(2t\partial_t - 2M - k_3 + 1) \quad (26)$$

$$M_1 + M_2 + M_3 = E \quad (27)$$

$$L_1 = \frac{1}{2} \left(\frac{1}{t} R^\alpha R + 8t(M - t\partial_t)(M - t\partial_t - k_3) \mathbf{L}_\alpha \mathbf{L} + \frac{1}{2\alpha^2} M_2 M_3 + \frac{1}{2} \right) \quad (28)$$

$$L_2 = \frac{1}{2} \left(\frac{1}{t} L_\beta L_\gamma + 8t(M - t\partial_t)(M - t\partial_t - k_3) \mathbf{R}^\beta \mathbf{R}^\gamma + \frac{1}{2\alpha^2} M_1 M_3 + \frac{1}{2} \right) \quad (29)$$

$$L_3 = -4y^2 + k_1^2 + k_2^2 \quad (30)$$

Here R, L, R^α, \dots are the difference operators for dual Hahn polynomials defined in Section II C. An operator in bold face, e.g., \mathbf{R}^β indicates that in the expression for the y -difference operator the parameter $\alpha = -N - (k_1 + k_2 + 1)/2$ is replaced by the differential operator $-t\partial_t - (k_1 + k_2 + 1)/2$. Note that the dual Hahn polynomials $\Omega_n^{(\alpha, \beta, \gamma)}(y^2)t^N$ are simultaneous eigenfunctions of operators M_1 and M_3 in this model.

E. Spherical case: A quantum model with $L_1 + L_2 + L_3, L_3$ diagonal

For the model of the finite dimensional irreducible representation (15) the basis eigenfunctions of L_3 and $L_1 + L_2 + L_3$ have the form $\phi_{N,j} = C_{N,j} \delta(y - \lambda_j) \delta(z - \lambda_N)$ where the support of the finite measure is $\lambda_j = j + \frac{k_1+k_2+1}{2}$ $\lambda_N = N + 1 - \frac{k_1+k_2-k_3}{2}$. We use the difference operators $T^\tau(f(y, z)) = f(y + \tau, z)$ and $Z^\tau(f(y, z)) = f(y, z + \tau)$ to construct our model operators. Setting

$$\alpha = z + \frac{k_3 + 1}{2} \quad \delta = -z + \frac{k_3 + 1}{2} \quad \beta = \frac{k_1 - k_2 + 1}{2} \quad \gamma = \frac{1 - k_1 - k_2}{2},$$

we find the model

$$L_1 = 2LR + (\alpha + \delta)(\beta + \gamma) \quad (31)$$

$$L_3 = -4y^2 + 2(\gamma^2 - \gamma + \beta^2 - \beta) + 1, \quad (32)$$

$$L_1 + L_2 + L_3 = 4z^2 + k_1^2 - k_2^2 - k_3 + \frac{3}{2} = -4(\alpha\delta - \gamma\beta) + 2(\alpha + \delta + \gamma + \beta) - \frac{1}{2}, \quad (33)$$

$$\frac{M_1}{2ia} = \frac{2(\alpha + \delta - 5 - 2M + 2\gamma)}{(\alpha - \delta - 1)(\alpha - \delta + 1)} (LR + 1 + 2\gamma\beta - \beta - \gamma(\gamma + \beta - 1)(\alpha + \delta) + 2\alpha\delta) \quad (34)$$

$$+ 2Z \frac{2 - \delta + M - \gamma}{(\alpha - \delta)(\alpha - \delta - 1)} L_{\alpha\beta} L_{\alpha\gamma} + 2Z^{-1} \frac{2 - \delta - \gamma + M}{(\alpha - \delta)(\alpha - \delta + 1)} R^{\alpha\beta} R^{\alpha\gamma},$$

$$\frac{M_3}{2ia} = \frac{\alpha^3 + \delta^2 - 2(\alpha^2 + \delta^2)(M - \gamma + 2) + 3(\alpha^2\delta + \alpha\delta^2 - 2\alpha\delta) + (\alpha + \delta)(2M + 2\gamma + 7) - 2}{(\alpha - \delta + 1)(\alpha - \delta - 1)} \quad (35)$$

$$- \frac{2(y - \alpha + 1)(y + \alpha - 1)(2 - \delta + M - \gamma)}{(\alpha - \delta)(\alpha - \delta - 1)} T - \frac{2(y - \delta + 1)(y + \delta - 1)(2 - \delta + M - \gamma)}{(\alpha - \delta)(\alpha - \delta - 1)} T^{-1}$$

$$M_1 + M_2 + M_3 = E \quad (36)$$

where the difference operators L, R , etc., are defined in section II B. Note that the Wilson polynomials (actually Racah polynomials for finite dimensional representations) $\Phi_n^{(\alpha, \beta, \gamma, \delta)}(y^2)\delta(z - \lambda_N)$ are simultaneous eigenfunctions of operators L_1 and $L_1 + L_2 + L_3$ in this model.

III. DISCUSSION AND OUTLOOK

We exhibited, for the first time, three models of the quadratic algebra for the 3D singular isotropic oscillator superintegrable system: a differential-differential, a differential-difference and a difference-difference operator model. We have presented the final results in the simplest form possible, to show the recurrence relation structure of Wilson (Racah) and dual Hahn polynomials. The models are associated with diagonalization of the operators responsible for separation of the Schrödinger eigenvalue problem in Cartesian, cylindrical and spherical coordinates, respectively. The derivation of these results, particularly the spherical model, is complicated; complete details will appear in [21]. First of all, although the general structure of the quadratic algebra follows from general theorems, the exact structure equations (4)-(6) are nontrivial to derive. A model must satisfy identically all of the quadratic algebra structure equations up to 8th order. The possibility and general form of each model follows from analysis of the Poisson algebra generated by the classical superintegrable system, just

as in the 2D cases, [10],[11]. Then the classical model must be quantized. For example it is possible to construct a classical two variable Cartesian model for which $M_1 = A, M_2 = B$ and $M_3 = E - A - B$ and

$$\begin{aligned} L_1 &= \frac{1}{4\alpha^2} [(4\delta\alpha - C_2) \exp(4i\alpha P_B) + (4\gamma\alpha - B_2) \exp(-4i\alpha P_B) + 2BC], \\ L_2 &= \frac{1}{4\alpha^2} [(4\delta\alpha - C_2) \exp(4i\alpha P_A) + (4\beta\alpha - A_2)] \exp(-4i\alpha P_A) + 2AC], \\ L_3 &= \frac{1}{4\alpha^2} [(4\beta\alpha - A_2) \exp(4i\alpha(P_A - P_B)) + (4\gamma\alpha - B_2)] \exp(-4i\alpha(P_A - P_B)) + 2AB] \end{aligned}$$

with P_A and P_B the canonical momenta conjugate to A and B . Here, all terms except the variables A, B, P_A, P_B are constants. This model then suggests the quantum Cartesian model via the quantisation rules $A \rightarrow \partial_A$ and $B \rightarrow \partial_B$. The deep connection between algebras generated by recurrences of families of orthogonal polynomials and the hidden algebras of quantum superintegrable systems may seem somewhat surprising, since the polynomials do not appear as eigenfunctions of the original quantum Hamiltonian. Rather they appear as connection coefficients between eigenfunctions of two different sets of commuting symmetry operators.

Much remains to be done. First, there are obvious connections between our models and Hecke algebras [22–24] and these relations have yet to be explored. Also, we have exhibited only the finite dimensional irreducible representations (associated with Racah polynomials) but there are also classes of infinite dimensional irreducible representations (associated with general Wilson polynomials). Intertwining operators mapping the models to the original quantum system remain to be constructed. Other nondegenerate 3D systems remain to be studied, particularly the generic potential on the 3-sphere. In that case the model will be expressible in terms of the recurrences for a family of two-variable Wilson polynomials. There are also families of degenerate 3D superintegrable systems. These have yet to be completely classified and their symmetry algebras studied. Then the study must be extended to n D second order superintegrable systems on conformally flat spaces. The formulation of a q version of superintegrable systems remains open.

-
- [1] E. G. Kalnins, W. Miller, Jr., G. C. Williams and G. S. Pogosyan, *J. Phys. A: Math Gen.* **35**, 4655 (2002).

- [2] N. W. Evans, *Phys. Rev. A.* **41**, 5666 (1990).
- [3] M. F. Rañada, *J. Math. Phys.* **38**, 4165 (1997).
- [4] *Superintegrability in Classical and Quantum Systems*, P. Tempesta, P. Winternitz, W. Miller, G. Pogosyan editors, AMS, 37, 2005.
- [5] E. G. Kalnins, J. M. Kress and W. Miller, Jr., *J. Math. Phys.* **46**, 053509 (2005).
- [6] E. G. Kalnins, J. M. Kress and W. Miller, Jr., *J. Math. Phys.* **47**, 043514 (2006).
- [7] C. Daskaloyannis and Y. Tanoudis, *J. Math Phys.* **48**, 072108 (2007)
- [8] E. G. Kalnins, J. M. Kress, W. Miller, Jr. and P. Winternitz, *J. Math. Phys.* **44**, 5811 (2003).
- [9] J. M. Kress, *Phys. Atomic Nuclei* **70**, 560 (2007).
- [10] E. G. Kalnins, W. Miller, Jr. and S. Post, *J. Phys. A: Math. Theor.* **40**, 11525 (2007).
- [11] E. G. Kalnins, W. Miller, Jr. and S. Post, *SIGMA* **4**, 008 arXiv:0801.2848 (2008).
- [12] E. G. Kalnins, W. Miller, Jr. and S. Post, *Models of quadratic quantum algebras and their relation to classical superintegrable systems*, (submitted) (2008)
- [13] D. Bonatos, C. Daskaloyannis and K. Kokkotas, *Phys. Rev. A.* **50**, 3700 (1994).
- [14] C. Daskaloyannis, *J. Math. Phys.* **42**, 1100 (2001).
- [15] Y. I. Granovskii, A. S. Zhedanov and I. M. Lutsenko, *Theoret. and Math. Phys.* **91**, 604 (1992).
- [16] C. Daskaloyannis C., *J. Math. Phys* **42**, 1100 math-ph/0003017 (2001).
- [17] C. Daskaloyannis and K. Ypsilantis, *J. Math Phys.* **47**, 042904 (2006).
- [18] C. Quesne, *SIGMA* **3**, 067 (2007).
- [19] W. Miller, Jr. *SIAM J. Math. Anal.* **18**, 1221 (1987).
- [20] G. E. Andrews, R. Askey and R. Roy, *Special Functions. Encyclopedia of Mathematics and its Applications*, *Cambridge University Press*, Cambridge, UK, 1999.
- [21] S. Post, *University of Minnesota Ph.D. Thesis*, Second order superintegrable systems, (in preparation), 2009.
- [22] S. Sahi, *Raising and Lowering Operators for Askey-Wilson Polynomials*, *SIGMA*, **3**, 002, (2007).
- [23] T. H. Koornwinder, *The relationship between Zhedanovs algebra AW (3) and the double affine Hecke algebra in the rank one case*, *SIGMA*, **3**, 063, (2007).
- [24] T. H. Koornwinder, *Zhedanovs Algebra AW(3) and the Double Affine Hecke Algebra in the Rank One Case. II. The Spherical Subalgebra* *SIGMA*, **4**, 052, (2008).