


# Mapping Groundwater Flow in Kieffer Valley, Winona County, MN


Bethany S. Ladd<sup>1</sup> and Dr. E. Calvin Alexander Jr.<sup>2</sup>

<sup>1</sup>Brown University, <sup>2</sup>Department of Geology and Geophysics, University of Minnesota



## INTRODUCTION

The Kieffer Valley Creek is a major tributary to the South Branch of the Whitewater River (SBWR), a designated trout stream in Winona County, Minnesota. Trout streams are sustained by the constant, cool, and clear water supplied by springs, which are under increasing pressure from human activities. Greater withdrawals of groundwater may be decreasing spring flow, while domestic and agricultural pollutants are affecting water quality. These pressures are especially important in the Kieffer Valley because it is an area of active karst, in which soluble carbonate material dissolves as groundwater flows through fractures, bedding planes, and conduits. Documenting groundwater flow paths is essential in such an area because the groundwater can flow rapidly through subsurface karst features. Here, the groundwater flows through Ordovician and Cambrian sandstone and carbonate aquifers, which feed the Crystal Springs State Fish Hatchery (CSSFH) via two major springs. Because it is very difficult to establish protective measures for trout streams without understanding the layout of the groundwater flow system, this study aims to begin the delineation of the springshed surrounding the CSSFH.



## BACKGROUND

Two previous dye traces have been conducted in the Kieffer Valley. The first, in 1980, occurred after sinkholes developed in the storage pond of the Altura Waste Water Treatment Facility, causing the sewage water to drain into the ground. This study traced the effluent from the two remaining aeration lagoons, which flowed on the surface into the Kieffer Valley and sank. That water then resurged into four springs in the Kieffer Valley in about one week. Originally, the researchers thought the sink point was in the Jordan Sandstone, but as knowledge improved in the subsequent 25 years, they realized it to be in the St. Lawrence Formation. In 1980, the St. Lawrence Formation was thought to be an aquitard. The second trace took place in 2008, this time pouring the dye directly into the same stream sink in the Kieffer Valley. The dye reappeared in the same four springs again within a week, confirming that the St. Lawrence Formation functions as an aquifer. The current study injected Rhodamine WT dye in a different Kieffer Valley stream sink approximately one half mile upstream of the 2008 input site, in the lower Jordan Sandstone.

The Crystal Springs State Fish Hatchery is 1.25 miles southwest of the Kieffer Valley dye input location, although it is on the other side of the SBWR. The CSSFH is supplied by two large springs in the St. Lawrence Formation. Paul Wotzka's monitoring of the CSSFH springs has shown significant evidence of varying concentrations of herbicides and nitrates. Approximately 1.25 farther to the southwest, surface water sinks into the Coon Valley Member of the Oneota Dolomite. The current study injected Uranine HS into this sink.

## MATERIALS AND METHODS


Monitored 11 locations in and around Kieffer Valley

- Teabag-sized, integrating charcoal detectors ("bugs") anchored underwater to adsorb fluorescent dye changed about once/week from June 22, 2010 – current
- Background bugs collected for two weeks prior to dye input, June 22-July 9
- Poured two types of dye in two different stream sinks, July 9, 2010
- D11006 Chromatint Uranine HS in stream sink 1.25 mi southwest of the Crystal Springs State Fish Hatchery
- D13800 Chromatint Rhodamine WT in a Kieffer Valley stream sink alongside County Rd 114

Analyzed charcoal detectors at the University of Minnesota

- Fluorescent dye extracted using eluent solution of 70% isopropyl alcohol, 30% deionized water, and 5 grams sodium hydroxide per half liter solution
- Samples analyzed by Shimadzu RF5000U scanning spectrofluorophotometer to detect fluorescent spectra
- Spectra analyzed using PeakFit® software

### 2010 Kieffer Valley trace using Light Detection and Ranging (LiDAR) imaging data




Left, Calvin Alexander and Paul Wotzka pouring Uranine HS into a stream sink southwest of CSSFH



Right, pouring Rhodamine WT into a Kieffer Valley stream sink

### Bedrock of the area around the Kieffer Valley





### 2008 Kieffer Valley trace



## RESULTS AND DISCUSSION

### Stratigraphic dye input locations



To date, no dye has been detected from either trace. It has been one month since the dye injection. There are four potential explanations for the lack of detection.


- 1) The dye has not yet reached the detection points. We have not waited long enough. The flow is slower than hypothesized.
- 2) The dye was diluted below the detection limit. We did not use enough dye.
- 3) The dye flowed to unmonitored locations. We did not look in the right places.
- 4) The dye may have been adsorbed or chemically changed. Something happened to the dye.

In the Kieffer Valley, dye was detected in the two previous traces in about a week. Whatever the reason, this dye trace is different from the 1980 and 2008 traces. One obvious difference is the input location. The input point for the first two traces was in the St. Lawrence Formation, whereas the input point for this trace is in the lower Jordan Sandstone. This may be a different flow system, either with much slower flow or flow in a different direction.


The Uranine HS dye input was the first attempt to trace to the CSSFH springs and was the first trace in Minnesota through the Jordan Sandstone. The above four potential explanations apply to this trace as well.

Although I will be returning to my home university in August, my advisor plans to continue monitoring these springs into the fall.

### Example comparison of 2008 and 2010 fluorescent spectra



Spectrum for Spring A7 in Kieffer Valley on Oct. 15, 2008, two weeks after dye input, showing pulse of Uranine dye



Spectrum for Spring A7 in Kieffer Valley on Aug. 2, 2010, three weeks after dye input, showing no Uranine dye

## LITERATURE CITED

Alexander, E.C. Jr. and Book, P.R. (1984). "Altura Minnesota Lagoon Collapses," in (Beck, B.F. ed.) *Proceedings of the First Multidisciplinary Conference on Sinkholes*, Orlando, FL, 15-17 Oct. 1984. A.A. Balkema pp. 311-318.

Alexander, S. "Spectral Deconvolution and Quantification of Natural Organic Material and Fluorescent Tracer Dyes," in (Beck B.F.) *Proceedings of the Tenth Multidisciplinary Conference on Sinkholes*, San Antonio, TX, 24-28 Sept. 2005.

Mossler, J.H. (2008). "Paleozoic stratigraphic nomenclature for Minnesota: Minnesota Geological Survey Report of Investigations" 65, 76 p., 1 pl.

## ACKNOWLEDGEMENTS

This research was made possible by the National Science Foundation Research Experience for Undergraduates program and the Department of Geology and Geophysics of the University of Minnesota. Sincere appreciation goes to the Crystal Springs State Fish Hatchery, Mike and Kaila Fish, and Paul Wotzka for their help in Winona County. I would also like to thank Scott Alexander and Andrew Luhmann for their assistance throughout this project. Calvin Alexander acknowledges support for the logistics of these traces by a grant provided by the Minnesota Environment and Natural Resources Trust Fund as recommended by the Legislative Citizen Commission on Minnesota Resources.