

**THE GLOBAL DYNAMICS OF DISCRETE
SEMILINEAR PARABOLIC EQUATIONS**

By

C.M. Elliott

and

A.M. Stuart

IMA Preprint Series # 973

May 1992

THE GLOBAL DYNAMICS OF DISCRETE SEMILINEAR PARABOLIC EQUATIONS

C.M. Elliott¹, and A.M. Stuart^{2 3}

Abstract A class of scalar semilinear parabolic equations possessing absorbing sets, a Lyapunov functional and a global attractor is considered. The gradient structure of the problem implies that, provided all steady states are isolated, solutions approach a steady state as $t \rightarrow \infty$. The dynamical properties of various finite difference and finite element schemes for the equations are analysed. The existence of absorbing sets, bounded independently of the mesh size, is proved for the numerical methods. Discrete Lyapunov functions are constructed to show that, under appropriate conditions on the mesh parameters, numerical orbits approach steady state solutions as discrete time increases. However, it is shown that insufficient spatial resolution can introduce deceptively smooth spurious steady solutions and cause the stability properties of the true steady solutions to be incorrectly represented. Furthermore, it is also shown that the explicit Euler scheme introduces spurious solutions with period 2 in the time-step. As a result, the absorbing set is destroyed and there is initial data leading to blow-up of the scheme, however small the mesh parameters are taken. To obtain stabilization to a steady state for this scheme, it is necessary to restrict the time step in terms of the initial data and the space step. Implicit schemes are constructed for which absorbing sets and Lyapunov functions exist under restrictions on the time-step which are independent of initial data and of the space step; both one-step and multi-step (BDF) methods are studied.

Key Words: Attractors, Absorbing Sets, Lyapunov Functions

AMS(MOS) Subject Classification: 65M99, 35K57

For Submission to SIAM Journal on Numerical Analysis

¹School of Mathematical and Physical Sciences, University of Sussex, Brighton BN1 9QH, UK.

²School of Mathematical Sciences, University of Bath, Bath BA2 7AY, UK.

³Current Address: Program in Scientific Computation and Computational Mathematics, Durand 252, Stanford University, CA 94305-4040, USA

1. Introduction

In this paper we consider the following semilinear parabolic initial value problem:

$$u_t = \gamma \Delta u - f(u), \quad x \in \Omega, \quad t > 0, \quad (1.1)$$

$$u = 0, \quad x \in \partial\Omega, \quad t \geq 0, \quad (1.2)$$

$$u(x, 0) = u_0(x), \quad x \in \Omega. \quad (1.3)$$

Here Ω is a bounded domain in \mathcal{R}^d ($d \leq 3$) with a Lipschitz boundary $\partial\Omega$ and $\gamma > 0$. We shall assume that the initial data $u_0 \in K_i$, $i = 1$ or 2 where $K_1 = \{\eta \in L^2(\Omega)\}$ and $K_2 = \{\eta \in L^2(\Omega) : \|\eta\|_\infty \leq M\}$. The assumptions on the nonlinear function $f(u)$ are:

(F1) $f(\bullet) \in C^2(\mathcal{R}, \mathcal{R})$ and $f(0) = 0$.

(F2) $\exists \bar{u} > 0$ such that $\frac{f(r)}{r} > 0$ for any $r : |r| > \bar{u}$.

(F3) $F''(u) \geq -C_F$ where

$$F(u) = \int f(u) du. \quad (1.4)$$

Assumptions (F1), (F2) and (F3) are made throughout the paper unless otherwise stated. Under these assumptions the following global existence result may be proved. For initial data in K_1 this result requires an additional growth assumption such as

(G1) $|f(r)| \leq c_0 |r|^{2p-1} + c_1$.

(see [Temam, 1988]) whereas for initial data in K_2 the result utilises the theory of invariant regions (see [Smoller, 1983]).

(R1) For each $u_0 \in K_i$ there exists a unique solution of (1.1)-(1.3) which satisfies, $\forall T > 0$,

$$u \in L^2(0, T; H_0^1(\Omega)) \cap C[0, T; L^2(\Omega)]$$

and the mapping $u_0 \rightarrow u(t)$ is continuous for each $t > 0$. Hence the family of solution operators $\{S(t)\}_{t \geq 0}$ defined by $S(t)u_0 \equiv u(t)$ forms a continuous semigroup on $L^2(\Omega)$. In addition, if $u_0 \in H_0^1(\Omega) \cap K_i$ then $\forall T > 0$

$$u \in C([0, T]; H_0^1(\Omega)) \cap L^2(0, T; H^2(\Omega)).$$

It is possible to show the following results about the semigroup $S(t)$ (see [Henry, 1981], [Hale, 1988], [Matano, 1979], [Zelenyak, 1968], [Temam, 1988]):

(R2) There exist balls $B_1 = \{v \in L^2(\Omega) : |v| \leq \rho_1\}$ and $B_2 = \{v \in H_0^1(\Omega) : \|v\| \leq \rho_2\}$ which are absorbing sets for the semi-group $\{S(t)\}_{t \geq 0}$; that is there exist t_1 and t_2 depending on $\{|u_0|, \rho_1\}$ and $\{|u_0|, \rho_2\}$ respectively such that $S(t)u_0 \in B_i \forall t \geq t_i$ ($i = 1, 2$). Here $|\bullet|$ and $\|\bullet\|$ denote the standard L^2 and H_0^1 norms respectively.

(R3) There exists a global attractor $\mathcal{A} \subset L^2(\Omega)$ for the semi-group $\{S(t)\}_{t \geq 0}$; that is \mathcal{A} attracts the bounded sets of $L^2(\Omega)$, and \mathcal{A} is bounded in $H_0^1(\Omega) \cup L^\infty(\Omega)$ and compact and connected in $L^2(\Omega)$.

(R4) The functional

$$I(v) := \int_{\Omega} \left[\frac{\gamma}{2} |\nabla v(x)|^2 + F(v(x)) \right] dx, \quad (1.5)$$

where $F(u)$ is defined by (1.4), is a Lyapunov functional for $\{S(t)\}_{t \geq 0}$, that is

I(i) $\exists c_l$ such that $I(v) \geq c_l \forall v \in H_0^1(\Omega)$, and for every c there exists $\rho = \rho(c)$ such that the diameter of $\{v : I(v) \leq c\}$ is bounded by ρ .

I(ii) $I(S(t)v) \leq I(S(t')v) \forall t \geq t' > 0, \forall v \in H_0^1(\Omega)$

I(iii) If $I(S(t)v) = I(v) \forall t > 0$ then $v \in \mathcal{E}$ where \mathcal{E} is the set of equilibrium points v satisfying

$$-\gamma \Delta v + f(v) = 0, \quad x \in \Omega, \quad v = 0, \quad x \in \partial\Omega. \quad (1.6)$$

Furthermore, for any $u_0 \in L^2(\Omega)$ the ω -limit set

$$\omega(u_0) = \{v \in L^2(\Omega) : \exists t_n \rightarrow \infty : \lim_{n \rightarrow \infty} S(t_n)u_0 = v\} \quad (1.7)$$

is contained in the set of equilibrium solutions (1.6), denoted E , and, if the equilibria are isolated, then

$$\mathcal{A} = \bigcup_{v \in \mathcal{E}} W^u(v)$$

where $W^u(v)$ denotes the unstable manifold of v . \square

It is our purpose in this paper to consider discrete dynamical systems generated by temporal and spatial discretisation of (1.1). We study the existence of absorbing sets, Lyapunov functions and attractors for the approximations and, in addition, describe various spurious features introduced by both spatial and temporal discretisation. We suppose that semi-discretisation in space yields a system of ordinary differential equations in time of the form:

$$M \frac{d\mathbf{U}}{dt} + \gamma A \mathbf{U} + M \mathbf{f}(\mathbf{U}) = 0, \quad t > 0, \quad (1.8)$$

$$\mathbf{U}(0) = \mathbf{U}_0. \quad (1.9)$$

Here $\mathbf{U} = (U_1, \dots, U_J)^T$ and $\{\mathbf{f}(\mathbf{U})\}_i = f(U_i)$. We use the notation that, for any $\phi: \mathcal{R} \rightarrow \mathcal{R}$

$$\{\phi(\mathbf{U})\}_i = \phi(\mathbf{U}_i).$$

M is a diagonal matrix with positive entries and A is a symmetric positive definite matrix.

If $\Omega = (0, 1)^d$ then such a system arises from finite difference approximations on a uniform grid $\{(ih, jh, kh) : i, j, k \in [0, N], h = 1/N\}$ with the elements of M being h^d and $\frac{1}{h^d} A$ being the matrix arising from a central difference approximation to $-\Delta$. The system also arises from a finite element approximation based on piecewise linear functions on a triangulation of Ω . Denoting by $\{\psi_i\}_{i=1}^{N_V}$ the set of piecewise linear basis functions associated with the internal vertices of the elements, then

$$M_{ii} = \langle \psi_i, 1 \rangle^h, \quad A_{ij} = \langle \nabla \psi_i, \nabla \psi_j \rangle^h,$$

where $\langle \cdot, \cdot \rangle^h$ denotes a discrete L^2 inner product defined by lumped mass integration.

Section 2 contains various general results on dissipative dynamical systems which will be required throughout the paper. In section 3 we analyse the semi-discrete problem (1.8)–(1.9). Semi-discrete analogues of (R1)–(R4) are proved. However, it is also shown that

insufficient spatial resolution can lead to the existence of deceptively smooth spurious steady solutions and also to incorrect stability properties for the true steady states – there exist many more discrete stable steady states than true stable steady states. In section 4 we study the spurious dynamics of the explicit Euler scheme. It is shown that the method can produce period 2 solutions in n , the time-step, for arbitrarily small Δt . Furthermore it is shown that, however small Δt and Δx are taken, the scheme blows-up with appropriate choice of initial data. Thus a global attractor cannot exist for the Euler method considered as a dynamical system on the whole of \mathcal{R}^J . We derive a restriction on the time step in terms of both the spatial grid and the magnitude of the initial data, under which discrete analogues of (R1)–(R4) hold. In section 5, various implicit schemes are considered. Discrete analogues of (R1)–(R4) are proved under far less restrictive conditions on the time step than for the explicit Euler scheme; in particular schemes are constructed for which the time-step restriction is independent of initial data and of h . Section 6 is concerned with multistep backward differentiation formulae and similar results to those in section 5 are proved. Finally, in section 7, numerical results are presented to illustrate the theory.

Similar questions to those addressed in this paper have been discussed for spectral approximations of (1.1) in [Shen, 1989] and for finite difference approximations to the Kuramoto-Sivashinsky equation in [Foias *et al*, 1991]. The question of spurious steady states for such problems has been simultaneously studied in [Bence *et al*, 1991]. The treatment of discrete Lyapunov functionals, which are used extensively in this paper, is motivated by studies of the Cahn-Hilliard equation [Elliott, 1989].

2. Dissipative Dynamical Systems

In this section we review the theory of dissipative dynamical systems sufficient for our needs. It is convenient to consider discrete and continuous dynamical systems simultaneously. Let H be a complete metric space and t be a real parameter taking values in \mathcal{R} or \mathcal{Z} . Let $\{S(t)\}_{t \geq 0}$ be a family of operators from H into itself satisfying the semigroup properties

$$S(t+s) = S(t)S(s), \forall s, t \geq 0, \\ S(0) = I$$

and $S(t) : \mathcal{R}_+ \times H \rightarrow H$ is continuous. In the case of a discrete dynamical system then $t \in \mathcal{Z}$ and $S(t) \equiv S^t$ where $S : H \rightarrow H$ is a continuous operator so that the semigroup properties are automatic.

For each $v \in H$ we denote the positive orbit

$$\bigcup_{t \geq 0} S(t)v$$

by $\gamma^+(v)$ and by $\omega(v)$ the ω -limit set defined by

$$\omega(v) = \bigcap_{s \geq 0} \overline{\bigcup_{t \geq s} S(t)v}$$

which is equivalent to definition (1.7) with $L^2(\Omega)$ replaced by the general metric space H . For a set $B \in H$ we define the ω -limit set $\omega(B)$ by

$$\omega(B) = \bigcap_{s \geq 0} \overline{\bigcup_{t \geq s} S(t)B}$$

which is equivalent to

$$\{w \in H : \exists t_j \rightarrow \infty \text{ and } v_j \in B : w = \lim_{j \rightarrow \infty} S(t_j)v_j\}.$$

A set B is said to be *invariant* if $S(t)B = B, \forall t \geq 0$. A set \mathcal{A} is said to *attract* a set B under S if, for any $\epsilon > 0$ there exists $t_0 = t_0(\epsilon, B, \mathcal{A})$ such that, for $t \geq t_0$ $S(t)B \subset N^\epsilon(\mathcal{A})$, an ϵ -neighbourhood of \mathcal{A} . A set $B \subset U$, where U is an open set, is said to be *absorbing* in U under S if for any bounded set $B_0 \subset U$ there exists $t_0 = t_0(B_0)$ such that $S(t)B_0 \subset B$ for $t \geq t_0$. A set \mathcal{A} is said to be a *global attractor* for an open set $U \subset H$ under S if

- a) \mathcal{A} is a compact invariant set.
- b) Every compact invariant set of S belongs to \mathcal{A}
- c) \mathcal{A} attracts every bounded set $B \subset U$ under S .

We say that v is an *equilibrium point* of S if $S(t)v = v, \forall t \geq 0$. The set of all equilibria is denoted by \mathcal{E} . A continuous functional $I : H \rightarrow \mathcal{R}$ is said to be a *Lyapunov functional* on U under S if

- I(i) $\exists c_l$ such that $I(v) \geq c_l \forall v \in U$, and for every c there exists $\rho = \rho(c)$ such that the diameter of $\{v : I(v) \leq c\}$ is bounded by ρ .
- I(ii) $I(S(t)v) \leq I(S(t')v) \forall t \geq t' > 0, \forall v \in U$
- I(iii) If $I(S(t)v) = I(v) \forall t > 0$ then $v \in \mathcal{E}$.

S is said to be *point dissipative* on an open set $U \subset H$ if there exists a bounded set $B \subset U$ which attracts each point of U . S is said to be *uniformly compact* on an open set $U \subset H$ if for each bounded set $B \subset U$ there exists $t_0 = t_0(B)$ such that

$$\bigcup_{t \geq t_0} S(t)B$$

is relatively compact.

Lemma 2.1 *Let H be finite dimensional and $U \subset H$ be open. Then each of the properties*

- (i) *There exists a bounded set B which is absorbing in U under S ;*
- (ii) *There exists a Lyapunov functional, I , on U under S*

implies that S is uniformly compact on U .

Proof Since H is finite dimensional it is sufficient to show that there exists $t_0 = t_0(B)$ such that, for each bounded set $B_0 \subset U$, $\bigcup_{t \geq t_0} S(t)B_0$ is bounded. If (i) holds then this is immediate since $\bigcup_{t > t_0} S(t)B_0 \subset B$ is a bounded set.

If (ii) holds, by I(iii), we have that for each $v \in B_0$, $I(S(t)v) \leq I(v) \leq M(B_0) \forall t > 0$ and by I(i) it holds that $S(t)B_0$ is bounded $\forall t > 0$. \square

Lemma 2.2 [Temam, 1989, p28] *Let $B \subset H$ be non-empty and $\bigcup_{t \geq t_0} S(t)B$ be relatively compact for some t_0 . Then $\omega(B)$ is non-empty, compact and invariant. \square*

Proposition 2.3 [Temam, 1988, p23] *Let there exist an open set $U \subset H$ and a bounded set $B \subset U$ such that B is absorbing in U under S . Suppose that S is uniformly compact on U . Then $\mathcal{A} = \omega(B)$ is a global attractor for U under S . Furthermore, if H is a Banach space and U is convex and connected, then \mathcal{A} is connected. \square*

Proposition 2.4 *Suppose that $\gamma^+(u_0)$ is relatively compact for each $u_0 \in U$ and that there exists a Lyapunov functional I on U under S . Then $\omega(u_0) \subset \mathcal{E}$ for each $u_0 \in U$. Furthermore, if \mathcal{E} is bounded then S is point dissipative on U .*

Proof By I(i) and I(iii), $I(S(t)u_0)$ is a bounded decreasing continuous functional for $t \in (0, \infty)$ (or sequence for $t \in \mathcal{Z}$) and so has a unique limit

$$e = \lim_{t \rightarrow \infty} I(S(t)u_0).$$

Since $\gamma^+(u_0)$ is relatively compact there exists a cluster point $v \in \omega(u_0) = \lim_{k \rightarrow \infty} S(t_k)u_0$ and, because I is continuous, $I(v) = \lim_{k \rightarrow \infty} I(S(t_k)u_0) = e$. Thus $I(v) = e \forall v \in \omega(u_0)$. The invariance of $\omega(u_0)$ (Lemma 2.2) implies that for $v \in \omega(u_0)$ $S(t)v \in \omega(u_0) \forall t \geq 0$. Therefore $I(S(t)v) = e = I(v)$ for all $t \geq 0$ and, by I(iii) it follows that $v \in \mathcal{E}$.

We have for each $u_0 \in U$ that

$$\lim_{t \rightarrow \infty} I(S(t)u_0) = e$$

which implies that for $t \geq t_0 = t_0(u_0)$, $I(S(t)u_0) \leq e + 1$. Therefore by I(i) there exists a bounded set B such that $S(t)u_0 \in B \forall t \geq t_0(u_0)$. Thus S is point dissipative. \square

Proposition 2.5 *Suppose that*

- (i) *S has a Lyapunov functional I on U .*
- (ii) *S is uniformly compact on U .*
- (iii) *\mathcal{E} is bounded.*

Then there exists a global attractor \mathcal{A} for U under S . Furthermore, if H is a Banach space and U is convex and connected, then \mathcal{A} is connected.

Proof It follows from Proposition 2.4 that S is point dissipative on U . Furthermore, by I(ii) and I(iii) the orbits of bounded sets are bounded. The existence of a connected global attractor is then a consequence of Hale [1988; p39 Theorem 3.4.6]. \square .

3. The Semi-Discrete Approximation: Spurious Steady Solutions

In this section we study the ordinary differential equations (1.8)–(1.9). We start by introducing some notation and making explicit some assumptions about the structure of the spatial discretization. We introduce a discrete L^2 inner product

$$\langle \mathbf{U}, \mathbf{V} \rangle = \mathbf{U}^T \mathbf{M} \mathbf{V}$$

and a discrete Dirichlet inner product

$$\langle \mathbf{U}, \mathbf{V} \rangle_A = \mathbf{U}^T \mathbf{A} \mathbf{V},$$

together with the associated norms

$$|\mathbf{U}| = \langle \mathbf{U}, \mathbf{U} \rangle^{\frac{1}{2}}, \quad \|\mathbf{U}\| = \langle \mathbf{U}, \mathbf{U} \rangle_A^{\frac{1}{2}}.$$

It is convenient to let L_2^h be the normed vector space $\{\mathcal{R}^J, |\bullet|\}$. We assume that the eigenvalue problem

$$A\psi_i = \lambda_i M\psi_i$$

has eigenvalues λ_i satisfying

$$C_p^{-2} \leq \lambda_i \leq C_0 h^{-2}.$$

Under this assumption it is possible to show (see Appendix 1) that the following Poincare and inverse inequalities hold for all $\mathbf{U} \in \mathcal{R}^J$, where C_p and C_0 are independent of h ,

$$|\mathbf{U}| \leq C_p \|\mathbf{U}\|, \quad (3.1)$$

$$\|\mathbf{U}\|^2 \leq \frac{C_0}{h^2} |\mathbf{U}|^2, \quad (3.2)$$

$$\|\mathbf{U}\| \leq C_p |M^{-1} \mathbf{A} \mathbf{U}|, \quad (3.3)$$

and

$$|M^{-1} \mathbf{A} \mathbf{U}|^2 \leq \frac{C_0}{h^2} \|\mathbf{U}\|^2. \quad (3.4)$$

We set

$$L = M^{-1} \mathbf{A}, \quad \mathbf{E} = (1, 1, \dots, 1)^T$$

and we assume that

$$0 < L_{ii} \leq \frac{C_1}{h^2}, \quad L_{ij} \leq 0, \quad j \neq i, \quad (3.5)$$

where C_1 is independent of h . In addition, we assume that

$$\{\mathbf{A} \mathbf{E}\}_i \geq 0, \quad \text{for } i = 1, \dots, J. \quad (3.6)$$

The assumptions (3.5) and (3.6) imply that

$$\lambda_\phi \langle \mathbf{U}, \phi(\mathbf{U}) \rangle_A \geq \|\phi(\mathbf{U})\|^2, \quad (3.7)$$

[Nochetto, 1991] for monotone ϕ with Lipschitz constant λ_ϕ satisfying $\phi(0) = 0$.

The assumptions that we make about the matrices M and A are reasonable and will be satisfied by standard second order finite difference approximations and piecewise linear finite element discretisations on ‘‘acute triangulations’’ of Ω .

3.1 Equilibrium Problem and Steady Solutions

We say that $\mathbf{U}^* \in L^2(\Omega)$ is an equilibrium of (1.8) provided

$$\gamma \mathbf{A} \mathbf{U}^* + M \mathbf{f}(\mathbf{U}^*) = 0, \quad (3.8)$$

and we let \mathcal{E}^h denote the set of all solutions to (3.8). Clearly $0 \in \mathcal{E}^h$ so that \mathcal{E}^h is nonempty. Furthermore, if $\mathbf{U}^* \in \mathcal{E}^h$ then \mathbf{U}^* is a critical point of the functional

$$I^h(\mathbf{V}) := \frac{\gamma}{2} \langle \mathbf{V}, \mathbf{V} \rangle_A + \langle \mathbf{F}(\mathbf{V}), \mathbf{E} \rangle. \quad (3.9)$$

By (F2) there exists, for each $\epsilon > 0$ a $C_\epsilon \geq 0$ such that

$$F(r) \geq -\epsilon r^2 - C_\epsilon, \quad \forall r \quad (3.10)$$

(see Hale[1988, p76]) and hence it follows that

$$I^h(\mathbf{V}) \geq \frac{\gamma}{2} \langle \mathbf{V}, \mathbf{V} \rangle_A - \epsilon |\mathbf{V}|^2 - C_\epsilon C(\Omega) \geq \frac{\gamma}{4} \|\mathbf{V}\|^2 - C_\gamma \quad \forall \mathbf{V} \in L_2^h \quad (3.11)$$

where $C(\Omega)$ is an h -independent upper bound for $|\mathbf{E}|^2$. Therefore, since (F1) implies that $I^h(\bullet)$ is continuous, classical arguments show that there exists at least one $\mathbf{U}^* \in \mathcal{E}^h$ such that

$$I^h(\mathbf{U}^*) = \inf_{\mathbf{V} \in L_2^h} I^h(\mathbf{V}).$$

Lemma 3.1 *The set \mathcal{E}^h is uniformly bounded, independent of h , in the discrete H_0^1 and in the maximum norm.*

Proof Taking the inner product of (3.8) and \mathbf{U}^* gives

$$\gamma \|\mathbf{U}^*\|^2 + \langle \mathbf{U}^*, \mathbf{f}(\mathbf{U}^*) \rangle = 0$$

and, since by (F1),(F2)

$$f(r)r \geq \min_{|r| \leq \bar{u}} f(r)r = -C_f, \quad (3.12)$$

it follows that \mathcal{E}^h is bounded in the discrete H_0^1 norm, viz

$$\|\mathbf{U}^*\| \leq \sqrt{(C_f C(\Omega)/\gamma)}, \quad \forall \mathbf{U}^* \in \mathcal{E}^h.$$

Furthermore, taking the inner product of (3.8) with $(\mathbf{U}^* - \bar{u}\mathbf{E})_+$, where the $+$ denotes positive part, and using (3.7), it follows that

$$\|(\mathbf{U}^* - \bar{u}\mathbf{E})_+\| = 0$$

since

$$f(r)(r - \bar{u})_+ \geq 0 \quad \forall r.$$

Similarly

$$\|(\mathbf{U}^* + \bar{u}\mathbf{E})_-\| = 0.$$

and hence we have that \mathcal{E}^h is uniformly bounded in the maximum norm

$$|\mathbf{U}^*|_\infty \leq \bar{u}, \quad \forall \mathbf{U}^* \in \mathcal{E}^h. \quad \square$$

Our next result concerns the number of steady solutions (equilibria) of (1.8)–(1.9). We make an additional assumption on $f(u)$, namely (F4).

(F4) $f(-1) = f(0) = f(1) = 0$ and $f'(-1), f'(1) > 0$ and $f'(0) = -1$.

The canonical example of a function with such properties is $f(u) = u^3 - u$. With this choice of f equation (1.1) is sometimes known as the Ginzburg-Landau or Allen-Cahn equation. Similar results can be proved if the number of zeros of f is different, with the obvious modification. In the case $f(u) = u^2 - u$ [Iserles, 1990] has numerical examples which illustrate an analogous result. Simultaneously, a result similar to Theorem 3.2 has been proved by [Bence *et al*, 1991]. Their proof of existence is identical, but their proof of stability is different, relying on the structure of the discrete Laplacian, leading to explicit estimates for $\gamma_0(h)$. A similar method was also used by [Stephens and Shubin, 1981] to study the existence of spurious solutions of the viscous Burgers' equation when the dissipation is small.

In the following, a steady state \mathbf{U}^* is said to be *asymptotically stable* if it is stable in the sense of Lyapunov and, in addition, $\mathbf{U}(t) \rightarrow \mathbf{U}^*$ as $t \rightarrow \infty$ for all $\mathbf{U}(0)$ in a neighbourhood of \mathbf{U}^* .

Theorem 3.2 *Under conditions (F1) and (F4) $\exists \gamma_0(h) > 0$ such that, for $\gamma \in [0, \gamma_0(h))$ (1.8) has at least 3^J steady solutions and 2^J of these are asymptotically stable.*

Proof Let \mathbf{W} be one of the 3^J solutions of (3.8) with $\gamma = 0$ which are generated by choosing

$$W_j = +1, -1 \text{ or } 0.$$

To show that each \mathbf{W} generates an equilibrium solution for γ sufficiently small, we employ the implicit function theorem. Set

$$G(\gamma, \mathbf{U}) := \gamma A\mathbf{U} + M\mathbf{f}(\mathbf{U}).$$

Since $f'(W_j) \neq 0$ and since M has positive elements we deduce that the Jacobian $D_{\mathbf{U}}G$ evaluated at $(0, \mathbf{W})$ is a diagonal matrix with non-zero diagonal entries and so is non-singular. Hence, by the implicit function theorem, we generate a family of solutions $\mathbf{U}^*(\gamma)$ parameterised by γ for $\gamma \in [0, \gamma_0)$ with $\mathbf{U}^*(0) = \mathbf{W}$.

The stability of these steady solutions is governed by the eigenvalues of $D_{\mathbf{U}}G(\gamma, \mathbf{U})$. Now consider the 2^J steady solutions of (1.8) generated by the choice $U_j^*(0) = \pm 1$. It follows from (F4) that $D_{\mathbf{U}}G(0, \mathbf{W})$ is a diagonal matrix with strictly positive entries defining its eigenvalues. By the implicit function theorem $\mathbf{U}(\gamma)$ is C^1 , and so the eigenvalues of $D_{\mathbf{U}}G(\gamma, \mathbf{U}(\gamma))$ depend continuously on γ and thus are positive for γ sufficiently small. This demonstrates that 2^J solutions are stable. \square

Many of these steady states constructed for $\gamma \ll 1$ are spurious and the stability properties of approximations of genuine steady solutions are often incorrect; for example the Ginzburg-Landau equation has only two stable steady states (for γ small) whereas the numerical approximation has 2^J for γ sufficiently small. This issue is discussed in detail in section 7, where numerical results are presented.

However, as $h \rightarrow 0$ for fixed γ the structure of the discrete steady solution set converges to that of the continuous steady solution set; we refer to [Crouzeix and Rappaz, 1990] for an account of the numerical solution of bifurcation problems. In Appendix 3 we show

that when (1.1)–(1.3) has N isolated solutions then the same is also true of (3.8) for h sufficiently small. See also [Khalsa, 1986].

3.2 Existence of Absorbing Sets, Attractors and Stabilization.

The appropriate analogues of (R1)–(R4), which we establish in Theorem 3.4 are stated below; note that B_2^h is an absorbing set in the discrete H_0^1 norm even though the initial data is only bounded uniformly in h in $\{L_2^h, |\bullet|\}$:-

(R1)^h For each $\mathbf{U}_0 \in L_2^h$ there exists a unique solution of (1.8)–(1.9) $\mathbf{U} \in C^2([0, T]; L_2^h)$ for all $T > 0$. The mapping $\mathbf{U}_0 \rightarrow \mathbf{U}(t)$ is continuous in L_2^h for each $t > 0$. Hence the family of solution operators $\{S^h(t)\}_{t \geq 0}$, defined by $S^h(t)\mathbf{U}_0 \equiv \mathbf{U}(t)$, forms a continuous semigroup on L_2^h . \square

(R2)^h There exist constants $\{\bar{\rho}_i\}_{i=1}^2$ independent of h and J such that the balls

$$B_1^h = \{\mathbf{V} \in L_2^h : |\mathbf{V}| \leq \bar{\rho}_1\}$$

$$B_2^h = \{\mathbf{V} \in L_2^h : \|\mathbf{V}\| \leq \bar{\rho}_2\}$$

are absorbing sets for the semigroup $\{S^h(t)\}_{t \geq 0}$; that is, for each $\mathbf{U}_0 \in L_2^h$ there exists $\{t_i\}_{i=1}^2$ (depending on $\{\mathbf{U}_0, \rho_1\}$ and $\{\mathbf{U}_0, \rho_2\}$ respectively) such that

$$S^h(t)\mathbf{U}_0 \in B_i^h, \forall t \geq t_i, (i = 1, 2). \quad \square$$

(R3)^h There exists a global attractor $\mathcal{A}^h \subset L_2^h$ for the semigroup $\{S^h(t)\}_{t \geq 0}$. Furthermore \mathcal{A}^h is connected and there exists a constant $\bar{\rho}_A$ independent of h such that

$$\max\{|\mathbf{V}|, \|\mathbf{V}\|, |\mathbf{V}|_\infty\} \leq \bar{\rho}_A \quad \forall \mathbf{V} \in \mathcal{A}^h. \quad \square$$

(R4)^h The functional defined by (3.9) is a Lyapunov functional for $\{S^h(t)\}_{t > 0}$. In addition, for any $\mathbf{U}_0 \in L_2^h$, the ω -limit set is contained in \mathcal{E}^h . \square

We now prove these results.

Lemma 3.3 *Let $m \geq \bar{u}$. If $\mathbf{U}(t)$ solves (1.8)–(1.9) then, for $t > t_0 \geq 0$*

$$|(\mathbf{U}(t) - m\mathbf{E})_+| \leq |(\mathbf{U}(t_0) - m\mathbf{E})_+| \exp(-(t - t_0)\gamma/C_p^2), \quad (3.13)$$

$$|(\mathbf{U}(t) + m\mathbf{E})_-| \leq |(\mathbf{U}(t_0) + m\mathbf{E})_-| \exp(-(t - t_0)\gamma/C_p^2), \quad (3.14)$$

and

$$|\mathbf{U}(t)|_\infty \leq \max(\bar{u}, |\mathbf{U}_0|_\infty). \quad (3.15)$$

Proof Set

$$\mathbf{Z}(t) = \mathbf{U}(t) - m\mathbf{E}$$

and define $\mathbf{Z}(t)_+$ by $\{\mathbf{Z}(t)_+\}_i = Z_i(t)_+$. It follows that

$$\left\langle \frac{d\mathbf{Z}}{dt}, \mathbf{Z}_+ \right\rangle + \gamma \langle \mathbf{Z}, \mathbf{Z}_+ \rangle_A + \langle \mathbf{f}(\mathbf{U}), \mathbf{Z}_+ \rangle = -m \langle \mathbf{E}, \mathbf{Z}_+ \rangle_A.$$

Since $m \geq \bar{u}$ we have

$$f(r)(r - m)_+ \geq 0, \forall r.$$

Combining with the assumptions (3.6),(3.7) on the matrix A yields

$$\frac{1}{2} \frac{d}{dt} |\mathbf{Z}_+|^2 + \gamma \|\mathbf{Z}_+\|^2 \leq 0. \quad (3.16)$$

Inequality (3.13) is an immediate consequence of (3.16) and the Gronwall inequality. The proof of (3.14) is very similar. We obtain the maximum norm bound (3.15) by taking $m = \max(\bar{u}, |\mathbf{U}_0|_\infty)$ and $t_0 = 0$ in (3.13),(3.14). \square

Theorem 3.4 $(R1)^h$ - $(R4)^h$ hold for (1.8)-(1.9).

Proof

$(R1)^h$:- local existence and uniqueness follows from the standard theory of ordinary differential equations, by (F1). Existence for any $T > 0$ follows from the a priori estimate (3.15). In order to prove continuity with respect to initial data, it is sufficient to consider $\mathbf{U}(t) - \mathbf{V}(t)$ where $\mathbf{V}(t)$ solves (1.8) with initial data \mathbf{V}_0 . Clearly

$$\frac{1}{2} \frac{d}{dt} |\mathbf{U} - \mathbf{V}|^2 + \gamma \|\mathbf{U} - \mathbf{V}\|^2 + \langle \mathbf{f}(\mathbf{U}) - \mathbf{f}(\mathbf{V}), \mathbf{U} - \mathbf{V} \rangle = 0$$

and the inequality

$$|\mathbf{U}(t) - \mathbf{V}(t)| \leq C(T, \mathbf{U}_0, \mathbf{V}_0) |\mathbf{U}_0 - \mathbf{V}_0|$$

follows from the a priori estimate (3.15) in Lemma 3.2 and Gronwall's inequality.

$(R2)^h$:- taking the inner product of (3.8) with $\mathbf{U}(t)$ implies that

$$\frac{1}{2} \frac{d}{dt} |\mathbf{U}(t)|^2 + \gamma \|\mathbf{U}\|^2 + \langle \mathbf{U}, \mathbf{f}(\mathbf{U}) \rangle = 0$$

and, by (3.12),

$$\frac{d}{dt} |\mathbf{U}|^2 + 2\gamma \|\mathbf{U}\|^2 \leq 2C_f C(\Omega). \quad (3.17)$$

Using Poincaré's inequality (3.1),

$$\frac{d}{dt} |\mathbf{U}|^2 + \frac{2\gamma}{C_p^2} |\mathbf{U}|^2 \leq 2C_f C(\Omega).$$

From this we obtain, using Gronwall's inequality,

$$|\mathbf{U}(t)|^2 \leq |\mathbf{U}_0|^2 \exp(-2\gamma t/C_p^2) + C_f C(\Omega) C_p^2 [1 - \exp(-2\gamma t/C_p^2)]/\gamma.$$

Therefore,

$$\limsup_{t \rightarrow \infty} |\mathbf{U}(t)| \leq \delta_1, \delta_1^2 = C_f C(\Omega) C_p^2/\gamma$$

and we can take $\bar{\rho}_1$ to be any number larger than δ_1 to define the absorbing set B_1^h .

In order to show that there is an absorbing set uniformly bounded in the discrete H_0^1 norm we pre-multiply equation (1.8) by M^{-1} and take the $\langle \bullet, \bullet \rangle_A$ inner product with \mathbf{U} to obtain

$$\frac{1}{2} \frac{d}{dt} \|\mathbf{U}\|^2 + \gamma |M^{-1} A \mathbf{U}|^2 + \langle \mathbf{f}(\mathbf{U}), \mathbf{U} \rangle_A = 0,$$

so that

$$\frac{1}{2} \frac{d}{dt} \|\mathbf{U}\|^2 + \langle \mathbf{f}(\mathbf{U}) + C_F \mathbf{U}, \mathbf{U} \rangle_A - \langle C_F \mathbf{U}, \mathbf{U} \rangle_A = 0.$$

Using the monotonicity of $\mathbf{f}(\mathbf{U}) + C_F \mathbf{U}$ implied by (F3) and (3.7) we obtain

$$\frac{1}{2} \frac{d}{dt} \|\mathbf{U}\|^2 \leq C_F \|\mathbf{U}\|^2.$$

Integrating (3.17) we obtain

$$\int_t^{t+r} \|\mathbf{U}(s)\|^2 ds \leq \frac{C_f C(\Omega) r}{\gamma} + \frac{1}{2\gamma} |\mathbf{U}(t)|^2.$$

Taking t sufficiently large and using the absorbing set in L_2^h this gives

$$\int_t^{t+r} \|\mathbf{U}(s)\|^2 ds \leq \frac{C_f C(\Omega) r}{\gamma} + \frac{\bar{\rho}_1^2}{2\gamma}.$$

Thus, applying the uniform Gronwall lemma [Temam 1989, p89] we obtain

$$\|\mathbf{U}(t+r)\|^2 \leq \left[\frac{C_f C(\Omega)}{\gamma} + \frac{\bar{\rho}_1}{2\gamma r} \right] \exp(2C_F r).$$

Since t may be arbitrarily large, this yields the absorbing set in the $\|\bullet\|$ norm.

(R3)^h:- We set ourselves in the framework of section 2. Since L_2^h is finite dimensional, the existence of a global attractor $\mathcal{A}^h = \omega^h(B_1^h)$ is an immediate consequence of Proposition 2.3. The existence of $\bar{\rho}_A$ is an immediate consequence of the bounds obtained in the proof of (R2)^h and from the following maximum norm bound: since

$$|(\mathbf{U}(t_0) - m\mathbf{E})_+| \leq |\mathbf{U}(t_0)| + mC(\Omega),$$

it follows from (3.13),(3.14) in Lemma 3.3 with $m = \bar{u}$ and the previous estimate on the absorbing set in the L_2^h norm that

$$\limsup_{t \rightarrow \infty} |\mathbf{U}(t)|_\infty \leq \bar{u}.$$

(R4)^h:- Taking the inner product of (1.8) with $d\mathbf{U}/dt$ yields

$$\left| \frac{d\mathbf{U}}{dt} \right|^2 + \frac{d}{dt} I^h(\mathbf{U}) = 0. \quad (3.18)$$

Facts I(i)–I(iii) concerning $I^h(\bullet)$ follow immediately from (3.11) and (3.18). Thus Proposition 2.3 yields $\omega(\mathbf{U}_0) \subset \mathcal{E}^h$.

4. The Explicit Euler Approximation

In this section we consider the Euler discretisation of (1.8)–(1.9). Let \mathbf{U}_n denote the approximation to $\mathbf{U}(n\Delta t)$. We obtain, for $n \geq 1$,

$$M\partial\mathbf{U}_{n+1} + \gamma A\mathbf{U}_n + M\mathbf{f}(\mathbf{U}_n) = 0, \quad (4.1)$$

where

$$\partial \mathbf{V}_n := (\mathbf{V}_n - \mathbf{V}_{n-1})/\Delta t.$$

The convergence of solutions of (4.1) to solutions of (1.1) on a finite time interval as $h, \Delta t \rightarrow 0$ is well-understood; see [Richtmyer and Morton, 1967]. Such results require that Δt be restricted by the square of the space step h . Here we are concerned with the long time dynamics of (4.1), that is $n \rightarrow \infty$, with fixed Δt and h . We show that the temporal discretisation (4.1) introduces spurious solutions with period 2 in the time-step and that, furthermore, under certain assumptions on $f(u)$, these periodic solutions can be found for Δt arbitrarily small. We then discuss the effect of these periodic solutions on the dynamics of the problem. If Δt is restricted in terms of the initial data then the effect of the periodic solutions is avoided, a gradient structure ensues and the existence of absorbing sets and attractors is proved. However, if Δt is not restricted sufficiently in terms of the initial data the scheme may be shown to blow-up, indicating that the unstable manifolds of the spurious periodic solutions are connected with infinity.

4.1 Spurious Dynamics

Period 2 solutions of (4.1) are pairs $\{\mathbf{U}, \mathbf{V}\}$ with $\mathbf{U} \neq \mathbf{V}$ which satisfy the equations

$$M(\mathbf{V} - \mathbf{U}) + \Delta t \gamma A \mathbf{U} + \Delta t M \mathbf{f}(\mathbf{U}) = 0, \quad (4.2)$$

$$M(\mathbf{U} - \mathbf{V}) + \Delta t \gamma A \mathbf{V} + \Delta t M \mathbf{f}(\mathbf{V}) = 0. \quad (4.3)$$

Existence of period 2 solutions can be established by a local bifurcation argument, treating Δt as a bifurcation parameter. See [Griffiths and Mitchell, 1988] and [Stuart, 1989]. Fixed points (i.e. equilibria of (4.1)) are solutions of (4.2)–(4.3) with the symmetry $\mathbf{U} = \mathbf{V}$. Genuine period 2 solutions can be constructed as local bifurcations which break this symmetry as Δt varies.

Suppose that \mathbf{U}^* is an equilibrium solution and let $d\mathbf{f}(\bullet)$ denote the Jacobian of \mathbf{f} so that

$$d\mathbf{f}(\mathbf{U}^*) = \text{diag}(f'(U_i^*)).$$

Applying Theorem 4.1 in [Stuart, 1989] we obtain the following result.

Lemma 4.1 *Let \mathbf{U}^* be an equilibrium point of (4.1) (i.e. a solution of (3.8)) such that all the eigenvalues of $\gamma L + d\mathbf{f}(\mathbf{U}^*)$ are non-zero. Assume that (F1) holds and $d = 1$. Then the eigenvectors $\{\phi_k\}_{k=1}^J$ and distinct eigenvalues $\{\eta_k\}_{k=1}^J$ of $\gamma L + d\mathbf{f}(\mathbf{U}^*)$ generate period 2 solutions of (4.1) with the form*

$$\mathbf{U}(\mu) := \mathbf{U}^* + \mu \phi_k + \mathcal{O}(|\mu|^2),$$

$$\mathbf{V}(\mu) := \mathbf{U}^* - \mu \phi_k + \mathcal{O}(|\mu|^2),$$

$$\Delta t_k(\mu) := 2/\eta_k + \mathcal{O}(|\mu|),$$

for μ sufficiently small which are C^1 in μ . \square

Note that most of the η_k are positive since the largest eigenvalues of $L = M^{-1}A$ scale like h^{-2} . Here μ parameterises the branches of period 2 solutions locally in the

neighbourhood of the bifurcation points. Having constructed these branches of spurious solutions locally in the neighbourhood of the bifurcation points Δt_k , it is natural to ask what happens to these branches of solutions far from the bifurcation points. In particular, it is of interest to know whether such solutions exist for Δt arbitrarily small. For simplicity we consider the case of $f(\bullet)$ being odd. It is then possible to seek period 2 solutions with the symmetry $\mathbf{U} = -\mathbf{V}$ under which (4.2),(4.3) become

$$2\mathbf{U} - \Delta t f(\mathbf{U}) - \Delta t \gamma L\mathbf{U} = 0. \quad (4.4)$$

Setting $\gamma = 0$, equation (4.4) admits solutions \mathbf{W} with the form

$$\{\mathbf{W}\}_i = \pm R(\Delta t), \quad (4.5)$$

where $R(\Delta t)$ solves

$$r = \Delta t f(r)/2. \quad (4.6)$$

Suppose that f satisfies (F1), (F2) and that, in addition,

$$(F5) |f(r)/r| \rightarrow \infty \text{ as } r \rightarrow \infty,$$

then (4.6) has solutions $\pm R(\Delta t)$ for arbitrarily small Δt which satisfy

$$\lim_{\Delta t \rightarrow 0} |R(\Delta t)| = \infty.$$

These two solutions generate 2^J solutions of (4.4), and hence 2^J period 2 solutions of (4.1) in the case of zero diffusion ($\gamma = 0$.) These solutions exist for Δt arbitrarily small but, in accordance with the results of [Humphries, 1991], they tend to infinity as $\Delta t \rightarrow 0$. We now show that these solutions persist for $\gamma > 0$ sufficiently small under appropriate conditions on $f(u)$. These conditions allow, for example, arbitrary superlinear polynomial power growth. In the following theorem the norm $|\bullet|$ on matrices denotes that subordinate to the related norm on vectors.

Theorem 4.2 *Let f satisfy (F1), (F2), (F5) and be odd. Assume that f satisfies the following two growth conditions: $\exists \bar{\Delta}t$ satisfying $0 < \bar{\Delta}t < 1$:*

(i) $|2 - \Delta t f'(R(\Delta t))| \geq d_1 > 0, \forall 0 \leq \Delta t \leq \bar{\Delta}t$, where d_1 is independent of Δt and $R(\Delta t)$ solves equation (4.6);

(ii) define

$$D = 2\text{Id} - \Delta t df(\mathbf{W})$$

where Id is the identity matrix and note that D^{-1} is defined for $0 \leq \Delta t \leq \bar{\Delta}t$ by (i). Let

$$\mathcal{B} := \{\mathbf{V} : |\mathbf{V} - \mathbf{W}| \leq \rho\}$$

where

$$\rho = \gamma |L| |D^{-1}| |\mathbf{W}|.$$

Then we assume $\Delta t K \leq d_2 \forall \Delta t \leq \bar{\Delta}t$ where K is the Lipschitz constant for df on \mathcal{B} .

Assume also that γ is chosen sufficiently small so that

$$d_2 |D^{-1}| \rho + \bar{\Delta}t (1 + \gamma |L| |D^{-1}|) \leq 1$$

and

$$2d_2|D^{-1}|\rho + \Delta t\gamma|L||D^{-1}| < 1.$$

Then, for each of the 2^J solutions \mathbf{W} defined by (4.5),(4.6), there exists a unique solution \mathbf{U}, \mathbf{V} to equations (4.2), (4.3) satisfying $\mathbf{V} = -\mathbf{U}$ and

$$|\mathbf{U} - \mathbf{W}| \leq \rho.$$

Proof Solutions of (4.4) are fixed points of the iteration

$$\mathbf{U}_{k+1} = \mathbf{U}_k - D^{-1}[2\mathbf{U}_k - \Delta t\mathbf{f}(\mathbf{U}_k) - \Delta t\gamma L\mathbf{U}_k].$$

By (i), this map is well-defined and $|D^{-1}|$ is bounded independently of Δt .

We first show that the map takes \mathcal{B} into itself. Let $\mathbf{U}_k \in \mathcal{B}$. Clearly

$$\mathbf{W} = \mathbf{W} - D^{-1}[2\mathbf{W} - \Delta t\mathbf{f}(\mathbf{W})].$$

Subtracting and exploiting the diagonal nature of \mathbf{f} we obtain

$$\mathbf{e}_{k+1} = -\Delta t D^{-1}[(d\mathbf{f}(\mathbf{W}) - d\mathbf{f}(\xi_k))\mathbf{e}_k - \gamma L\mathbf{U}_k],$$

where $\mathbf{e}_k = \mathbf{U}_k - \mathbf{W}$ and $\xi_k \in \mathcal{B}$. Using the Lipschitz constant for $d\mathbf{f}$ we find that

$$\begin{aligned} |\mathbf{e}_{k+1}| &\leq \Delta t|D^{-1}|K|\mathbf{e}_k|^2 + \Delta t\gamma|L||D^{-1}||\mathbf{W}| + \Delta t\gamma|L||D^{-1}||\mathbf{e}_k| \\ &\leq d_2|D^{-1}|\rho^2 + \Delta t\rho(1 + \gamma|L||D^{-1}|) \\ &\leq [d_2|D^{-1}|\rho + \Delta t(1 + \gamma|L||D^{-1}|)]\rho \end{aligned}$$

and hence, by assumption on γ the mapping takes \mathcal{B} into itself.

Now consider a second sequence of iterates \mathbf{V}_k satisfying the same iteration scheme and assume that \mathbf{U}_0 and $\mathbf{V}_0 \in \mathcal{B}$. Defining $\mathbf{d}_k = \mathbf{U}_k - \mathbf{V}_k$ we obtain

$$\mathbf{d}_{k+1} = -\Delta t D^{-1}[(d\mathbf{f}(\mathbf{W}) - d\mathbf{f}(\eta_k))\mathbf{d}_k - \gamma L\mathbf{d}_k],$$

where $\eta_k \in \mathcal{B}$. Thus

$$\begin{aligned} |\mathbf{d}_{k+1}| &\leq \Delta t|D^{-1}|K|\mathbf{d}_k|^2 + \Delta t\gamma|L||D^{-1}||\mathbf{d}_k| \\ &\leq [2d_2|D^{-1}|\rho + \Delta t\gamma|L||D^{-1}|]|\mathbf{d}_k|. \end{aligned}$$

Hence, by assumption on γ the mapping is a contraction on \mathcal{B} . By the contraction mapping theorem, the proof is complete. \square

We now consider the effect of these spurious period 2 solutions on the dynamics of the Euler scheme. It is clear that, if the scheme is to inherit the gradient structure of the underlying PDE, then some restriction will be required on Δt in terms of the magnitude of the initial data – this is because, for any Δt sufficiently small, it is possible to choose initial data in the domain of attraction of a period 2 solution which precludes stabilisation to the set of equilibria and hence a gradient structure cannot hold. Furthermore, we

conjecture that the unstable manifolds of the period 2 solutions are connected to infinity. To substantiate this conjecture, we prove that, for all Δt sufficiently small there is initial data which blows up under the Euler iteration (4.1). This should be contrasted with the underlying partial differential equation and its semi-discretisation, for which arbitrarily large initial data is mapped into an absorbing set under the evolution semigroup.

The following relationships are needed in the analysis:

$$|\mathbf{U}| \leq C(\Omega)|\mathbf{U}|_\infty, \quad \forall U \in \mathcal{R}^J \quad (4.7)$$

and

$$|\mathbf{U}|^2 \geq C_2 h^d |\mathbf{U}|_\infty^2, \quad \forall U \in \mathcal{R}^J. \quad (4.8)$$

Here C_2 is independent of h , and d is the spatial dimension of the problem. In the following set $r = \Delta t/h^2$.

Theorem 4.3 Consider (4.1) under conditions (F1), (F2) and (F5) with initial data \mathbf{U}_0 satisfying the property that $|f(u)| > C_3|u|$ for all $u : |u| \geq |\mathbf{U}_0|_\infty$, where C_3, h and r are chosen so that

$$\Delta t C_3 [\Delta t C_2 h^d C_3 - 2C(\Omega)^2 - 2r C_0 \gamma C(\Omega)^2] \geq C(\Omega)^2 (1+k) - (1 - 2r\gamma C_0 + \frac{\Delta t^2 \gamma^2}{C_p^4}) C_2 h^d,$$

for some $k > 0$ and

$$\Delta t C_3 [\Delta t C_2 h^d C_3 - 2C(\Omega)^2 - 2r C_0 \gamma C(\Omega)^2] \geq 0.$$

Then

$$|\mathbf{U}_n|_\infty^2 \geq (1+k)^n |\mathbf{U}_0|_\infty^2.$$

Proof The proof is by induction. Re-arranging (4.1) gives

$$M\mathbf{U}_{n+1} = M\mathbf{U}_n - \Delta t \gamma A \mathbf{U}_n - \Delta t M \mathbf{f}(\mathbf{U}_n).$$

Taking the inner product of each side of this equation with itself yields

$$\begin{aligned} |\mathbf{U}_{n+1}|^2 &= |\mathbf{U}_n|^2 - 2\Delta t \gamma \|\mathbf{U}_n\|^2 - 2\Delta t \langle \mathbf{U}_n, \mathbf{f}(\mathbf{U}_n) \rangle + \\ &2\Delta t^2 \gamma \langle \mathbf{f}(\mathbf{U}_n), \mathbf{U}_n \rangle_A + \Delta t^2 \gamma^2 |M^{-1} A \mathbf{U}_n|^2 + \Delta t^2 |\mathbf{f}(\mathbf{U}_n)|^2. \end{aligned}$$

Using (3.1),(3.2),(3.3), (4.7) and (4.8) we obtain

$$\begin{aligned} |\mathbf{U}_{n+1}|^2 &\geq (1 - 2r\gamma C_0 + \frac{\Delta t^2 \gamma^2}{C_p^4}) |\mathbf{U}_n|^2 - 2\Delta t |\mathbf{U}_n| |\mathbf{f}(\mathbf{U}_n)| - 2\Delta t^2 \gamma \|\mathbf{U}_n\| \|\mathbf{f}(\mathbf{U}_n)\| + \Delta t^2 |\mathbf{f}(\mathbf{U}_n)|^2 \\ &\geq (1 - 2r\gamma C_0 + \frac{\Delta t^2 \gamma^2}{C_p^4}) |\mathbf{U}_n|^2 - 2\Delta t |\mathbf{U}_n| |\mathbf{f}(\mathbf{U}_n)| - 2\Delta t r C_0 \gamma |\mathbf{U}_n| |\mathbf{f}(\mathbf{U}_n)| + \Delta t^2 |\mathbf{f}(\mathbf{U}_n)|^2 \\ &\geq (1 - 2r\gamma C_0 + \frac{\Delta t^2 \gamma^2}{C_p^4}) |\mathbf{U}_n|^2 - 2\Delta t C(\Omega)^2 |\mathbf{U}_n|_\infty |\mathbf{f}(\mathbf{U}_n)|_\infty \end{aligned}$$

$$\begin{aligned}
& -2\Delta t r C_0 \gamma C(\Omega)^2 |\mathbf{U}_n|_\infty |\mathbf{f}(\mathbf{U}_n)|_\infty + \Delta t^2 C_2 h^d |\mathbf{f}(\mathbf{U}_n)|_\infty^2 \\
& = (1 - 2r\gamma C_0 + \frac{\Delta t^2 \gamma^2}{C_p^4}) |\mathbf{U}_n|^2 + \Delta t |\mathbf{f}(\mathbf{U}_n)|_\infty [\Delta t C_2 h^d |\mathbf{f}(\mathbf{U}_n)|_\infty \\
& \quad - 2C(\Omega)^2 |\mathbf{U}_n|_\infty - 2r C_0 \gamma C(\Omega)^2 |\mathbf{U}_n|_\infty].
\end{aligned}$$

Assume for the purposes of induction that $|\mathbf{U}_n|_\infty$ satisfies

$$|\mathbf{U}_n|_\infty \geq (1+k)^{n/2} |\mathbf{U}_0|_\infty. \quad (4.9)$$

Now

$$|\mathbf{f}(\mathbf{U}_n)|_\infty \geq |f(v)|$$

where $|v| = |\mathbf{U}_n|_\infty$ and hence

$$|\mathbf{f}(\mathbf{U}_n)|_\infty > C_3 |\mathbf{U}_n|_\infty.$$

By assumption on C_3 it follows that

$$\Delta t C_2 h^d C_3 - 2C(\Omega)^2 - 2r C_0 \gamma C(\Omega)^2 > 0.$$

Thus we have

$$\begin{aligned}
|\mathbf{U}_{n+1}|^2 & \geq (1 - 2r\gamma C_0 + \frac{\Delta t^2 \gamma^2}{C_p^4}) |\mathbf{U}_n|^2 + \Delta t C_3 [\Delta t C_2 h^d C_3 - 2C(\Omega)^2 - 2r C_0 \gamma C(\Omega)^2] |\mathbf{U}_n|_\infty^2 \\
& \geq C_2 h^d (1 - 2r\gamma C_0 + \frac{\Delta t^2 \gamma^2}{C_p^4}) |\mathbf{U}_n|_\infty^2 + \Delta t C_3 [\Delta t C_2 h^d C_3 - 2C(\Omega)^2 - 2r C_0 \gamma C(\Omega)^2] |\mathbf{U}_n|_\infty^2.
\end{aligned}$$

By assumption this yields

$$|\mathbf{U}_{n+1}|^2 \geq C(\Omega)^2 (1+k) |\mathbf{U}_n|_\infty^2.$$

Hence

$$|\mathbf{U}_{n+1}|_\infty^2 \geq (1+k) |\mathbf{U}_n|_\infty^2.$$

Thus, since $k > 0$, (4.9) holds for $n = n + 1$ and the induction is complete. \square

The magnitude of the data required to prove blow-up in Theorem 4.3 is very large: a rough estimate shows that $C_3 = \mathcal{O}(\Delta t^{-1} h^{-d/2})$. The analysis is very crude since it is based in the maximum norm and no attempt is made to determine the most unstable spatial structure required for blow-up. In the case of one-dimensional finite difference schemes, the analysis can be tightened to yield a critical $C_3 = \mathcal{O}(\Delta t^{-1})$. In this case a scheme of the form (4.1) may be written

$$U_j^{n+1} = r U_{j-1}^n + (1 - 2r) U_j^n + r U_{j+1}^n - \Delta t f(U_j^n), \quad \text{for } j = 1, \dots, J, \quad (4.10)$$

together with the boundary conditions

$$U_0^n = U_{J+1}^n = 0. \quad (4.11)$$

Consider initial data U_j^0 satisfying

$$U_l^0 U_{l+1}^0 < 0 \quad \forall l = 1, \dots, J-1. \quad (4.12)$$

This choice of initial data is very unstable and is motivated by the most unstable mode according to linear theory [Richtmyer and Morton, 1967]. Note that the condition we derive for blow-up may be re-formulated as

$$\Delta t > \frac{(2+k)h^2}{2+h^2C}.$$

where C is defined below. This should be contrasted with the bound (4.15) on Δt in Lemma 4.5 which ensures boundedness of the numerical solution as $n \rightarrow \infty$.

Theorem 4.4 Consider (4.10)–(4.11) under conditions (F1), (F2) and (F5) together with initial data satisfying (4.12). In addition, assume that the initial data satisfies the property that $f(u)/u > C$ for all $u : |u| \geq \min_j \{|\mathbf{U}_0\}_j\}$ and where C is chosen so that

$$C \geq \frac{2-2r+k}{\Delta t},$$

for some $k > 0$. Then

$$|\mathbf{U}_n|_\infty \geq (1+k)^n |\mathbf{U}_0|_\infty.$$

Proof The proof proceeds by induction. We assume that

$$U_l^n U_{l+1}^n < 0 \quad \forall l = 1, \dots, J-1. \quad (4.13)$$

and that

$$f(U_j^n)/U_j^n > C \quad \text{for } j = 1, \dots, J. \quad (4.14)$$

Both (4.13) and (4.14) hold by assumption for $n = 0$. Consider any $U_j^n > 0$. From (4.10) and (4.13) we deduce that

$$U_j^{n+1} \leq (1-2r)U_j^n - \Delta t f(U_j^n).$$

Hence, by (4.14),

$$\begin{aligned} U_j^{n+1} &\leq [1-2r - \Delta t \frac{f(U_j^n)}{U_j^n}] U_j^n \\ &\leq [1-2r - \Delta t C] U_j^n. \end{aligned}$$

By assumption on C we obtain

$$U_j^{n+1} \leq -(1+k)U_j^n.$$

Similarly, for $U_j^n < 0$, we obtain

$$U_j^{n+1} \geq -(1+k)U_j^n.$$

Thus (4.13)–(4.14) are true for $n = n + 1$. Furthermore we have

$$|U_j^{n+1}| \geq (1 + k)|U_j^n|$$

and the result follows by induction. \square

4.2 Existence of Absorbing Sets, Attractors and Stabilization

We now set (4.1) in the context of dissipative dynamical systems. For fixed h and Δt let $S_\Delta^h : L_2^h \rightarrow L_2^h$ be defined by

$$S_\Delta^h(\mathbf{V}) = (I - \Delta t \gamma L)\mathbf{V} - \Delta t \mathbf{f}(\mathbf{V}).$$

The continuity of S_Δ^h follows directly from (F2); the equilibria of S_Δ^h are given by the set \mathcal{E}^h of solutions to (3.8). In order for S_Δ^h to yield a dissipative dynamical system it is necessary to restrict the initial condition to the set

$$K_2^h = \{\mathbf{V} \in L_2^h : |\mathbf{V}|_\infty \leq a\},$$

for any $a \geq \bar{u}$. In the following we set

$$K = \max_{|r| \leq a} \{|f'(r)|\}.$$

We derive the following discrete analogues of (R1)–(R4) *under appropriate restrictions on the time step which are dependent upon initial data and h* and detailed in Theorem 4.10.

(RE1) ^{h} For each $\mathbf{U}_0 \in K_2^h$ there exists a unique solution of (4.1) $\mathbf{U}_n \in L_2^h$ for all $n > 0$. The mapping $\mathbf{U}_0 \rightarrow \mathbf{U}_n$ is continuous in L_2^h for each $n > 0$ and hence the family of solution operators $\{(S_\Delta^h)^n\}_{n \geq 0}$, defined by $(S_\Delta^h)^n \mathbf{U}_0 \equiv \mathbf{U}_n$, forms a continuous semigroup on K_2^h . \square

(RE2) ^{h} There exist constants $\{\bar{\rho}_i\}_{i=1}^2$ independent of h and Δt such that the balls

$$B_1^h = \{\mathbf{V} \in L_2^h : |\mathbf{V}| \leq \bar{\rho}_1\}$$

$$B_2^h = \{\mathbf{V} \in L_2^h : \|\mathbf{V}\| \leq \bar{\rho}_2\}$$

are absorbing sets for the semigroup $\{(S_\Delta^h)^n\}_{n \geq 0}$, that is, for each $\mathbf{U}_0 \in K_2^h$ there exists $\{n_i\}_{i=1}^2$ (depending on $\{\mathbf{U}_0, \rho_1\}$ and $\{\mathbf{U}_0, \rho_2\}$ respectively) such that

$$(S_\Delta^h)^n \mathbf{U}_0 \in B_i^h, \forall n \geq n_i, (i = 1, 2). \quad \square$$

(RE3) ^{h} There exists a global attractor $\mathcal{A}^h \subset L_2^h$ for the semigroup $\{(S_\Delta^h)^n\}_{n \geq 0}$. Furthermore \mathcal{A}^h is connected and there exists a constant $\bar{\rho}_A$ independent of h and J such that

$$\max\{|\mathbf{V}|, \|\mathbf{V}\|, |\mathbf{V}|_\infty\} \leq \bar{\rho}_A \quad \forall \mathbf{V} \in \mathcal{A}^h.$$

(RE4) ^{h} The functional defined by (3.9) is a Lyapunov functional for $\{(S_\Delta^h)^n\}_{t > 0}$. In addition, for any $\mathbf{U}_0 \in K_2^h$, the ω -limit set is contained in \mathcal{E}^h .

We proceed by obtaining some estimates of $\mathbf{U}_n = (S_\Delta^h)^n \mathbf{U}_0$. We begin by proving a time discrete version of the maximum norm bounds of Lemma 2.1. The first bound is a special case of Theorem 3.3 in [Hoff, 1978].

Lemma 4.5 *If the initial data and mesh parameters satisfy*

$$\mathbf{U}_0 \in K_2^h$$

and

$$\Delta t \leq \frac{h^2}{\gamma C_1 + K h^2}, \quad (4.15)$$

then

$$|\mathbf{U}_n|_\infty \leq a, \quad \forall n > 0.$$

If, for $\delta \in (0, 2)$

$$\Delta t \leq \min\left\{\frac{h^2}{\gamma C_1 + K h^2}, \frac{(2 - \delta)h^2}{(C_0 \gamma)}\right\} \quad (4.16)$$

and $a \geq m \geq \bar{u}$ then

$$|(\mathbf{U}_n - m\mathbf{E})_+|^2 \leq (1 + \Delta t \gamma \delta / C_p^2)^{-1} |(\mathbf{U}_0 - m\mathbf{E})_+|^2$$

and

$$|(\mathbf{U}_n + m\mathbf{E})_-|^2 \leq (1 + \Delta t \gamma \delta / C_p^2)^{-1} |(\mathbf{U}_0 + m\mathbf{E})_-|^2$$

Proof We write (4.1) componentwise as

$$U_i^{n+1} - a = U_i^n - a - \Delta t \gamma \sum_{j=1}^J L_{ij}(U_j^n - a) - \Delta t \gamma (LE)_{i,a} - \Delta t f(U_i^n).$$

Let us assume that $|\mathbf{U}_n|_\infty \leq a$. Re-writing we obtain

$$U_i^{n+1} - a = [1 - \Delta t \gamma L_{ii} - \Delta t f'(\xi_i^n)](U_i^n - a) - \Delta t \gamma \sum_{j \neq i}^J L_{ij}(U_j^n - a) - \Delta t \gamma (LE)_{i,a} - \Delta t f(a).$$

Since $|\xi_j^n| \leq a$ it follows from the fact that $f(a) > 0$, the assumptions (3.5), (3.6) on L and the restriction (4.15) on the time step that

$$U_i^{n+1} - a \leq 0$$

and it follows that

$$U_i^n \leq a$$

by induction. Similarly it may be shown that

$$U_i^n \geq -a.$$

For $a \geq m \geq \bar{u}$, it holds that

$$\begin{aligned} U_i^{n+1} - m &= (1 - \Delta t \gamma L_{ii})(U_i^n - m) - \Delta t \gamma \sum_{j \neq i}^J L_{ij}(U_j^n - m) - \Delta t \gamma (LE)_i m - \Delta t f(U_i^n) \\ &\leq (1 - \Delta t \gamma L_{ii})(U_i^n - m) - \Delta t \gamma \sum_{j \neq i}^J L_{ij}(U_j^n - m) - \Delta t f(U_i^n) \\ &\leq (1 - \Delta t \gamma L_{ii} - \Delta t f'(\xi_i^n))(U_i^n - m) - \Delta t \gamma \sum_{j \neq i}^J L_{ij}(U_j^n - m) - \Delta t f(m) \end{aligned}$$

where $|\xi_i^n| \leq a$. Thus, if $U_i^n \leq m$, then by (4.15) and (3.5)

$$\begin{aligned} U_i^{n+1} - m &\leq (1 - \Delta t \gamma L_{ii} - \Delta t f'(\xi_i^n))(U_i^n - m)_+ - \Delta t \gamma \sum_{j \neq i} L_{ij}(U_j^n - m)_+ \\ &= (U_i^n - m)_+ - \Delta t \gamma \sum_j L_{ij}(U_j^n - m)_+. \end{aligned}$$

Alternatively, if $U_i^n > m$ then $f(U_i^n) > 0$ and we again obtain

$$U_i^{n+1} - m \leq (U_i^n - m)_+ - \Delta t \gamma \sum_j L_{ij}(U_j^n - m)_+.$$

Multiplying by $(U_i^{n+1} - m)_+$ and summing gives

$$\langle \mathbf{Z}_{n+1} - \mathbf{Z}_n, \mathbf{Z}_{n+1} \rangle + \Delta t \gamma \langle \mathbf{Z}_n, \mathbf{Z}_{n+1} \rangle_A \leq 0$$

where

$$\mathbf{Z}_n = (\mathbf{U}_n - m\mathbf{E})_+.$$

This is the time discrete analogue of (3.16). Re-arranging we obtain, by (3.2),

$$\begin{aligned} |\mathbf{Z}_{n+1}|^2 - |\mathbf{Z}_n|^2 + |\mathbf{Z}_{n+1} - \mathbf{Z}_n|^2 + 2\Delta t \gamma \|\mathbf{Z}_{n+1}\|^2 &\leq 2\Delta t \gamma \|\mathbf{Z}_{n+1} - \mathbf{Z}_n\| \|\mathbf{Z}_{n+1}\| \\ &\leq \frac{\gamma \Delta t}{2 - \delta} \frac{C_0}{h^2} |\mathbf{Z}_{n+1} - \mathbf{Z}_n|^2 + \gamma \Delta t (2 - \delta) \|\mathbf{Z}_{n+1}\|^2 \end{aligned}$$

and, provided that (4.16) holds, we obtain

$$|\mathbf{Z}_{n+1}|^2 (1 + \delta \gamma \Delta t / C_p^2) \leq |\mathbf{Z}_n|^2.$$

Similarly we estimate $(\mathbf{U}_n + m\mathbf{E})_-$ and obtain the desired results. \square

Remark The preceding lemma is also true with the same proof if we set

$$a = \max\{|\mathbf{U}_0|_\infty, \bar{u}\}.$$

Lemma 4.6 *If, for $\delta \in (0, 1)$*

$$\Delta t \leq \min\{2(1 - \delta)\gamma h^2 / (\gamma C_0^{\frac{1}{2}} + hKC_p)^2, h^2 / (\gamma C_1 + Kh^2)\} \quad (4.17)$$

then

$$|\mathbf{U}_n|^2 \leq |\mathbf{U}_0|^2 (1 + 2\Delta t \gamma \delta / C_p^2)^{-n} + \frac{C_f C(\Omega) C_p^2}{\gamma \delta} [1 - (1 + 2\Delta t \gamma \delta / C_p^2)^{-n}].$$

Proof Taking the inner product of (4.1) with \mathbf{U}_{n+1} yields

$$\begin{aligned} & \frac{1}{2} \partial |\mathbf{U}_{n+1}|^2 + \frac{\Delta t}{2} |\partial \mathbf{U}_{n+1}|^2 + \gamma \|\mathbf{U}_{n+1}\|^2 + \langle \mathbf{f}(\mathbf{U}_{n+1}), \mathbf{U}_{n+1} \rangle \\ &= \langle \mathbf{f}(\mathbf{U}_{n+1}) - \mathbf{f}(\mathbf{U}_n), \mathbf{U}_{n+1} \rangle + \gamma \langle \mathbf{U}_{n+1} - \mathbf{U}_n, \mathbf{U}_{n+1} \rangle_A \\ &\leq K \Delta t |\partial \mathbf{U}_{n+1}| \|\mathbf{U}_{n+1}\| + \frac{\Delta t C_0^{\frac{1}{2}} \gamma}{h} |\partial \mathbf{U}_{n+1}| \|\mathbf{U}_{n+1}\|. \end{aligned}$$

Here we have used Lemma 4.5 and (3.2). Using Poincaré's inequality (3.1) and (3.12) we find that

$$\begin{aligned} & \frac{1}{2} \partial |\mathbf{U}_{n+1}|^2 + \frac{\Delta t}{2} |\partial \mathbf{U}_{n+1}|^2 + \gamma \|\mathbf{U}_{n+1}\|^2 \\ &\leq C_f C(\Omega) + (K \Delta t C_p + \frac{\Delta t C_0^{\frac{1}{2}} \gamma}{h}) |\partial \mathbf{U}_{n+1}| \|\mathbf{U}_{n+1}\| \end{aligned}$$

and completing the square on the left-hand side of this inequality yields

$$\begin{aligned} & \frac{1}{2} \partial |\mathbf{U}_{n+1}|^2 + \sqrt{(2\Delta t \gamma (1 - \delta))} |\partial \mathbf{U}_{n+1}| \|\mathbf{U}_{n+1}\| + \gamma \delta \|\mathbf{U}_{n+1}\|^2 \\ &\leq C_f C(\Omega) + (K \Delta t C_p + \frac{\Delta t C_0^{\frac{1}{2}} \gamma}{h}) |\partial \mathbf{U}_{n+1}| \|\mathbf{U}_{n+1}\|. \end{aligned}$$

Under the time-step restriction (4.17) we have that

$$\partial |\mathbf{U}_{n+1}|^2 + 2\gamma \delta \|\mathbf{U}_{n+1}\|^2 \leq 2C_f C(\Omega) \quad (4.18)$$

so that

$$(1 + 2\Delta t \gamma \delta / C_p^2) |\mathbf{U}_{n+1}|^2 \leq |\mathbf{U}_n|^2 + 2\Delta t C_f C(\Omega)$$

and application of the standard discrete Gronwall lemma gives the result. \square

For any $\epsilon > 0$, let

$$\rho_1 = C_f C(\Omega) C_p^2 / (\gamma \delta) + \epsilon.$$

It follows from Lemma 4.6 that there exists n_0 depending only on $|\mathbf{U}_0|$ and $|\mathbf{U}_0|_\infty$ such that

$$|\mathbf{U}_n| \leq \rho_1, \quad \forall n \geq n_0$$

provided that (4.17) holds. Thus the following statement about absorbing sets has been established.

Proposition 4.7 *There exists a constant ρ_1 independent of h and J such that the ball*

$$B_1^h = \{\mathbf{V} \in L_2^h : |\mathbf{V}| \leq \rho_1\}$$

is absorbing in \mathbf{U}^h under S_Δ^h provided the time-step restriction (4.17) holds. \square

We now prove the existence of an absorbing set in the $\|\bullet\|$ norm. To do this we employ a discrete version of the uniform Gronwall lemma, which is stated and proved in Appendix 2.

Proposition 4.8 *There exists a constant ρ_2 independent of h such that the ball*

$$B_2^h = \{\mathbf{V} \in L_2^h : \|\mathbf{V}\| \leq \rho_2\}$$

is absorbing in K_2^h under S_Δ^h provided the time-step restriction (4.17) and

$$\Delta t \leq \min\left\{\frac{2\gamma h^2}{(\gamma C_0^{\frac{1}{2}} + K C_\rho h)^2}, \frac{1 - \delta_1}{2C_F}\right\} \quad (4.19)$$

holds for $\delta_1 \in (0, 1)$. \square

Proof Taking the $\langle \bullet, \bullet \rangle_A$ inner product of M^{-1} times (4.1) with \mathbf{U}_{n+1} gives

$$\langle \mathbf{U}_{n+1}, \partial \mathbf{U}_{n+1} \rangle_A + \gamma \langle M^{-1} A \mathbf{U}_n, M^{-1} A \mathbf{U}_{n+1} \rangle + \langle \mathbf{f}(\mathbf{U}_n), \mathbf{U}_{n+1} \rangle_A = 0.$$

Hence

$$\begin{aligned} & \frac{1}{2} \partial \|\mathbf{U}_{n+1}\|^2 + \frac{\Delta t}{2} \|\partial \mathbf{U}_{n+1}\|^2 + \gamma |M^{-1} A \mathbf{U}_{n+1}|^2 + \langle \mathbf{f}(\mathbf{U}_{n+1}), \mathbf{U}_{n+1} \rangle_A \\ &= \gamma \Delta t \langle M^{-1} A \partial \mathbf{U}_{n+1}, M^{-1} A \mathbf{U}_{n+1} \rangle + \langle \mathbf{f}(\mathbf{U}_{n+1}) - \mathbf{f}(\mathbf{U}_n), \mathbf{U}_{n+1} \rangle_A \\ &\leq \gamma \Delta t |M^{-1} A \partial \mathbf{U}_{n+1}| |M^{-1} A \mathbf{U}_{n+1}| + K \Delta t \|\partial \mathbf{U}_{n+1}\| \|\mathbf{U}_{n+1}\|. \end{aligned}$$

Using (3.3), (3.4) and splitting \mathbf{f} as the difference of two monotone functions $f(u) = f(u) + C_F u - C_F u$ and applying (3.7) we obtain

$$\begin{aligned} & \frac{1}{2} \partial \|\mathbf{U}_{n+1}\|^2 + \frac{\Delta t}{2} \|\partial \mathbf{U}_{n+1}\|^2 + \gamma |M^{-1} A \mathbf{U}_{n+1}|^2 \\ &\leq C_F \|\mathbf{U}_{n+1}\|^2 + \frac{\gamma C_0^{\frac{1}{2}} \Delta t}{h} \|\partial \mathbf{U}_{n+1}\| |M^{-1} A \mathbf{U}_{n+1}| + K C_p \Delta t \|\partial \mathbf{U}_{n+1}\| |M^{-1} A \mathbf{U}_{n+1}|. \end{aligned}$$

Under restriction (4.19) we obtain

$$\frac{1}{2} \partial \|\mathbf{U}_{n+1}\|^2 \leq C_F \|\mathbf{U}_n\|^2. \quad (4.20)$$

From equation (4.18) we have that

$$2\gamma \delta \sum_{n=K_0}^{N+K_0} \Delta t \|\mathbf{U}_{n+1}\|^2 \leq 2C_f C(\Omega) r + |\mathbf{U}_{K_0}|^2$$

where

$$r = \sum_{n=K_0}^{N+K_0} \Delta t = (N+1)\Delta t.$$

By taking K_0 sufficiently large and using Proposition 4.7 we obtain

$$2\gamma\delta \sum_{K_0}^{N+K_0} \Delta t \|\mathbf{U}_{n+1}\|^2 \leq 2C_f C(\Omega)r + \rho_1^2.$$

Application of the discrete uniform Gronwall lemma to (4.20) (Appendix 2) gives

$$\|\mathbf{U}_{N+K_0+1}\|^2 \leq \exp(2C_F r / \delta_1) \left[\frac{2C_f C(\Omega)r + \rho_1^2}{2\gamma r \delta} \right]$$

for all K_0 sufficiently large. This completes the proof. \square

Proposition 4.9 *Let the mesh parameters satisfy*

$$\Delta t \leq \min \left\{ \frac{h^2}{\gamma C_1 + K h^2}, \frac{2(1-\delta)h^2}{\gamma C_0 + K h^2} \right\}, \quad (4.21)$$

for $\delta \in (0, 1)$ then $I^h(\bullet)$ is a Lyapunov functional for S_Δ^h on K_2^h .

Proof Since

$$f(s)(r-s) = F(r) - F(s) - \frac{1}{2}f'(\xi)(r-s)^2$$

it follows that, after taking the inner product of (4.1) with $\partial \mathbf{U}_{n+1}$ and using Lemma 4.5,

$$\Delta t |\partial \mathbf{U}_{n+1}|^2 + I^h(\mathbf{U}_{n+1}) \leq I^h(\mathbf{U}_n) + \frac{\gamma \Delta t^2}{2} \|\partial \mathbf{U}_{n+1}\|^2 + \frac{K \Delta t^2}{2} |\partial \mathbf{U}_{n+1}|^2.$$

Therefore, using (3.2) and the time-step restriction (4.21) we obtain

$$\delta \Delta t |\partial \mathbf{U}_{n+1}|^2 + I^h(\mathbf{U}_{n+1}) \leq I^h(\mathbf{U}_n). \quad (4.22)$$

Clearly $I^h(\mathbf{U}_n)$ is non-increasing in n and if $I^h(\mathbf{U}_{n+1}) = I^h(\mathbf{U}_n)$ then $|\partial \mathbf{U}_{n+1}| = 0$ by (4.22) which implies that $\mathbf{U}_{n+1} = \mathbf{U}_n$ and by (4.1) $\mathbf{U}_n = \mathbf{U}^* \in \mathcal{E}^h$. Thus $I^h(\bullet)$ is a Lyapunov functional for $S_\Delta^h(\bullet)$. \square

It follows from Lemma 2.1, Propositions 2.3, 4.7, 4.8, 4.9 that the following Theorem holds, noting that the boundedness of the attractor in $|\bullet|_\infty$ follows from Lemma 4.5.

Theorem 4.10 Setting

$$\mathcal{A}^h = \omega(B_1^h)$$

(RE1)^h–(RE4)^h hold under restrictions (4.15), (4.16), (4.17), (4.19), (4.21). \square

5. One Step Implicit Approximations: Existence of Absorbing Sets, Attractors and Stabilization

Given $\mathbf{U}_{n-1} \in L_2^h$ we consider the following schemes to find $\mathbf{U}_n \in L_2^h$ for $n \geq 1$:

$$(S1) \quad M \partial \mathbf{U}_n + \gamma A \mathbf{U}_n + M \mathbf{f}(\mathbf{U}_n) = 0. \quad (5.1)$$

$$(S2) \quad M\partial\mathbf{U}_n + \gamma A\mathbf{U}_n + M\mathbf{f}(\mathbf{U}_{n-1}) = 0. \quad (5.2)$$

$$(S3) \quad M\partial\mathbf{U}_n + \frac{\gamma}{2}A(\mathbf{U}_n + \mathbf{U}_{n-1}) + M\hat{\mathbf{f}}(\mathbf{U}_n, \mathbf{U}_{n-1}) = 0. \quad (5.3)$$

$$(S4) \quad M\partial\mathbf{U}_n + \gamma A\mathbf{U}_n + M\tilde{\mathbf{f}}(\mathbf{U}_n, \mathbf{U}_{n-1}) = 0. \quad (5.4)$$

where

$$\begin{aligned} \{\hat{\mathbf{f}}(\mathbf{U}, \mathbf{V})\}_i &:= \hat{f}(U_i, V_i) \\ \hat{f}(a, b) &:= (F(a) - F(b))/(a - b) \quad a \neq b \\ \tilde{f}(a, b) &:= f(a) \quad a = b \end{aligned}$$

and

$$\begin{aligned} \{\tilde{\mathbf{f}}(\mathbf{U}, \mathbf{V})\}_i &:= \mathbf{f}_0(\mathbf{U}) - \mathbf{f}_1(\mathbf{V}) \\ f(r) &:= f_0(r) - f_1(r). \end{aligned}$$

Under condition (F3) we may choose $f_0(\bullet)$ and $f_1(\bullet)$ to be monotone increasing; for example $f_0(r) = f(r) + rC_F$, $f_1(r) = rC_F$. We assume that such a choice has been made throughout this section. As in section 4,

$$K_2^h = \{\mathbf{V} \in L_2^h : |\mathbf{V}|_\infty \leq a\},$$

where $a \geq \bar{u}$ and we set

$$K = \max_{|r| \leq a} |f'(r)|.$$

We also set

$$K_1^h \equiv L_2^h$$

for notational convenience.

Replacing \mathbf{U}_{n-1} by \mathbf{V} and \mathbf{U}_n by \mathbf{W} in each of (5.1)–(5.4) yields systems of algebraic equations for \mathbf{W} , given \mathbf{V} . Provided they have unique solution then we write $\mathbf{W} = S_\Delta^h(\mathbf{V})$. Under appropriate restrictions (see Theorems 5.6 and 5.10) we show below that schemes (S1) and (S2) satisfy (RI1)^h–(RI4)^h with $j = 1$ for (S1) and $j = 2$ for (S2). We show that schemes (S3) and (S4) satisfy (RI1)^h, (RI3*)^h and (RI4)^h with $j = 1$.

(RI1)^h For each $\mathbf{U}_0 \in K_j^h$ there exists a unique solution $\mathbf{U}_n \in L_2^h$ for all $n > 0$. The mapping $\mathbf{U}_0 \rightarrow \mathbf{U}_n$ is continuous in L_2^h and hence the family of solution operators $\{(S_\Delta^h)^n\}_{n \geq 0}$, defined by $(S_\Delta^h)^n \mathbf{U}_0 \equiv \mathbf{U}_n$, forms a continuous semigroup on K_j^h . \square

(RI2)^h There exist constants $\{\bar{\rho}_i\}_{i=1}^2$ independent of h and J such that the balls

$$B_1^h = \{\mathbf{V} \in L_2^h : |\mathbf{V}| \leq \bar{\rho}_1\}$$

$$B_2^h = \{\mathbf{V} \in L_2^h : \|\mathbf{V}\| \leq \bar{\rho}_2\}$$

are absorbing sets for the semigroup $\{(S_\Delta^h)^n\}_{n \geq 0}$, that is, for each $\mathbf{U}_0 \in K_j^h$ there exists $\{n_i\}_{i=1}^2$ (depending on $\{\mathbf{U}_0, \rho_1\}$ and $\{\mathbf{U}_0, \rho_2\}$ respectively) such that

$$(S_\Delta^h)^n \mathbf{U}_0 \in B_i^h, \forall n \geq n_i, (i = 1, 2). \quad \square$$

(RI3)^h There exists a global attractor $\mathcal{A}^h \subset L_2^h$ for the semigroup $\{(S_\Delta^h)^n\}_{n \geq 0}$. Furthermore \mathcal{A}^h is connected and there exists a constant $\bar{\rho}_A$ independent of h and J such that

$$\max\{|\mathbf{V}|, \|\mathbf{V}\|, |\mathbf{V}|_\infty\} \leq \bar{\rho}_A \forall \mathbf{V} \in \mathcal{A}^h. \quad \square$$

(RI3*)^h Let H^h be the normed vector space $\{\mathcal{R}^J, \|\bullet\|\}$. Then $\{(S_\Delta^h)^n\}_{n \geq 0}$ is point dissipative on H^h and there exists a connected global attractor $\mathcal{A}^h \subset L_2^h$ for the semigroup $\{(S_\Delta^h)^n\}_{n \geq 0}$. \square

(RI4)^h The functional defined by (3.9) is a Lyapunov functional for $\{(S_\Delta^h)^n\}_{t > 0}$. In addition, for any $\mathbf{U}_0 \in K_j^h$, the ω -limit set is contained in \mathcal{E}^h . \square

5.1 Schemes (S1) and (S2)

We establish that (S1) and (S2) define dynamical systems and prove the existence of absorbing sets, a global attractor and a Lyapunov functional.

Lemma 5.1 *Suppose that the time-step restriction*

$$\Delta t \leq 1/C_F \quad (5.5)$$

holds for (S1) and the restriction

$$\Delta t \leq 1/K \quad (5.6)$$

holds for (S2). Then (S1) and (S2) generate continuous solution operators S_Δ^h and unique sequences $\{\mathbf{U}_n\}_{n \geq 0}$ such that

$$|\mathbf{U}_n|_\infty \leq a, \forall n \geq 0. \quad (5.7)$$

Furthermore, for $\bar{u} \leq m \leq a$ and any integer $k > 1$

$$|(\mathbf{U}_{n+2k} - m\mathbf{E})_+| \leq (1 + 2\gamma\Delta t/C_p^2)^{-k} |(\mathbf{U}_n - m\mathbf{E})_+| \quad (5.8)$$

and

$$|(\mathbf{U}_{n+2k} + m\mathbf{E})_-| \leq (1 + 2\gamma\Delta t/C_p^2)^{-k} |(\mathbf{U}_n + m\mathbf{E})_+|. \quad (5.9)$$

Proof First we show that (S1) generates a continuous operator S_Δ^h under the restriction (5.5). Existence of a solution follows from a well known variational argument based on minimising the functional

$$\frac{1}{2\Delta t} |\mathbf{W} - \mathbf{V}|^2 + \frac{\gamma}{2} \|\mathbf{W}\|^2 + \langle \mathbf{F}(\mathbf{W}), \mathbf{E} \rangle$$

over L_2^h given $\mathbf{V} \in L_2^h$ (cf. [Elliott, 1989]).

Let

$$M\mathbf{W}^i + \Delta t \gamma A\mathbf{W}^i + \Delta t M\mathbf{f}(\mathbf{W}^i) = M\mathbf{V}^i, \quad i = 1, 2,$$

and

$$\mathbf{E}_w = \mathbf{W}^1 - \mathbf{W}^2, \quad \mathbf{E}_v = \mathbf{V}^1 - \mathbf{V}^2.$$

It follows that

$$|\mathbf{E}_w|^2 + \Delta t \gamma \|\mathbf{E}_w\|^2 + \Delta t \langle \mathbf{f}(\mathbf{W}^1) - \mathbf{f}(\mathbf{W}^2), \mathbf{E}_w \rangle = \langle \mathbf{E}_v, \mathbf{E}_w \rangle.$$

Using

$$\langle \mathbf{f}(\mathbf{W}^1) - \mathbf{f}(\mathbf{W}^2), \mathbf{E}_w \rangle \geq -C_F |\mathbf{E}_w|^2$$

and (5.5) we have

$$\Delta t \gamma \|\mathbf{E}_w\|^2 \leq |\mathbf{E}_v| |\mathbf{E}_w|.$$

Hence, under the restriction (5.5) the scheme (S1) generates a unique sequence $\{\mathbf{U}_n\}$ and defines a continuous solution operator S_Δ^h . We now turn to the proof of (5.7)–(5.9) for (S1).

For $n \geq 0$ set $\mathbf{Z}_n = \mathbf{U}_n - m\mathbf{E}$. It follows that

$$\langle \partial \mathbf{Z}_n, (\mathbf{Z}_n)_+ \rangle + \gamma \langle \mathbf{Z}_n, (\mathbf{Z}_n)_+ \rangle_A + \langle \mathbf{f}(\mathbf{U}_n), (\mathbf{Z}_n)_+ \rangle = -m \langle \mathbf{E}, (\mathbf{Z}_n)_+ \rangle_A.$$

Noting (F2), (3.6), (3.7) and the inequality

$$(b - c)b_+ \geq (|b_+|^2 - |c_+|^2)/2, \quad \forall b, c \in \mathcal{R}$$

we have

$$\frac{1}{2} \partial |(\mathbf{Z}_n)_+|^2 + \gamma |(\mathbf{Z}_n)_+|^2 / C_p^2 \leq 0, \quad (5.10)$$

or

$$(1 + 2\Delta t \gamma / C_p^2) |(\mathbf{Z}_n)_+|^2 \leq |(\mathbf{Z}_{n-1})_+|^2$$

and (5.8) is an immediate consequence. (5.9) is proved similarly by consideration of $\mathbf{Z}_n = \mathbf{U}_n + m\mathbf{E}$ and then (5.7) is a consequence of (5.8), (5.9) thus completing the proof of the Lemma for (S1).

Since $M + \Delta t A$ is positive definite the system of equations (5.2) always has a unique solution so that (S2) trivially generates a unique sequence $\{\mathbf{U}_n\}$ and the continuity of f implies that S_Δ^h is continuous because of the finite dimensionality of L_2^h . Again, introducing $\mathbf{Z}_n = \mathbf{U}_n - m\mathbf{E}$ it follows from (3.6) that

$$\langle \partial \mathbf{Z}_n, (\mathbf{Z}_n)_+ \rangle + \gamma \langle \mathbf{Z}_n, (\mathbf{Z}_n)_+ \rangle_A + \langle \mathbf{f}(\mathbf{U}_{n-1}), (\mathbf{Z}_n)_+ \rangle \leq 0. \quad (5.11)$$

Either

$$Z_i^{n-1} = U_i^{n-1} - m \geq 0$$

and so

$$(-Z_i^{n-1} + \Delta t f(U_i^{n-1}))(U_i^n - m)_+ \geq -(Z_i^{n-1})(Z_i^n)_+ = -(Z_i^{n-1})_+(Z_i^n)_+$$

or

$$Z_i^{n-1} = U_i^{n-1} - m < 0$$

and

$$(-Z_i^{n-1} + \Delta t f(U_i^{n-1}))(U_i^n - m)_+$$

$\geq (-Z_i^{n-1} + \Delta t(f(U_i^{n-1}) - f(m)))(U_i^n - m)_+ = (-1 + \Delta t f'(\xi))Z_i^{n-1}(Z_i^n)_+ \geq 0 = -(Z_i^{n-1})_+(Z_i^n)_+$
provided

$$\|\mathbf{U}_{n-1}\| \leq a \quad (5.12)$$

since (5.6) holds and $f(m) \geq 0$. Thus we have shown that

$$\Delta t \langle \mathbf{f}(\mathbf{U}_{n-1}), (\mathbf{Z}_n)_+ \rangle \geq - \langle (\mathbf{Z}_{n-1})_+, (\mathbf{Z}_n)_+ \rangle + \langle \mathbf{Z}_{n-1}, (\mathbf{Z}_n)_+ \rangle.$$

Hence (5.10) follows from (5.11) provided (5.12) holds. By induction we deduce that (5.7) holds and then (5.8),(5.9) by arguments similar to those used for (S1). \square

Remark As for the explicit scheme, the preceding Lemma is also true (with an identical proof) if $a = \max\{\|\mathbf{U}_0\|_\infty, \bar{u}\}$.

From Lemma 5.1 we have

Proposition 5.2 For each $\mathbf{U}_0 \in K_1^h$ (resp. K_2^h) there exists a unique solution of (5.1) (resp. (5.2)) $\mathbf{U}_n \in L_2^h$ provided (5.5) holds (resp. (5.6)). The mapping $\mathbf{U}_0 \rightarrow \mathbf{U}_n$ is continuous in L_2^h for each $n > 0$ and hence the family of solution operators $\{(S_\Delta^h)^n\}_{n \geq 0}$ defined by $(S_\Delta^h)^n \mathbf{U}_0 = \mathbf{U}_n$ forms a discrete continuous semi-group on K_1^h (resp. K_2^h).

Lemma 5.3 For schemes (S1) and (S2)

$$|\mathbf{U}_n|^2 \leq |\mathbf{U}_0|^2 (1 + 2\Delta t \gamma \delta / C_p^2)^{-n} + \frac{C_f C(\Omega) C_p^2}{\gamma \delta} [1 - (1 + 2\Delta t \gamma \delta / C_p^2)^{-n}], \quad (5.13)$$

where $\delta = 1$ for (S1) and $\delta \in (0, 1)$ for (S2), provided that the time-step restriction

$$\Delta t \leq \min\{1/K, 2\gamma(1 - \delta)/(K^2 C_p^2)\} \quad (5.14)$$

holds for (S2).

Proof Taking the inner product of (5.1) with \mathbf{U}_n and using (3.1) yields

$$\begin{aligned} \frac{1}{2} \partial |\mathbf{U}_n|^2 + \frac{1}{2} \Delta t |\partial \mathbf{U}_n|^2 + \gamma \|\mathbf{U}_n\|^2 &\leq C_f C(\Omega) \\ \Rightarrow \frac{1}{2} \partial |\mathbf{U}_n|^2 + \frac{1}{2} \Delta t |\partial \mathbf{U}_n|^2 + \gamma |\mathbf{U}_n|^2 / C_p^2 &\leq C_f C(\Omega) \end{aligned} \quad (5.15)$$

where we have used (3.12) and Poincaré's inequality. The bound (5.13) follows from application of Gronwall's inequality. Similarly for (S2) we find, using (5.7),

$$\frac{1}{2} \partial |\mathbf{U}_n|^2 + \frac{1}{2} \Delta t |\partial \mathbf{U}_n|^2 + \gamma \|\mathbf{U}_n\|^2 \leq C_f C(\Omega) + K \Delta t |\partial \mathbf{U}_n| |\mathbf{U}_n| \quad (5.16)$$

so that

$$\frac{1}{2} \partial |\mathbf{U}_n|^2 + \frac{1}{2} \Delta t |\partial \mathbf{U}_n|^2 + \gamma |\mathbf{U}_n|^2 / C_p^2 \leq C_f C(\Omega) + K \Delta t |\partial \mathbf{U}_n| |\mathbf{U}_n|$$

and under the time-step restriction, completing the square on the left hand side it follows that

$$\frac{1}{2} \partial |\mathbf{U}_n|^2 + \delta \gamma |\mathbf{U}_n|^2 / C_o^2 \leq C_f C(\Omega).$$

Applying the Gronwall inequality, the proof of the Lemma is complete. \square

We may now prove

Proposition 5.4 *For scheme (S1) (resp. (S2)) there exist constants ρ_1, ρ_2 independent of h such that the balls*

$$B_1^h = \{\mathbf{V} \in L_2^h : |\mathbf{V}| \leq \rho_1\}, \quad B_2^h = \{\mathbf{V} \in L_2^h : \|\mathbf{V}\| \leq \rho_2\}$$

are absorbing in K_1^h (resp. K_2^h) for S_Δ^h provided the time-step restriction

$$\Delta t \leq \frac{1 - \delta}{2C_F}, \delta \in (0, 1), \quad (5.17)$$

holds (resp.

$$\Delta t \leq \min\left\{\frac{1}{K}, \frac{1 - \delta_1}{2C_F}, \frac{2\gamma(1 - \delta_2)}{K^2 C_p^2}\right\}, \delta_i \in (0, 1)). \quad (5.18)$$

Proof The absorbing sets in $|\bullet|$ follow from Lemma 5.3. Taking the $\langle \bullet, \bullet \rangle_A$ inner product of M^{-1} times (5.1) with $\partial \mathbf{U}_n$ yields, using the monotonicity of $f(u) + C_F u$

$$\|\mathbf{U}_{n+1}\|^2 - \|\mathbf{U}_n\|^2 \leq 2\Delta t C_F \|\mathbf{U}_{n+1}\|^2.$$

By (5.15) we have

$$2\Delta t \gamma \sum_{n=K_0}^{N+K_0} \|\mathbf{U}_{n+1}\|^2 \leq 2r C_f C(\Omega) + |\mathbf{U}_{K_0}|^2,$$

where $r = (N + 1)\Delta t$. Hence, for K_0 sufficiently large, Lemma 5.3 gives

$$\sum_{n=K_0}^{N+K_0} \Delta t \|\mathbf{U}_{n+1}\|^2 \leq \frac{C_f C(\Omega) r}{\gamma} + \frac{\rho_1^2}{2\gamma}.$$

Applying the discrete uniform Gronwall Lemma (Appendix 2) we obtain

$$\|\mathbf{U}_{N+K_0+1}\|^2 \leq \exp\left(\frac{2C_F r}{\delta}\right) \left[\frac{C_f C(\Omega)}{\gamma} + \frac{\rho_1^2}{2r\gamma}\right]$$

for arbitrarily large K_0 . The statement of the Lemma follows for (S1).

For (S2) a similar analysis as for (S1) yields

$$\begin{aligned} \frac{1}{2} \partial \|\mathbf{U}_n\|^2 + \frac{\Delta t}{2} \|\partial \mathbf{U}_n\|^2 + \gamma |M^{-1} A \mathbf{U}_n|^2 + \langle \mathbf{f}(\mathbf{U}_n), \mathbf{U}_n \rangle_A &= \langle \mathbf{f}(\mathbf{U}_n) - \mathbf{f}(\mathbf{U}_{n-1}), \mathbf{U}_n \rangle \\ &\leq K \Delta t \|\partial \mathbf{U}_n\| \|\mathbf{U}_n\|. \end{aligned}$$

Using the monotonicity of $f(u) + C_F u$ and inequality (3.3) we obtain, under restriction (5.18),

$$\frac{1}{2} \partial \|\mathbf{U}_n\|^2 \leq C_F \|\mathbf{U}_n\|^2.$$

From (5.16) we have

$$\frac{1}{2}\partial|\mathbf{U}_n|^2 + \frac{\Delta t}{2}|\partial\mathbf{U}_n|^2 + \frac{\gamma(1-\delta)}{C_p^2}|\mathbf{U}_n|^2 + \frac{\gamma}{\delta}\|\mathbf{U}_n\|^2 \leq C_f C(\Omega) + K\Delta t|\partial\mathbf{U}_n||\mathbf{U}_n|.$$

Completing the square on the left-hand side under restriction (5.18) we obtain

$$\frac{1}{2}\partial|\mathbf{U}_n|^2 + \gamma\delta_2\|\mathbf{U}_n\|^2 \leq C_f C(\Omega)$$

and the argument now proceeds as for (S1) to obtain

$$\|\mathbf{U}_{N+K_0+1}\|^2 \leq \exp\left(\frac{2C_F r}{\delta}\right)\left[\frac{C_f C(\Omega)}{\gamma\delta_2} + \frac{\rho_1^2}{2r\gamma\delta_2}\right]$$

for arbitrarily large K_0 . The statement of the Lemma follows for (S2). \square

Proposition 5.5 *Suppose that the time-step restrictions (5.5) (resp. (5.6)) hold for (S1) (resp. (S2)). Then for (S1) (resp. (S2)) $I^h(\bullet)$ is a Lyapunov functional for S_Δ^h on K_1^h (resp. K_2^h).*

Proof Taking the inner product of (5.1) with $\partial\mathbf{U}_n$ yields

$$\Delta t|\partial\mathbf{U}_n|^2 + \frac{\gamma}{2}[\|\mathbf{U}_n\|^2 - \|\mathbf{U}_{n-1}\|^2 + \|\mathbf{U}_n - \mathbf{U}_{n-1}\|^2] + \langle \mathbf{f}(\mathbf{U}_n), \mathbf{U}_n - \mathbf{U}_{n-1} \rangle = 0.$$

Note that

$$f(r)(r-s) = F(r) - F(s) + \frac{f'(\xi)}{2}(r-s)^2.$$

Hence for (S1),

$$|\partial\mathbf{U}_n|^2 + \frac{\Delta t\gamma}{2}\|\partial\mathbf{U}_n\|^2 + \partial I^h(\mathbf{U}_n) \leq \frac{C_F\Delta t}{2}|\partial\mathbf{U}_n|^2.$$

Under the restriction (5.5) S_Δ^h defines a dynamical system and, in addition,

$$\partial I^h(\mathbf{U}_n) \leq -\frac{1}{2}|\partial\mathbf{U}_2|^2$$

and the statement of the Theorem follows.

An identical argument holds for (S2) if we note that

$$f(r)(r-s) \geq F(r) - F(s) - \frac{K}{2}(r-s)^2$$

for $|r|, |s| \leq a$. \square

By Lemma 2.1 and Propositions 2.3, 5.2 and 5.5 we have proved the following

Theorem 5.6 *Let $\mathcal{A} = \omega(B_1^h)$ for (S1) and (S2). Then, under the restrictions (5.5), (5.17) for (S1) and (5.6), (5.18) for (S2) $(RI1)^h - (RI4)^h$ hold with $j = 1$ for (S1) and $j = 2$ for (S2).*

5.2 Schemes (S3) and (S4)

We establish that (S3) and (S4) define dynamical systems on L_2^h and prove the existence of a global attractor by establishing a Lyapunov functional structure. We note that the proof of uniqueness can be used to prove continuity of S_Δ^h as for the scheme (S1).

Lemma 5.7 *There exist unique sequences generated by (S3) and (S4) for any $U_0 \in L_2^h$ provided, in the case of (S3), that for some $\delta \in (0, 1)$,*

$$\Delta t \leq \max\left\{\frac{8\gamma(1-\delta)}{(C_F C_p)^2}, \frac{2}{C_F}\right\}. \quad (5.19)$$

Proof We consider (S3) first. The analysis of existence and uniqueness is identical to that used by [Elliott,1989] for a similar scheme applied to the Cahn-Hilliard equation. Set

$$\{\mathbf{F}^n(\mathbf{W})\}_i = \int_0^{W_i} \hat{f}(U_i^n, s) ds + c_i$$

where c_i is chosen so that $\{\mathbf{F}^n(\mathbf{W})\}_i \geq 0$. A standard argument in the calculus of variations yields a minimiser of the functional

$$\frac{1}{2\Delta t} |\mathbf{W} - \mathbf{U}_{n-1}|^2 + \frac{\gamma}{4} \|\mathbf{W}\|^2 + \frac{\gamma}{2} \langle \mathbf{U}_{n-1}, \mathbf{W} \rangle_A + \langle \mathbf{E}, \mathbf{F}^n(\mathbf{W}) \rangle$$

which satisfies the Euler-Lagrange equations (5.3). Furthermore, from [Elliott, 1989]

$$(\hat{f}(r, s_1) - \hat{f}(r, s_2))(s_1 - s_2) = F[s_1, r, s_2](s_1 - s_2)^2 \geq -C_F(s_1 - s_2)^2/2$$

where $F[s_1, r, s_2]$ denotes the second divided difference. Thus, denoting the difference of two solutions to (5.3) by \mathbf{Y} for given \mathbf{U}_{n-1} we obtain

$$|\mathbf{Y}|^2 + \frac{\Delta t \gamma}{2} \|\mathbf{Y}\|^2 \leq \Delta t C_F |\mathbf{Y}|^2 / 2 \leq \Delta t C_F C_p \|\mathbf{Y}\| |\mathbf{Y}| / 2.$$

Uniqueness under the condition (5.19) on Δt follows.

Because of (F3), $f_0(\bullet)$ is monotone increasing and the existence and uniqueness of solutions to (5.4) is an immediate consequence of the method of monotonicity. \square

Thus we have established

Proposition 5.8 *For each $U_0 \in L_2^h$ there exists a unique solution of (5.3),(5.4) $U_n \in L_2^h$ provided (5.19) holds for (S3). The mapping $U_0 \rightarrow U_n$ is continuous in L_2^h for each $n > 0$ and hence the family of solution operators $\{(S_\Delta^h)^n\}_{n \geq 0}$ defined by $(S_\Delta^h)^n U_0 = U_n$ forms a discrete continuous semi-group on L_2^h .*

Proposition 5.9 *Under the restriction (5.19) for scheme (S3) and no restriction for (S4), $I^h(\bullet)$ is a Lyapunov functional on L_2^h for the dynamical systems generated by schemes (S3) and (S4).*

Proof Taking the inner product of (S3) with $\Delta t \partial U_n$ yields

$$\Delta t |\partial U_n|^2 + I^h(U_n) - I^h(U_{n-1}) = 0.$$

This is the discrete analogue of (3.18) and the conclusion of the Lemma is evident for (S3). For (S4) we use the inequality

$$(f_0(r) - f_1(s))(r - s) = F_0(r) - F_0(s) + \frac{f'_0(\xi)}{2}(r - s)^2 + F_1(s) - F_1(r) + \frac{f'_1(\eta)}{2}(r - s)^2 \geq F(r) - F(s),$$

which holds as a result of (F3) and the monotonicity of f_0 and f_1 . Taking the inner product of (5.4) with $\Delta t \partial \mathbf{U}_n$, yields

$$\Delta t |\partial \mathbf{U}_n|^2 + I^h(\mathbf{U}_n) - I^h(\mathbf{U}_{n-1}) + \frac{\gamma \Delta t^2}{2} \|\partial \mathbf{U}_n\|^2 \leq 0,$$

and the conclusion follows for (S4). \square

Applying Lemmas 2.1 and 2.4 and Propositions 2.5, 5.8 and 5.9 we establish the following

Theorem 5.10 *Then, under the restriction (5.19) for (S3) and no restriction for (S4), (RI1)^h, (RI3*)^h and (RI4)^h hold with $j = 1$.*

6. Multi-Step Backward Differentiation Methods

In this section we consider backward differentiation formulae. Given $\{\mathbf{U}_j \in L_2^h, j = 0, 1, \dots, q - 1\}$ we define the sequence $\{\mathbf{U}_n \in L_2^h\}$ by

$$M \left(\sum_{j=1}^q \frac{\Delta t^{j-1}}{j} \partial^j \mathbf{U}_n \right) + \gamma A \mathbf{U}_n + M \mathbf{f}(\mathbf{U}_n) = 0, \quad (6.1)$$

where $q \geq 1$ is a given integer and ∂^j is defined recursively by

$$\partial^j \mathbf{V}_n := \frac{\partial^{j-1} \mathbf{V}_n - \partial^{j-1} \mathbf{V}_{n-1}}{\Delta t}, \quad j \geq 1.$$

This scheme is the result of applying the q -step backward differentiation formula to (1.8). For $q \in [1, 6]$ the formula is known to be $A(\alpha)$ -stable [Lambert, 1991]. Since in order to find \mathbf{U}_n one solves a system of the form

$$(\alpha_q M + \Delta t \gamma A) \mathbf{W} + \Delta t M \mathbf{f}(\mathbf{W}) = \mathbf{b}$$

it is straightforward to apply the proof of Lemma 5.1 to show

Lemma 6.1 *Suppose that the time-step restriction*

$$\Delta t \leq \alpha_q / C_F \quad (6.2)$$

holds. Then there exists a unique sequence $\{\mathbf{U}_n\}$ satisfying (6.1) for given $\{\mathbf{U}_j\}_0^{q-1}$. \square

Since knowing $\{\partial^j \mathbf{U}_n\}_{j=0}^{q-1}$ is equivalent to knowing $\{\mathbf{U}_{n-j}\}_{j=0}^{q-1}$ the system (6.1) is equivalent to

$$M \left(\sum_{j=0}^{q-1} \frac{\Delta t^j}{j+1} \partial^j \mathbf{U}_n \right) + \Delta t \gamma A \mathbf{U}_n + \Delta t M \mathbf{f}(\mathbf{U}_n) = M \left(\sum_{j=0}^{q-1} \frac{\Delta t^j}{j+1} \partial^j \mathbf{U}_{n-1} \right), \quad (6.3)$$

$$\Delta t \partial^j \mathbf{U}_n - \partial^{j-1} \mathbf{U}_n = -\partial^{j-1} \mathbf{U}_{n-1}, \quad j = 1, 2, \dots, q-1. \quad (6.4)$$

We let $\hat{\mathbf{V}} = (\mathbf{V}^{(1)}, \dots, \mathbf{V}^{(q)})$ be an arbitrary element of $\hat{L}_2^h := (L_2^h)^q$ and set

$$\hat{\mathbf{U}}_n := (\mathbf{U}_n, \Delta t \partial \mathbf{U}_n, \dots, \Delta t^{q-1} \partial^{q-1} \mathbf{U}_n).$$

It follows that, under the conditions of Lemma 6.1, there exists a continuous mapping $\hat{S}_\Delta^h : \hat{L}_2^h \rightarrow \hat{L}_2^h$ defined by

$$\hat{\mathbf{W}} := \hat{S}_\Delta^h(\hat{\mathbf{V}})$$

where

$$M\left(\sum_{j=1}^q \frac{\mathbf{W}^{(j)}}{j}\right) + \Delta t \gamma A \mathbf{W}^{(1)} + \Delta t M \mathbf{f}(\mathbf{W}^{(1)}) = M\left(\sum_{j=1}^q \frac{\mathbf{V}^{(j)}}{j}\right), \quad (6.5)$$

$$\mathbf{W}^{(j)} - \mathbf{W}^{(j-1)} = -\mathbf{V}^{(j-1)}, \quad j = 2, \dots, q, \quad (6.6)$$

and

$$\hat{\mathbf{U}}_n = \hat{S}_\Delta^h(\hat{\mathbf{U}}_{n-1}), \quad n \geq 1. \quad (6.7)$$

Thus (6.1) defines a discrete dynamical system on \hat{L}_2^h via (6.7). We now proceed to show the properties of the dynamical system. Clearly a fixed point $\hat{\mathbf{U}}^*$ of \hat{S}_Δ^h satisfies

$$\gamma A \mathbf{U}^{*(1)} + M \mathbf{f}(\mathbf{U}^{*(1)}) = 0, \quad \mathbf{U}^{*(j)} = 0 \quad j = 2, \dots, q. \quad (6.8)$$

It follows that the set of equilibria $\hat{\mathcal{E}}^h$ is given by $\hat{\mathbf{U}}^* = (\mathbf{U}^*, 0, 0, \dots, 0)$ where $\mathbf{U}^* \in \mathcal{E}^h$ is defined by (3.8). Thus $\hat{\mathcal{E}}^h$ is bounded and it follows that, by Proposition 2.5, the existence of a Lyapunov functional is sufficient to ensure the existence of a global attractor. In the case $q = 1$ we set

$$\hat{I}_1(\bullet) = I^h(\bullet).$$

For $q = 2, 3$ we set

$$\hat{I}_2(\hat{\mathbf{V}}) = \frac{1}{4\Delta t} |\mathbf{V}^{(2)}|^2 + I^h(\mathbf{V}^{(1)}) \quad (6.9)$$

and

$$\hat{I}_3(\hat{\mathbf{V}}) = \frac{1}{6\Delta t} |\mathbf{V}^{(3)}|^2 + \frac{5}{12\Delta t} |\mathbf{V}^{(2)}|^2 + I^h(\mathbf{V}^{(1)}). \quad (6.10)$$

We introduce the following norms on \hat{L}_2^h :

$$|\hat{\mathbf{V}}| = \left(\sum_{j=1}^q |\mathbf{V}^{(j)}|^2\right)^{\frac{1}{2}}$$

and

$$\|\hat{\mathbf{V}}\| = \left(\sum_{j=1}^q \|\mathbf{V}^{(j)}\|^2\right)^{\frac{1}{2}}$$

It is clear that $I(i)$ of section 2 is satisfied. Taking the inner product of (6.1) with $\Delta t \partial \mathbf{U}_n$ yields

$$\sum_{j=1}^q \frac{\Delta t^j}{j} \langle \partial^j \mathbf{U}_n, \partial \mathbf{U}_n \rangle + \frac{\gamma}{2} \Delta t^2 \|\partial \mathbf{U}_n\|^2 + I^h(\mathbf{U}_n) \leq I^h(\mathbf{U}_{n-1}) + \frac{C_F \Delta t^2}{2} |\partial \mathbf{U}_n|^2, \quad (6.11)$$

where we have used the argument of Proposition 5.5. It remains to evaluate the first term on the left-hand side of (6.11) which we call $\Gamma_q/\Delta t$. We set $\mathbf{Z}_n = \mathbf{U}_n - \mathbf{U}_{n-1}$ and consider the cases $q = 2$ and $q = 3$ in turn.

q=2 It may be shown that

$$\begin{aligned}\Gamma_2 &= \langle \mathbf{U}_n - \mathbf{U}_{n-1} + \frac{1}{2}(\mathbf{U}_n - 2\mathbf{U}_{n-1} + \mathbf{U}_{n-2}), \mathbf{U}_n - \mathbf{U}_{n-1} \rangle \\ &= \langle \mathbf{Z}_n + \frac{1}{2}(\mathbf{Z}_n - \mathbf{Z}_{n-1}), \mathbf{Z}_n \rangle = |\mathbf{Z}_n|^2 + \frac{1}{4}(|\mathbf{Z}_n|^2 - |\mathbf{Z}_{n-1}|^2 + |\mathbf{Z}_n - \mathbf{Z}_{n-1}|^2).\end{aligned}$$

Hence (6.11) becomes

$$\Delta t |\partial \mathbf{U}_n|^2 + \frac{\Delta t^3}{4} |\partial^2 \mathbf{U}_n|^2 + \frac{\gamma}{2} \Delta t^2 \|\partial \mathbf{U}_n\|^2 + \hat{I}_2^h(\hat{\mathbf{U}}_n) \leq \hat{I}_2^h(\hat{\mathbf{U}}_{n-1}) + \frac{C_F \Delta t^2}{2} |\partial \mathbf{U}_n|^2. \quad (6.12)$$

q=3 It is a calculation to show that

$$\begin{aligned}\Gamma_3 &= \langle \mathbf{Z}_n + \frac{1}{2}(\mathbf{Z}_n - \mathbf{Z}_{n-1}) + \frac{1}{3}(\mathbf{Z}_n - 2\mathbf{Z}_{n-1} + \mathbf{Z}_{n-2}), \mathbf{Z}_n \rangle \\ &= |\mathbf{Z}_n|^2 + \frac{1}{4}(|\mathbf{Z}_n|^2 - |\mathbf{Z}_{n-1}|^2 + |\mathbf{Z}_n - \mathbf{Z}_{n-1}|^2) \\ &+ \frac{1}{3} \left[\frac{1}{2}(|\mathbf{Z}_n|^2 - |\mathbf{Z}_{n-1}|^2 + |\mathbf{Z}_n - \mathbf{Z}_{n-1}|^2) - \frac{1}{2}|\mathbf{Z}_n|^2 - \frac{1}{2}|\mathbf{Z}_{n-1} - \mathbf{Z}_{n-2}|^2 + \frac{1}{2}|\mathbf{Z}_n + \mathbf{Z}_{n-2} - \mathbf{Z}_{n-1}|^2 \right] \\ &= \frac{5}{6}|\mathbf{Z}_n|^2 + \frac{5}{12}(|\mathbf{Z}_n|^2 - |\mathbf{Z}_{n-1}|^2) + \frac{1}{4}|\mathbf{Z}_n - \mathbf{Z}_{n-1}|^2 \\ &+ \frac{1}{6}(|\mathbf{Z}_n - \mathbf{Z}_{n-1}|^2 - |\mathbf{Z}_{n-1} - \mathbf{Z}_{n-2}|^2) + \frac{1}{6}|\mathbf{Z}_n + \mathbf{Z}_{n-2} - \mathbf{Z}_{n-1}|^2.\end{aligned}$$

Hence (6.11) becomes

$$\begin{aligned}\frac{5}{6} \Delta t |\partial \mathbf{U}_n|^2 + \frac{\Delta t^3}{4} |\partial^2 \mathbf{U}_n|^2 + \frac{\Delta t}{6} |\partial \mathbf{U}_n - \Delta t \partial^2 \mathbf{U}_{n-1}|^2 + \frac{\gamma \Delta t^2}{2} \|\partial \mathbf{U}_n\|^2 \\ \hat{I}_3(\hat{\mathbf{U}}_n) \leq \hat{I}_3(\hat{\mathbf{U}}_{n-1}) + \frac{C_F \Delta t^2}{2} |\partial \mathbf{U}_n|^2.\end{aligned} \quad (6.13)$$

Setting $\tilde{\alpha}_1 = \alpha_1 = 1$, $\tilde{\alpha}_2 = \alpha_2 = \frac{3}{2}$ and $\tilde{\alpha}_3 = \frac{5}{3}$ we have the following proposition.

Theorem 6.2 *Provided the time step restriction*

$$\Delta t \leq \tilde{\alpha}_q / C_F, \quad (6.14)$$

holds then, for $q = 1, 2, 3$ $\hat{I}_q(\bullet)$ is a Lyapunov function for the dynamical system \hat{S}_Δ^h on $\{\hat{L}_2^h, \|\bullet\|\}$. Furthermore, \hat{S}_Δ^h is point dissipative on $\{\hat{L}_2^h, \|\bullet\|\}$ and there exists a global attractor $\mathcal{A}^h \subset \hat{L}_2^h$ for the semigroup \hat{S}_Δ^h .

Proof The case $q = 1$ is considered in Proposition 5.5. Clearly we just need to check condition I(iii) of section 2 because I(ii) follows immediately from (6.12) and (6.13). We

detail the case $q = 3$ since the other case is treated in an identical fashion. It follows from (6.13) and the definition of \hat{S}_Δ^h that, if

$$\hat{I}_q(\hat{S}_\Delta^h(\hat{\mathbf{V}})) = \hat{I}_q(\hat{\mathbf{V}})$$

then, with $\hat{\mathbf{W}} = \hat{S}_\Delta^h(\hat{\mathbf{V}})$,

$$\mathbf{W}^{(2)} = \mathbf{W}^{(3)} = \mathbf{V}^{(3)} = 0.$$

Furthermore from (6.6) we have that $\mathbf{V}^{(2)} = 0$ and $\mathbf{V}^{(1)} = \mathbf{W}^{(1)}$. Hence, by (6.5), it holds that

$$A\mathbf{W}^{(1)} + Mf(\mathbf{W}^{(1)}) = 0,$$

and $\hat{\mathbf{V}} = \hat{S}_\Delta^h(\hat{\mathbf{V}}) \in \mathcal{E}^h$. By Lemma 2.1 the existence of a Lyapunov functional implies that \hat{S}_Δ^h is uniformly compact and hence that individual positive orbits are relatively compact; thus, by Proposition 2.4, \hat{S}_Δ^h is point dissipative on $\{\hat{L}_2^h, \|\bullet\|\}$. The existence of a global attractor follows from Proposition 2.5. \square

7. Numerical Results

In this section we present numerical results which illustrate the material in section 3. Numerical results illustrating the material in sections 4 and 5 may be found in [Stuart and Peplow, 1991].

We illustrate the spurious steady states and spurious stability properties implied by Theorem 3.2. For simplicity we assume, in addition to (F1) and (F4) on $f(u)$, that the equation is posed on the interval $\Omega = (0, 1)$ and that $\text{sgn}(f''(u)) = \text{sgn}(u)$. (The canonical example is still $f(u) = u^3 - u$.) It is proved in [Chafee and Infante, 1974] that (1.1)–(1.3) has $2n + 1$ steady solutions for $\gamma \in [((n+1)\pi)^{-2}, (n\pi)^{-2}]$. Thus the number of steady solutions approaches infinity as $\gamma \rightarrow 0_+$. Hence Theorem 2.1 does not seem to be worrying, at first glance – the method just resolves as many solutions as possible on the grid. However, a closer inspection shows that most of the numerical steady states are spurious: the true solutions have zeros which are equidistributed in the interval $[0, 1]$ whereas most of the numerical steady states do not share this property. A rough calculation based on the number of zeros shows that, for γ small, we would expect the numerical method to resolve approximately $2J + 1$ steady solutions (because of the equidistributed zeros property.) This implies the existence of approximately $3^J - 2J - 1$ spurious solutions.

At this point it is important to understand the dynamics of the true equation, for γ small. On a short time scale diffusion is negligible and the initial data rapidly evolves towards the stable zeros of f , ± 1 . Diffusion layers join together regions in which $u \approx 1$ and $u \approx -1$. Typically, these diffusion layers are not equidistributed and so they start to move towards a steady state configuration in which they are equidistributed. Furthermore, it is shown in [Chafee and Infante, 1974] that, for $\gamma < \pi^{-2}$, only two of the steady states are stable and that these are one-signed in the interval $(0, 1)$ – one negative and one positive. (Note that, in contrast, the numerical method has 2^J stable steady states.) The analysis of [Carr and Pego, 1989] shows that the evolution of interfaces occurs very slowly – on a

time-scale of $\mathcal{O}(\exp(-\gamma^{-\frac{1}{2}}))$. The states which evolve on such a slow time-scale are known as *meta-stable states*. Figures 7.1 and 7.2 show such an evolution starting from initial data $u_0(x) = \sin(7\pi x)$. The first figure shows the solution at times $t = 0, 20, 40$ and 80 . Notice the development of an interface separating regions in which $u > 0$ (to the left) and $u < 0$ (to the right). The interface propagates to the boundary $x = 0$ and, at $t = 80$ the unique negative stable steady solution has been reached. The propagation is very slow and this can be seen in Figure 7.2 where a three-dimensional projection of the solution $u(x, t)$ is shown.

Figure 7.1. Solutions of (1.1)–(1.3). $\Omega = [0, 1], \gamma = 0.0025, u_0(x) = \sin(7\pi x)$. For a) $t = 0$; b) $t = 20$; c) $t = 40$; d) $t = 80$. Implicit Euler applied to (1.5) with $\Delta t = 0.1, \Delta x = 0.01$.

Figure 7.2. Solution of (1.1)–(1.3). $\Omega = [0, 1], \gamma = 0.0025, u_0(x) = \sin(7\pi x)$. $0 \leq t \leq 80$. Implicit Euler applied to (1.5) with $\Delta t = 0.1, \Delta x = 0.01$.

The spurious steady states introduced by the discretisation are close in form to the meta-stable states; they are $\approx \pm 1$ everywhere except in transition layers. The resolution of the numerical method is insufficient to capture the tiny propagation speeds for the transition layers and hence these meta-stable states are stabilised by the discretisation and become steady solutions of the numerical method. Figures 7.3 and 7.4 show the numerical solution of (1.1)–(1.3) posed on the unit interval with the same initial data. In both cases the implicit Euler scheme is applied to (1.5). In Figure 7.3 a value of $\Delta x = 0.05$ is taken and the solution approaches a spurious steady state. The computed solution is smooth and there is nothing obvious which tells us that it is spurious – it is only the fact that we know *a priori* that genuine steady solutions have equidistributed zeros that enables to rule out the computation as spurious. In Figure 2.4 the value of Δx is decreased to 0.01 . Notice that the computed solution now tends to a true steady solution; however, the density of the profiles indicates that the solution evolves to form a meta-stable state before finally approaching the negative stable steady state. The meta-stable state is clearly closely related to the spurious steady state in Figure 7.3.

Figure 7.3. Solutions of (1.1)–(1.3). $\Omega = [0, 1], \gamma = 0.0025, u_0(x) = 1, x < 0.3; u_0(x) = -1, x \geq 0.3$. Implicit Euler applied to (1.5) with $\Delta t = 0.1, \Delta x = 0.05$. Profiles at intervals of 10 seconds for $0 \leq t \leq 500$.

Figure 7.4. Solutions of (1.1)–(1.3). $\Omega = [0, 1], \gamma = 0.0025, u_0(x) = 1, x < 0.3; u_0(x) = -1, x \geq 0.3$. Implicit Euler applied to (1.5) with $\Delta t = 0.1, \Delta x = 0.01$. Profiles at intervals of 10 seconds for $0 \leq t \leq 500$.

Figure 7.5 shows a trajectory approaching a steady solution which is unstable for the underlying PDE (since it changes sign) but which is artificially stabilised by the discretisation. To check that the solution is stable for the discretisation we took the computed steady state from Figure 7.5, denoted $u_s(x)$ in the caption of Figure 7.6, and perturbed it, taking this as initial data. The results are shown in Figure 7.6 which demonstrates the (spurious) stability of the steady solution.

Figure 7.5. Solutions of (1.1)–(1.3). $\Omega = [0, 1], \gamma = 0.000784, u_0(x) = 1, x < 0.3, x > 0.7; u_0(x) = -1, 0.3 \leq x \leq 0.7$. Implicit Euler applied to (1.5) with $\Delta t = 0.5, \Delta x =$

0.02. Profiles at intervals of 50 seconds for $0 \leq t \leq 5000$.

Figure 7.6. Solutions of (1.1)–(1.3). $\Omega = [0, 1]$, $\gamma = 0.000784$, $u_0(x) = u_s(x) + 0.2 * \sin(7\pi x)$. Implicit Euler applied to (1.5) with $\Delta t = 0.2$, $\Delta x = 0.02$. Profiles at intervals of 0.2 seconds for $0 \leq t \leq 10.0$

Our numerical experiments are purely one dimensional but analogous properties hold in dimensions greater than one.

Acknowledgement: A preliminary version of this work was presented at the IMA (UK) “Dynamics of Numerics and Numerics of Dynamics” conference, July 1990, Bristol. The work of CM Elliott was partially supported by the Institute of Mathematics and its Applications (Minneapolis) with funds provided by the National Science Foundation during the 1990/91 program on “Phase Transitions and Free Boundaries.” The authors are grateful to Arieh Iserles for helpful discussions regarding the material in section 3.1.

REFERENCES

- Bence, J., Merriman, B. and Osher, S. *On spurious numerical solutions of fast reaction, slow diffusion equations*. In preparation.
- Carr, J. and Pego, R. *Metastable patterns in solutions of $u_t = \epsilon^2 u_{xx} - f(u)$* . Comm. Pure. Appl. Math. **42**(1989),
- Chafee, N. and Infante, E. *A bifurcation problem for a nonlinear parabolic problem*. J. Applic. Anal. **4**(1974), 17-37.
- Crouzeix, M. and Rappaz, J. *On numerical approximation in bifurcation theory*. Masson-Springer-Verlag, Paris, 1990.
- Elliott, C.M. *The Cahn-Hilliard model for the kinetics of phase separation*. Appears in "Mathematical Models for Phase Change Problems," ed J.F.Rodrigues, Birkhauser-Verlag, 1989.
- Foias, C., Jolly, M.S., Kevrekidies, I.G. and Titi, E.S. *Dissipativity of numerical schemes*. Nonlinearity **4**(1991), 591-613.
- Griffiths, D.F. and Mitchell, A.R. *Stable periodic bifurcations of an explicit discretisation of a nonlinear partial differential equation in reaction diffusion*. IMA J. Num. Anal. **8**(1988), 435-454.
- Hale, J. *Asymptotic behaviour of dissipative systems*. AMS Mathematical Surveys and Monographs, 25. 1988.
- Henry, D. *Geometric Theory of Semilinear Parabolic Equations*. Lecture Notes in Mathematics 840, Springer, New York, 1981.
- Hoff, D. *Stability and convergence of finite difference methods for systems of nonlinear reaction-diffusion equations*. SIAM J. Num. Anal. **15**(1978), 1161-1177.
- Humphries, A.R. *Spurious solutions of numerical methods for initial value problems*. To appear IMA J. Num. Anal.
- Iserles, A. *Private Communication*. 1990.
- Khalsa, S.N.S *Finite element approximation of a reaction diffusion equation. Part I: application of a topological technique to the analysis of asymptotic behaviour of the semi-discrete equation*, Quart. Appl. Math. **44**(1986), 375-386.
- Lambert, J.D. *Numerical Methods for Ordinary Differential Equations*. Wiley, London, 1991.

Matano, H. *Asymptotic behaviour and stability of solutions of semilinear diffusion equations.* Publ. RIMS, Kyoto Univ. **15**(1979), 401-454.

Nochetto, R. *Finite element methods for parabolic free boundary problems.* Appears in "Advances in Numerical Analysis, Vol. 1" ed W.Light, Oxford University Press, 1991.

Richtmyer, R.D. and Morton, K.W., *Difference Methods for Initial Value Problems,* Academic Press, New York, 1967.

Shen, J. *Convergence of approximate attractors for a fully discrete system for reaction-diffusion equations.* Num. Func. Anal. Opt. **10**(1989), 1213-1234.

Stephens, A.B. and Shubin, G.R. *Multiple solutions and bifurcations of finite difference approximations to some steady problems of fluid mechanics.* SIAM J. Sci. Stat. Comp. **2**(1981), 404-415.

Stuart, A.M. *Linear instability implies spurious periodic solutions.* IMA J. Num. Anal. **9**(1989), 465-486.

Stuart, A.M. and Peplow, A.T. *The dynamics of the theta method.* SIAM J. Sci. Stat. Comp. **12**(1991), 1351-1372.

Temam, R. *Infinite Dimensional Dynamical Systems In Mechanics and Physics.* Springer-Verlag, New York, 1988.

Zelenyak, T.J. *Stabilization of solutions of boundary value problems for a second order parabolic equation with one space variable.* Differential Equations **4**(1968), 17-22.

APPENDIX 1

Poincare and Inverse Inequalities

Consider a diagonal matrix M with positive diagonal entries and a positive definite symmetric matrix A with eigenvectors and eigenvalues $\{\psi_i, \lambda_i\}$ satisfying

$$A\psi_i = \lambda_i M\psi_i$$

where

$$C_p^{-2} \leq \lambda_i \leq C_0/h^2.$$

Under these assumptions the following inequalities hold for all $U \in \mathcal{R}^J$:

$$|U| \leq C_p \|U\|,$$

$$\|U\|^2 \leq \frac{C_0}{h^2} |U|^2,$$

$$\|U\| \leq C_p |M^{-1}AU|,$$

and

$$|M^{-1}AU|^2 \leq \frac{C_0}{h^2} \|U\|^2,$$

where C_p and C_0 are independent of h and where $\langle \bullet, \bullet \rangle$, $|\bullet|$ and $\|\bullet\|$ are as defined in section 3.

Proof Normalise the eigenvectors so that

$$\langle \psi_i, \psi_j \rangle = \delta_{ij}.$$

Then any $U \in \mathcal{R}^J$ may be written as

$$U = \sum_{i=1}^J a_i \psi_i$$

and

$$AU = \sum_{i=1}^J a_i \lambda_i M\psi_i.$$

$$|U|^2 = \sum_{i=1}^J a_i^2$$

and

$$\|U\|^2 = \sum_{i=1}^J \lambda_i a_i^2 \geq C_p^{-2} \sum_{i=1}^J a_i^2.$$

Also

$$\|U\|^2 \leq \frac{C_0}{h^2} \sum_{i=1}^J a_i^2.$$

Hence the first pair of inequalities follow. Furthermore

$$M^{-1}AU = \sum_{i=1}^J a_i \lambda_i \psi_i$$

and hence

$$|M^{-1}AU|^2 = \sum_{i=1}^J a_i^2 \lambda_i^2$$

and the second pair of inequalities follow. \square

APPENDIX 2

A Discrete Uniform Gronwall Lemma Let G_n, P_n, Y_n be three positive sequences satisfying, $\forall n \geq n_0$ and $K_0 \geq n_0$

$$\frac{Y_{n+1} - Y_n}{\Delta t_n} \leq G_n Y_{n+1} + P_n, \quad \sum_{n=K_0}^{N+K_0} \Delta t_n G_n \leq a_1(r), \quad \sum_{n=K_0}^{N+K_0} \Delta t_n P_n \leq a_2(r), \quad \sum_{n=K_0}^{N+K_0} \Delta t_n Y_{n+1} \leq a_3(r),$$

where

$$\sum_{n=K_0}^{N+K_0} \Delta t_n = r.$$

If $\Delta t_n G_n \leq 1 - \delta \quad \forall n \geq n_0$ then

$$Y_{N+K_0+1} \leq \exp(a_1/\delta) \left[\frac{a_3}{r} + a_2 \right], \quad \forall K_0 \geq n_0.$$

Proof We have

$$(1 - \Delta t_n G_n) Y_{n+1} - Y_n \leq \Delta t_n P_n.$$

Let

$$Q_{n+1} = (1 - \Delta t_n G_n) Q_n, \quad n \geq n_0, \quad Q_{n_0} = 1$$

so that

$$Q_{n+1} Y_{n+1} - Q_n Y_n \leq \Delta t_n P_n Q_n.$$

Summing we obtain

$$Q_{N+K_0+1} Y_{N+K_0+1} - Q_M Y_M \leq \sum_{n=M}^{N+K_0} \Delta t_n P_n Q_n \leq Q_M \left(\sum_{n=M}^{N+K_0} \Delta t_n P_n \right),$$

for $M \geq K_0$. Now

$$Q_{N+K_0+1} = Q_M \prod_{n=M}^{N+K_0} (1 - \Delta t_n G_n).$$

Hence

$$Y_{N+K_0+1} \leq (Y_M + \sum_{n=M}^{N+K_0} \Delta t_n P_n) \prod_{n=M}^{N+K_0} (1 - \Delta t_n G_n)^{-1}.$$

Now

$$(1 - x)^{-1} \leq \exp(x/(1 - x)), \quad |x| < 1$$

so that

$$(1 - \Delta t_n G_n)^{-1} \leq \exp(\Delta t_n G_n / \delta).$$

Hence

$$\prod_{n=M}^{N+K_0} (1 - \Delta t_n G_n)^{-1} \leq \exp\left(\sum_{n=M}^{N+K_0} \Delta t_n G_n / \delta \right) \leq \exp(a_1/\delta).$$

Thus

$$Y_{N+K_0+1} \leq (Y_M + a_2) \exp(a_1/\delta).$$

Hence

$$\sum_{M=K_0+1}^{N+K_0+1} \Delta t_{M-1} Y_{N+K_0+1} \leq \exp(a_1/\delta) \left[\sum_{M=K_0+1}^{N+K_0+1} \Delta t_{M-1} Y_M + a_2 \left(\sum_{M=K_0+1}^{N+K_0+1} \Delta t_{M-1} \right) \right]$$

$$\Rightarrow r(Y_{N+K_0+1}) \leq \exp(a_1/\delta) [a_3 + a_2 r]$$

and division by r gives the result. \square

APPENDIX 3

Convergence of the number of discrete solutions

Throughout this appendix we use C to denote a constant independent of h . By \mathcal{G} we denote the Greens' operator for the negative Laplacian with homogeneous boundary data on Ω and we write the steady state equation for (1.1) as

$$\gamma u = \mathcal{G}f(u). \quad (A1)$$

Let $S^h \subset C(\bar{\Omega})$ denote the finite element space consisting of piecewise linear functions on an acute triangulation of Ω which vanish on $\partial\Omega$. The finite element approximation to (1.1) can be written as

$$\gamma u^h = \mathcal{G}^h f(u^h),$$

rather than the matrix version (3.8). Here $\mathcal{G}^h \in \mathcal{L}(C(\bar{\Omega}), S^h)$ is defined by

$$\langle \nabla \mathcal{G}^h \eta, \nabla \xi \rangle = \langle \eta, \xi \rangle^h$$

where $\langle \bullet, \bullet \rangle^h$ is the discrete L^2 inner product arising from lumped mass integration. It is known that (see [Crouzeix and Rappaz, 1990])

$$\|(\mathcal{G} - \mathcal{G}^h)\eta\| \leq Ch\|\eta\|_{W^{1,q}(\Omega)}, \quad q \geq 2$$

and

$$\|\mathcal{G}^h \eta\|_{W^{1,p}(\Omega)} \leq c|I^h \eta|, \quad p \geq 1$$

where I^h is the interpolation operator from $C(\bar{\Omega})$ into S^h and we use $\|\bullet\|$ and $|\bullet|$ for the $H_0^1(\Omega)$ and $L^2(\Omega)$ norms.

We suppose that (A1) has exactly Q solutions $\{u_i\}$ for a given γ and that there exists $\delta > 0$ such that in the balls

$$B_i^\delta = \{\eta : \|\eta - u_i\| \leq \delta\}$$

there is only one solution of (A1). Thus, in the closed set

$$V^\delta = H_0^1(\Omega) \setminus \left(\bigcup_i B_i^\delta \right)$$

there are no solutions and

$$\epsilon = \inf_{\eta \in V^\delta} \|\gamma \eta - \mathcal{G}f(\eta)\| > 0.$$

Hence, if $u^h \in V^\delta$ then

$$\begin{aligned} \epsilon &\leq \|\gamma u^h - \mathcal{G}f(u^h)\| = \|\gamma u^h - \mathcal{G}^h f(u^h) + (\mathcal{G}^h - \mathcal{G})f(u^h)\| \\ &\leq \|(\mathcal{G}^h - \mathcal{G})f(u^h)\| \leq Ch\|f(u^h)\|_{W^{1,q}(\Omega)}. \end{aligned}$$

Using the L^∞ and H_0^1 bounds on u^h from Lemma 3.1 we find that

$$\|f(u^h)\|_{W^{1,q}(\Omega)} \leq C$$

and so $u^h \in V^\delta$ implies that $\epsilon \leq Ch$. Thus for $h < h_0 = \epsilon/C$ we have that u^h belongs to one of the balls B_i^δ .

If (γ, u_j) is a regular point of (A1) then the analysis in [Crouziex and Rappaz, 1990, Chapter 3] implies that for h sufficiently small there is a unique solution u^h in the ball B_j^δ .




Figure 7.1. Solutions of (1.1)–(1.3). $\Omega = [0, 1], \gamma = 0.0025, u_0(x) = \sin(7\pi x)$. For a) $t = 0$; b) $t = 20$; c) $t = 40$; d) $t = 80$. Implicit Euler applied to (1.5) with $\Delta t = 0.1, \Delta x = 0.01$.

Solution Profiles




Figure 7.2. Solution of (1.1)–(1.3). $\Omega = [0, 1]$, $\gamma = 0.0025$, $u_0(x) = \sin(7\pi x)$. $0 \leq t \leq 80$. Implicit Euler applied to (1.5) with $\Delta t = 0.1$, $\Delta x = 0.01$.




Figure 7.3. Solutions of (1.1)–(1.3). $\Omega = [0, 1]$, $\gamma = 0.0025$, $u_0(x) = 1$, $x < 0.3$; $u_0(x) = -1$, $x \geq 0.3$. Implicit Euler applied to (1.5) with $\Delta t = 0.1$, $\Delta x = 0.05$. Profiles at intervals of 10 seconds for $0 \leq t \leq 500$.




Figure 7.4. Solutions of (1.1)–(1.3). $\Omega = [0, 1], \gamma = 0.0025, u_0(x) = 1, x < 0.3; u_0(x) = -1, x \geq 0.3$. Implicit Euler applied to (1.5) with $\Delta t = 0.1, \Delta x = 0.01$. Profiles at intervals of 10 seconds for $0 \leq t \leq 500$.




Figure 7.5. Solutions of (1.1)–(1.3). $\Omega = [0, 1]$, $\gamma = 0.000784$, $u_0(x) = 1$, $x < 0.3$, $x > 0.7$; $u_0(x) = -1$, $0.3 \leq x \leq 0.7$. Implicit Euler applied to (1.5) with $\Delta t = 0.5$, $\Delta x = 0.02$. Profiles at intervals of 50 seconds for $0 \leq t \leq 5000$.




Figure 7.6. Solutions of (1.1)–(1.3). $\Omega = [0, 1]$, $\gamma = 0.000784$, $u_0(x) = u_s(x) + 0.2 * \sin(7\pi x)$. Implicit Euler applied to (1.5) with $\Delta t = 0.2$, $\Delta x = 0.02$. Profiles at intervals of 0.2 seconds for $0 \leq t \leq 10.0$

#	Author/s	Title
892	E.G. Kalnins, Willard Miller, Jr. and Sanchita Mukherjee,	Models of q -algebra representations: the group of plane motions
893	T.R. Hoffend Jr. and R.K. Kaul,	Relativistic theory of superpotentials for a nonhomogeneous, spatially isotropic medium
894	Reinhold von Schwerin,	Two metal deposition on a microdisk electrode
895	Vladimir I. Oliker and Nina N. Uraltseva,	Evolution of nonparametric surfaces with speed depending on curvature, III. Some remarks on mean curvature and anisotropic flows
896	Wayne Barrett, Charles R. Johnson, Raphael Loewy and Tamir Shalom,	Rank incrementation via diagonal perturbations
898	Mingxiang Chen, Xu-Yan Chen and Jack K. Hale,	Structural stability for time-periodic one-dimensional parabolic equations
899	Hong-Ming Yin,	Global solutions of Maxwell's equations in an electromagnetic field with the temperature- dependent electrical conductivity
900	Robert Grone, Russell Merris and William Watkins,	Laplacian unimodular equivalence of graphs
901	Miroslav Fiedler,	Structure-ranks of matrices
902	Miroslav Fiedler,	An estimate for the nonstochastic eigenvalues of doubly stochastic matrices
903	Miroslav Fiedler,	Remarks on eigenvalues of Hankel matrices
904	Charles R. Johnson, D.D. Olesky, Michael Tsatsomeros and P. van den Driessche,	Spectra with positive elementary symmetric functions
905	Pierre-Alain Gremaud,	Thermal contraction as a free boundary problem
906	K.L. Cooke, Janos Turi and Gregg Turner,	Stabilization of hybrid systems in the presence of feedback delays
907	Robert P. Gilbert and Yongzhi Xu,	A numerical transmutation approach for underwater sound propagation
908	LeRoy B. Beasley, Richard A. Brualdi and Bryan L. Shader,	Combinatorial orthogonality
909	Richard A. Brualdi and Bryan L. Shader,	Strong hall matrices
910	Håkan Wennerström and David M. Anderson,	Difference versus Gaussian curvature energies; monolayer versus bilayer curvature energies applications to vesicle stability
911	Shmuel Friedland,	Eigenvalues of almost skew symmetric matrices and tournament matrices
912	Avner Friedman, Bei Hu and J.L. Velazquez,	A Free Boundary Problem Modeling Loop Dislocations in Crystals
913	Ezio Venturino,	The Influence of Diseases on Lotka-Volterra Systems
914	Steve Kirkland and Bryan L. Shader,	On Multipartite Tournament Matrices with Constant Team Size
915	Richard A. Brualdi and Jennifer J.Q. Massey,	More on Structure-Ranks of Matrices
916	Douglas B. Meade,	Qualitative Analysis of an Epidemic Model with Directed Dispersion
917	Kazuo Murota,	Mixed Matrices Irreducibility and Decomposition
918	Richard A. Brualdi and Jennifer J.Q. Massey,	Some Applications of Elementary Linear Algebra in Combinations
919	Carl D. Meyer,	Sensitivity of Markov Chains
920	Hong-Ming Yin,	Weak and Classical Solutions of Some Nonlinear Volterra Integrodifferential Equations
921	B. Leinkuhler and A. Ruehli,	Exploiting Symmetry and Regularity in Waveform Relaxation Convergence Estimation
922	Xinfu Chen and Charles M. Elliott,	Asymptotics for a Parabolic Double Obstacle Problem
923	Yongzhi Xu and Yi Yan,	An Approximate Boundary Integral Method for Acoustic Scattering in Shallow Oceans
924	Yongzhi Xu and Yi Yan,	Source Localization Processing in Perturbed Waveguides
925	Kenneth L. Cooke and Janos Turi,	Stability, Instability in Delay Equations Modeling Human Respiration
926	F. Bethuel, H. Brezis, B.D. Coleman and F. Hélein,	Bifurcation Analysis of Minimizing Harmonic Maps Describing the Equilibrium of Nematic Phases Between Cylinders
927	Frank W. Elliott, Jr.,	Signed Random Measures: Stochastic Order and Kolmogorov Consistency Conditions
928	D.A. Gregory, S.J. Kirkland and B.L. Shader,	Pick's Inequality and Tournaments
929	J.W. Demmel, N.J. Higham and R.S. Schreiber,	Block LU Factorization
930	Victor A. Galaktionov and Juan L. Vazquez,	Regional Blow-Up in a Semilinear Heat Equation with Convergence to a Hamilton-Jacobi Equation
931	Bryan L. Shader,	Convertible, Nearly Decomposable and Nearly Reducible Matrices
932	Dianne P. O'Leary,	Iterative Methods for Finding the Stationary Vector for Markov Chains
933	Nicholas J. Higham,	Perturbation theory and backward error for $AX - XB = C$
934	Z. Strakos and A. Greenbaum,	Open questions in the convergence analysis of the lanczos process for the real symmetric eigenvalue problem
935	Zhaojun Bai,	Error analysis of the lanczos algorithm for the nonsymmetric eigenvalue problem
936	Pierre-Alain Gremaud,	On an elliptic-parabolic problem related to phase transitions in shape memory alloys
937	Bojan Mohar and Neil Robertson,	Disjoint essential circuits in toroidal maps

- 939 **Bojan Mohar and Svatopluk Poljak** Eigenvalues in combinatorial optimization
- 940 **Richard A. Brualdi, Keith L. Chavey and Bryan L. Shader**, Conditional sign-solvability
- 941 **Roger Fosdick and Ying Zhang**, The torsion problem for a nonconvex stored energy function
- 942 **René Ferland and Gaston Giroux**, An unbounded mean-field intensity model:
 Propagation of the convergence of the empirical laws and compactness of the fluctuations
- 943 **Wei-Ming Ni and Izumi Takagi**, Spike-layers in semilinear elliptic singular Perturbation Problems
- 944 **Henk A. Van der Vorst and Gerard G.L. Sleijpen**, The effect of incomplete decomposition preconditioning
 on the convergence of conjugate gradients
- 945 **S.P. Hastings and L.A. Peletier**, On the decay of turbulent bursts
- 946 **Apostolos Hadjidimos and Robert J. Plemmons**, Analysis of p -cyclic iterations for Markov chains
- 947 **ÅBjörck, H. Park and L. Eldén**, Accurate downdating of least squares solutions
- 948 **E.G. Kalnins, Willard Miller, Jr. and G.C. Williams**, Recent advances in the use of separation of
 variables methods in general relativity
- 949 **G.W. Stewart**, On the perturbation of LU, Cholesky and QR factorizations
- 950 **G.W. Stewart**, Gaussian elimination, perturbation theory and Markov chains
- 951 **G.W. Stewart**, On a new way of solving the linear equations that arise in the method of least squares
- 952 **G.W. Stewart**, On the early history of the singular value decomposition
- 953 **G.W. Stewart**, On the perturbation of Markov chains with nearly transient states
- 954 **Umberto Mosco**, Composite media and asymptotic dirichlet forms
- 955 **Walter F. Mascarenhas**, The structure of the eigenvectors of sparse matrices
- 956 **Walter F. Mascarenhas**, A note on Jacobi being more accurate than QR
- 957 **Raymond H. Chan, James G. Nagy and Robert J. Plemmons**, FFT-based preconditioners for
 Toeplitz-Block least squares problems
- 958 **Zhaojun Bai**, The CSD, GSVD, their applications and computations
- 959 **D.A. Gregory, S.J. Kirkland and N.J. Pullman**, A bound on the exponent of a primitive matrix using
 Boolean rank
- 960 **Richard A. Brualdi, Shmuel Friedland and Alex Pothén**, Sparse bases, elementary vectors and nonzero
 minors of compound matrices
- 961 **J.W. Demmel**, Open problems in numerical linear algebra
- 962 **James W. Demmel and William Gragg**, On computing accurate singular values and eigenvalues of acyclic
 matrices
- 963 **James W. Demmel**, The inherent inaccuracy of implicit tridiagonal QR
- 964 **J.J.L. Velázquez**, Estimates on the $(N - 1)$ -dimensional Hausdorff measure of the blow-up set
 for a semilinear heat equation
- 965 **David C. Dobson**, Optimal design of periodic antireflective structures for the Helmholtz equation
- 966 **C.J. van Duijn and Joseph D. Fehribach**, Analysis of planar model for the molten carbonate fuel cell
- 967 **Yongzhi Xu, T. Craig Poling and Trent Brundage**, Source localization in a waveguide with unknown
 large inclusions
- 968 **J.J.L. Velázquez**, Higher dimensional blow up for semilinear parabolic equations
- 969 **E.G. Kalnins and Willard Miller, Jr.**, Separable coordinates, integrability and the Niven equations
- 970 **John M. Chadam and Hong-Ming Yin**, A diffusion equation with localized chemical reactions
- 971 **A. Greenbaum and L. Gurvits**, Max-min properties of matrix factor norms
- 972 **Bei Hu**, A free boundary problem arising in smoulder combustion
- 973 **C.M. Elliott and A.M. Stuart**, The global dynamics of discrete semilinear parabolic equations
- 974 **Avner Friedman and Jianhua Zhang**, Swelling of a rubber ball in the presence of good solvent
- 975 **Avner Friedman and Juan J.L. Velázquez**, A time-dependence free boundary problem modeling
 the visual image in electrophotography
- 976 **Richard A. Brualdi, Hyung Chan Jung and William T. Trotter, Jr.**, On the poset of all posets on
 n elements
- 977 **Ricardo D. Fierro and James R. Bunch**, Multicollinearity and total least squares
- 978 **Adam W. Bojanczyk, James G. Nagy and Robert J. Plemmons**, Row householder transformations for
 rank- k Cholesky inverse modifications
- 979 **Chaocheng Huang**, An age-dependent population model with nonlinear diffusion in R^n
- 980 **Emad Fatemi and Faroukh Odeh**, Upwind finite difference solution of Boltzmann equation applied to
 electron transport in semiconductor devices
- 981 **Esmond G. Ng and Barry W. Peyton**, A tight and explicit representation of Q in sparse QR
 factorization
- 982 **Robert J. Plemmons**, A proposal for FFT -based fast recursive least-squares
- 983 **Anne Greenbaum and Zdenek Strakos**, Matrices that generate the same Krylov residual spaces
- 984 **Alan Edelman and G.W. Stewart**, Scaling for orthogonality
- 985 **G.W. Stewart**, Note on a generalized sylvester equation
- 986 **G.W. Stewart**, Updating URV decompositions in parallel