

MN 2500 AGFO-3332

AG-FO-3332

1987

MANAGEMENT OF SOILS  
IN SOUTHWESTERN MINNESOTA  
A Correspondence Course


FEB 8 1988  
ST. PAUL COLLEGE

# Unit 1: Soil Testing

George Rehm

## Objectives

- Gain a better understanding of the importance of soil testing.
- Become acquainted with the techniques involved in obtaining accurate and representative soil samples.
- Become familiar with, in a general way, the principles involved in arriving at fertilizer recommendations.



## THE IMPORTANCE OF A SOIL TEST

There is general agreement among those who work with fertilizer recommendations and fertilizer use that there is no substitute for a regular soil testing program as a management tool for crop production. Routine soil testing provides a basis for adequate, but not excessive, fertilizer recommendations.

There are, of course, other methods for arriving at fertilizer recommendations. For example, you could use the same fertilizer program as your neighbor, replace the amounts of nutrients that the crop removes each year, or guess at the amount of fertilizer needed. All of these approaches can be expensive. Excessive fertilizer use leads to added cost without added return. Inadequate fertilizer use usually produces lower yields, reducing the economic return.

Fertilizer does not supply all of the nutrients used by the growing crop. The soil is capable of supplying some, and in many cases all, of the essential nutrients. If a nutrient in the soil is not adequate, the soil test provides a basis for determining the amount of that nutrient that should be applied in a fertilizer program.

Sometimes we expect a soil test to do more than what it is intended to do. A soil test *does not* measure the total amount of plant nutrients in soils. Except for extreme cases, it *does not* tell us if a "balanced" fertilizer program is being used. A test for a given nutrient *does not* measure the amount of that nutrient available to crops. A soil test is simply intended to give a grower an index of the nutrient status of the soil. It's a gauge to tell us if nutrient levels are low, medium, or high. If we know the relative level of a nutrient, we can develop a fertilizer program that will produce optimum yields.

If the relative level of a plant nutrient is known, the grower has some idea of the probability of getting an economic response if that nutrient is added in a fertilizer program (see Table 1).

**Table 1. Relative soil test levels and probabilities of crop response.**

Relative Level of P or K	Probability of a Response (%)
Very Low	95-100
Low	70-95
Medium	40-70
High	10-40
Very High	0-10

Source: Purdue

There is, however, no general agreement among those who make fertilizer recommendations about what the relative levels should be. For example, some may consider a 25 lb/A phosphorus (P) test to be medium, while others might consider this value to be low. This lack of agreement is one of the causes for different fertilizer recommendations by different soil testing laboratories. Relative levels for both P and potassium (K) will be discussed in future lessons.

## COLLECTING SOIL SAMPLES

Collecting soil samples is not a complicated task, but it is one that should be taken seriously. The results of any soil test can be only as good as the sample submitted for analy-

sis. In collecting soil samples, give special attention to soil type, sampling depth, and time of sample collection.

In the past, an adequate soil sample consisted of 15 to 20 cores for each 20-acre field. For today's agriculture, it is suggested that fields be sampled according to soil type. Cores from similar soils are combined into one sample using this technique, and one soil sample can represent few or many acres. In any given field, sample cores are collected from areas where the soils have similar properties. If there is more than one major soil type in a field, each major soil type should be sampled.

A copy of a modern USDA—SCS Soil Survey is needed before fields can be sampled by soil type. The old sampling procedure should be followed in counties where the modern soil survey has not been completed.

It's important to collect samples from the same depth each year to prevent wide fluctuations in fertilizer recommendations. Where subsoil samples are not taken, the sampling depth for a surface sample can vary from 0-to-6 to 0-to-8 inches without causing problems.

There are frequent complaints that soil test values fluctuate widely from year to year, creating a general lack of confidence in soil testing as a management tool. Wide fluctuations can be caused by several factors, but sampling to different depths each year is one of the major sources of error. Be consistent. Don't take samples to a depth of 6 inches one year and sample to a depth of 12-14 inches the following year.

Samples can be collected in either fall or spring. It's important, however, to be consistent in the time of sampling. There are several advantages for collecting samples in the fall. Special attention given to depth of sampling and consistent time of sampling will result in consistent, cost-effective fertilizer recommendations.

In many cases the local fertilizer dealer has the equipment and experience necessary for collecting soil samples. Generally, fertilizer dealers are not committed to sending soil samples to a specific soil testing laboratory. The grower may choose the laboratory for analysis and fertilizer recommendations.

## **Conservation Tillage Systems**

Collection of soil samples where corn is planted in a ridge-till system presents a special situation. There is no widespread agreement on the best sampling procedure to use for this planting system. We suggest that samples be taken in the ridge six inches from the row. The depth of sampling should remain at 0-to-6 or 0-to-8 inches.

## **Frequency of Sample Collection**

In southwestern Minnesota there is no need to collect soil samples each year. Soil test values for P and K do not change dramatically in a short time.

Fields should be sampled at least once in three years. This sampling schedule makes it easy to monitor any changes in soil test levels and make necessary corrections in the fertilizer program. The nitrate soil test is an important management tool for southwestern Minnesota. Samples for this test should be collected before corn and/or small grain crops are grown in the rotation.

It is important to collect samples about one year before seeding alfalfa to allow time to broadcast and incorporate lime. Annual topdress applications of P and/or K for alfalfa

also can be based on the results of analysis of soil samples collected at this time.

## **The Information Sheet**

Every laboratory that provides a chemical analysis of soil samples and corresponding fertilizer recommendations requests an information sheet for the field represented by the soil sample. Although some growers may think that the information is not necessary, it is needed before accurate fertilizer recommendations can be made. This is especially true if soil samples were taken from problem areas in a field.

The information sheet also gives the grower the opportunity to select the tests that will be performed on the soil sample. For southwestern Minnesota, it's not necessary to have each soil sample analyzed for all nutrients.

Most soil testing laboratories offer a basic test that includes a measurement of pH and organic matter content as well as the concentration of P and K. This basic test should be completed for all soil samples. The soil test for sulfur(S) is appropriate only for sandy soils. This test does not accurately predict the need for S on fine-textured soils that are dominant in southwestern Minnesota. Zinc (Zn) is the only micronutrient that may be needed for crop production in this part of the state. You may want to have a soil sample analyzed for Zn, especially if the sample was taken from a problem area. Soils in southwestern Minnesota have an adequate supply of the other micronutrients. Soils naturally have high levels of magnesium (Mg) and adequate Mg will be supplied with dolomitic lime if lime is ever needed for alfalfa production, so there is no reason to test for Mg.

As mentioned previously, the soil nitrate test is important in southwestern Minnesota. For this test, soil should be collected from 0-to-6 inches and from 6-to-24 inches. These samples are analyzed for nitrate and carry-over N is calculated. Nitrogen fertilizer recommendations are then adjusted for the amount of carry-over N.

The soil test for chloride (Cl) is new. Chloride use in a fertilizer program has increased small grain yields in South Dakota. Chloride may improve small grain yields in southwestern Minnesota. Use the sampling procedure that is used for the soil nitrate test.

## **SOIL TESTING AND DIFFERENT FERTILIZER RECOMMENDATIONS**

In recent years, many growers evaluated the usefulness of soil testing by collecting soil from a field, dividing the sample in two, and sending the two samples to different soil testing laboratories. The fertilizer recommendations returned didn't agree. So, the growers logically questioned the usefulness of soil testing in determining the amount of fertilizer to use.

Several universities in the north-central United States have taken the evaluation one step further, using the widely different recommendations and recording fertilizer costs and yields. After five years with this study in Minnesota, there are some general conclusions.

First of all, the quality of the chemical analysis is high for nearly all soil testing laboratories. The laboratory results are comparable, although values are rarely identical for a particular soil sample since there is always a small amount of variability in laboratory procedures.

There are, however, large differences in the fertilizer

recommendations that come from the analysis of identical soil samples. In the Minnesota studies, fertilizer recommendations coming from some commercial soil testing laboratories headquartered in other states would cost a corn grower \$20 to \$40 per acre per year more than recommendations from the University of Minnesota (Table 2). Thus, the soil testing laboratory that the grower chooses can have a major impact on fertilizer costs.

**Table 2. Effect of fertilizer recommendations on total crop value, total fertilizer cost, and return on sites testing very high and medium high in P and K at Waseca, six-year total (1980-1985).**

Laboratory	Very High Testing Site			Medium High Testing Site		
	Crop Value	Fertilizer Cost	Return	Crop Value	Fertilizer Cost	Return
\$/A						
A & L	1929	337	- 101	2154	371	+ 180
Harris	1946	352	- 96	2159	435	+ 121
MVTL	2009	248	+ 71	2159	283	+ 273
Cropmate*	1999	407	- 98	2159	432	+ 124
U of M	1962	206	+ 66	2165	255	+ 307
No fertilizer	1690	0	—	1603	0	—

\*Cropmate no longer has a soil testing laboratory.

The results of this comparison study emphasize the difference between two general approaches used in making fertilizer recommendations. The approach that is usually associated with the highest fertilizer costs has two objectives, to *build up* the nutrient status (primarily P and K) to a desired level, and to *maintain* the nutrient status at these optimum levels.

With this approach, more fertilizer is applied than is needed to achieve optimum yields until the nutrient status of the soil reaches the "optimum" level. There is, however, no universal agreement as to what the "optimum" levels are. For example, a soil test value of 300 pounds of K per acre might be considered "optimum" by some but low or medium to others, who would recommend adding potash to build the K to a higher level. Thus, a lack of agreement on an "optimum" soil test value creates some differences in fertilizer recommendations.

The approach that is usually associated with the lower fertilizer costs focuses on providing nutrients that will result in optimum yields. Very little emphasis is placed on achieving an "optimum" level of a plant nutrient in the soil. In this approach, plant nutrients are applied only when the soil test level indicates that there is a high probability that yields will increase if the nutrient is applied. This approach requires a considerable amount of research in the field, laboratory, and greenhouse. Lower fertilizer costs without reduced yields demonstrate that there is no substitute for field research as a basis for making fertilizer recommendations.

Growers frequently ask for a recommendation regarding the soil testing laboratory that they should use. In Minnesota, there are several laboratories that offer good service and precise analytical results. The laboratory should be chosen on the basis of the fertilizer recommendations that it provides. Growers should choose a soil testing laboratory that bases fertilizer recommendations on research conducted in their state. Excessive fertilizer recommendations are costly for the grower and can create some concerns for environmental quality.

The information given in this publication is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Minnesota Extension Service is implied.

Copyright © 1987 by Minnesota Extension Service, University of Minnesota. All rights reserved. No part of these materials may be reproduced, transmitted, transcribed, stored in a retrieval system, or translated into any language or computer language, in any form or by any means, electronic, mechanical, magnetic, optical, chemical, manual, or otherwise, without the prior written permission of Minnesota Extension Service, Distribution, Room 3 Coffey Hall, 1420 Eckles Ave., University of Minnesota, St. Paul, MN 55108.

Issued in furtherance of cooperative extension work in agriculture and home economics, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Patrick J. Borich, Dean and Director of Minnesota Extension Service, University of Minnesota, St. Paul, Minnesota 55108. The University of Minnesota, including the Minnesota Extension Service, is committed to the policy that all persons shall have equal access to its programs, facilities, and employment without regard to race, religion, color, sex, national origin, handicap, age, veteran status, or sexual orientation.