

mn 2500 AG FO 3770-13

AG-FO-3770-B
Reviewed 1990

CLEAN WATER

*You Can
Make A Difference*

Understanding Nitrogen In Soils

by
Mike O'Leary, George Rehm, and Michael Schmitt

UNIVERSITY OF MINNESOTA
DOCUMENTS
AUG 28 1990
ST. PAUL CAMPUS
LIBRARIES

UNDERSTANDING NITROGEN IN SOILS

Environmental and economic issues combined have increased the need to better understand the role and fate of nitrogen (N) in crop production systems. Nitrogen is the nutrient most often deficient for crop production in Minnesota and its use can result in substantial economic return for farmers. However, when N inputs to the soil system exceed crop needs, there is a possibility that excessive amounts of nitrate (NO_3^-) may enter

either ground or surface water.


Managing N inputs to achieve a balance between profitable crop production and environmentally tolerable levels of NO_3^- in water supplies should be every grower's goal. The behavior of N in the soil system is complex, yet an understanding of these basic processes is essential for a more efficient N management program.

NITROGEN CYCLE

Nitrogen exists in the soil system in many forms and changes (transforms) very easily from one form to another. The route that N follows in and out of the soil system is collectively called the "nitrogen cycle" (figure 1) and is biologically influenced. Biological processes, in turn, are influenced

by prevailing climatic conditions along with the physical and chemical properties of a particular soil. Both climate and soils vary greatly across Minnesota and affect the N transformations for the different areas.

Figure 1



SOURCES OF N FOR PLANT GROWTH

Nitrogen can be supplied for plant growth from several sources:

- The atmosphere
 - Biological fixation
 - Atmospheric fixation
 - Precipitation
- Commercial fertilizers
- Soil organic matter
- Crop residues
- Animal manures

Atmospheric N is the major reservoir for N in the N cycle (air is 79% N_2 gas). Although unavailable to most plants, large amounts of N_2 can be used by leguminous plants via **N fixation**. In this biological process, nodule-forming *Rhizobium* bacteria inhabit the roots of leguminous plants and

through a symbiotic relationship convert atmospheric N_2 to a form the plant can use. The amount of N_2 fixed by legumes into usable N can be substantial, with a potential for several hundred lbN/acre/year to be fixed in an alfalfa crop. Any portion of a legume crop, that is left after harvest, including roots and nodules, supplies N to the soil system. When the plant material is decomposed, N is released. Several non-symbiotic organisms exist that fix N, but N additions from these organisms are quite low (1-5 lb/acre/year). In addition small amounts of N are added to soil from **precipitation**. The amount of N supplied from precipitation averages 5-10 lb/acre/year in Minnesota.

Commercial N fertilizers are also derived from the atmospheric N pool. The major step is to combine N_2 with hydrogen (H_2) to form ammonia (NH_3). Anhydrous ammonia is then used as a starting point in the manufacture of other nitrogen fertilizers. Anhydrous ammonia or other N products derived from NH_3 can then supplement other N sources for crop nutrition.

Nitrogen can also become available for plant use from **organic N sources** which must be converted to inorganic forms before they are available to plants. Nitrogen is available to plants as either ammonium (NH_4^+) or nitrate (NO_3^-). Animal ma-

nures and other organic wastes can be important sources of N for plant growth. The amount of N supplied by manure will vary with the type of livestock, handling, rate applied, and method of application. Since the N form and content of manures varies widely, an analysis of manure is recommended to improve N management.

Crop residues from non-leguminous plants also contain N, but in relatively small amounts compared with legumes. Nitrogen exists in crop residues in complex organic forms and the residue must decay (a process that can take several years) before N is made available for plant use.

Soil organic matter is also a major source of N used by crops. Organic matter is composed primarily of rather stable material called humus that has collected over a long period of time. Easily decomposed portions of organic material disappear relatively quickly, leaving behind residues more resistant to decay. Soils contain approximately 2,000 pounds N in organic forms for each percent of organic matter. Decomposition of this portion of organic matter proceeds at a rather slow rate and releases about 20 lbN/acre/year for each percent of organic matter. A credit for the amount of N released by organic matter is built into current University of Minnesota N recommendations.

NITROGEN TRANSFORMATIONS

Nitrogen, present or added to the soil, is subject to several changes (transformations) that dictate the availability of N to plants and influence the potential movement of NO_3^- to water supplies.

Organic N that is present in soil organic matter, crop residues, and manure is converted to inorganic N through the process of **mineralization**. In this process, bacteria digest organic material and release ammonium (NH_4^+) nitrogen. Formation of NH_4^+ increases as microbial activity increases. Bacterial growth is directly related to soil temperature and water content. The ammonium supplied from fertilizers is the same as the ammonium supplied from organic matter.

Ammonium-N has properties that are of practical importance for N management. Plants can absorb NH_4^+ -N. Ammonium also has a positive

charge and, therefore, is attracted or held by negatively charged soil and soil organic matter. This means that NH_4^+ does not move downward in soils. Nitrogen in the ammonium form that is not taken up by plants is subject to other changes in the soil system.

Nitrification is the conversion of NH_4^+ -N to NO_3^- -N. Nitrification is a biological process and proceeds rapidly in warm, moist, well-aerated soils. Nitrification slows at soil temperatures below 50° F—thus, the general recommendation is that ammoniacal (NH_4^+ forming) fertilizers should not be fall-applied until soils are below 50° F. Nitrate is a negatively charged ion and is not attracted to soil particles or soil organic matter like NH_4^+ . Nitrate-N is water soluble and can move below the crop rooting zone under certain conditions.

Denitrification is a process by which bacteria convert NO_3^- to N gases that are lost to the atmosphere. Denitrifying bacteria use NO_3^- instead of oxygen in the metabolic processes. Denitrification takes place where there is waterlogged soil and where there is ample organic matter to provide energy for bacteria. For these reasons, denitrification is generally limited to topsoil. Denitrification can proceed rapidly when soils are warm and become saturated for 2 or 3 days.

A temporary reduction in the amount of plant-available N can occur from **immobilization** (tie

up) of soil N. Bacteria that decompose high carbon-low N residues, such as corn stalks or small grain straw, need more N to digest the material than is present in the residue. Immobilization occurs when nitrate and/or ammonium present in the soil is used by the growing microbes to build proteins. The actively growing bacteria that immobilize some soil N also break down soil organic matter to release available N during the growing season. There is often a net gain of N during the growing season because the additional N in the residue will be the net gain after immobilization-mineralization processes.

NITROGEN LOSS FROM THE SOIL SYSTEM

Nitrogen is lost from the soil system in several ways:

- Leaching
- Denitrification
- Volatilization
- Crop removal
- Soil erosion and runoff

In contrast to the biological transformations previously described, loss of nitrate by **leaching** is a physical event. Leaching is the loss of soluble NO_3^- as it moves with soil water, generally excess water, below the root zone. Nitrate that moves below the root zone has potential to enter either groundwater or surface water through tile drainage systems.

Coarse-textured soils have a lower water-holding capacity and, therefore, a higher potential to lose nitrate from leaching when compared with fine-textured soils. Some sandy soils, for instance, may retain only 1/2 inch of water per foot of soil while some silt loam or clay loam soils may retain up to 2 inches of water per foot. Nitrate can be leached from any soil if rainfall or irrigation moves water through the root zone.

Denitrification can be a major loss mechanism of NO_3^- when soils are saturated with water for 2 or 3 days. Nitrogen in the NH_4^+ form is not subject to this loss. Management alternatives are

available if denitrification losses are a potential problem.

Significant losses from some surface-applied N sources can occur through the process of **volatilization**. In this process, N is lost as the ammonia (NH_3) gas. Nitrogen can be lost in this way from manure and fertilizer products containing urea. Ammonia is an intermediate form of N during the process in which urea is transformed to NH_4^+ . Incorporation of these N sources will virtually eliminate volatilization losses. Loss of N from volatilization is greater when soil pH is higher than 7.3, the air temperature is high, the soil surface is moist, and there is a lot of residue on the soil.

Substantial amounts of N are lost from the soil system through **crop removal**. A 150 bu/acre corn crop, for example, removes approximately 135 pounds of N with the grain. Crop removal accounts for a majority of the N that leaves the soil system.

Nitrogen can be lost from agricultural lands through soil **erosion and runoff**. Losses through these events do not normally account for a large portion of the soil N budget, but should be considered for surface water quality issues. Incorporation or injection of manure and fertilizer can help to protect against N loss through erosion or runoff. Where soils are highly erodible, conservation tillage can reduce soil erosion and runoff, resulting in less surface loss of N.

AVOID MISCONCEPTIONS

In considering the many transformations and reactions of N in soils, there are some major points to keep in mind. Although N can be added to soil in either organic or inorganic forms, plants take up only inorganic N (that is, NO_3^- and NH_4^+). One form is not more important than the other and all sources of N can be converted to nitrate. Commercial N fertilizers, legumes, manures, and crop residues are all initial sources of NO_3^- and NH_4^+ and once in the plant or in the water supply it is impossible to identify the initial source.

Nitrate is always present in the soil solution and will move with the soil water. Inhibiting the conversion of NH_4^+ to NO_3^- can result in less N loss and more plant uptake; however, it is not possible to totally prevent nitrification. There is no way to totally prevent the movement of some NO_3^- to water supplies, but sound management practices can keep losses within acceptable limits.

SUMMARY

This publication discusses several factors that are key to understanding N behavior in a soil system. Numerous sources of N exist and must be considered when evaluating the N budget for any field or region. Nitrogen's mobility factor in the soil must be considered when developing N programs and evaluating environmental effects. Nitrogen loss from the soil system is greatly affected by soil type and climate. Sandy soils may lose N through leaching while on heavy, poorly drained soils it

may be lost through denitrification. Because Minnesota has such diverse soils and climate, interpretation of the N cycle should be site specific.

The following publications which discuss several aspects of N management in more detail can be requested through the county offices of the Minnesota Extension Service or from the Distribution Center, 3 Coffey Hall, 1420 Eckles Ave., University of Minnesota, St. Paul, MN 55108.

AG-FO-2392, *Managing Nitrogen for Corn Production on Irrigated Sandy Soils*

AG-FO-3425, *Potato Fertilization on Irrigated Soils*

AG-FO-2613, *Utilization of Animal Manure as a Fertilizer*

AG-FO-0636, *Fertilizer Urea*

AG-FO-3073, *Using Anhydrous Ammonia in Minnesota*

AG-FO-3110, *Management of Soils in Southeastern Minnesota Unit 5: Nitrogen Management*

Mike O'Leary, former assistant extension specialist, soil science, is now with the Minnesota Department of Agriculture. George Rehm and Michael Schmitt are extension soil scientists, soil fertility.

"This material is based upon work supported by the U.S. Department of Agriculture, Extension Service, under special project number 89-EWQI-1-9180."

Issued in furtherance of cooperative extension work in agriculture and home economics, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Patrick J. Borich, Dean and Director of Minnesota Extension Service, University of Minnesota, St. Paul, Minnesota 55108. The University of Minnesota, including the Minnesota Extension Service is committed to the policy that all persons shall have equal access to its programs, facilities and employment without regard to race, religion, color, sex, national origin, handicap, age, veteran status, or sexual orientation.