

**STABILITY OF MATTER THROUGH AN ELECTROSTATIC
INEQUALITY**

By

Gian Michele Graf

IMA Preprint Series # 1342

October 1995

INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS

UNIVERSITY OF MINNESOTA

**514 Vincent Hall
206 Church Street S.E.
Minneapolis, Minnesota 55455**

Stability of matter through an electrostatic inequality

Gian Michele Graf
 Theoretische Physik
 ETH Honggerberg
 CH-8093 Zurich, Switzerland

Abstract. Stability of matter is proved using an electrostatic inequality which is a manifestation of screening.

1. Introduction

Nonrelativistic matter is described by the Hamiltonian

$$H = - \sum_{i=1}^N \Delta^{(i)} + \sum_{\substack{i,j=1 \\ i < j}}^{N+M} \frac{e_i e_j}{|x_i - x_j|},$$

accounting for N fermionic electrons $i = 1, \dots, N$ and M nuclei $i = N + 1, \dots, N + M$ with positions $x_i \in \mathbb{R}^3$ and charges $e_i = -1$, resp. $1 \leq e_i \leq \text{const.}$ Stability of matter is the statement:

Theorem 1. *There is a constant C such that*

$$H \geq -C(N + M). \tag{1}$$

This result has first been proved by Dyson and Lenard [3] and subsequently by Lenard [10], Federbush [5], Eckmann [4], Lieb and Thirring [14] and Fefferman [6]. We refer to [11] for the implications of this result. More recently, stability of matter in magnetic field has been proved [7] (but see [13] for another proof), thereby extending previous results [8, 12]. Here we propose a fairly direct proof of (1) based on an electrostatic inequality: Essentially, Coulomb energies are lowered as \mathbb{R}^3 is decomposed into simplices and the interaction is restricted to pairs belonging to the same simplex. This procedure is then repeated until only a few nuclei are left in each simplex.

2. Inequalities

Let \mathcal{L} be a lattice in \mathbb{R}^3 with unit cell of unit volume: $|\mathbb{R}^3/\mathcal{L}| = 1$. An open simplex, i.e., a tetrahedron, is a bounded set

$$\Delta = \{x \in \mathbb{R}^3 \mid a_i x < c_i, i = 1, \dots, 4\} \quad (2)$$

with $a_i \in \mathbb{R}^3$, $c_i \in \mathbb{R}$. A periodic tiling of \mathbb{R}^3 is a collection $T_0 = \{\Delta_\alpha\}$ of disjoint simplices, finitely many up to congruences, such that

$$\bigcup_{\alpha \in T_0} \overline{\Delta}_\alpha = \mathbb{R}^3$$

$$T_0 + u := \{\Delta_\alpha + u\} = T_0 \quad (u \in \mathcal{L}).$$

An example is the tiling given by the \mathbb{Z}^3 -translations of the simplices obtained by cutting the unit cube $W = [0, 1]^3$ with all planes passing through the centre and an edge or a face diagonal of W . This tiling contains just one simplex up to congruences.

We now regard \mathcal{L} , T_0 as fixed and define a tiling T of scale $l > 0$ to be one congruent to lT_0 . Its simplices are also said to be of scale l . Given a tiling T (of any scale) let

$$\delta_T(x_1, x_2) = \begin{cases} 1 & \text{if } x_1, x_2 \text{ belong to the same simplex of } T, \\ 0 & \text{otherwise.} \end{cases}$$

The average of a function $f(T)$ of the tilings T of scale l is defined as

$$\langle f \rangle = \int_{\text{SO}(3) \times \mathbb{R}^3 / \mathcal{L}} d\mu(R) dy f(lR(T_0 + y)),$$

where $d\mu(R)$ is the Haar measure on $R \in \text{SO}(3)$. This definition is Euclidean invariant in the sense that it is not affected if \mathcal{L} , T_0 are replaced by $R\mathcal{L}$, $R(T_0 + y)$ for some $R \in \text{SO}(3)$, $y \in \mathbb{R}^3$.

Theorem 2. *There is $C > 0$ such that for any $N \in \mathbb{N}$, any $x_i \in \mathbb{R}^3$, $e_i \in \mathbb{R}$, ($i = 1, \dots, N$) and any $l > 0$*

$$\sum_{\substack{i,j=1 \\ i < j}}^N \frac{e_i e_j}{|x_i - x_j|} \geq \left\langle \sum_{\substack{i,j=1 \\ i < j}}^N \frac{e_i e_j}{|x_i - x_j|} \delta_T(x_i, x_j) \right\rangle - \frac{C}{l} \sum_{i=1}^N e_i^2, \quad (3)$$

where the average is over tilings T of scale l .

This result, although not explicitly stated, is contained in [9]. Previously, similar inequalities were derived in [1] and in [2]: There the tiling is made of cubes, the average is over translations and the interaction on the r.h.s. is of Yukawa type but, as here, the two sides of the inequality differ by an interaction of positive type.

The proof of (3) is based on the following two lemmas, whose proof is given in the Appendix. The spherical average of a function $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ is the function $\bar{f} : [0, +\infty) \rightarrow \mathbb{R}$ given by

$$\bar{f}(|x|) = \int_{\text{SO}(3)} d\mu(R) f(R^{-1}x). \quad (4)$$

Lemma 3. *Let Δ be a simplex with characteristic function χ . Set $\chi_-(x) = \chi(-x)$ and let $h(r)$ be the spherical average of $\chi * \chi_-$. Then $h \in C_0^2[0, +\infty)$, $h(0) = |\Delta|$ and $h''(r)$ is non-increasing in r .*

Lemma 4. *Let $h \in C^2[0, +\infty)$ with $\lim_{r \rightarrow +\infty} h(r) = 0$ and let $h''(r)$ be non-increasing. Then*

$$w(x) = \frac{h(0) - h(|x|)}{|x|}, \quad (x \in \mathbb{R}^3)$$

has positive Fourier transform: $\widehat{w}(p) \geq 0$.

Proof of Theorem 2. By scaling it suffices to prove (3) for $l = 1$. In this case it follows from the fact that the function $w(x)$ given by

$$w(x_1 - x_2) = \frac{1}{|x_1 - x_2|} (1 - \langle \delta_T(x_1, x_2) \rangle)$$

is of positive type, and that $C = w(0)/2 < +\infty$. The proof of these properties is as follows: T_0 consists of finitely many simplices $\Delta^{(i)}$, ($i = 1, \dots, n$) up to \mathcal{L} -translations. Let $\chi^{(i)}$ (resp. χ_α) be the characteristic function of $\Delta^{(i)}$, (resp. Δ_α). Using $\delta_{T_0}(x_1, x_2) = \sum_{\alpha \in T_0} \chi_\alpha(x_1) \chi_\alpha(x_2)$ and $\delta_{R(T_0+y)}(x_1, x_2) = \delta_{T_0}(R^{-1}x_1 - y, R^{-1}x_2 - y)$ we get

$$\begin{aligned} \langle \delta_T(x_1, x_2) \rangle &= \int_{\text{SO}(3) \times \mathbb{R}^3 / \mathcal{L}} d\mu(R) dy \sum_{\alpha \in T_0} \chi_\alpha(R^{-1}x_1 - y) \chi_\alpha(R^{-1}x_2 - y) \\ &= \sum_{i=1}^n \int_{\text{SO}(3) \times \mathbb{R}^3} d\mu(R) dy \chi^{(i)}(R^{-1}x_1 - y) \chi^{(i)}(R^{-1}x_2 - y) \\ &= \sum_{i=1}^n \int_{\text{SO}(3)} d\mu(R) \chi^{(i)} * \chi_-^{(i)}(R^{-1}(x_1 - x_2)). \end{aligned}$$

The claim now follows from the above lemmas, together with $\langle \delta_T(x, x) \rangle = 1$. ■

For the sake of simplicity we shall from now on assume that all Δ_α , $\alpha \in T$ are congruent to a single one $\Delta = l\Delta_0$, as in the example previously mentioned. We fix an open cube $Q_0 \supset \overline{\Delta_0}$ and let Q (resp. Q_α) be a cube in a fixed relative position to Δ (resp. Δ_α), i.e., if $\Delta_\alpha = lR(\Delta_0 + y)$ then $Q_\alpha = lR(Q_0 + y)$.

Lemma 5. *Let Δ_B be the Neumann-Laplacian for an open set $B \subset \mathbb{R}^3$. Then*

$$-\Delta_B \geq \left\langle \frac{|\Delta_0|}{|Q_0|} \sum_{\alpha \in T} (-\Delta_{Q_\alpha \cap B}) \right\rangle, \quad (5)$$

where the average is over tilings T of scale l .

Proof. This follows from integrating $|\nabla\psi|^2$ against

$$\left\langle \sum_{\alpha \in T} \chi_{Q_\alpha \cap B}(x) \right\rangle = \left\langle \sum_{\alpha \in T} \chi_{Q_\alpha}(x) \right\rangle \chi_B(x) = \frac{|Q_0|}{|\Delta_0|} \chi_B(x). \quad \blacksquare$$

Let Δ be the simplex of scale l . We shall consider Hamiltonians of the form

$$H_{\Delta,S} = \kappa l^2 K_Q + lV_S,$$

where $\kappa > 0$ will be fixed later, $S \subset \Delta$ and

$$K_Q = \sum_i^* (-\Delta_Q^{(i)}), \quad V_S = \sum_{i < j} \frac{e_i e_j}{|x_i - x_j|} \chi_S(x_i) \chi_S(x_j),$$

with the $*$ indicating that the sum is over electrons only. Scaling yields the unitary equivalence $H_{l\Delta_0, lS} \cong H_{\Delta_0, S}$. We also set $H_\Delta = H_{\Delta, \Delta}$. Moreover, let M_B , resp. N_B be the number of nuclei, resp. electrons in $B \subset \mathbb{R}^3$.

From (3, 5) we obtain a decoupling inequality for tilings T of scale l ,

$$H \geq l^{-1} \left(\left\langle \sum_{\alpha \in T} H_{\Delta_\alpha} \right\rangle - C(N + M) \right), \quad (6)$$

provided $\kappa l \leq |\Delta_0|/|Q_0|$.

3. Stability of matter

We shall prove that the ‘energy in finite volume’ is bounded below:

Proposition 6. *There are $\kappa, C > 0$ such that*

$$H_{\Delta_0} \geq -C. \quad (7)$$

Proof of Theorem 1. By scaling, the bound (7) holds for Δ of any scale and can be replaced by $H_\Delta \geq -CM_\Delta$, since $H_\Delta \geq 0$ if $M_\Delta = 0$. Together with (6) this proves stability of matter (1). \blacksquare

i) One electron, one nucleus: The uncertainty principle. For the simplex Δ_0 we have

$$-\Delta_{Q_0} - \frac{e}{|x - y|} \chi_S(x) \geq -C \quad (8)$$

for some $C > 0$, uniformly in $y \in \mathbb{R}^3$, $e \leq \text{const}$ and $S \subset \Delta_0$. To show this [6], let $V(x) = |x - y|^{-1} \chi_S(x)$ and $\psi, \nabla\psi \in L^2(Q_0)$. Set $\bar{\psi} = |Q_0|^{-1} \int_{Q_0} \psi$. Then, for any $\varepsilon > 0$,

$$\begin{aligned} (V\psi, \psi) &\leq 2\varepsilon^{-1} (V\bar{\psi}, \bar{\psi}) + 2\varepsilon (V(\psi - \bar{\psi}), \psi - \bar{\psi}) \leq 2\varepsilon^{-1} \|V\|_1 |\bar{\psi}|^2 + 2\varepsilon \|V\|_{3/2} \|\psi - \bar{\psi}\|_6^2 \\ &\leq 2\varepsilon^{-1} |Q_0|^{-1} \|V\|_1 \|\psi\|_2^2 + 2C'\varepsilon \|V\|_{3/2} \|\nabla\psi\|_2^2 \end{aligned}$$

by Hölder and Sobolev inequalities, where the p -norms are those of $L^p(Q_0)$. This proves (8).

ii) *Many electrons, no nuclei: The Pauli principle.* For the cube Q_0 we have

$$K_{Q_0} \geq c(N_{Q_0} - 1)_+^{5/3} \quad (9)$$

for some $c > 0$. This follows from the Pauli principle by filling one-particle levels.

iii) *Many electrons, many nuclei: Screening.*

Lemma 7. *Given $C_0, K > 0$ there are $C_1, \kappa > 0$ depending respectively on C_0 and K only, such that*

$$H_{\Delta_0, S} \geq C_0(N_S + M_S) - C_1 \quad (10)$$

for any $S \subset \Delta_0$, provided

$$M_S \leq K. \quad (11)$$

Proof. Let $y_j, j = 1, \dots, M_S$ be the positions of the nuclei in S . By dropping the repulsion between electrons we have for $\kappa \geq 2K$

$$\begin{aligned} H_{\Delta_0, S} &\geq \frac{\kappa}{2} K_{Q_0} + \sum_{j=1}^{M_S} \sum_i^* \left(-\frac{\kappa}{2M_S} \Delta_{Q_0}^{(i)} - \frac{e_j}{|x_i - y_j|} \chi_S(x_i) \right) + \sum_{j < k} \frac{e_j e_k}{|y_j - y_k|} \\ &\geq \frac{c\kappa}{2} (N_{Q_0} - 1)_+^{5/3} - C M_S N_{Q_0} + \text{const } M_S (M_S - 1), \end{aligned}$$

by using (8, 9) and $\text{diam}(\Delta_0) < +\infty$. Due to $M_S N_{Q_0} \leq M_S + K(N_{Q_0} - 1)_+^{5/3}$ we see that if $c\kappa/2 \geq CK + 1$ in addition then

$$H_{\Delta, S} - C_0(N_{Q_0} + M_S) \geq (N_{Q_0} - 1)_+^{5/3} - C_0 N_{Q_0} + \text{const } M_S (M_S - 1) - (C + C_0) M_S.$$

The r.h.s. has a lower bound $-C_1$ depending on C_0 only. Clearly, $N_S \leq N_{Q_0}$. ■

Lemma 8. *There is $\kappa > 0$ such that (10) holds for some $C_0, C_1 > 0$ without the restriction (11).*

Remark. This implies Proposition 6.

Proof. We choose $0 < l < \min(|\Delta_0|/|Q_0|, 1/2)$ such that if Δ is any simplex of scale l intersecting Δ_0 then $Q \subset Q_0$. Let ω be the maximal number of such Δ 's occurring in any tiling of scale l . Using (3, 5) once more we have

$$H_{\Delta_0, S} \geq l^{-1} \left(\left\langle \sum_{\substack{\alpha \in T \\ Q_\alpha \subset Q_0}} H_{\Delta_\alpha, S_\alpha} \right\rangle - C(N_S + M_S) \right), \quad (12)$$

where $S_\alpha = S \cap \Delta_\alpha$ and the average is over tilings $T = \{\Delta_\alpha\}$ of scale l . Here we dropped any $\alpha \in T$ with $Q_\alpha \not\subset Q_0$ since their contribution is purely kinetic.

We shall prove the lemma by induction in $n = 0, 1, \dots$, the induction assumption being: (10) holds provided $M_{S \cap \Delta^{(n)}} \leq K$ for any simplex $\Delta^{(n)}$ of scale l^n , where K will be

fixed below. Clearly any given configuration of non-coinciding nuclei satisfies this condition for some $n \in \mathbb{N}$. The case $n = 0$ corresponds to Lemma 7. We may thus assume $M_S > K$. If $n \geq 1$ the induction assumption applies to the simplices $\Delta_\alpha \supset S_\alpha$ of scale l , after scaling them to scale 1. The r.h.s. of (12) is thus the average of

$$\begin{aligned} l^{-1} \sum_{\substack{\alpha \in T \\ Q_\alpha \subset Q_0}} \left(H_{\Delta_\alpha, S_\alpha} - C(N_{S_\alpha} + M_{S_\alpha}) \right) &\geq l^{-1} \left((C_0 - C)(N_S + M_S) - C_1 \omega \right) \\ &= 2C_0(N_S + M_S) - \frac{C_1 \omega}{l}, \end{aligned}$$

where we set $C_0 = C(1 - 2l)^{-1}$. Now C_1 is fixed by the previous lemma, but not K . By taking it large enough (independent of n) we have

$$\frac{C_1 \omega}{l M_S} \leq \frac{C_1 \omega}{l K} \leq C_0$$

and hence $H_{\Delta_0, S} \geq C_0(N_S + M_S)$. ■

Appendix

For the convenience of the reader we include the proofs [9] of Lemma 3 and 4. We remark that $a_i \in \mathbb{R}^3$, $c_i \in \mathbb{R}$ in (2) can be normalized as

$$\sum_{i=1}^4 a_i = 0, \tag{13}$$

$$|\det(a_i, a_j, a_k)| = \frac{1}{6}. \tag{14}$$

Here i, j, k is some (and, by (13), any) triple of distinct integers in $\{1, \dots, 4\}$. Elementary considerations show that the volume of Δ is

$$|\Delta| = \left(\sum_{i=1}^4 c_i \right)_+^3, \tag{15}$$

where $x_+ = \max(x, 0)$. The spherical average (4) of a function $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ is also given by $\bar{f}(r) = \int_{S^2} d\omega f(r\omega)$, where $d\omega$ is the normalized surface measure on the unit sphere $S^2 = \{\omega \in \mathbb{R}^3 \mid |\omega| = 1\}$.

Proof of Lemma 3. We begin with

$$\chi * \chi_-(x) = \int dy \chi(x-y) \chi(-y) = \int dy \chi(y) \chi(y+x) = |\Delta \cap (\Delta - x)|.$$

Note that $y \in \Delta \cap (\Delta - x)$ iff $a_i y < c_i$ and $a_i y < c_i - a_i x$ for $i = 1, \dots, 4$, i.e., iff

$$a_i y < \min(c_i, c_i - a_i x) = c_i - (a_i x)_+ \quad (i = 1, \dots, 4).$$

Hence $\Delta \cap (\Delta - x)$ is again a simplex. According to (15) its volume is

$$|\Delta \cap (\Delta - x)| = \left(\sum_{i=1}^4 c_i - (a_i x)_+ \right)_+^3 = |\Delta| (1 - k(\omega)r)_+^3,$$

where we set $x = r\omega$ ($r \geq 0, \omega \in S^2$) and $k(\omega) = |\Delta|^{-1/3} \sum_{i=1}^4 (a_i \omega)_+$. This last function is continuous on S^2 and has a positive minimum there. Indeed, if $k(\omega) = 0$ for some $\omega \in S^2$ then $a_i \omega = 0$ for $i = 1, \dots, 4$ because of (13). Together with (14), this would imply that the four vectors $a_1, a_2, a_3, \omega \in \mathbb{R}^3$ are linearly independent, which is impossible. As a result,

$$h(r) = |\Delta| \int_{S^2} d\omega (1 - k(\omega)r)_+^3$$

has compact support. Its second derivative $h''(r) = 6|\Delta| \int_{S^2} d\omega k(\omega)^2 (1 - k(\omega)r)_+$ is continuous and non-increasing in r . ■

Proof of Lemma 4. We note that $h, -h', h'' \geq 0$. Passing to spherical coordinates we find for $p \neq 0$

$$\begin{aligned} \widehat{w}(p) &= \lim_{\varepsilon \downarrow 0} \int dx e^{-ipx} e^{-\varepsilon|x|} w(x) = \frac{4\pi}{|p|} \lim_{\varepsilon \downarrow 0} \int_0^\infty dr \sin(|p|r) e^{-\varepsilon r} (h(0) - h(r)) \\ &= - \lim_{\varepsilon \downarrow 0} \frac{4\pi}{\varepsilon^2 + |p|^2} \int_0^\infty dr (\cos(|p|r) + \frac{\varepsilon}{|p|} \sin(|p|r)) e^{-\varepsilon r} h'(r) = - \frac{4\pi}{|p|^2} \int_0^\infty dr \cos(|p|r) h'(r) \\ &= \frac{4\pi}{|p|^3} \int_0^\infty dr \sin(|p|r) h''(r) = \frac{4\pi}{|p|^4} \sum_{k=0}^\infty (-1)^k \int_0^\pi dt \sin t h''\left(\frac{k\pi + t}{|p|}\right) \geq 0. \end{aligned}$$

The third and fifth equalities are obtained by partial integration. The series above is alternating because $h'' \geq 0$ is non-increasing. Hence the final inequality. ■

Acknowledgements. I thank L. Erdős and J. Fröhlich for interesting discussions and A. Friedman and R. Gulliver for hospitality at the Institute for Mathematics and its Applications of the University of Minnesota, Minneapolis, where part of this work was done.

References.

- [1] J.G. Conlon, E.H. Lieb, H.T. Yau, The $N^{7/5}$ law for charged bosons, *Commun. Math. Phys.* **116**, 417-448 (1988).
- [2] J.G. Conlon, E.H. Lieb, H.T. Yau, The Coulomb gas at low temperature and low density, *Commun. Math. Phys.* **125**, 153-180 (1989).
- [3] F.J. Dyson, A. Lenard, Stability of matter, I, *J. Math. Phys.* **8**, 423-434 (1967).
- [4] J.P. Eckmann, Sur la stabilité de la matière, Lecture notes (unpublished).
- [5] P. Federbush, A new approach to the stability of matter problem, *J. Math. Phys.* **16**, 347-351, 706-709 (1975).
- [6] C.L. Fefferman, The uncertainty principle, *Bull. A.M.S.* **9**, 129-206 (1983).
- [7] C.L. Fefferman, unpublished.
- [8] J. Fröhlich, E.H. Lieb, M. Loss, Stability of Coulomb systems with magnetic fields I. The one-electron atom, *Commun. Math. Phys.* **104**, 251-270 (1986).
- [9] G.M. Graf, D. Schenker, On the molecular limit of Coulomb gases, *Commun. Math. Phys.*, to appear.
- [10] A. Lenard, in *Statistical Mechanics and Mathematical Problems*, A. Lenard ed., Springer (1973).
- [11] E.H. Lieb, The stability of matter, *Rev. Mod. Phys.* **48**, 553-569 (1976).
- [12] E.H. Lieb, M. Loss, Stability of Coulomb systems with magnetic fields II. The many-electron atom and the one-electron molecule, *Commun. Math. Phys.* **104**, 271-282 (1986).
- [13] E.H. Lieb, M. Loss, J.P. Solovej, Stability of matter in magnetic fields, preprint.
- [14] E.H. Lieb, W. Thirring, Bound for the kinetic energy of fermions which proves the stability of matter, *Phys. Rev. Lett.* **35**, 687-689 (1975). *Errata* **35**, 1116 (1975).

Recent IMA Preprints

#	Author/s	Title
1260	Mary Ann Horn & I. Lasiecka,	Nonlinear boundary stabilization of parallelly connected Kirchhoff plates
1261	B. Cockburn & H. Gau,	A posteriori error estimates for general numerical methods for scalar conservation laws
1262	B. Cockburn & P-A. Gremaud,	A priori error estimates for numerical methods for scalar conservation laws. Part I: The general approach
1263	R. Spigler & M. Vianello,	Convergence analysis of the semi-implicit euler method for abstract evolution equations
1264	R. Spigler & M. Vianello,	WKB-type approximation for second-order differential equations in C^* -algebras
1265	M. Menshikov & R.J. Williams,	Passage-time moments for continuous non-negative stochastic processes and applications
1266	C. Mazza,	On the storage capacity of nonlinear neural networks
1267	Z. Chen, R.E. Ewing & R. Lazarov,	Domain decomposition algorithms for mixed methods for second order elliptic problems
1268	Z. Chen, M. Espedal & R.E. Ewing,	Finite element analysis of multiphase flow in groundwater hydrology
1269	Z. Chen, R.E. Ewing, Y.A. Kuznetsov, R.D. Lazarov & S. Maliassov,	Multilevel preconditioners for mixed methods for second order elliptic problems
1270	S. Kichenassamy & G.K. Srinivasan,	The structure of WTC expansions and applications
1271	A. Zinger,	Positiveness of Wigner quasi-probability density and characterization of Gaussian distribution
1272	V. Malkin & G. Papanicolaou,	On self-focusing of short laser pulses
1273	J.N. Kutz & W.L. Kath,	Stability of pulses in nonlinear optical fibers using phase-sensitive amplifiers
1274	S.K. Patch,	Recursive recovery of a family of Markov transition probabilities from boundary value data
1275	C. Liu,	The completeness of plane waves
1276	Z. Chen & R.E. Ewing,	Stability and convergence of a finite element method for reactive transport in ground water
1277	Z. Chen & Do Y. Kwak,	The analysis of multigrid algorithms for nonconforming and mixed methods for second order elliptic problems
1278	Z. Chen,	Expanded mixed finite element methods for quasilinear second order elliptic problems II
1279	M.A. Horn & W. Littman,	Boundary control of a Schrödinger equation with nonconstant principal part
1280	R.E. Ewing, Yu.A. Kuznetsov, R.D. Lazarov & S. Maliassov,	Substructuring preconditioning for finite element approximations of second order elliptic problems. I. Nonconforming linear elements for the Poisson equation in a parallelepiped
1281	S. Maliassov,	Substructuring preconditioning for finite element approximations of second order elliptic problems. II. Mixed method for an elliptic operator with scalar tensor
1282	V. Jakšić & C.-A. Pillet,	On model for quantum friction II. Fermi's golden rule and dynamics at positive temperatures
1283	V. M. Malkin,	Kolmogorov and nonstationary spectra of optical turbulence
1284	E.G. Kalnins, V.B. Kuznetsov & W. Miller, Jr.,	Separation of variables and the XXZ Gaudin magnet
1285	E.G. Kalnins & W. Miller, Jr.,	A note on tensor products of q -algebra representations and orthogonal polynomials
1286	E.G. Kalnins & W. Miller, Jr.,	q -algebra representations of the Euclidean, pseudo-Euclidean and oscillator algebras, and their tensor products
1287	L.A. Pastur,	Spectral and probabalistic aspects of matrix models
1288	K. Kastella,	Discrimination gain to optimize detection and classification
1289	L.A. Peletier & W.C. Troy,	Spatial patterns described by the Extended Fisher-Kolmogorov (EFK) equation: Periodic solutions
1290	A. Friedman & Y. Liu,	Propagation of cracks in elastic media
1291	A. Friedman & C. Huang,	Averaged motion of charged particles in a curved strip
1292	G. R. Sell,	Global attractors for the 3D Navier-Stokes equations
1293	C. Liu,	A uniqueness result for a general class of inverse problems
1294	H-O. Kreiss,	Numerical solution of problems with different time scales II
1295	B. Cockburn, G. Gripenberg, S-O. Londen,	On convergence to entropy solutions of a single conservation law
1296	S-H. Yu,	On stability of discrete shock profiles for conservative finite difference scheme
1297	H. Behncke & P. Rejto,	A limiting absorption principle for separated Dirac operators with Wigner Von Neumann type potentials
1298	R. Lipton B. Vernescu,	Composites with imperfect interface
1299	E. Casas,	Pontryagin's principle for state-constrained boundary control problems of semilinear parabolic equations
1300	G.R. Sell,	References on dynamical systems
1301	J. Zhang,	Swelling and dissolution of polymer: A free boundary problem
1302	J. Zhang,	A nonlinear nonlocal multi-dimensional conservation law
1303	M.E. Taylor,	Estimates for approximate solutions to acoustic inverse scattering problems
1304	J. Kim & D. Sheen,	A priori estimates for elliptic boundary value problems with nonlinear boundary conditions
1305	B. Engquist & E. Luo,	New coarse grid operators for highly oscillatory coefficient elliptic problems

- 1306 **A. Boutet de Monvel & I. Egorova**, On the almost periodicity of solutions of the nonlinear Schrödinger equation with the cantor type spectrum
- 1307 **A. Boutet de Monvel & V. Georgescu**, Boundary values of the resolvent of a self-adjoint operator: Higher order estimates
- 1308 **S.K. Patch**, Diffuse tomography modulo Graßmann and Laplace
- 1309 **A. Friedman & J.J.L. Velázquez**, Liouville type theorems for fourth order elliptic equations in a half plane
- 1310 **T. Aktosun, M. Klaus & C. van der Mee**, Recovery of discontinuities in a nonhomogeneous medium
- 1311 **V. Bondarevsky**, On the global regularity problem for 3-dimensional Navier-Stokes equations
- 1312 **M. Cheney & D. Isaacson**, Inverse problems for a perturbed dissipative half-space
- 1313 **B. Cockburn, D.A. Jones & E.S. Titi**, Determining degrees of freedom for nonlinear dissipative equations
- 1314 **B. Engquist & E. Luo**, Convergence of a multigrid method for elliptic equations with highly oscillatory coefficients
- 1315 **L. Pastur & M. Shcherbina**, Universality of the local eigenvalue statistics for a class of unitary invariant random matrix ensembles
- 1316 **V. Jakšić, S. Molchanov & L. Pastur**, On the propagation properties of surface waves
- 1317 **J. Nečas, M. Ružička & V. Šverák**, On self-similar solutions of the Navier-Stokes equations
- 1318 **S. Stojanovic**, Remarks on $W^{2,p}$ -solutions of bilateral obstacle problems
- 1319 **E. Luo & H-O. Kreiss**, Pseudospectral vs. Finite difference methods for initial value problems with discontinuous coefficients
- 1320 **V.E. Grikurov**, Soliton's rebuilding in one-dimensional Schrödinger model with polynomial nonlinearity
- 1321 **J.M. Harrison & R.J. Williams**, A multiclass closed queueing network with unconventional heavy traffic behavior
- 1322 **M.E. Taylor**, Microlocal analysis on Morrey spaces
- 1323 **C. Huang**, Homogenization of biharmonic equations in domains perforated with tiny holes
- 1324 **C. Liu**, An inverse obstacle problem: A uniqueness theorem for spheres
- 1325 **M. Luskin**, Approximation of a laminated microstructure for a rotationally invariant, double well energy density
- 1326 **Rakesh & P. Sacks**, Impedance inversion from transmission data for the wave equation
- 1327 **O. Lafitte**, Diffraction for a Neumann boundary condition
- 1328 **E. Sobel, K. Lange, J.R. O'Connell & D.E. Weeks**, Haplotyping algorithms
- 1329 **B. Cockburn, D.A. Jones & E.S. Titi**, Estimating the number of asymptotic degrees of freedom for nonlinear dissipative systems
- 1330 **T. Aktosun**, Inverse Schrödinger scattering on the line with partial knowledge of the potential
- 1331 **T. Aktosun & C. van der Mee**, Partition of the potential of the one-dimensional Schrödinger equation
- 1332 **B. Engquist & E. Luo**, Convergence of the multigrid method with a wavelet coarse grid operator
- 1333 **V. Jakšić & C.-A. Pillet**, Ergodic properties of the Spin-Boson system
- 1334 **S.K. Patch**, Recursive solution for diffuse tomographic systems of arbitrary size
- 1335 **J.C. Bronski**, Semiclassical eigenvalue distribution of the non self-adjoint Zakharov-Shabat eigenvalue problem
- 1336 **J.C. Cockburn**, Bitangential structured interpolation theory
- 1337 **S. Kichenassamy**, The blow-up problem for exponential nonlinearities
- 1338 **F.A. Grünbaum & S.K. Patch**, How many parameters can one solve for in diffuse tomography?
- 1339 **R. Lipton**, Reciprocal relations, bounds and size effects for composites with highly conducting interface
- 1340 **H.A. Levine & J. Serrin**, A global nonexistence theorem for quasilinear evolution equations with dissipation
- 1341 **A. Boutet de Monvel & R. Purice**, The conjugate operator method: Application to DIRAC operators and to stratified media
- 1342 **G. Michele Graf**, Stability of matter through an electrostatic inequality
- 1343 **G. Avalos**, Sharp regularity estimates for solutions of the wave equation and their traces with prescribed Neumann data
- 1344 **G. Avalos**, The exponential stability of a coupled hyperbolic/parabolic system arising in structural acoustics
- 1345 **G. Avalos & I. Lasićka**, A differential Riccati equation for the active control of a problem in structural acoustics
- 1346 **G. Avalos**, Well-posedness for a coupled hyperbolic/parabolic system seen in structural acoustics
- 1347 **G. Avalos & I. Lasićka**, The strong stability of a semigroup arising from a coupled hyperbolic/parabolic system
- 1348 **A.V. Fursikov**, Certain optimal control problems for Navier-Stokes system with distributed control function
- 1349 **F. Gesztesy, R. Nowell & W. Pötz**, One-dimensional scattering theory for quantum systems with nontrivial spatial asymptotics
- 1350 **F. Gesztesy & H. Holden**, On trace formulas for Schrödinger-type operators
- 1351 **X. Chen**, Global asymptotic limit of solutions of the Cahn-Hilliard equation
- 1352 **X. Chen**, Lorenz equations, Part I: Existence and nonexistence of homoclinic orbits
- 1353 **X. Chen**, Lorenz equations Part II: "Randomly" rotated homoclinic orbits and chaotic trajectories
- 1354 **X. Chen**, Lorenz equations, Part III: Existence of hyperbolic sets
- 1355 **R. Abeyaratne, C. Chu & R.D. James**, Kinetics of materials with wiggly energies: Theory and application to the evolution of twinning microstructures in a Cu-Al-Ni shape memory alloy
- 1356 **C. Liu**, The Helmholtz equation on Lipschitz domains