Redwood County

Pregnancy and Birth (8)

<u></u>						
	2017					
P	Pregnancy Rates per 1,000			Birth Rates per 1,000		
	Redwood Co.	MN		Redwood Co.	MN	
15-17 years	10.3	6.6	15-17 years	10.3	4.5	
18-19 years	32.3	31.3	18-19 years	25.8	23.5	
15-19 years	17.9	16.4	15-19 years	15.7	12.1	

 15-17 years
 10.3
 4.5

 18-19 years
 25.8
 23.5

 15-19 years
 15.7
 12.1

In 2017, 2,877 teens aged 15-19 and

Number of Pregnancies and Births, 2017

Redwood Co. MN

Number of Pregnancies 19 years and under * 2,902

Number of Births 19 years and under * 2,125

* Numbers not reported for counties with <20 births or pregnancies


25 teens under age 15 became


pregnant in Minnesota. 2,113 teens
aged 15-19 and 12 teens under age
15 gave birth in Minnesota.

State Comparison: Since 1990, teen

pregnancy and birth rates have

Every day in 2017, approximately 8 adolescents became pregnant and 6 gave birth in Minnesota.





Birth Rate per 1,000 by Minnesota County among

National Comparison: From 1991 to 2017, the birth rate among adolescents age 15–19 in the United States dropped 70%, reaching a record low of 18.8 births per 1,000 (2). The decline in the adolescent birth rate over the past two decades is due to a combination of improved contraceptive use and delayed initiation of sexual activity (3). More recent declines have mainly been driven by increased use of highly effective contraceptive methods (IUDs and implants) and dual methods (4,5).

Sexually Transmitted Infections (STIs) (10)


STI rates, 2018					
(aged 15-19 per 100,000 population)					
	Redwood Co.	MN			
Chlamydia rate	573	1,624			
Gonorrhea rate	96	339			


State Comparison: Although they account for only 7% of the
population in Minnesota, adolescents aged 15-19 accounted for 25%
of the chlamydia and 17% of the gonorrhea cases in Minnesota in
2018 (9,11).

Adolescents Who Talked with Partner(s) About Protecting Themselves from STIs/HIV, 2016 Redwood Co.				
	Males	Females		
Never				
9th grade	45%	50%		
11th grade	45%	14%		
At least once per partner				
9th grade	55%	50%		
11th grade	40%	71%		

Adolescents experience a disproportionately high rate of sexually transmitted infections. This is likely to due to a combination of biological, behavioral, and cultural factors, including young women's increased anatomical vulnerability to infection, barriers to health services and peer and media influences (12).

Chlamydia and gonorrhea are at an all-time high among Minnesota youth





Gonorrhea Rate per 100,000 by Minnesota County

Prenatal Care and Low Birth Weight

important factor for improving the birth outcomes fo adolescent mothers (23).

Infants born at a low birth weight are more likely than infants born at a normal


Access to prenatal care and support from family members may be the most

weight to die within the first year of life. There are many factors associated with giving birth to a low birth weight infant, such as maternal age, health, nutrition, access to prenatal care, education level and contact with alcohol, cigarette smoke and lead (24).

No Prenatal Care or Care only in 3rd

Trimester			% Low Birth Weight		
	Redwood Co.	MN		Redwood Co.	MN
Under 15 years	0.00	16.67	Under 15 years	0.00	0
15-19 years	14.29	8.10	15-19 years	0.00	8.15
20-29 years	0.00	4.04	20-29 years	4.76	6.65
30-39 years	2.30	2.74	30-39 years	7.61	6.58
40+ years	0.00	3.91	40+ years	0.00	10.56

Percentage Low Birth Weight Among Babies Born to Mothers Aged 15-19



Sexual Behavior

The 2016 Minnesota Student Survey was administered to public school students in grades 5, 8, 9 and 11. Sexual health questions are only asked in grades 9 and 11. Approximately 71% of 9th graders and 61% of 11th graders participated in the 2016 Minnesota Student Survey (20).

