

**STABILITY AND CONVERGENCE OF A FINITE ELEMENT
METHOD FOR REACTIVE TRANSPORT IN GROUND WATER**

By

Zhangxin Chen

and

Richard E. Ewing

IMA Preprint Series # 1276

December 1994

**INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
UNIVERSITY OF MINNESOTA**

**514 Vincent Hall
206 Church Street S.E.
Minneapolis, Minnesota 55455**

STABILITY AND CONVERGENCE OF A FINITE ELEMENT METHOD FOR REACTIVE TRANSPORT IN GROUND WATER

ZHANGXIN CHEN AND RICHARD E. EWING

ABSTRACT. An explicit finite element method is used to solve the convection-diffusion-reaction equations governing contaminant transport in ground water flowing through an adsorbing porous medium. The use of discontinuous finite elements for the convective part of the equations combined with mixed finite elements for the diffusive part renders the method for the concentration solution, which displays strong gradients, trivially conservative and fully parallelizable. We carry out a stability and convergence analysis. In particular, the method is proven to satisfy a maximum principle, to be total variation bounded, and to converge to the unique weak solution of the equations. Special attention is paid to the convective part of the equations. Numerical simulations are presented and discussed.

1. Introduction. In this paper we propose and analyze a finite element method for solving the convection-diffusion-reaction equation:

$$(1.1) \quad \frac{\partial}{\partial t}(\Phi u) + \operatorname{div}(Vu - D\nabla u) = -Ku,$$

which describes the transport of a solute in a fluid phase flowing through a porous medium [1], [14]. In this case, u is the concentration of the solute in the fluid phase, V is the Darcy velocity of fluid, Φ is the volume fraction-dependent constant, D is the diffusion constant, and $K \geq 0$ is the first-order chemical reaction rate. This equation, while formally parabolic, is more nearly hyperbolic in practice. In recent years many finite element methods have been proposed to solve this important partial differential equation. The classes of optimal spatial methods and characteristic methods have been extensively studied [2], [7], [13], [15], [16], for example. However, all these finite element methods are defined by taking advantage of the parabolicity of the equation for the concentration u . As a result, the solution of the differential equation is required very smooth in the derivation of error estimates, and the constants for the error estimates blow up as the coefficient of the diffusion term goes to zero.

In this paper we propose and analyze a finite element method for numerically solving (1.1). It is similar to a finite element method introduced in [3], [4], [5], [8], [9] in that we

1991 *Mathematics Subject Classification.* 65N30, 65N10, 76S05, 76T05.

Key words and phrases. convection-diffusion-reaction equation, finite element and volume method, conservation law, stability, convergence, mixed method.

approximate the convective part of the equation using a upwinding discontinuous finite element method or a upwinding finite volume method [19], [17]. We use, however, a mixed finite element method for the diffusive part of (1.1). The main advantages of this method are that it is trivially conservative and fully parallelizable, and that it can capture discontinuities within a couple of elements without producing spurious oscillations.

A stability and convergence analysis is carried out here for the finite element method for equation (1.1) in two space dimensions. While a stability analysis was completed for the similar approach for the two-dimensional semiconductor device equations in [5], we are here able to prove much stronger results than those obtained in [5]. Namely, besides a strong maximum principle, the boundedness of the total variation and the modulus of continuity in time of the approximate solution is proven here; only an estimate on the weak derivatives of the approximate solution is given in [5]. These properties suffice to show that the numerical method converges to the weak solution of the differential equation; in [5], however, convergence of the approximate solutions to the weak solution is proven under the assumption that there is a convergent subsequence. It is also emphasized that this paper contains the first stability analysis for the two-dimensional equation (1.1) with the diffusion term included and the first convergence analysis for (1.1) with the boundary conditions. The properties derived in this paper will be exploited in a forthcoming paper where error estimates will be obtained with minimum requirements on the solution and with the property that the constant for the error estimates does not involve the small diffusion coefficient. Especially, the error estimates apply to the case of D equal to zero.

The equation (1.1) is completed by specifying the boundary and initial conditions:

$$(1.2a) \quad \partial u / \partial \nu = 0, \quad (x, y) \in \partial\Omega_1, \quad t \in J,$$

$$(1.2b) \quad u = u_D, \quad (x, y) \in \partial\Omega_2, \quad t \in J.$$

$$(1.2c) \quad u(0, x, y) = u_{\text{init}}(x, y), \quad (x, y) \in \Omega.$$

where $J = (0, T)$, $\Omega = (0, 1)^2$, $\partial\Omega = \partial\Omega_1 \cup \partial\Omega_2$ with $\partial\Omega_1 \cap \partial\Omega_2 = \emptyset$ and Ω_1 containing the endpoints of its segments, and ν denotes the normal unit-vector to $\partial\Omega$. The boundary conditions need to be modified properly in the case of $D = 0$. Namely, only the inflow boundary condition is imposed for the concentration (see (3.9) below). Moreover, in this case, note that, while the equation (1.1) is analogous to a classical conservation law, the value of the Darcy velocity V at a point (t, x, y) contains the information of all the values of the solution $u(t, \cdot, \cdot)$ on Ω . Hence a perturbation of the solution u at any given point of the domain has a global effect immediately. This is in sharp contrast with the classical conservation laws where local perturbations of the solution have a local effect in finite time.

The rest of the paper is organized as follows. The finite element method is defined in the next section. Then, in §3 we state and discuss our main results on a maximum principle (Theorem 3.1), a total variation boundedness of the scheme (Theorem 3.2), continuity with respect to data (Theorem 3.3), and convergence to the weak solution (Theorem 3.4). The proofs of these properties are carried out in §4, §5, §6, and §7, respectively. Numerical results are displayed in §8. These numerical results are devised to test the performance

of the method and to indicate the order of convergence. Finally, a concluding remark is given in §9.

2. The finite element method. In this section we define the finite element method for approximating the solution of the differential system (1.1). Toward that end, let $\{x_{i+1/2}\}_{i=0}^{n_x} \times \{y_{j+1/2}\}_{j=0}^{n_y}$ be a partition of Ω with $x_{1/2} = y_{1/2} = 0$ and $x_{n_x+1/2} = y_{n_y+1/2} = 1$ and let $\{t^n\}_{n=0}^{n_T}$ be a partition of $[0, T]$ with $t^0 = 0$ and $t^{n_T} = T$. Then, set $I_i^x = (x_{i-1/2}, x_{i+1/2})$, $I_j^y = (y_{j-1/2}, y_{j+1/2})$, $\Delta x_i = x_{i+1/2} - x_{i-1/2}$, $\Delta y_j = y_{j+1/2} - y_{j-1/2}$, $J^n = [t^n, t^{n+1})$, and $\Delta t^n = t^{n+1} - t^n$. Let $\Delta x = \max_{1 \leq i \leq n_x} \Delta x_i$, $\Delta y = \max_{1 \leq j \leq n_y} \Delta y_j$, $\Delta t = \max_{0 \leq n \leq n_T} \Delta t^n$, and $h = \max\{\Delta x, \Delta y\}$. We tacitly assume that each exterior edge has imposed on it either Dirichlet or Neumann conditions, but not both. Associated with these partitions, we introduce the spaces

$$Q_h = \{v \in H(\text{div}; \Omega) : v|_{I_i^x \times I_j^y} = (a_{i,j}^1 + a_{i,j}^2 x + a_{i,j}^3 y + a_{i,j}^4 xy), a_{i,j}^k \in \mathbb{R}, \\ i = 1, \dots, n_x, j = 1, \dots, n_y, v \cdot \nu|_{\partial\Omega_1} = 0\},$$

$$W_h = \{w \in L^\infty(\Omega) : w|_{I_i^x \times I_j^y} \in P^0(I_i^x \times I_j^y), i = 1, \dots, n_x, j = 1, \dots, n_y\},$$

$$W_{\Delta t} = \{w \text{ right continuous} : w|_{J^n} \in P^0(J^n), n = 0, \dots, n_T - 1\}.$$

If $v \in Q_h$, $v_{i+1/2,j}$ and $v_{i,j+1/2}$ denote $v(x_{i+1/2}, y_j)$ and $v(x_i, y_{j+1/2})$, respectively. If $w \in W_h$, then $w_{i,j}$ represents the constant value $w(x, y)$, $(x, y) \in I_i^x \times I_j^y$. w^n indicates the constant $w(t)$, $t \in J^n$, if $w \in W_{\Delta t}$. For notational and expositional convenience, let $\Delta x_0 = \Delta x_1$, $\Delta x_{n_x+1} = \Delta x_{n_x}$, $\Delta y_0 = \Delta y_1$, $\Delta y_{n_y+1} = \Delta y_{n_y}$, $\Delta x_{i+1/2} = (\Delta x_i + \Delta x_{i+1})/2$, $i = 1, \dots, n_x$, $\Delta y_{j+1/2} = (\Delta y_j + \Delta y_{j+1})/2$, $j = 1, \dots, n_y$, and $\Phi = 1$. Finally, define the notation $v^+ = \max\{v, 0\}$ and $v^- = \min\{v, 0\}$.

Let P_{Q_h} , P_{W_h} , and $P_{W_{\Delta t}}$ denote the L^2 -projections into Q_h , W_h , and $W_{\Delta t}$, respectively. To discretize (1.1), we first discretize the data as follows:

$$(2.1a) \quad u_{\text{init},h} = P_{W_h} u_{\text{init}},$$

$$(2.1b) \quad u_{D,\Delta t} = P_{W_{\Delta t}} u_D,$$

$$(2.1c) \quad V_h = P_{Q_h} V.$$

The subscript h is omitted below when no ambiguity occurs. Then the approximate solution $u_h \in W_{\Delta t} \otimes W_h$ is required to satisfy the equation, for $n = 0, \dots, n_T - 1$, $i = 1, \dots, n_x$, and $j = 1, \dots, n_y$:

$$(2.2a) \quad \frac{u_{i,j}^{n+1} - u_{i,j}^n}{\Delta t^n} + \frac{f_{1,i+1/2,j}^n - f_{1,i-1/2,j}^n}{\Delta x_i} + \frac{f_{2,i,j+1/2}^n - f_{2,i,j-1/2}^n}{\Delta y_j} \\ - \frac{D}{\Delta x_i} (q_{1,i+1/2,j}^n - q_{1,i-1/2,j}^n) - \frac{D}{\Delta y_j} (q_{2,i,j+1/2}^n - q_{2,i,j-1/2}^n) = -K_{i,j}^n u_{i,j}^n,$$

where

$$(2.2b) \quad f_{1,i-1/2,j}^n = u_{i-1,j}^n V_{1,i-1/2,j}^{n+} + u_{i,j}^n V_{1,i-1/2,j}^{n-},$$

$$(2.2c) \quad f_{2,i,j-1/2}^n = u_{i,j-1}^n V_{2,i,j-1/2}^{n+} + u_{i,j}^n V_{2,i,j-1/2}^{n-},$$

and the function $q_h = (q_1, q_2) \in W_{\Delta t} \otimes Q_h$ is the solution of

$$(q_h(t^n), v_h) = -(u_h(t^n), \operatorname{div} v_h) + \langle u_{D,\Delta t}, v_h \cdot \nu \rangle_{\partial\Omega_D}, \quad \forall v_h \in Q_h.$$

After the mass matrix has been mass-lumped [21], the expression for the degrees of freedom of q_h is taken as follows:

$$(2.2d) \quad q_{1,i-1/2,j}^n = (u_{i,j}^n - u_{i-1,j}^n) / \Delta x_{i-1/2},$$

$$(2.2e) \quad q_{2,i,j-1/2}^n = (u_{i,j}^n - u_{i,j-1}^n) / \Delta y_{j-1/2}.$$

Finally, the Neumann boundary condition (1.2a) is discretized by the usual reflection principle, and on $\partial\Omega_2$ u_h is defined by $u_{D,\Delta t}$.

Note that the lowest-order Raviart-Thomas mixed method [20] over rectangles has been used in (2.2a). Since the elements in Q_h have continuous normal components on interelement edges, the numerical fluxes $f_{1,i-1/2,j}^n$ and $f_{2,i,j-1/2}^n$ in (2.2b) and (2.2c) are well defined. Furthermore, if appropriate approximations of the coefficient V_h are introduced and the mass-lumping technique is used as in (2.2d) and (2.2e), the conservative scheme (2.2a) can be deduced from the discontinuous finite element method [6], [10] or from the finite volume method [19], [17] combined with the mixed finite element method [21]. Finally, the scheme applies to the case of $D = 0$.

The following approximation properties are used later [12], [20]:

$$(2.3a) \quad \|V_h^n\|_{L^\infty(\Omega)} \leq C_0 \|V^n\|_{L^\infty(\Omega)},$$

$$(2.3b) \quad \|\operatorname{div} V_h^n\|_{BV(\Omega)} \leq C_0 \|\operatorname{div} V^n\|_{BV(\Omega)},$$

$$(2.3c) \quad \|V^n - V_h^n\|_{L^\infty(\Omega)} \leq C_0 h \|V^n\|_{W^{1,\infty}(\Omega)},$$

for each n . Moreover, the operator P_{Q_h} is locally defined. □

3. Stability and convergence results. In this section we state and discuss the stability and convergence results of the scheme (2.2). Let $Q_T = T \times \Omega$. We assume that the data satisfy the following conditions:

$$(3.1a) \quad u_{\text{init}}, u_D \in [0, u^*],$$

$$(3.1b) \quad V, \nabla V \in L^\infty(Q_T),$$

$$(3.1c) \quad u_D \in L^\infty(J; BV(\partial\Omega_2)),$$

$$(3.1d) \quad u_D \in L^1(\partial\Omega_2; BV(J)),$$

$$(3.1e) \quad \operatorname{div} V \in L^\infty(J; BV(\Omega)),$$

$$(3.1f) \quad K \in [0, K^*].$$

For expositional convenience, let

$$V_1^* = C_0 \|V_1\|_{L^\infty(Q_T)}, \quad V_2^* = C_0 \|V_2\|_{L^\infty(Q_T)}, \quad V_D^* = C_0 \|\operatorname{div} V\|_{L^\infty(Q_T)}.$$

Theorem 3.1 (Stability). *Suppose that (3.1a), (3.1b), (3.1f), and for $n = 0, \dots, n_T - 1$ the following CFL condition are satisfied:*

$$(3.2) \quad \Delta t^n \leq \frac{1}{D_{ij}^* + 2V_1^*/\Delta x_i + 2V_2^*/\Delta y_j}, \quad i = 1, \dots, n_x, j = 1, \dots, n_y,$$

where $D_{ij}^* = \frac{D}{\Delta x_i} \left(\frac{1}{\Delta x_{i+1/2}} + \frac{1}{\Delta x_{i-1/2}} \right) + \frac{D}{\Delta y_j} \left(\frac{1}{\Delta y_{j+1/2}} + \frac{1}{\Delta y_{j-1/2}} \right) + K^*$. Then

$$(3.3) \quad 0 \leq u_h(t, x, y) \leq e^{tV_D^*} u^*, \quad (t, x, y) \in Q_T.$$

In addition, if

$$(3.4) \quad (\operatorname{div} V_h^n)_{ij} + K_{i,j}^n \geq 0,$$

then we have

$$(3.5) \quad 0 \leq u_h(t, x, y) \leq u^*, \quad (t, x, y) \in Q_T.$$

Obviously, since $K \geq 0$, (3.4) is satisfied if V_1 is nondecreasing in x and V_2 is nondecreasing in y , or $\operatorname{div} V$ is uniformly positive by the definition of P_{Q_h} [20].

Define, for $n = 0, \dots, n_T$,

$$\|u_h^n\|_{BV(\Omega)} = \sum_{i=0}^{n_x} \sum_{j=0}^{n_y} (|u_{i+1,j}^n - u_{i,j}^n| \Delta y_j + |u_{i,j+1}^n - u_{i,j}^n| \Delta x_i),$$

and set

$$\Delta x_\star = \min\{\Delta x_i, i = 1, \dots, n_x\}. \quad \Delta y_\star = \min\{\Delta y_j, j = 1, \dots, n_y\}.$$

Theorem 3.2 (TVB). *Assume that (3.1) and for $n = 0, \dots, n_T - 1$ the following CFL condition are satisfied:*

$$(3.6) \quad \Delta t^n \leq \frac{1}{D_{ij}^* + 3V_1^*/\Delta x_\star + 3V_2^*/\Delta y_\star}, \quad i = 1, \dots, n_x, j = 1, \dots, n_y.$$

Then there is a constant C_1 depending solely on the data and T such that

$$(3.7) \quad \|u_h\|_{L^\infty(J; BV(\Omega))} \leq C_1 \left\{ 1 + \|K\|_{L^\infty(J; BV(\Omega))} + D \left(\frac{1}{\Delta x_\star} + \frac{1}{\Delta y_\star} \right) \|u_D\|_{L^\infty(J; BV(\partial\Omega_2))} \right\}.$$

We remark that either in the case of $D = 0$ or in the case of u_D being constant in space, (3.7) shows that the total variation of the solution u_h is bounded.

In the following, v_h stands for the approximate solution of (1.1) and (1.2) with the data v_{init} and v_D satisfying the conditions (3.1a), (3.1c), and (3.1d).

Theorem 3.3 (Continuity with respect to data). *Assume that the hypotheses of Theorem 3.1 are satisfied for both sets of data. Then there exists a constant C_2 depending only on the data and T such that*

$$(3.8) \quad \|u_h - v_h\|_{L^\infty(J; L^1(\Omega))} \leq C_2 \left\{ \left(1 + D \left(\frac{1}{\Delta x_*} + \frac{1}{\Delta y_*} \right) \right) \right. \\ \left. \times \|u_D - v_D\|_{L^\infty(J; L^1(\partial\Omega_2))} + \|u_{\text{init}} - v_{\text{init}}\|_{L^1(\Omega)} \right\}.$$

As for the convergence result, we now consider a simple case where $D = 0$. In this case Theorem 3.2 implies the total variation boundedness of the numerical scheme as remarked above, which together with Theorem 3.3 yields the following convergence result (see §7). For nonzero D , concrete error estimates for the numerical scheme (2.2) will be obtained in the work mentioned earlier.

In the simple case the boundary condition (1.2a) and (1.2a) is replaced by the following inflow boundary condition:

$$(3.9) \quad u = u_D, \quad (x, y) \in \partial\Omega_-, \quad t \in J.$$

where $\partial\Omega_- = \{(x, y) \in \partial\Omega : (V \cdot \nu)(x, y) < 0\}$. We now extend the numerical flux introduced in (2.2) to the general setting:

$$f(u_{\text{left}}, u_{\text{right}}; \alpha) = u_{\text{left}}\alpha^+ + u_{\text{right}}\alpha^-.$$

Also, we define

$$\mathcal{C}_0^1([0, T] \times \bar{\Omega}) = \{\varphi \in \mathcal{C}^1(\bar{J} \times \bar{\Omega}) : \varphi(T, x, y) = 0, (x, y) \in \bar{\Omega}\}.$$

Then a weak solution of the differential equation given by (1.1) with $D = 0$, (3.9), and (1.2c) is defined to be a function $u \in L^\infty(J; BV(\Omega))$ satisfying the weak formulation:

$$(3.10) \quad (u, \varphi_t)_{Q_T} + (uV, \nabla\varphi)_{Q_T} + (u_{\text{init}}, \varphi)_{\{t=0\} \times \Omega} \\ - (f(u, u_D; V \cdot \nu), \varphi)_{J \times (\partial\Omega)} - (Ku, \varphi)_{Q_T} = 0, \quad \forall \varphi \in \mathcal{C}_0^1([0, T] \times \bar{\Omega}),$$

where $(\cdot, \cdot)_S$ denotes the inner product in $L^2(S)$ for some set S . Note that the role of the flux f is to select the correct boundary value for u , and that the smoothness hypothesis on V guarantees the uniqueness of weak solution to (3.10) [18].

Theorem 3.4 (Convergence). *Assume that the hypotheses of Theorem 3.2 are satisfied. Then the sequence $\{u_h\}_{h>0}$ produced by the scheme (2.2) converges in $L^\infty(J; L^1(\Omega))$ to the unique solution of (3.10). Moreover, $u \in L^\infty(J; BV(\Omega))$.*

4. Proof of the maximum principle. In this section we prove Theorem 3.1. Let

$$U^n = \max\{u_n^{i,j}, 0 \leq i \leq n_x + 1, 0 \leq j \leq n_y + 1\}.$$

Lemma 4.1. Suppose that

$$(4.1) \quad 1 - \frac{\Delta t^n}{\Delta x_i} (V_{n+}^{1,i+1/2,j} - V_{n-}^{1,i-1/2,j}) - \frac{\Delta t^n}{\Delta y_j} (V_{n+}^{2,i,j+1/2} - V_{n-}^{2,i,j-1/2}) - \frac{\Delta t^n}{D \Delta x_i} \left(\frac{\Delta x_{i+1/2}}{1} + \frac{\Delta x_{i-1/2}}{1} \right) \left(\frac{\Delta y_{j+1/2}}{1} + \frac{\Delta y_{j-1/2}}{1} \right) - K_{n,j}^{i,j} \Delta t^n \geq 0.$$

Then, if

$$(4.2) \quad 0 \leq u_n^{i,j}, \quad 0 \leq i \leq n_x + 1, 0 \leq j \leq n_y + 1,$$

we have, for $0 \leq i \leq n_x + 1$ and $0 \leq j \leq n_y + 1$,

$$(4.3) \quad 0 \leq u_{n+1}^{i,j} \leq U^{n+1} \leq 1 + \Delta t^n \max_{I_i^j \times I_j^i} \{ |\operatorname{div} V_n^h| \} \{ U^n \}.$$

In addition, if

$$(4.4) \quad 1 - \Delta t^n (\operatorname{div} V_n^h)^{i,j} - K_{n,j}^{i,j} \Delta t^n \leq 1,$$

we have

$$(4.5) \quad 0 \leq u_{n+1}^{i,j} \leq U^n, \quad 0 \leq i \leq n_x + 1, 0 \leq j \leq n_y + 1.$$

Proof. For $i = 1, \dots, n_x$ and $j = 1, \dots, n_y$, it follows from (2.2) that

$$u_{n+1}^{i,j} = A_n^{i+1,j} u_n^{i+1,j} + A_n^{i,j+1} u_n^{i,j+1} + B_n^{i,j} u_n^{i,j} + E_n^{i-1,j} u_n^{i-1,j} + E_n^{i,j-1} u_n^{i,j-1},$$

where

$$\begin{aligned} A_n^{i+1,j} &= \frac{\Delta t^n}{D \Delta x_i} V_{n-}^{1,i+1/2,j} + \frac{\Delta x_i}{D \Delta t^n} \Delta x_i \Delta x_{i+1/2}, \\ A_n^{i,j+1} &= \frac{\Delta t^n}{D \Delta y_j} V_{n-}^{2,i,j+1/2} + \frac{\Delta y_j}{D \Delta t^n} \Delta y_j \Delta y_{j+1/2}, \\ B_n^{i,j} &= 1 - \frac{\Delta t^n}{D \Delta x_i} \left(V_{n+}^{1,i+1/2,j} - V_{n-}^{1,i-1/2,j} \right) - \frac{\Delta t^n}{D \Delta y_j} \left(V_{n+}^{2,i,j+1/2} - V_{n-}^{2,i,j-1/2} \right) \\ &\quad - \frac{D \Delta t^n}{1} \frac{\Delta x_i}{1} \left(\frac{\Delta x_{i+1/2}}{1} + \frac{\Delta x_{i-1/2}}{1} \right) \left(\frac{\Delta y_{j+1/2}}{1} + \frac{\Delta y_{j-1/2}}{1} \right) - K_{n,j}^{i,j} \Delta t^n, \\ E_n^{i-1,j} &= \frac{\Delta t^n}{D \Delta x_i} V_{n+}^{1,i-1/2,j} + \frac{\Delta x_i}{D \Delta t^n} \Delta x_i \Delta x_{i-1/2}, \\ E_n^{i,j-1} &= \frac{\Delta t^n}{D \Delta y_j} V_{n+}^{2,i,j-1/2} + \frac{\Delta y_j}{D \Delta t^n} \Delta y_j \Delta y_{j-1/2}. \end{aligned}$$

Then, by (4.1), we see that

$$A_{i+1,j}^n, A_{i,j+1}^n, B_{i,j}^n, E_{i-1,j}^n, E_{i,j-1}^n \geq 0,$$

so that, by (4.2),

$$u_{i,j}^{n+1} \geq 0, \quad i = 1, \dots, n_x, \quad j = 1, \dots, n_y.$$

Furthermore, by the definition of Q_h and (4.2),

$$\begin{aligned} u_{i,j}^{n+1} &\leq \left(1 - \frac{\Delta t^n}{\Delta x_i} (V_{1,i+1/2,j}^n - V_{1,i-1/2,j}^n) - \frac{\Delta t^n}{\Delta y_j} (V_{2,i,j+1/2}^n - V_{2,i,j-1/2}^n) - K_{i,j}^n \Delta t^n \right) U^n \\ &= (1 - \Delta t^n (\operatorname{div} V_h^n)_{i,j} - K_{i,j}^n \Delta t^n) U^n, \end{aligned}$$

which implies (4.3) immediately since $K \geq 0$, and together with (4.4) yields (4.5). \square

Lemma 4.2. *If for $i = 1, \dots, n_x$ and $j = 1, \dots, n_y$*

$$(4.6) \quad \Delta t^n \leq \frac{1}{D_{i,j}^* + 2\|V_{h1}^n\|_{L^\infty(\Omega)}/\Delta x_i + 2\|V_{h2}^n\|_{L^\infty(\Omega)}/\Delta y_j},$$

where $D_{i,j}^*$ is defined as in Theorem 3.1, then (4.1) is satisfied.

The lemma follows obviously from the inequality (4.1) and the definition of $D_{i,j}^*$.

We are now ready to prove Theorem 3.1 by means of induction on n .

Proof of Theorem 3.1. For $n = 0$, the results (3.3) and (3.5) follow trivially from the assumption (3.1a). Let the results be true up to n . By Lemma 4.2 and (2.3a), (3.2) clearly implies (4.1). Then iterating (4.3) on n and using the induction hypothesis and (2.3b) yield that

$$0 \leq u_{i,j}^{n+1} \leq e^{t^{n+1} V_D^*} U^0, \quad i = 1, \dots, n_x, \quad j = 1, \dots, n_y.$$

Consequently, by (3.1a), (3.3) follows.

If (3.4) is true, so is (4.4). Then, in this case, it follows from (4.5) and the induction hypothesis that

$$0 \leq u_{i,j}^{n+1} \leq U^n \leq U^0, \quad i = 1, \dots, n_x, \quad j = 1, \dots, n_y,$$

which implies (3.5) by (3.1a). \square

5. Proof of total variation boundedness. In this section we prove Theorem 3.2. In order to fix ideas, let

$$\begin{aligned} \partial\Omega_2 &= \{(x, y) : x = 0, 0 < y < 1\} \cup \{(x, y) : x = 1, 0 < y < 1\}, \\ \partial\Omega_1 &= \partial\Omega \setminus \partial\Omega_2; \end{aligned}$$

other cases can be treated similarly.

Lemma 5.1. For $i = 0$ and $j = 1, \dots, n_y$,

$$\begin{aligned}
u_{1,j}^{n+1} - u_{0,j}^{n+1} &= u_{0,j}^n - u_{0,j}^{n+1} + \left(-\frac{\Delta t^n}{\Delta x_1} V_{1,3/2,j}^{n-} + \frac{D\Delta t^n}{\Delta x_{3/2}\Delta x_1} \right) (u_{2,j}^n - u_{1,j}^n) \\
&+ \left(-\frac{\Delta t^n}{\Delta y_j} V_{2,1,j+1/2}^{n-} + \frac{D\Delta t^n}{\Delta y_{j+1/2}\Delta y_j} \right) (u_{1,j+1}^n - u_{0,j+1}^n) \\
&+ \left\{ 1 - \frac{\Delta t^n}{\Delta x_1} V_{1,1/2,j}^{n+} + \frac{\Delta t^n}{\Delta y_j} V_{2,1,j+1/2}^{n-} - \frac{\Delta t^n}{\Delta y_j} V_{2,1,j-1/2}^{n+} \right. \\
&\quad \left. - \frac{D\Delta t^n}{\Delta x_0\Delta x_{1/2}} - \frac{D\Delta t^n}{\Delta y_j\Delta y_{j+1/2}} - \frac{D\Delta t^n}{\Delta y_j\Delta y_{j-1/2}} \right\} (u_{1,j}^n - u_{0,j}^n) \\
&+ \left(\frac{\Delta t^n}{\Delta y_j} V_{2,1,j-1/2}^{n+} + \frac{D\Delta t^n}{\Delta y_j\Delta y_{j-1/2}} \right) (u_{1,j-1}^n - u_{0,j-1}^n) \\
&- \left(\frac{\Delta t^n}{\Delta x_1} (V_{1,3/2,j}^n - V_{1,1/2,j}^n) + \frac{\Delta t^n}{\Delta y_j} (V_{2,1,j+1/2}^n - V_{2,1,j-1/2}^n) \right) u_{0,j}^n \\
&+ \left(-\frac{\Delta t^n}{\Delta y_j} V_{2,1,j+1/2}^{n-} + \frac{D\Delta t^n}{\Delta y_j\Delta y_{j+1/2}} \right) (u_{0,j+1}^n - u_{0,j}^n) \\
&- \left(\frac{\Delta t^n}{\Delta y_j} V_{2,1,j-1/2}^{n+} + \frac{D\Delta t^n}{\Delta y_j\Delta y_{j-1/2}} \right) (u_{0,j}^n - u_{0,j-1}^n) - K_{1,j}^n \Delta t^n u_{1,j}^n,
\end{aligned}$$

for $i = 1, \dots, n_x - 1$ and $j = 1, \dots, n_y$,

$$\begin{aligned}
u_{i+1,j}^{n+1} - u_{i,j}^{n+1} &= \left(-\frac{\Delta t^n}{\Delta x_{i+1}} V_{1,i+3/2,j}^{n-} + \frac{D\Delta t^n}{\Delta x_{i+1}\Delta x_{i+3/2}} \right) (u_{i+2,j}^n - u_{i+1,j}^n) \\
&+ \left(-\frac{\Delta t^n}{\Delta y_j} V_{2,i+1,j+1/2}^{n-} + \frac{D\Delta t^n}{\Delta y_j\Delta y_{j+1/2}} \right) (u_{i+1,j+1}^n - u_{i,j+1}^n) \\
&+ \left\{ 1 - \frac{\Delta t^n}{\Delta x_{i+1}} (V_{1,i+3/2,j}^n - V_{1,i+1/2,j}^n) - \frac{\Delta t^n}{\Delta y_j} (V_{2,i+1,j+1/2}^n - V_{2,i+1,j-1/2}^n) \right. \\
&\quad + \frac{\Delta t^n}{\Delta x_i} V_{1,i+1/2,j}^{n-} - \frac{\Delta t^n}{\Delta x_{i+1}} V_{1,i+1/2,j}^{n+} + \frac{\Delta t^n}{\Delta y_j} (V_{2,i+1,j+1/2}^{n-} - V_{2,i+1,j-1/2}^{n+}) \\
&\quad \left. - \frac{D\Delta t^n}{\Delta x_{i+1/2}} \left(\frac{1}{\Delta x_i} + \frac{1}{\Delta x_{i+1}} \right) - \frac{D\Delta t^n}{\Delta y_j} \left(\frac{1}{\Delta y_{j+1/2}} + \frac{1}{\Delta y_{j-1/2}} \right) \right. \\
&\quad \left. - K_{i+1,j}^n \Delta t^n \right\} (u_{i+1,j}^n - u_{i,j}^n) \\
&+ \left(\frac{\Delta t^n}{\Delta x_i} V_{1,i-1/2,j}^{n+} + \frac{D\Delta t^n}{\Delta x_i\Delta x_{i-1/2}} \right) (u_{i,j}^n - u_{i-1,j}^n) \\
&+ \left(\frac{\Delta t^n}{\Delta y_j} V_{2,i+1,j-1/2}^{n+} + \frac{D\Delta t^n}{\Delta y_j\Delta y_{j-1/2}} \right) (u_{i+1,j-1}^n - u_{i,j-1}^n)
\end{aligned}$$

$$\begin{aligned}
& - \frac{\Delta t^n}{\Delta y_j} \left(V_{2,i+1,j+1/2}^{n-} - V_{2,i,j+1/2}^{n-} \right) (u_{i,j+1}^n - u_{i,j}^n) \\
& + \frac{\Delta t^n}{\Delta y_j} \left(V_{2,i+1,j-1/2}^{n+} - V_{2,i,j-1/2}^{n+} \right) (u_{i,j-1}^n - u_{i,j}^n) \\
& - \Delta t^n \left((\operatorname{div} V_h^n)_{i+1,j} - (\operatorname{div} V_h^n)_{i,j} \right) u_{i,j}^n \\
& + \Delta t^n (K_{i,j}^n - K_{i+1,j}^n) u_{i,j}^n,
\end{aligned}$$

and for $i = n_x$ and $j = 1, \dots, n_y$,

$$\begin{aligned}
u_{n_x+1,j}^{n+1} - u_{n_x,j}^{n+1} &= u_{n_x+1,j}^{n+1} - u_{n_x+1,j}^n \\
& + \left\{ 1 + \frac{\Delta t^n}{\Delta x_{n_x}} V_{1,n_x+1/2,j}^{n-} + \frac{\Delta t^n}{\Delta y_j} V_{2,n_x,j+1/2}^{n-} - \frac{\Delta t^n}{\Delta y_j} V_{2,n_x,j-1/2}^{n+} \right. \\
& \quad \left. - \frac{D\Delta t^n}{\Delta x_{n_x} \Delta x_{n_x+1/2}} - \frac{D\Delta t^n}{\Delta y_j \Delta y_{j+1/2}} - \frac{D\Delta t^n}{\Delta y_j \Delta y_{j-1/2}} \right\} (u_{n_x+1,j}^n - u_{n_x,j}^n) \\
& + \left(\frac{\Delta t^n}{\Delta x_{n_x}} V_{1,n_x-1/2,j}^{n+} + \frac{D\Delta t^n}{\Delta x_{n_x} \Delta x_{n_x-1/2}} \right) (u_{n_x,j}^n - u_{n_x-1,j}^n) \\
& + \left(-\frac{\Delta t^n}{\Delta y_j} V_{2,n_x,j+1/2}^{n-} + \frac{D\Delta t^n}{\Delta y_j \Delta y_{j+1/2}} \right) (u_{n_x+1,j+1}^n - u_{n_x,j+1}^n) \\
& + \left(\frac{\Delta t^n}{\Delta y_j} V_{2,n_x,j-1/2}^{n+} + \frac{D\Delta t^n}{\Delta y_j \Delta y_{j-1/2}} \right) (u_{n_x+1,j-1}^n - u_{n_x,j-1}^n) \\
& + \left(\frac{\Delta t^n}{\Delta x_{n_x}} \left(V_{1,n_x+1/2,j}^n - V_{1,n_x-1/2,j}^n \right) \right. \\
& \quad \left. + \frac{\Delta t^n}{\Delta y_j} \left(V_{1,n_x,j+1/2}^n - V_{1,n_x,j-1/2}^n \right) \right) u_{n_x,j}^n \\
& - \left(-\frac{\Delta t^n}{\Delta y_j} V_{2,n_x,j+1/2}^{n-} + \frac{D\Delta t^n}{\Delta y_j \Delta y_{j+1/2}} \right) (u_{n_x+1,j+1}^n - u_{n_x+1,j}^n) \\
& + \left(\frac{\Delta t^n}{\Delta y_j} V_{2,n_x,j-1/2}^{n+} + \frac{D\Delta t^n}{\Delta y_j \Delta y_{j-1/2}} \right) (u_{n_x+1,j}^n - u_{n_x+1,j-1}^n) \\
& + K_{n_x,j}^n \Delta t^n u_{n_x,j}^n.
\end{aligned}$$

Similar expressions hold for $u_{i,j+1}^{n+1} - u_{i,j}^{n+1}$.

Proof. From (2.2), we see that

$$\begin{aligned}
u_{i,j}^{n+1} &= \left(-\frac{\Delta t^n}{\Delta x_i} V_{1,i+1/2,j}^{n-} + \frac{D\Delta t^n}{\Delta x_i \Delta x_{i+1/2}} \right) (u_{i+1,j}^n - u_{i,j}^n) \\
& + \left(-\frac{\Delta t^n}{\Delta y_j} V_{2,i,j+1/2}^{n-} + \frac{D\Delta t^n}{\Delta y_j \Delta y_{j+1/2}} \right) (u_{i,j+1}^n - u_{i,j}^n)
\end{aligned}$$

$$\begin{aligned}
& + \left(\frac{\Delta t^n}{\Delta x_i} V_{1,i-1/2,j}^{n+} + \frac{D\Delta t^n}{\Delta x_i \Delta x_{i-1/2}} \right) (u_{i-1,j}^n - u_{i,j}^n) \\
& + \left(\frac{\Delta t^n}{\Delta y_j} V_{2,i,j-1/2}^{n+} + \frac{D\Delta t^n}{\Delta y_j \Delta y_{j-1/2}} \right) (u_{i,j-1}^n - u_{i,j}^n) - K_{i,j}^n \Delta t^n u_{i,j}^n \\
& + \left(1 - \frac{\Delta t^n}{\Delta x_i} (V_{1,i+1/2,j}^n - V_{1,i-1/2,j}^n) - \frac{\Delta t^n}{\Delta y_j} (V_{2,i,j+1/2}^n - V_{2,i,j-1/2}^n) \right) u_{i,j}^n.
\end{aligned}$$

Then, the proof is completed by simple algebraic manipulations on $u_{i+1,j}^{n+1} - u_{i,j}^{n+1}$. \square

Lemma 5.2. *Assume that*

$$(5.1a) \quad 0 \leq u_{i,j}^n, \quad i = 0, \dots, n_x + 1, j = 0, \dots, n_y + 1,$$

and, for $i = 1, \dots, n_x$ and $j = 1, \dots, n_y$,

$$(5.1b) \quad \Delta t^n \leq \frac{1}{D_{i,j}^* + 3\|V_{h1}^n\|_{L^\infty(\Omega)}/\Delta x_* + 3\|V_{h2}^n\|_{L^\infty(\Omega)}/\Delta y_*}.$$

Then there is a constant $C_3 = C_3(C_0)$ such that

$$\begin{aligned}
\|u_h^{n+1}\|_{BV(\Omega)} & \leq (1 + C_3 \Delta t^n \|\nabla V^n\|_{L^\infty(\Omega)}) \|u_h^n\|_{BV(\Omega)} + \Delta t^n \|\operatorname{div} V_h^n\|_{BV(\Omega)} U^n \\
& + \Delta t^n C_3 (\|\nabla V^n\|_{L^\infty(\Omega)} + K^*) U^n + \Delta t^n \|K^n\|_{BV(\Omega)} U^n \\
& + 2\Delta t^n \sum_{j=0}^{n_y} \left(\|V_{h2}^n\|_{L^\infty(\Omega)} + \frac{D}{\Delta y_{j+1/2}} \right) |u_{0,j+1}^n - u_{0,j}^n| \\
& + 2\Delta t^n \sum_{j=0}^{n_y} \left(\|V_{h2}^n\|_{L^\infty(\Omega)} + \frac{D}{\Delta y_{j+1/2}} \right) |u_{n_x+1,j+1}^n - u_{n_x+1,j}^n| \\
& + \sum_{j=0}^{n_y} |u_{0,j}^{n+1} - u_{0,j}^n| \Delta y_j + \sum_{j=0}^{n_y} |u_{n_x+1,j}^{n+1} - u_{n_x+1,j}^n| \Delta y_j.
\end{aligned}$$

Proof. From (5.1b) we see that the coefficients of the terms between the brackets $\{\}$ in the expressions of Lemma 5.1 are nonnegative. Then simple algebraic manipulations and use of (2.3c) yield the desired result. \square

We are now in a position to prove Theorem 3.2.

Proof of Theorem 3.2. Note that the CFL condition (3.6) implies (5.1b) by (2.3a). Then the result (3.7) follows by iterating on n the inequality in Lemma 5.2 and using Theorem 3.1 and (2.3b). \square

6. Proof of continuity with respect to data. In this section we prove Theorem 3.3 and a result on equicontinuity in time of the approximate solution, Proposition 6.4 below. We recall that v_h stands for the solution of (2.2) with the data v_D and v_{init} .

Lemma 6.1. For $n = 0, \dots, n_T$, $i = 1, \dots, n_x$, and $j = 1, \dots, n_y$, we have

$$\begin{aligned} u_{i,j}^{n+1} - v_{i,j}^{n+1} &= A_{i+1,j}^n (u_{i+1,j}^n - v_{i+1,j}^n) + A_{i,j+1}^n (u_{i,j+1}^n - v_{i,j+1}^n) \\ &\quad + B_{i,j}^n (u_{i,j}^n - v_{i,j}^n) + E_{i-1,j}^n (u_{i-1,j}^n - v_{i-1,j}^n) \\ &\quad + E_{i,j-1}^n (u_{i,j-1}^n - v_{i,j-1}^n), \end{aligned}$$

where $A_{i+1,j}^n$, $A_{i,j+1}^n$, $B_{i,j}^n$, $E_{i-1,j}^n$, and $E_{i,j-1}^n$ are defined as in the proof of Lemma 4.1.

The result easily follows from (2.2).

Lemma 6.2. Supposed that (4.1) is satisfied. Then

$$\begin{aligned} \|u_h^{n+1} - v_h^{n+1}\|_{L^1(\Omega)} &\leq \Delta t^n \left(\|V_{h1}^n\|_{L^\infty(\Omega)} + \|V_{h2}^n\|_{L^\infty(\Omega)} + D \left(\frac{1}{\Delta x_\star} + \frac{1}{\Delta y_\star} \right) \right) \\ &\quad \times \|u_D - v_D\|_{L^1(\partial\Omega_2)} + \|u_h^n - v_h^n\|_{L^1(\Omega)}. \end{aligned}$$

Proof. Since, by (4.1), the coefficients in the equality of Lemma 6.1 are nonnegative, we see that

$$\begin{aligned} |u_{i,j}^{n+1} - v_{i,j}^{n+1}| &\leq A_{i+1,j}^n |u_{i+1,j}^n - v_{i+1,j}^n| + A_{i,j+1}^n |u_{i,j+1}^n - v_{i,j+1}^n| \\ &\quad + B_{i,j}^n |u_{i,j}^n - v_{i,j}^n| + E_{i-1,j}^n |u_{i-1,j}^n - v_{i-1,j}^n| \\ &\quad + E_{i,j-1}^n |u_{i,j-1}^n - v_{i,j-1}^n|. \end{aligned}$$

Then multiplying by $\Delta x_i \Delta y_j$, adding over i, j , and rearranging terms imply the desired result. \square

Now Theorem 3.3 can be easily seen from Lemma 6.2.

Lemma 6.3. Assume that the CFL condition (4.1) is satisfied. Then

$$\begin{aligned} \|u_h^1 - u_h^0\|_{L^1(\Omega)} &\leq 2\Delta t^0 \left(\|V_{h1}^0\|_{L^\infty(\Omega)} + \|V_{h2}^0\|_{L^\infty(\Omega)} + D \left(\frac{1}{\Delta x_\star} + \frac{1}{\Delta y_\star} \right) \right) \|u_{\text{init}}\|_{BV(\Omega)} \\ &\quad + \Delta t^0 \left(\|V_{h1}^0\|_{BV(\Omega)} + \|V_{h2}^0\|_{BV(\Omega)} + K^\star \right) u^\star. \end{aligned}$$

Proof. By (2.2), we observe that

$$\begin{aligned} |u_{i,j}^1 - u_{i,j}^0| &\leq \left(-\frac{\Delta t^0}{\Delta x_i} V_{1,i+1/2,j}^{0-} + \frac{D\Delta t^0}{\Delta x_i \Delta x_{i+1/2}} \right) |u_{i+1,j}^0 - u_{i,j}^0| \\ &\quad + \left(-\frac{\Delta t^0}{\Delta y_j} V_{2,i,j+1/2}^{0-} + \frac{D\Delta t^0}{\Delta y_j \Delta y_{j+1/2}} \right) |u_{i,j+1}^0 - u_{i,j}^0| \\ &\quad + \left(\frac{\Delta t^0}{\Delta x_i} V_{1,i-1/2,j}^{0+} + \frac{D\Delta t^0}{\Delta x_i \Delta x_{i-1/2}} \right) |u_{i-1,j}^0 - u_{i,j}^0| \\ &\quad + \left(\frac{\Delta t^0}{\Delta y_j} V_{2,i,j-1/2}^{0+} + \frac{D\Delta t^0}{\Delta y_j \Delta y_{j-1/2}} \right) |u_{i,j-1}^0 - u_{i,j}^0| + \Delta t^0 K_{i,j}^0 u_{i,j}^0 \\ &\quad + \Delta t^0 \left| \frac{1}{\Delta x_i} (V_{1,i+1/2,j}^0 - V_{1,i-1/2,j}^0) + \frac{1}{\Delta y_j} (V_{2,i,j+1/2}^0 - V_{2,i,j-1/2}^0) \right| u_{i,j}^0. \end{aligned}$$

Then the lemma follows by multiplying this inequality by $\Delta x_i \Delta y_j$ and adding the resulting one over i, j . \square

Proposition 6.4 (Equicontinuity in time). *Under the assumptions of Theorem 3.1, there is a constant C_4 depending only on the data and T such that for $n = 0, \dots, n_T$*

$$\begin{aligned} \|u_h^{n+1} - u_h^n\|_{L^1(\Omega)} &\leq C_4 \Delta t \left(1 + D \left(\frac{1}{\Delta x_\star} + \frac{1}{\Delta y_\star} \right) \right) \\ &\quad \times \left(\|u_{\text{init}}\|_{BV(\Omega)} + \|u_D\|_{L^1(\partial\Omega_2; BV(J))} \right). \end{aligned}$$

Proof. Take $v_h^{n+1} = u_h^n$ in Lemma 6.2 and use Lemma 6.3 to obtain the result. \square

7. A convergence analysis. In this section we prove Theorem 3.4 by applying the ideas used in [8] for analyzing the one-dimensional drift-diffusion semiconductor device equations. We point out that the analysis here is much simpler than that given in [8]. The reason is that we are here using the standard entropy $|\cdot|$, while a smoother entropy has been used there, which requires much work to estimate the distance between the smooth entropy and the standard one. We also emphasize the difference between the present analysis and that used in classical conservation laws; in the present case the delicate part is how to handle the boundary terms in the ‘entropy form’ Θ (see (7.3) below), while an unbounded domain is treated in the classical conservation laws.

The proof of Theorem 3.4 proceeds as follows. First, we prove that there is a subsequence $\{u_{h'}\}_{h'>0}$ converging to a limit u . Then, we show that

$$(7.1a) \quad \lim_{h' \rightarrow 0} \mathcal{R}(u_{h'}, \varphi) = \mathcal{R}(u, \varphi),$$

$$(7.1b) \quad \mathcal{R}(u, \varphi) = 0,$$

for $\varphi \in C_0^1([0, T] \times \bar{\Omega})$, where $\mathcal{R}(\cdot, \cdot)$ defines the left-hand side of (3.10). Since the weak solution of (3.10) is assumed unique, this completes the proof of Theorem 3.4.

As in classical conservation laws, (7.1) follows from the following result [11]: \blacktriangleright

$$(7.2a) \quad \lim_{h' \rightarrow 0} \Theta(u_{h'}, c; V_{h'}; \varphi) = \Theta(u, c; V; \varphi), \quad \forall c \in \mathbb{R}, \varphi \in C^1(\bar{Q}_T),$$

$$(7.2b) \quad \Theta(u, c; V; \varphi) \leq 0, \quad \forall c \in \mathbb{R}, 0 \leq \varphi \in C^1(\bar{Q}_T),$$

where Θ is defined in (7.3) below. Most part of this section is devoted to proving this result.

7.1. The entropy form. The entropy form $\Theta(u, c; V; \varphi)$ with boundary terms included is defined as follows:

$$\begin{aligned} (7.3) \quad \Theta(u, c; V; \varphi) &= -(|u - c|, \varphi_t)_{Q_T} - (|u - c|V, \nabla \varphi)_{Q_T} \\ &\quad + (|u - c|, \varphi)|_{\{t=T\} \times \Omega} - (|u_{\text{init}} - c|, \varphi)|_{\{t=0\} \times \Omega} \\ &\quad + (G(u - c, u_D - c; V \cdot \nu), \varphi)_{J \times (\partial\Omega)} \\ &\quad - (H(u, c)(\text{div } V + K), \varphi)_{Q_T} + (K|u - c|, \varphi)_{Q_T}, \end{aligned}$$

where $c \in \mathbb{R}$, $\varphi \in \mathcal{C}^1(\overline{Q}_T)$, and the ‘entropy flux’ G and the function H are defined by

$$\begin{aligned} G(u_{\text{left}}, u_{\text{right}}; V \cdot \nu) &= |u_{\text{left}}|(V \cdot \nu)^+ + |u_{\text{right}}|(V \cdot \nu)^-, \\ H(u, c) &= |u - c| - u \operatorname{sign}(u - c). \end{aligned}$$

The motivation of the form Θ can be given as in the one-dimensional case [8].

7.2. A convergent subsequence. In this subsection we prove the existence of a convergent subsequence $\{u_{h'}\}_{h'>0}$.

Lemma 7.1. *Assume that the hypotheses of Theorem 3.2 are satisfied. Then there exists a subsequence $\{u_{h'}\}_{h'>0}$ converging in $L^\infty(J; L^1(\Omega))$ to a limit u in $L^\infty(J; BV(\Omega)) \cap \mathcal{C}^0(J; L^1(\Omega))$.*

Proof. We note that the ideas in [11] can be used to prove the lemma. In [11], a discrete version of Azcoli-Arzelá Theorem was used. In the present case with $D = 0$, the equicontinuity in time is provided by Proposition 6.4. and the compactness of the range is given by Theorem 3.2. Also, the regularity result on u follows from the convergence and Theorem 3.2. \square

7.3. Proof of (7.2a). Here we prove (7.2a) under a condition.

Lemma 7.2. *Suppose that for $c \in \mathbb{R}$ and nonnegative $\varphi \in \mathcal{C}^1(\overline{Q}_T)$,*

$$(7.4) \quad \lim_{h' \rightarrow 0} \Theta(u_{h'}, c; V_{h'}; \varphi) \leq 0.$$

Then

$$(7.5) \quad \lim_{h' \rightarrow 0} \Theta(u_{h'}, c; V_{h'}; \varphi) = \Theta(u, c; V; \varphi).$$

Proof. First, for every nonnegative $\varphi \in \mathcal{C}_0^1([0, T] \times \Omega)$, (7.5) follows from Lemma 7.1 and the standard argument in the classical conservation laws [11]. Also, since $u \in \mathcal{C}^0(J; L^1(\Omega))$ by Lemma 7.1, the same result holds for $\varphi \in \mathcal{C}_0^1([0, T] \times \Omega)$.

We now consider the case where $\varphi \in \mathcal{C}_0^1(J \times [0, 1] \times (0, 1))$. Since we are mainly concerned with the boundary term associated with the edge $\{x = 0, 0 < y < 1\}$, it suffices to consider $\varphi(t, x, y)$ of this form $\omega(t, y)\xi(x)$. Then, set

$$\begin{aligned} g_{h'}(x) &= \int_0^T \int_0^1 |u_{h'} - c| V_{h'1} \omega(t, y) dy dt, \\ g_{h'}(0-) &= \int_0^T \int_0^1 \left\{ |u_{D, \Delta t} - c| V_{h'1}^+(t, 0, y) + |u_{h'}(t, 0+, y) - c| V_{h'1}^-(t, 0, y) \right\} \omega dt dy. \end{aligned}$$

and rewrite $\Theta(u_{h'}, c; V_{h'}; \varphi)$ as follows:

$$\begin{aligned} \Theta(u_{h'}, c; \varphi) &= - \left(|u_{h'} - c|, \xi \frac{\partial \omega}{\partial t} \right)_{Q_T} - \left(|u_{h'} - c| V_{h',2}, \xi \frac{\partial \omega}{\partial y} \right)_{Q_T} \\ &\quad - \int_0^1 g_{h'}(x) \xi'(x) dx - g_{h'}(0-) \xi(0) \\ &\quad - (H(u_{h'}, c)(\operatorname{div} V_{h'} + K), \xi \omega)_{Q_T} + (K|u_{h'} - c|, \xi \omega)_{Q_T}. \end{aligned}$$

Since the sequence $\{u_{h'}(\cdot, 0+, \cdot)\}_{h' > 0}$ is bounded in $L^\infty(J \times (0, 1))$ by Theorem 3.1, there is a subsequence $\{u_{h''}(\cdot, 0+, \cdot)\}_{h'' > 0}$ converging in $L^\infty(J \times (0, 1))$ -weak* to a limit \tilde{u} . Let $\gamma_{t,y}$ be the Young measure corresponding to \tilde{u} . Then, by Lemma 7.1 and (2.3a), we see that

$$\begin{aligned} (7.6) \quad \lim_{h'' \rightarrow 0} \Theta(u_{h''}, c; V_{h''}; \varphi) &= - \left(|u - c|, \xi \frac{\partial \omega}{\partial t} \right)_{Q_T} - \left(|u - c| V_2, \xi \frac{\partial \omega}{\partial y} \right)_{Q_T} \\ &\quad - \int_0^1 g(x) \xi'(x) dx - g_0 \xi(0) \\ &\quad - (H(u, c)(\operatorname{div} V + K), \xi \omega)_{Q_T} + (K|u - c|, \xi \omega)_{Q_T}, \end{aligned}$$

where

$$(7.7a) \quad g(x) = \int_0^T \int_0^1 |u - c| V_1 \omega(t, y) dy dt,$$

$$(7.7b) \quad g_0 = \int_0^T \int_0^1 \left\{ |u_D - c| V_1^+(t, 0, y) + w(t, y) V_1^-(t, 0, y) \right\} \omega dt dy,$$

where

$$(7.8a) \quad w(t, y) = \int_0^{u^{**}} |\lambda - c| d\gamma_{t,y}(\lambda),$$

$$(7.8b) \quad u^{**} = e^{TV_D^*} u^*.$$

Thus, to prove (7.5), it suffices to prove that $g_0 = g^*$ where

$$(7.9) \quad g^* = \int_0^T \int_0^1 \left\{ |u_D - c| V_1^+(t, 0, y) + |u(t, 0+, y) - c| V_1^-(t, 0, y) \right\} \omega dt dy.$$

Take ξ such that its support is contained in $[0, \epsilon]$. Then, by (7.4) and Theorem 3.1, it follows from (7.6) and (7.7) that

$$- \int_0^1 g(x) \xi'(x) dx - g_0 \xi(0) \leq C \epsilon \|\xi\|_{L^1(0,1)}.$$

Since ϵ is arbitrary, this inequality yields that

$$(7.10) \quad g(0+) - g_0 \leq 0.$$

Choose $c \in \mathbb{R}$ such that $|u - c| = \alpha(u - c)$ for some $\alpha \in \mathbb{R}$. Then, by (7.8) and the definition of $\gamma_{t,y}$,

$$w(t, y) = \alpha(\tilde{u} - c),$$

so that, by (7.7), (7.10) becomes

$$\alpha \left\{ \int_0^T \int_0^1 (u(t, 0+, y) - u_D) V_1^+(t, 0, y) \omega dy dt \right. \\ \left. + (u(t, 0+, y) - \tilde{u}) V_1^-(t, 0, y) \omega dy dt \right\} \leq 0.$$

Since the sign of α is arbitrary and this inequality is true for any nonnegative $\omega \in C_0^1(T \times (0, 1))$, we have

$$(7.11a) \quad V_1^+(t, 0, y)(u(t, 0+, y) - u_D) = 0, \quad \text{a.e. in } J \times (0, 1),$$

$$(7.11b) \quad V_1^-(t, 0, y)(u(t, 0+, y) - \tilde{u}) = 0, \quad \text{a.e. in } J \times (0, 1).$$

Finally, by (7.7)–(7.9) and (7.11), we see that

$$\begin{aligned} g(0+) - g_0 &= g^* - g_0 \\ &= \int_0^T \int_0^1 (|u(t, 0+, y) - c| - w(t, y)) V_1^-(t, 0, y) \omega dy dt \\ &= \int_0^T \int_0^1 (|\tilde{u}(t, y) - c| - w(t, y)) V_1^-(t, 0, y) \omega dy dt \\ &= \int_0^T \int_0^1 \left(\left| \int_0^{u^{**}} (\lambda - c) d\gamma_{t,y}(\lambda) \right| - w(t, y) \right) V_1^-(t, 0, y) \omega dy dt \\ &\geq \int_0^T \int_0^1 \left(\int_0^{u^{**}} |\lambda - c| d\gamma_{t,y}(\lambda) - w(t, y) \right) V_1^-(t, 0, y) \omega dy dt \\ &= 0. \end{aligned}$$

which together with (7.10) implies that $g_0 = g^*$. This completes the proof of the case where $\varphi \in C_0^1(J \times [0, 1) \times (0, 1))$. The same argument applies to the remaining three cases. \square

It is now clear that it suffices to prove (7.4). This is done in the next two subsections.

7.4. A discrete entropy inequality. The following discrete entropy inequality will be needed for obtaining an upper bound for $\Theta(u_{h'}, c; V_{h'}; \varphi)$.

Lemma 7.3. *Under the CFL condition (3.2), we have, for $c \in \mathbb{R}$,*

$$\begin{aligned} & |u_{i,j}^{n+1} - c| - (1 - K_{i,j}^n \Delta t^n) |u_{i,j}^n - c| + \frac{\Delta t^n}{\Delta x_i} \left(G_{i+1/2,j}^n - G_{i-1/2,j}^n \right) \\ & + \frac{\Delta t^n}{\Delta y_j} \left(G_{i,j+1/2}^n - G_{i,j-1/2}^n \right) - H(u_{i,j}^{n+1}, c) \{ (\operatorname{div} V_h^n)_{i,j} + K_{i,j}^n \} \Delta t^n \leq 0, \end{aligned}$$

where

$$\begin{aligned} G_{i+1/2,j}^n &= V_{1,i+1/2,j}^{n+} |u_{i,j}^n - c| + V_{1,i+1/2,j}^{n-} |u_{i+1,j}^n - c|, \\ G_{i,j+1/2}^n &= V_{2,i,j+1/2}^{n+} |u_{i,j}^n - c| + V_{2,i,j+1/2}^{n-} |u_{i,j+1}^n - c|. \end{aligned}$$

Proof. From (2.2) and the definition of the mixed finite element space Q_h , we have, for $c \in \mathbb{R}$,

$$\begin{aligned} u_{i,j}^{n+1} - c &= \left(-\frac{\Delta t^n}{\Delta x_i} V_{1,i+1/2,j}^{n-} \right) (u_{i+1,j}^n - c) + \left(-\frac{\Delta t^n}{\Delta y_j} V_{2,i,j+1/2}^{n-} \right) (u_{i,j+1}^n - c) \\ &+ \left\{ 1 - \frac{\Delta t^n}{\Delta x_i} (V_{1,i+1/2,j}^{n+} - V_{1,i-1/2,j}^{n-}) - \frac{\Delta t^n}{\Delta y_j} (V_{2,i,j+1/2}^{n+} - V_{2,i,j-1/2}^{n-}) \right. \\ &\quad \left. - K_{i,j}^n \Delta t^n \right\} (u_{i,j}^n - c) \\ &+ \left(\frac{\Delta t^n}{\Delta x_i} V_{1,i-1/2,j}^{n+} \right) (u_{i-1,j}^n - c) + \left(\frac{\Delta t^n}{\Delta y_j} V_{2,i,j-1/2}^{n+} \right) (u_{i,j-1}^n - c) \\ &- \Delta t^n [(\operatorname{div} V_h^n)_{i,j} + K_{i,j}^n] c. \end{aligned}$$

Note that the term between the brackets is nonnegative by (3.2). Thus the lemma follows by multiplying this expression by $\operatorname{sign}(u_{i,j}^{n+1} - c)$. \square

7.5. An upper bound of entropy form. In this section we obtain an upper bound for $\Theta(u_{h'}, c; V_{h'}; \varphi)$, which implies the inequality (7.4). We first have the following decomposition of $\Theta(u_{h'}, c; V_{h'}; \varphi)$.

Let, for $\varphi \in C^1(\overline{Q}_T)$,

$$\begin{aligned} \varphi_{i,j}^n &= \frac{1}{\Delta x_i} \frac{1}{\Delta y_j} \int_{x_{i-1/2}}^{x_{i+1/2}} \int_{y_{j-1/2}}^{y_{j+1/2}} \varphi(t^n, x, y) dx dy, \\ \varphi_{i,j}^{n+1/2} &= \frac{1}{\Delta t^n} \frac{1}{\Delta x_i} \frac{1}{\Delta y_j} \int_{t^n}^{t^{n+1}} \int_{x_{i-1/2}}^{x_{i+1/2}} \int_{y_{j-1/2}}^{y_{j+1/2}} \varphi(t, x, y) dt dx dy, \\ \varphi_{i+1/2,j}^{n+1/2} &= \frac{1}{\Delta t^n} \frac{1}{\Delta y_j} \int_{t^n}^{t^{n+1}} \int_{y_{j-1/2}}^{y_{j+1/2}} \varphi(t, x_{i+1/2}, y) dt dy, \\ \varphi_{i,j+1/2}^{n+1/2} &= \frac{1}{\Delta t^n} \frac{1}{\Delta x_i} \int_{t^n}^{t^{n+1}} \int_{y_{j-1/2}}^{y_{j+1/2}} \varphi(t, x, y_{j+1/2}) dt dx. \end{aligned}$$

Lemma 7.4. (decomposition of Θ). *We have*

$$\Theta(u_h, c; V_h'; \varphi) = \Theta_{ent}(u_h, c; V_h'; \varphi) + \Theta_{com}(u_h, c; V_h'; \varphi),$$

where (with arguments omitted)

$$\begin{aligned} \Theta_{ent} = & \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} \left\{ |u_{i,j}^{n+1} - c| - (1 - K_{i,j}^n \Delta t^n) |u_{i,j}^n - c| + \frac{\Delta t^n}{\Delta x_i} \left(G_{i+1/2,j}^n - G_{i-1/2,j}^n \right) \right. \\ & \left. + \frac{\Delta t^n}{\Delta y_j} \left(G_{i,j+1/2}^n - G_{i,j-1/2}^n \right) - H(u_{i,j}^{n+1}, c) [(\operatorname{div} V_h^n)_{i,j} + K_{i,j}^n] \Delta t^n \right\} \varphi_{i,j}^{n+1} \Delta x_i \Delta y_j, \end{aligned}$$

and

$$\begin{aligned} \Theta_{com} = & \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} \left\{ (|u_{i+1,j}^n - c| - |u_{i,j}^n - c|) (-V_{1,i+1/2,j}^{n-}) (\varphi_{i,j}^{n+1} - \varphi_{i+1/2,j}^{n+1/2}) \right. \\ & \left. + (|u_{i-1,j}^n - c| - |u_{i,j}^n - c|) (V_{1,i-1/2,j}^{n+}) (\varphi_{i,j}^{n+1} - \varphi_{i-1/2,j}^{n+1/2}) \right\} \Delta t^n \Delta y_j \\ & + \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} \left\{ (|u_{i,j+1}^n - c| - |u_{i,j}^n - c|) (-V_{2,i,j+1/2}^{n-}) (\varphi_{i,j}^{n+1} - \varphi_{i,j+1/2}^{n+1/2}) \right. \\ & \left. + (|u_{i,j-1}^n - c| - |u_{i,j}^n - c|) (V_{2,i,j-1/2}^{n+}) (\varphi_{i,j}^{n+1} - \varphi_{i,j-1/2}^{n+1/2}) \right\} \Delta t^n \Delta x_i \\ & - \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} (\operatorname{div} V_h^n)_{i,j} u_{i,j}^n \operatorname{sign}(u_{i,j}^n - c) (\varphi_{i,j}^{n+1} - \varphi_{i,j}^{n+1/2}) \Delta t^n \Delta x_i \Delta y_j \\ & + \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} K_{i,j}^n H(u_{i,j}^{n+1}, c) (\varphi_{i,j}^{n+1} - \varphi_{i,j}^{n+1/2}) \Delta t^n \Delta x_i \Delta y_j \\ & + \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} K_{i,j}^n |u_{i,j}^n - c| (\varphi_{i,j}^n - \varphi_{i,j}^{n+1}) \Delta t^n \Delta x_i \Delta y_j \\ & + \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} (\operatorname{div} V_h^n)_{i,j} (H(u_{i,j}^{n+1}, c) - H(u_{i,j}^n, c)) \varphi_{i,j}^{n+1} \Delta t^n \Delta x_i \Delta y_j \\ & + \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} K_{i,j}^n (H(u_{i,j}^{n+1}, c) - H(u_{i,j}^n, c)) \varphi_{i,j}^{n+1/2} \Delta t^n \Delta x_i \Delta y_j. \end{aligned}$$

Proof. From the definition of Θ and the fact that $\operatorname{div} V_h$ is piecewise constant, we have

$$\Theta = \Psi_t + \Psi_x + \Psi,$$

where

$$\begin{aligned}
\Psi_t &= - \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} |u_{i,j}^n - c| (\varphi_{i,j}^{n+1} - \varphi_{i,j}^n) \Delta x_i \Delta y_j + \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} |u_{i,j}^{n_T} - c| \varphi_{i,j}^{n_T} \Delta x_i \Delta y_j \\
&\quad - \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} |u_{i,j}^0 - c| \varphi_{i,j}^0 \Delta x_i \Delta y_j, \\
\Psi_x &= - \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} |u_{i,j}^n - c| \left(V_{1,i+1/2,j}^n \varphi_{i+1/2,j}^{n+1/2} - V_{1,i-1/2,j}^n \varphi_{i-1/2,j}^{n+1/2} \right) \Delta t^n \Delta y_j \\
&\quad - \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} |u_{i,j}^n - c| \left(V_{2,i,j+1/2}^n \varphi_{i,j+1/2}^{n+1/2} - V_{2,i,j-1/2}^n \varphi_{i,j-1/2}^{n+1/2} \right) \Delta t^n \Delta x_i \\
&\quad + \sum_{n=0}^{n_T-1} \sum_{j=1}^{n_y} G_{n_x+1/2,j}^n \varphi_{n_x+1/2,j}^{n+1/2} \Delta t^n \Delta y_j - \sum_{n=0}^{n_T-1} \sum_{j=1}^{n_y} G_{1/2,j}^n \varphi_{1/2,j}^{n+1/2} \Delta t^n \Delta y_j \\
&\quad + \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} G_{i,n_y+1/2}^n \varphi_{i,n_y+1/2}^{n+1/2} \Delta t^n \Delta x_i - \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} G_{i,1/2}^n \varphi_{i,1/2}^{n+1/2} \Delta t^n \Delta x_i, \\
\Psi &= - \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} (\operatorname{div} V_h^n)_{i,j} u_{i,j}^n \operatorname{sign}(u_{i,j}^n - c) \varphi_{i,j}^{n+1/2} \Delta t^n \Delta x_i \Delta y_j \\
&\quad - \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} K_{i,j}^n H(u_{i,j}^n, c) \varphi_{i,j}^n \Delta t^n \Delta x_i \Delta y_j \\
&\quad + \sum_{n=0}^{n_T-1} \sum_{i=1}^{n_x} \sum_{j=1}^{n_y} K_{i,j}^n |u_{i,j}^n - c| \varphi_{i,j}^n \Delta t^n \Delta x_i \Delta y_j.
\end{aligned}$$

Then simple algebraic manipulations yield the desired result. \square

Lemma 7.5. (Upper bound of Θ). *Suppose that the conditions of Theorem 3.2 are satisfied. Then there is a constant C_5 depending solely on the data and T such that*

$$\begin{aligned}
\Theta_{ent} &\leq 0, \\
\Theta_{com} &\leq C_5 (1 + |c|) \left\{ \Delta x \|\varphi_x\|_{L^\infty(Q_T)} + \Delta y \|\varphi_y\|_{L^\infty(Q_T)} \right. \\
&\quad \left. + \Delta t (\|\varphi_t\|_{L^\infty(Q_T)} + \|\varphi\|_{L^\infty(Q_T)}) \right\}.
\end{aligned}$$

Proof. The first inequality follows immediately from Lemmas 7.3 and 7.4. Also, observe

that

$$\begin{aligned} |\varphi_{i,j}^{n+1} - \varphi_{i+1/2,j}^{n+1/2}| &\leq \frac{1}{2} \Delta x_i \|\varphi_x\|_{L^\infty(J \times \Omega)} + \frac{1}{2} \Delta t^n \|\varphi_t\|_{L^\infty(J \times \Omega)}, \\ |\varphi_{i,j}^{n+1} - \varphi_{i,j+1/2}^{n+1/2}| &\leq \frac{1}{2} \Delta y_j \|\varphi_y\|_{L^\infty(J \times \Omega)} + \frac{1}{2} \Delta t^n \|\varphi_t\|_{L^\infty(J \times \Omega)}, \\ |\varphi_{i,j}^{n+1} - \varphi_{i,j}^{n+1/2}| &\leq \frac{1}{2} \Delta t^n \|\varphi_t\|_{L^\infty(J \times \Omega)}, \\ |\varphi_{i,j}^{n+1} - \varphi_{i,j}^n| &\leq \Delta t^n \|\varphi_t\|_{L^\infty(J \times \Omega)}. \end{aligned}$$

Then, if an integration by parts on n is applied to the last two terms in the expression of Θ_{com} , the second inequality follows from Theorems 3.1 and 3.2, Proposition 6.4, Lemma 7.4, and (2.3a). \square

We are now in a position to prove Theorem 3.4.

Proof of Theorem 3.4. From Lemma 7.1 there exists a subsequence $\{u_{h'}\}_{h'>0}$ converging in $L^\infty(J; L^1(\Omega))$ to a limit u . Now, by Lemma 7.5, we have

$$\lim_{h' \rightarrow 0} \Theta(u_{h'}, c; V_{h'}; \varphi) \leq 0,$$

for every $c \in \mathbb{R}$ and nonnegative $\varphi \in \mathcal{C}^1(\overline{Q}_T)$. Thus, by Lemma 7.2, we see that

$$\lim_{h' \rightarrow 0} \Theta(u_{h'}, c; V_{h'}; \varphi) = \Theta(u, c; V; \varphi) \leq 0.$$

This implies that u is the unique solution of (3.10). Consequently, the whole sequence $\{u_h\}_{h>0}$ converges to u , and thus Theorem 3.4 is proven. \square

8. Numerical results. This section reports on numerical results with the finite element method (2.2) for three problems. They are designed to show the performance of the method and to indicate the convergence properties. In all examples the CFL condition (3.6) is required to satisfy.

Example 1. In this example we consider a convecting Gaussian hill in one space dimension. Specifically, we solve equation (1.1) with $\Phi = 1$, $V = 10$, $D = 0.1$, and $K = 0$ on the interval $[0, 6]$. The initial datum u_{init} is given by

$$u_{\text{init}}(x) = e^{-\pi x^2}.$$

As a pure initial-value problem, this leads to the analytical solution

$$u_a(t, x) = \frac{1}{\sqrt{1 + 4\pi Dt}} e^{-\frac{\pi(x - Vt)^2}{1 + 4\pi Dt}}.$$

We obtain an initial-boundary-value problem with the same solution by imposing the Dirichlet boundary condition:

$$u(t, 0) = u_a(t, 0), \quad u(t, 6) = u_a(t, 6).$$

In Figure 1 we display the analytical solution u_a and the approximate solution u_h at time $T = 0.25$. In Table 1 we display the errors and their respective order of convergence at the same time. From the table we see that the scheme is first-order accurate both in L^1 and in L^∞ for the concentration. This shows that the scheme (2.2) is first-order accurate in both spaces when the solution of the differential equation is smooth. Also, Figure 1 justifies the stability property given in Theorem 3.1.

Example 2. In this example we consider a problem whose solution displays a discontinuity. The problem has the data: $\Phi = 1$, $V = -0.5$, and $K = 0$. The boundary and initial conditions are given by

$$\begin{aligned} u(t, 0) &= 0, & u(t, 1) &= 1, & t > 0, \\ u_{\text{init}}(x, 0) &= 0. & x &\in [0, 1]. \end{aligned}$$

The exact and approximate ‘nonviscous’ solution (i.e., in the case of $D = 0$) and the ‘viscous’ solution with $D = 10^{-3}$ at $T = 0.5$ are displayed in Figure 2. Notice that the biggest error in the approximation of u occurs around the location of the discontinuity $x = 0.75$. In Table 2 we show the errors and their convergence orders. Note that the orders of convergence in L^1 and L^∞ are nearly $1/2$. This implies that the presence of discontinuity has an effect on the convergence. Finally, from Figure 2, we see that the ‘nonviscous’ solution provides a good approximation to the ‘viscous’ solution.

Example 3. In the final example we consider a two-dimensional problem which has a shock. The data are set as follows: $\Omega = (0, 1)^2$, $\Phi = 1$, $V = (\cos(\frac{8}{\pi}), \sin(\frac{8}{\pi}))$, $k = 0$, and $D = 10^{-3}$. The Neumann and Dirichlet boundaries $\partial\Omega_1$ and $\partial\Omega_2$ are defined by

$$\begin{aligned} \partial\Omega_1 &= \{(x, y) : 0 \leq x \leq 1, y = 1\}, \\ \partial\Omega_2 &= \partial\Omega \setminus \partial\Omega_1, \end{aligned}$$

and the boundary and initial data by

$$\begin{aligned} u_D &= \begin{cases} 1, & x = 0, \frac{1}{2} < y < 1, \\ 0, & \text{elsewhere,} \end{cases} \\ u_{\text{init}} &= \begin{cases} 1, & 0 \leq x \leq 1, \frac{1}{2} < y \leq 1, \\ 0, & \text{elsewhere.} \end{cases} \end{aligned}$$

The approximate solution of this problem obtained using the method (2.2) with $\Delta x = \Delta y = 10^{-2}$ at time $T = 2$ is shown in Figure 3. The graph clearly shows that the method can capture the shock around the location $y = 1/2$.




FIG. 1. The ‘—’, ‘---’, and ‘...’ represent the exact u and approximate solution u_h with $h = .001$ and $h = .01$.

$1/\Delta x$	L^∞ -error ($\times 10^2$)	L^∞ -order	L^1 -error ($\times 10^2$)	L^1 -order
50	6.09	-	5.10	-
100	3.57	0.76	2.50	1.03
200	1.89	0.91	1.20	0.99
400	0.99	0.93	0.62	1.02
800	0.51	0.98	0.29	1.09

Table 1. Convergence of u_h in $(0, 6)$ at $T = .25$.




FIG. 2. The '—', '- - -', and 'xxx' represent u with $D = 0$ and the approximate solution u_h with $D = .0001$ and $D_{\text{ex}} = 0$.

$1/\Delta x$	L^∞ -error ($\times 10^2$)	L^∞ -order	L^1 -error ($\times 10^2$)	L^1 -order
50	8.65	-	9.57	-
100	6.15	0.48	8.09	0.26
200	5.02	0.29	6.32	0.37
400	3.76	0.43	4.36	0.54
800	2.49	0.60	2.77	0.65

Table 2. Convergence of u_h in $(0, 1)$ at $T = .5$.




FIG. 3. The approximate solution u_h on $(0, 1)^2$.

9. A concluding remark. A new finite element method for numerically solving the two-dimensional convection-dominated transport equation in ground water has been formulated and analyzed in this paper. The primary computational advantage of the method is that it is local and thus fully parallelizable, and is conservative. The stability properties of this method and its convergence in a suitable topology have been established. Moreover, the numerical results have shown that the method is first-order accurate when the solution is smooth and is one-half accurate when the solution has discontinuities, and that the method is non-oscillatory and shock-capturing. Future work will be devoted to obtaining error estimates for both cases of zero and nonzero coefficient D .

REFERENCES

1. M. B. Allen III, G. A. Behie, and J. A. Trangenstein, *Multiphase Flow in Porous Media. Lecture Notes in Engineering*. Springer-Verlag, 1988.
2. J. W. Barrett and K. W. Morton, *Approximate symmetrization and Petrov-Galerkin methods for diffusion-convection problems*, *Comp. Meth. Appl. Mech. Engng.* **45** (1984), 97-122.
3. Z. Chen and B. Cockburn, *Error estimates for a finite element method for the drift-diffusion semiconductor device equations*, to appear in *SIAM J. Numer. Anal.*
4. Z. Chen, *Finite element analysis of the 1D full drift diffusion semiconductor model*, to appear in *SIAM J. Numer. Anal.*
5. Z. Chen and B. Cockburn, *Convergence of a finite element method for the drift-diffusion semiconductor device equations: the multidimensional case*, to appear in *Numer. Math.*

6. Z. Chen and J. Douglas, Jr., *Approximation of coefficients in hybrid and mixed methods for nonlinear parabolic problems*, *Mat. Applic. Comp.* **10** (1991), 137–160.
7. I. Christie, D. F. Griffiths, and A. R. Mitchell, *Finite element methods for second order differential equations with significant first derivatives*, *Int. J. Num. Engrg.* **10** (1976), 1389–1396.
8. B. Cockburn and I. Triandaf, *Convergence of a finite element method for the drift diffusion semiconductor device equations*, *Math. Comp.* **59** (1992), 383–401.
9. B. Cockburn and I. Triandaf, *Error estimates for a finite element method for the drift diffusion semiconductor device equations: the zero-diffusion case*, to appear in *Math. Comp.*
10. B. Cockburn and C.W. Shu, *TVB Runge-Kutta local projection discontinuous Galerkin finite element method for scalar conservation laws II: general framework*, *Math. Comp.* **52** (1989), 411–435.
11. M. Crandall and A. Majda, *Monotone difference approximations for scalar conservation laws*, *Math. Comp.* **34** (1980), 1–21.
12. J. Douglas, Jr. and J. Roberts, *Global estimates for mixed methods for second order elliptic problems*, *Math. Comp.* **45** (1985), 39–52.
13. J. Douglas, Jr. and T. F. Russell, *Numerical methods for convection-dominated diffusion problems based on combining the method of characteristics with finite element or finite difference procedures*, *SIAM J. Num. Anal.* **19** (1982), 871–885.
14. R. E. Ewing, ed., *Research Frontiers in Applied Mathematics. Vol. 1*. SIAM, Philadelphia, 1983.
15. R. E. Ewing and T. F. Russell, *Multistep Galerkin methods along characteristics for convection-diffusion problems*. *Advances in Computer Methods for PDEs IV*, Vichnevetsky and Stepleman, (eds.) IMACS, Rutgers Univ. (1981). 28–36.
16. R. E. Ewing, T. F. Russell, and M. F. Wheeler, *Convergence analysis of an approximation of miscible displacement in porous media by mixed finite elements and a modified method of characteristics*, *Comp. Meth. Appl. Mech. Engrg.* **47** (1984), 73–92.
17. R. D. Lazarov, I. D. Mishev, and P. S. Vassilevski, *Finite volume methods with local refinement for convection-diffusion problems*, Technical Report 92-50, Department of Mathematics. UCLA, Los Angeles, 1992.
18. P. A. Markowich and P. Szmolyan, *A system of convection-diffusion equations with small diffusion coefficient arising in semiconductor physics*, *J. Diff. Equ.* **81** (1989), 234–254.
19. S. McCormick, *Multilevel Adaptive Methods for Partial Differential Equations*, SIAM, Philadelphia, 1989.
20. P.A. Raviart and J.M. Thomas, *A mixed finite element method for second order elliptic problems*. *Lecture Notes in Mathematics*, Springer-Verlag, 1977.
21. T. Russell and M. Wheeler, *Finite element and finite difference methods for continuous flows in porous media*, Chapter II, *The Mathematics of Reservoir Simulation*. R. Ewing, ed., *Frontiers in Applied Mathematics 1*, Society for Industrial and Applied Mathematics, Philadelphia, 1983. pp. 35–106.

DEPARTMENT OF MATHEMATICS AND THE INSTITUTE FOR SCIENTIFIC COMPUTATION, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843.

E-mail address: zchen@isc.tamu.edu, ewing@ewing.tamu.edu

Recent IMA Preprints

#	Author/s	Title
1189	Frank H. Shaw & Charles J. Geyer,	Constrained covariance component models
1190	Tomasz Luczaka,	A greedy algorithm estimating the height of random trees
1191	Timo Seppäläinen,	Maximum entropy principles for disordered spins
1192	Yuandan Lin, Eduardo D. Sontag & Yuan Wang,	Recent results on Lyapunov-theoretic techniques for nonlinear stability
1193	Svante Janson,	Random regular graphs: Asymptotic distributions and contiguity
1194	Rachid Ababou,	Random porous media flow on large 3-D grids: Numerics, performance, & application to homogenization
1195	Moshe Fridman,	Hidden Markov model regression
1196	Petr Klouček, Bo Li & Mitchell Luskin,	Analysis of a class of nonconforming finite elements for Crystalline microstructures
1197	Steven P. Lalley,	Random series in inverse Pisot powers
1198	Rudy Yaksick,	Expected optimal exercise time of a perpetual American option: A closed-form solution
1199	Rudy Yaksick,	Valuation of an American put catastrophe insurance futures option: A Martingale approach
1200	János Pach, Farhad Shahrokhi & Mario Szegedy,	Application of the crossing number
1201	Avner Friedman & Chaocheng Huang,	Averaged motion of charged particles under their self-induced electric field
1202	Joel Spencer,	The Erdős-Hanani conjecture via Talagrand's inequality
1203	Zhangxin Chen,	Superconvergence results for Galerkin methods for wave propagation in various porous media
1204	Russell Lyons, Robin Pemantle & Yuval Peres,	When does a branching process grow like its mean? Conceptual proofs of $L \log L$ criteria
1205	Robin Pemantle,	Maximum variation of total risk
1206	Robin Pemantle & Yuval Peres,	Galton-Watson trees with the same mean have the same polar sets
1207	Robin Pemantle,	A shuffle that mixes sets of any fixed size much faster than it mixes the whole deck
1208	Itai Benjamini, Robin Pemantle & Yuval Peres,	Martin capacity for Markov chains and random walks in varying dimensions
1209	Włodzimierz Bryc & Amir Dembo,	On large deviations of empirical measures for stationary Gaussian processes
1210	Martin Hildebrand,	Some random processes related to affine random walks
1211	Alexander E. Mazel & Yurii M. Suhov,	Ground states of a Boson quantum lattice model
1212	Roger L. Fosdick & Darren E. Mason,	Single phase energy minimizers for materials with nonlocal spatial dependence
1213	Bruce Hajek,	Load balancing in infinite networks
1214	Petr Klouček,	The transonic flow problems stability analysis and numerical results
1215	Petr Klouček,	On the existence of the entropic solutions for the transonic flow problem
1216	David A. Schmidt & Chjan C. Lim,	Full sign-invertibility and symplectic matrices
1217	Piermarco Cannarsa & Maria Elisabetta Tessitore,	Infinite dimensional Hamilton-Jacobi equations and Dirichlet boundary control problems of parabolic type
1218	Zhangxin Chen,	Multigrid algorithms for mixed methods for second order elliptic problems
1219	Zhangxin Chen,	Expanded mixed finite element methods for linear second order elliptic problems I
1220	Gang Bao,	A note on the uniqueness for an inverse diffraction problem
1221	Moshe Fridman,	A two state capital asset pricing model
1222	Paolo Baldi,	Exact asymptotics for the probability of exit from a domain and applications to simulation
1223	Carl Dou & Martin Hildebrand,	Enumeration and random random walks on finite groups
1224	Jaksa Cvitanic & Ioannis Karatzas,	On portfolio optimization under "drawdown" constraints
1225	Avner Friedman & Yong Liu,	A free boundary problem arising in magnetohydrodynamic system
1226	Dominic Welsh,	Randomised approximation schemes for Tutte-Gröthendieck invariants
1227	Zhangxin Chen, Bernardo Cockburn, Carl L. Gardner, & Joseph W. Jerome,	Quantum hydrodynamic simulation of hysteresis in the resonant tunneling diode
1228	E.G. Kalnins, G.C. Williams, & Willard Miller, Jr.,	Intrinsic characterisation of the separation constant for spin one and gravitational perturbations in Kerr geometry
1229	Zhangxin Chen,	Large-scale averaging analysis of multiphase flow in fractured reservoirs
1230	Bruce Hajek & Babu Narayanan,	Multigraphs with the most edge covers
1231	K.B. Athreya,	Entropy maximization
1232	F.I. Karpelevich & Yu.M. Suhov,	Functional equations in the problem of boundedness of stochastic branching dynamics
1233	E. Dibenedetto & V. Vespri,	On the singular equation $\beta(u)_t = \Delta u$
1234	M.Ya. Kelbert & Yu.M. Suhov,	The Markov branching random walk and systems of reaction-diffusion (Kolmogorov-Petrovskii-Piskunov) equations
1235	M. Hildebrand,	Random walks on random regular simple graphs

- 1236 **W.S. Don & A. Solomonoff**, Accuracy enhancement for higher derivatives using Chebyshev collocation and a mapping technique
- 1237 **D. Gurarie**, Symmetries and conservation laws of two-dimensional hydrodynamics
- 1238 **Z. Chen**, Finite element methods for the black oil model in petroleum reservoirs
- 1239 **G. Bao & A. Friedman**, Inverse problems for scattering by periodic structure
- 1240 **G. Bao**, Some inverse problems in partial differential equations
- 1241 **G. Bao**, Diffractive optics in periodic structures: The TM polarization
- 1242 **C.C. Lim & D.A. Schmidt**, On noneven digraphs and symplectic pairs
- 1243 **H.M. Soner, S.E. Shreve & J. Cvitanić**, There is no nontrivial hedging portfolio for option pricing with transaction costs
- 1244 **D.L. Russell & B-Yu Zhang**, Exact controllability and stabilizability of the Korteweg-de Vries equation
- 1245 **B. Morton, D. Enns & B-Yu Zhang**, Stability of dynamic inversion control laws applied to nonlinear aircraft pitch-axis models
- 1246 **S. Hansen & G. Weiss**, New results on the operator Carleson measure criterion
- 1247 **V.A. Malyshev & F.M. Spiessma**, Intrinsic convergence rate of countable Markov chains
- 1248 **G. Bao, D.C. Dobson & J.A. Cox**, Mathematical studies in rigorous grating theory
- 1249 **G. Bao & W.W. Symes**, On the sensitivity of solutions of hyperbolic equations to the coefficients
- 1250 **D.A. Huntley & S.H. Davis**, Oscillatory and cellular mode coupling in rapid directional solidification
- 1251 **M.J. Donahue, L. Gurvits, C. Darken & E. Sontag**, Rates of convex approximation in non-Hilbert spaces
- 1252 **A. Friedman & B. Hu**, A Stefan problem for multi-dimensional reaction diffusion systems
- 1253 **J.L. Bona & B-Y. Zhang**, The initial-value problem for the forced Korteweg-de Vries equation
- 1254 **A. Friedman & R. Gulliver**, **Organizers**, Mathematical modeling for instructors
- 1255 **S. Kichenassamy**, The prolongation formula for tensor fields
- 1256 **S. Kichenassamy**, Fuchsian equations in Sobolev spaces and blow-up
- 1257 **H.S. Dumas, L. Dumas, & F. Golse**, On the mean free path for a periodic array of spherical obstacles
- 1258 **C. Liu**, Global estimates for solutions of partial differential equations
- 1259 **C. Liu**, Exponentially growing solutions for inverse problems in PDE
- 1260 **Mary Ann Horn & I. Lasiecka**, Nonlinear boundary stabilization of parallelly connected Kirchhoff plates
- 1261 **B. Cockburn & H. Gau**, A posteriori error estimates for general numerical methods for scalar conservation laws
- 1262 **B. Cockburn & P-A. Gremaud**, A priori error estimates for numerical methods for scalar conservation laws. Part I: The general approach
- 1263 **R. Spigler & M. Vianello**, Convergence analysis of the semi-implicit euler method for abstract evolution equations
- 1264 **R. Spigler & M. Vianello**, WKB-type approximation for second-order differential equations in C^* -algebras
- 1265 **M. Menshikov & R.J. Williams**, Passage-time moments for continuous non-negative stochastic processes and applications
- 1266 **C. Mazza**, On the storage capacity of nonlinear neural networks
- 1267 **Z. Chen, R.E. Ewing & R. Lazarov**, Domain decomposition algorithms for mixed methods for second order elliptic problems
- 1268 **Z. Chen, M. Espedal & R.E. Ewing**, Finite element analysis of multiphase flow in groundwater hydrology
- 1269 **Z. Chen, R.E. Ewing, Y.A. Kuznetsov, R.D. Lazarov & S. Maliassov**, Multilevel preconditioners for mixed methods for second order elliptic problems
- 1270 **S. Kichenassamy & G.K. Srinivasan**, The structure of WTC expansions and applications
- 1271 **A. Zinger**, Positiveness of Wigner quasi-probability density and characterization of Gaussian distribution
- 1272 **V. Malkin & G. Papanicolaou**, On self-focusing of short laser pulses
- 1273 **J.N. Kutz & W.L. Kath**, Stability of pulses in nonlinear optical fibers using phase-sensitive amplifiers
- 1274 **S.K. Patch**, Recursive recovery of a family of Markov transition probabilities from boundary value data
- 1275 **C. Liu**, The completeness of plane waves
- 1276 **Z. Chen & R.E. Ewing**, Stability and convergence of a finite element method for reactive transport in ground water
- 1277 **Z. Chen & Do Y. Kwak**, The analysis of multigrid algorithms for nonconforming and mixed methods for second order elliptic problems
- 1278 **Z. Chen**, Expanded mixed finite element methods for quasilinear second order elliptic problems II
- 1279 **M.A. Horn & W. Littman**, Boundary control of a Schrödinger equation with nonconstant principal part
- 1280 **R.E. Ewing, Yu.A. Kuznetsov, R.D. Lazarov & S. Maliassov**, Substructuring preconditioning for finite element approximations of second order elliptic problems. I. Nonconforming linear elements for the Poisson equation in a parallelepiped
- 1281 **S. Maliassov**, Substructuring preconditioning for finite element approximations of second order elliptic problems. II. Mixed method for an elliptic operator with scalar tensor
- 1282 **V. Jakšić & C.-A. Pillet**, On model for quantum friction II. Fermi's golden rule and dynamics at positive temperatures
- 1283 **V. M. Malkin**, Kolmogorov and nonstationary spectra of optical turbulence

Recent IMA Preprints

- | # | Author/s | Title |
|------|--|---|
| 1189 | Frank H. Shaw & Charles J. Geyer, | Constrained covariance component models |
| 1190 | Tomasz Luczaka, | A greedy algorithm estimating the height of random trees |
| 1191 | Timo Seppäläinen, | Maximum entropy principles for disordered spins |
| 1192 | Yuandan Lin, Eduardo D. Sontag & Yuan Wang, | Recent results on Lyapunov-theoretic techniques for nonlinear stability |
| 1193 | Svante Janson, | Random regular graphs: Asymptotic distributions and contiguity |
| 1194 | Rachid Ababou, | Random porous media flow on large 3-D grids: Numerics, performance, & application to homogenization |
| 1195 | Moshe Fridman, | Hidden Markov model regression |
| 1196 | Petr Klouček, Bo Li & Mitchell Luskin, | Analysis of a class of nonconforming finite elements for Crystalline microstructures |
| 1197 | Steven P. Lalley, | Random series in inverse Pisot powers |
| 1198 | Rudy Yaksick, | Expected optimal exercise time of a perpetual American option: A closed-form solution |
| 1199 | Rudy Yaksick, | Valuation of an American put catastrophe insurance futures option: A Martingale approach |
| 1200 | János Pach, Farhad Shahrokhi & Mario Szegedy, | Application of the crossing number |
| 1201 | Avner Friedman & Chaocheng Huang, | Averaged motion of charged particles under their self-induced electric field |
| 1202 | Joel Spencer, | The Erdős-Hanani conjecture via Talagrand's inequality |
| 1203 | Zhangxin Chen, | Superconvergence results for Galerkin methods for wave propagation in various porous media |
| 1204 | Russell Lyons, Robin Pemantle & Yuval Peres, | When does a branching process grow like its mean? Conceptual proofs of $L \log L$ criteria |
| 1205 | Robin Pemantle, | Maximum variation of total risk |
| 1206 | Robin Pemantle & Yuval Peres, | Galton-Watson trees with the same mean have the same polar sets |
| 1207 | Robin Pemantle, | A shuffle that mixes sets of any fixed size much faster than it mixes the whole deck |
| 1208 | Itai Benjamini, Robin Pemantle & Yuval Peres, | Martin capacity for Markov chains and random walks in varying dimensions |
| 1209 | Włodzimierz Bryc & Amir Dembo, | On large deviations of empirical measures for stationary Gaussian processes |
| 1210 | Martin Hildebrand, | Some random processes related to affine random walks |
| 1211 | Alexander E. Mazel & Yurii M. Suhov, | Ground states of a Boson quantum lattice model |
| 1212 | Roger L. Fosdick & Darren E. Mason, | Single phase energy minimizers for materials with nonlocal spatial dependence |
| 1213 | Bruce Hajek, | Load balancing in infinite networks |
| 1214 | Petr Klouček, | The transonic flow problems stability analysis and numerical results |
| 1215 | Petr Klouček, | On the existence of the entropic solutions for the transonic flow problem |
| 1216 | David A. Schmidt & Chjan C. Lim, | Full sign-invertibility and symplectic matrices |
| 1217 | Piermarco Cannarsa & Maria Elisabetta Tessitore, | Infinite dimensional Hamilton-Jacobi equations and Dirichlet boundary control problems of parabolic type |
| 1218 | Zhangxin Chen, | Multigrid algorithms for mixed methods for second order elliptic problems |
| 1219 | Zhangxin Chen, | Expanded mixed finite element methods for linear second order elliptic problems I |
| 1220 | Gang Bao, | A note on the uniqueness for an inverse diffraction problem |
| 1221 | Moshe Fridman, | A two state capital asset pricing model |
| 1222 | Paolo Baldi, | Exact asymptotics for the probability of exit from a domain and applications to simulation |
| 1223 | Carl Dou & Martin Hildebrand, | Enumeration and random random walks on finite groups |
| 1224 | Jaksa Cvitanic & Ioannis Karatzas, | On portfolio optimization under “drawdown” constraints |
| 1225 | Avner Friedman & Yong Liu, | A free boundary problem arising in magnetohydrodynamic system |
| 1226 | Dominic Welsh, | Randomised approximation schemes for Tutte-Gröthendieck invariants |
| 1227 | Zhangxin Chen, Bernardo Cockburn, Carl L. Gardner, & Joseph W. Jerome, | Quantum hydrodynamic simulation of hysteresis in the resonant tunneling diode |
| 1228 | E.G. Kalnins, G.C. Williams, & Willard Miller, Jr., | Intrinsic characterisation of the separation constant for spin one and gravitational perturbations in Kerr geometry |
| 1229 | Zhangxin Chen, | Large-scale averaging analysis of multiphase flow in fractured reservoirs |
| 1230 | Bruce Hajek & Babu Narayanan, | Multigraphs with the most edge covers |
| 1231 | K.B. Athreya, | Entropy maximization |
| 1232 | F.I. Karpelevich & Yu.M. Suhov, | Functional equations in the problem of boundedness of stochastic branching dynamics |
| 1233 | E. Dibenedetto & V. Vespri, | On the singular equation $\beta(u)_t = \Delta u$ |
| 1234 | M.Ya. Kelbert & Yu.M. Suhov, | The Markov branching random walk and systems of reaction-diffusion (Kolmogorov-Petrovskii-Piskunov) equations |
| 1235 | M. Hildebrand, | Random walks on random regular simple graphs |

- 1236 **W.S. Don & A. Solomonoff**, Accuracy enhancement for higher derivatives using Chebyshev collocation and a mapping technique
- 1237 **D. Gurarie**, Symmetries and conservation laws of two-dimensional hydrodynamics
- 1238 **Z. Chen**, Finite element methods for the black oil model in petroleum reservoirs
- 1239 **G. Bao & A. Friedman**, Inverse problems for scattering by periodic structure
- 1240 **G. Bao**, Some inverse problems in partial differential equations
- 1241 **G. Bao**, Diffractive optics in periodic structures: The TM polarization
- 1242 **C.C. Lim & D.A. Schmidt**, On noneven digraphs and symplectic pairs
- 1243 **H.M. Soner, S.E. Shreve & J. Cvitanić**, There is no nontrivial hedging portfolio for option pricing with transaction costs
- 1244 **D.L. Russell & B-Yu Zhang**, Exact controllability and stabilizability of the Korteweg-de Vries equation
- 1245 **B. Morton, D. Enns & B-Yu Zhang**, Stability of dynamic inversion control laws applied to nonlinear aircraft pitch-axis models
- 1246 **S. Hansen & G. Weiss**, New results on the operator Carleson measure criterion
- 1247 **V.A. Malyshev & F.M. Spijksma**, Intrinsic convergence rate of countable Markov chains
- 1248 **G. Bao, D.C. Dobson & J.A. Cox**, Mathematical studies in rigorous grating theory
- 1249 **G. Bao & W.W. Symes**, On the sensitivity of solutions of hyperbolic equations to the coefficients
- 1250 **D.A. Huntley & S.H. Davis**, Oscillatory and cellular mode coupling in rapid directional solidification
- 1251 **M.J. Donahue, L. Gurvits, C. Darken & E. Sontag**, Rates of convex approximation in non-Hilbert spaces
- 1252 **A. Friedman & B. Hu**, A Stefan problem for multi-dimensional reaction diffusion systems
- 1253 **J.L. Bona & B-Y. Zhang**, The initial-value problem for the forced Korteweg-de Vries equation
- 1254 **A. Friedman & R. Gulliver**, **Organizers**, Mathematical modeling for instructors
- 1255 **S. Kichenassamy**, The prolongation formula for tensor fields
- 1256 **S. Kichenassamy**, Fuchsian equations in Sobolev spaces and blow-up
- 1257 **H.S. Dumas, L. Dumas, & F. Golse**, On the mean free path for a periodic array of spherical obstacles
- 1258 **C. Liu**, Global estimates for solutions of partial differential equations
- 1259 **C. Liu**, Exponentially growing solutions for inverse problems in PDE
- 1260 **Mary Ann Horn & I. Lasiecka**, Nonlinear boundary stabilization of parallelly connected Kirchhoff plates
- 1261 **B. Cockburn & H. Gau**, A posteriori error estimates for general numerical methods for scalar conservation laws
- 1262 **B. Cockburn & P-A. Gremaud**, A priori error estimates for numerical methods for scalar conservation laws. Part I: The general approach
- 1263 **R. Spigler & M. Vianello**, Convergence analysis of the semi-implicit euler method for abstract evolution equations
- 1264 **R. Spigler & M. Vianello**, WKB-type approximation for second-order differential equations in C^* -algebras
- 1265 **M. Menshikov & R.J. Williams**, Passage-time moments for continuous non-negative stochastic processes and applications
- 1266 **C. Mazza**, On the storage capacity of nonlinear neural networks
- 1267 **Z. Chen, R.E. Ewing & R. Lazarov**, Domain decomposition algorithms for mixed methods for second order elliptic problems
- 1268 **Z. Chen, M. Espedal & R.E. Ewing**, Finite element analysis of multiphase flow in groundwater hydrology
- 1269 **Z. Chen, R.E. Ewing, Y.A. Kuznetsov, R.D. Lazarov & S. Maliassov**, Multilevel preconditioners for mixed methods for second order elliptic problems
- 1270 **S. Kichenassamy & G.K. Srinivasan**, The structure of WTC expansions and applications
- 1271 **A. Zinger**, Positiveness of Wigner quasi-probability density and characterization of Gaussian distribution
- 1272 **V. Malkin & G. Papanicolaou**, On self-focusing of short laser pulses
- 1273 **J.N. Kutz & W.L. Kath**, Stability of pulses in nonlinear optical fibers using phase-sensitive amplifiers
- 1274 **S.K. Patch**, Recursive recovery of a family of Markov transition probabilities from boundary value data
- 1275 **C. Liu**, The completeness of plane waves
- 1276 **Z. Chen & R.E. Ewing**, Stability and convergence of a finite element method for reactive transport in ground water
- 1277 **Z. Chen & Do Y. Kwak**, The analysis of multigrid algorithms for nonconforming and mixed methods for second order elliptic problems
- 1278 **Z. Chen**, Expanded mixed finite element methods for quasilinear second order elliptic problems II
- 1279 **M.A. Horn & W. Littman**, Boundary control of a Schrödinger equation with nonconstant principal part
- 1280 **R.E. Ewing, Yu.A. Kuznetsov, R.D. Lazarov & S. Maliassov**, Substructuring preconditioning for finite element approximations of second order elliptic problems. I. Nonconforming linear elements for the Poisson equation in a parallelepiped
- 1281 **S. Maliassov**, Substructuring preconditioning for finite element approximations of second order elliptic problems. II. Mixed method for an elliptic operator with scalar tensor
- 1282 **V. Jakšić & C.-A. Pillet**, On model for quantum friction II. Fermi's golden rule and dynamics at positive temperatures
- 1283 **V. M. Malkin**, Kolmogorov and nonstationary spectra of optical turbulence