



United States Department of Agriculture

Animal and Plant  
Health Inspection  
Service

**Veterinary Services**

Science, Technology,  
and Analysis Services

Center for  
Epidemiology and  
Animal Health

2150 Centre Avenue  
Building B  
Fort Collins, CO 80526

# **An Assessment of the Risk Associated with the Movement Broilers to Market Into, Within, and Out of a Control Area During a Highly Pathogenic Avian Influenza Outbreak in the United States**

**February 2017 FIRST DRAFT**

**June 2017 SECOND REVIEW**

**October 2017 THIRD REVIEW**

**October 2018 FINAL REVIEW AND CLEARANCE**

A Collaboration between the Broiler Sector Working Group, the  
University of Minnesota's Secure Food Systems Team, and  
USDA:APHIS:VS:CEAH



**UNITED STATES DEPARTMENT OF AGRICULTURE**

**ANIMAL AND PLANT HEALTH INSPECTION SERVICE**

**VETERINARY SERVICES**

**CENTER FOR EPIDEMIOLOGY AND ANIMAL HEALTH**



**Suggested bibliographic citation for this report:**

Carol Cardona, Carie Alexander, Peter Bonney, Francesca Contadini, Marie Culhane, Timothy Goldsmith, David Halvorson, Eric Linskens, Sasidhar Malladi, Amos Ssematimba, Jamie Umber, Todd Weaver, Emily Walz. **An Assessment of the Risk Associated with the Movement of Broilers to Market Into, Within, and Out of a Control Area during a Highly Pathogenic Avian Influenza Outbreak in the United States.** Collaborative agreement between USDA:APHIS:VS and University of Minnesota Center for Secure Food Systems. Fort Collins, CO. October 2018. 221 pgs

This document was developed through the Continuity of Business / Secure Food Supply Plans / Secure Poultry Supply project initiative. Related documents can be found at:

[https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/sa\\_emergency\\_management/ct\\_fadprep\\_continuity\\_of\\_business](https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/sa_emergency_management/ct_fadprep_continuity_of_business)

**Contributors:**

Carol Cardona, Professor, University of Minnesota (UMN)  
Carie Alexander, Veterinary Public Health and Preventive Medicine Resident, UMN  
Peter Bonney, Epidemiologist, UMN  
Francesca Contadini, Veterinary Public Health and Preventive Medicine Resident, UMN  
Marie Culhane, Associate Professor, UMN  
Timothy Goldsmith, Associate Professor, UMN  
David Halvorson, Professor Emeritus, UMN  
Eric Linskens, Researcher, UMN  
Sasidhar Malladi, Risk Analyst, UMN  
Amos Ssematimba, Epidemiologist, UMN  
Emily Walz, Risk Analyst, UMN  
Todd Weaver, Epidemiologist, USDA:APHIS:VS:CEAH  
Jamie Umber, Risk Analyst, UMN

**Acknowledgments:**

We thank the Broiler Sector Working Group for their support during the development of this risk assessment.

**Keywords:** avian influenza, continuity of business, HPAI, market, outbreak, risk, broiler, chicken

Questions or comments on data analysis, contact: Carol Cardona, DVM, PhD, DACPV, University of Minnesota College of Veterinary Medicine, Secure Food Systems, 651-253-2870 Email: ccardona@umn.edu
--

## Contents

<b>1</b>	<b>Abbreviations and Definitions</b> .....	<b>8</b>
<b>2</b>	<b>Executive Summary</b> .....	<b>13</b>
2.1	Likelihood of Broiler Flock Becoming Infected with HPAI via Components of Local Area Spread Resulting in Infected but Undetected Movement to Market.....	14
2.2	Likelihood of Broiler Flock Becoming Infected with HPAI via Movements of People, Vehicles, or Equipment, Resulting in Infected but Undetected Movement to Market .....	16
2.3	Likelihood of Broiler Flock Becoming Infected with HPAI via Load-out Crews, Vehicles, or Equipment Resulting in an Infected but Undetected Movement to Market .....	18
<b>3</b>	<b>Introduction</b> .....	<b>20</b>
<b>4</b>	<b>Scope</b> .....	<b>22</b>
4.1	Facilities Covered under this Risk Assessment.....	22
4.2	Types of Movements Addressed under this Risk Assessment .....	22
<b>5</b>	<b>Overview of Data Analysis Approaches</b> .....	<b>23</b>
<b>6</b>	<b>Significant Assumptions Used in the Risk Assessment</b> .....	<b>25</b>
<b>7</b>	<b>Background</b> .....	<b>27</b>
7.1	Definition of the Grow-out and Harvest Process.....	27
7.2	Overview of Broiler Meat Production in the United States.....	27
7.2.1	Vertical Integration .....	28
7.2.2	Service Technicians and Poultry Health Monitoring.....	29
7.2.3	Broiler Distribution and Logistics .....	29
7.3	Overview of Major Steps in Production of Broilers during Routine Operations.....	31
7.3.1	Broiler Facility Operations.....	31
7.3.1.1	Broiler House Preparations.....	31
7.3.1.2	Grow-out Period Management.....	31
7.3.1.2.1	Receiving the Chicks .....	31
7.3.1.2.2	Brooding.....	32
7.3.1.2.3	Brood and Grow Ventilation .....	32
7.3.1.3	Load-out.....	33
7.3.1.4	Transportation of Broilers to Processing and Awaiting Processing.....	34
7.4	Overview of Current Disease Prevention and Biosecurity Efforts in Broiler Production .....	36
7.4.1	Current Disease Prevention and Containment Measures in Grow-out Operations during Normal (non-outbreak) Situations.....	36
7.4.2	Structural Biosecurity .....	36
7.4.2.1	Conceptual Biosecurity .....	36
7.4.2.2	Secured Farm Entry .....	37
7.4.3	Operational Biosecurity.....	37
7.4.3.1	Farm Personnel, Service Technicians, and Visitors.....	37
7.4.3.2	Feed Delivery .....	38
7.4.3.3	Sanitation Facilities on Farm.....	38
7.4.3.4	Cleaning & Disinfection (C&D).....	38
7.4.3.4.1	Vehicles and Drivers .....	38
7.4.3.4.2	Equipment .....	39
7.4.3.4.3	Water Supplies.....	39
7.4.3.4.4	Housing Area .....	39
7.4.3.4.5	Load-out and Slaughter Plant .....	39
7.4.3.5	Animal, Pest, Insect Control.....	39
7.4.3.6	Dead Bird Disposal .....	40
7.4.3.7	Manure and Litter Management.....	40

<b>8</b>	<b>Hazard Identification: HPAI overview .....</b>	<b>41</b>
8.1	Agent .....	41
8.1.1	Definition of Highly Pathogenic Notifiable Avian Influenza.....	41
8.1.2	Host Range .....	42
8.2	Geographic Distribution of H5 and H7 HPAI.....	43
8.3	Virus Shedding .....	44
8.4	Chemical and Physical Inactivation .....	44
8.5	Persistence of HPAI Virus in Manure and Other Media .....	45
8.6	Transmission .....	45
8.6.1	Vertical transmission in chickens and turkeys .....	45
8.6.2	Transmission via artificial insemination in turkeys.....	46
8.7	Dose Response.....	47
8.7.1	Dose Response in Turkeys.....	47
8.7.2	Dose Response in Chickens.....	47
8.7.3	Route of Entry and 50 Percent Infectious Dose Estimate Used in this Assessment .	49
8.8	Latently Infected and Infectious Periods.....	49
8.8.1	Latently Infected and Infectious Periods in Turkeys.....	49
8.8.2	Latently Infected and Infectious Periods in Chickens .....	50
8.9	Clinical Signs .....	50
8.10	Diagnosis.....	51
8.11	Differential Diagnosis .....	51
<b>9</b>	<b>Risk Evaluation .....</b>	<b>52</b>
9.1	Pathways for a Broiler Flock Becoming Infected with HPAI via Local Area Spread Components other than those Involving Movements of People, Vehicles, and Equipment..	52
9.1.1	Role of Local Spread Components in Previous AI Outbreaks .....	52
9.1.2	Role of Insects in the Transmission of HPAI Virus .....	53
9.1.2.1	Literature Review .....	53
9.1.2.2	Expert Opinion .....	55
9.1.2.3	Qualitative Analysis.....	56
9.1.2.4	Likelihood Rating and Conclusion .....	57
9.1.3	Role of Aerosol Transmission of HPAI Virus .....	58
9.1.3.1	Aerosol Transmission of AI Virus in Past Outbreaks.....	58
9.1.3.2	Experimental Studies of Aerosol Transmission of AI Virus .....	60
9.1.3.3	Expert Opinion .....	61
9.1.3.4	Qualitative analysis .....	61
9.1.3.5	Likelihood Rating and Conclusion .....	62
9.1.3.5.1	Likelihood of HPAI Spread to a Broiler Flock in a Control Area via Aerosol Transmission from a Known HPAI-Infected Flock.....	62
9.1.3.5.2	Likelihood of HPAI Spread to a Broiler Flock in a Control Area via Aerosol Transmission from an Infected but Undetected Flock .....	62
9.1.3.5.3	Conclusion .....	62
9.1.4	Role of HPAI Spread to a Broiler Flock in a Control Area via Wild Aquatic Birds in the Farm Vicinity.....	63
9.1.4.1	Literature Review .....	64
9.1.4.2	Qualitative Analysis.....	71
9.1.4.3	Likelihood Rating and Conclusion .....	72
9.1.5	Role of HPAI Virus Spread to a Broiler Flock via Wild Non-Aquatic Birds in Farm Vicinity .....	72
9.1.5.1	Likelihood of Infection via Passerine or Columbiforme Birds in Farm Vicinity .....	72
9.1.5.1.1	Literature Review .....	73

	9.1.5.1.2	Qualitative Analysis.....	78
	9.1.5.1.3	Likelihood Rating and Conclusion .....	79
9.1.5.2		Likelihood of Infection via Other Non-Aquatic Bird Species in Farm Vicinity.....	79
	9.1.5.2.1	Literature Review .....	80
	9.1.5.2.2	Qualitative Analysis.....	84
	9.1.5.2.3	Likelihood Rating and Conclusion .....	84
9.1.6		Role of HPAI Virus Spread to Broiler Grow-Out Premises near Poultry Live-Haul Routes via Feathers, Feces, and Other Fomites.....	85
9.1.6.1		Risk of HPAI Virus Transmission to Broiler Grow-out Premises near Poultry Live-haul Routes.....	85
9.1.6.2		Literature Review .....	86
9.1.6.3		Qualitative Analysis.....	88
9.1.6.4		Risk Rating and Conclusion.....	92
	9.1.6.4.1	Risk of HPAI Transmission to Broiler Grow-Out Premises in a Control Area near Route of Live-Haul Trailer .....	92
	9.1.6.4.2	Conclusion .....	93
9.2		Pathways for a Broiler Flock Becoming Infected with HPAI via Movements of People, Vehicles, or Equipment .....	93
9.2.1		Role of HPAI Virus Spread to a Broiler Flock via Critical Operational Visits during PMIP.....	93
9.2.1.1		Likelihood of Infection via Feeds .....	94
9.2.1.2		Likelihood of Infection via Feed Delivery or Other Critical Operations Visits.....	94
	9.2.1.2.1	Literature Review .....	95
	9.2.1.2.2	Qualitative Analysis.....	96
	9.2.1.2.3	Likelihood Rating and Conclusion .....	96
9.2.2		Role of HPAI Virus Spread to a Broiler Flock via Growers or Employees and their Vehicles Entering the Premises.....	97
9.2.2.1		Likelihood of Infection via Movement of Growers.....	97
	9.2.2.1.1	Literature Review .....	98
	9.2.2.1.2	Qualitative Analysis.....	99
9.2.2.2		Likelihood of Infection via Farm Employees .....	101
	9.2.2.2.1	Literature Review .....	101
	9.2.2.2.2	Qualitative Analysis.....	102
9.2.2.3		Likelihood Rating and Conclusion .....	103
9.2.3		Role of HPAI Virus Spread to a Broiler Flock via Machinery or Equipment Shared between Multiple Premises .....	103
9.2.3.1		Likelihood of Infection via Shared Machinery or Equipment .....	104
9.2.3.2		Literature Review .....	104
9.2.3.3		Qualitative Analysis.....	107
9.2.3.4		Likelihood Rating and Conclusion .....	110
9.2.4		Role of HPAI Virus Spread to a Broiler Flock via Dead Bird Disposal .....	111
9.2.4.1		Dead Bird Disposal during PMIP .....	111
9.2.4.2		Likelihood of a Broiler Flock Becoming Infected via On-farm Dead Bird Disposal and Scavengers during PMIP .....	113
	9.2.4.2.1	Literature Review .....	113
	9.2.4.2.2	Qualitative Analysis.....	116
	9.2.4.2.3	Likelihood Rating and Conclusion .....	118
9.2.4.3		Likelihood of a Broiler Flock Becoming Infected with HPAI via Dead Bird Disposal that Takes Place before PMIP .....	118
	9.2.4.3.1	Literature Review .....	120
	9.2.4.3.2	Qualitative Analysis.....	121

9.2.4.3.3	Likelihood Rating and Conclusion .....	124
9.2.5	Role of HPAI Virus Spread to a Broiler Flock due to Garbage Management .....	125
9.2.5.1	Likelihood of HPAI-Virus Infection via Garbage Management .....	125
9.2.5.2	Literature Review .....	125
9.2.5.3	Qualitative Analysis.....	127
9.2.5.4	Likelihood Rating and Conclusion .....	131
9.2.5.4.1	Likelihood of a Broiler Flock Becoming Infected with HPAI-Virus due to Garbage Management.....	131
9.2.5.4.2	Likelihood of a Broiler Flock Becoming Infected with HPAI Virus due to Garbage Management when a PMIP is Implemented .....	131
9.3	Pathways for a Broiler Flock Becoming Infected with HPAIV via Load-Out Operations.....	132
9.3.1	PMIP Measures for Moving Broilers to Slaughter .....	133
9.3.1.1	Additional Load-out Mitigation Measures for Permitted Movement of Poultry to Market from a Control Area .....	133
9.3.2	Literature Review .....	134
9.3.3	Qualitative Analysis.....	135
9.3.4	Risk Rating and Conclusion.....	139
9.4	Likelihood of Detecting HPAI in an Infected Broiler House .....	140
9.4.1	HPAI Surveillance Measures .....	140
9.4.1.1	Current Measures .....	140
9.4.1.2	Outbreak Measures.....	140
9.4.2	Quantitative Methods for the Estimated Likelihood of Detection prior to the Start of Load-out on a Premises.....	141
9.4.2.1	Overview of Disease Transmission and Active Surveillance Models.....	141
9.4.2.2	Model Scenarios .....	142
9.4.2.3	Estimated Likelihood of Detection under a Pre Movement Isolation Period (PMIP) .....	143
9.4.2.4	Overall Likelihood of not Detecting HPAI in a Broiler House prior to the Start of Load-out on the Premises.....	146
9.4.2.4.1	Estimation of the Probability of Infection via a Spatial Transmission Kernel .....	146
9.4.2.4.2	Estimated Overall Likelihood of not Detecting HPAI in a Broiler House Prior to the Start of Load-out .....	148
9.4.3	Likelihood of Moving Infectious but Undetected Broilers Following Exposure during Load-out.....	149
9.4.4	Conclusions.....	151
<b>10</b>	<b>Overall Conclusion.....</b>	<b>153</b>
<b>Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates.....</b>		<b>157</b>
<b>Appendix 2: Literature Review on the Role of Local Area Spread in Previous Outbreaks .....</b>		<b>168</b>
<b>Appendix 3: Estimating an Approximate Posterior Distribution for the Prevalence among Insects.....</b>		<b>171</b>
<b>Appendix 4: Expert Polling on Insect Transmission Routes .....</b>		<b>172</b>
<b>Appendix 5: Live Broiler Movement Aerosol Modeling .....</b>		<b>175</b>
<b>Appendix 6: Expert Polling on Aerosol Transmission Route .....</b>		<b>184</b>
<b>Appendix 7: Cross-Commodity Pre-Movement Isolation Period (PMIP).....</b>		<b>187</b>

**Appendix 8: Modeling Technical Details ..... 191**  
**Appendix 9: Poultry Industry Survey on Garbage Management Practices..... 198**  
**Appendix 10: Supplementary Testing Protocols ..... 212**  
**Appendix 11: Load-out Mitigation Measures ..... 217**  
**References ..... 222**

## 1 Abbreviations and Definitions

AC	Antigen capture (as in "AC testing")
AI	Avian influenza
APHIS	Animal and Plant Health Inspection Service (USDA:APHIS)
BWG	Broiler Working Group
CEAH	Centers for Epidemiology and Animal Health (USDA:APHIS:VS:CEAH)
CFR	U.S. Code of Federal Regulations
C&D	Cleaning and disinfection, or cleaned and disinfected
DPI	Days post-inoculation (or days post-infection)
EA/AM	Eurasian/American
EPA	U.S. Environmental Protection Agency
FAO	Food and Agriculture Organization of the United Nations
GIS	Geographic Information System
GLEWS	Global Early Warning System for Major Animal Diseases Including Zoonoses
HA	Hemagglutinin
HI	Hemagglutination inhibition
HPAI	Highly pathogenic avian influenza
IAV-S	Influenza A virus of swine
ICS	Incident Command System
ILT	Infectious laryngotracheitis
IP	Infected premises
IVPI	Intravenous pathogenicity index
LPAI	Low pathogenic avian influenza
NA	Neuraminidase
NAHLN	National Animal Health Laboratory Network
NAHMS	National Animal Health Monitoring System (USDA)
NDV	Newcastle disease virus
NPIP	National Poultry Improvement Plan
NVSL	National Veterinary Services Laboratory (USDA)
OIE	World Organization for Animal Health (formerly Office International des Epizooties)
PBA	Perimeter Buffer Area



PM	Particulate matter
PMIP	Pre-Movement Isolation Period
PPE	Personal protective equipment
PRRSV	Porcine reproductive and respiratory syndrome virus
RH	Relative humidity
rRT-PCR	Real-time reverse transcription polymerase chain reaction
SAHO	State animal health official
SPF	Specific Pathogen Free
U.S.	United States of America
USDA	United States Department of Agriculture
UV	Ultraviolet Light
VS	Veterinary Services (USDA:APHIS:VS)
WHO	World Health Organization

**AERMOD**

Aerosol dispersion model developed by the EPA and recommended to be used for regulatory decisions associated with air quality.

**BID<sub>50</sub>**

50 percent bird infectious dose. One BID<sub>50</sub> unit is the amount of virus that will infect 50 percent of inoculated birds.

**Biosecurity**

A comprehensive set of measures to prevent the introduction of disease agents into a specific area.

**Breeder farm**

Farms with multiplier broiler breeder flocks that produce broiler hatching eggs. The hatching eggs from a breeder farm are transported to a hatchery.

**Broiler sector working group (BWG)**

A working group, which is made up of representatives from the broiler industry, academia, SAHOs, and USDA:APHIS, to support permits for the movement of broiler hatching eggs, chicks, or birds during an HPAI outbreak.

**Brooder premises**

Premises with facilities that raise poults (young turkeys) during the first few weeks of production. Day-old poults from a hatchery are transported to a brooder farm, some of which may also include grow-out barns.

**Buffer Zone**

The zone immediately surrounding the infected zone; the Buffer Zone and the Infected Zone constitute the Control Area.

**CID<sub>50</sub>**

50 percent chicken infectious dose. One CID<sub>50</sub> unit is the amount of virus that will infect 50 percent of inoculated chickens.

**Control Area**

A Control Area, consisting of an Infected Zone and a Buffer Zone, will be established to ensure the rapid and effective containment of the disease. The potential modes of transmission of HPAI are considered when determining the minimum size and shape of a Control Area. Movement control—through the use of permits—should be maintained until the disease is eradicated.

**Downtime for visitors**

For purposes of this assessment, downtime refers to the time interval between when a visitor enters the hatchery and the time of last contact with other domestic poultry, other avian species, and/or related organic material from the Control Area.

**EID<sub>50</sub>**

50 percent chicken embryo infectious dose. One EID<sub>50</sub> unit is the amount of virus that will infect 50 percent of inoculated embryos.

**ELD<sub>50</sub>**

50 percent chicken embryo lethal dose. One ELD<sub>50</sub> unit is the amount of virus that will be lethal to 50 percent of inoculated embryos. Since most HPAI viruses are embryo lethal, the ELD<sub>50</sub> estimates would be similar to EID<sub>50</sub>.

**Fomite**

An inanimate object, such as boots, clothing, etc., that, when contaminated with a viable disease agent, can serve as a source of infection for a susceptible host.

**Free Area**

Any area outside of the Control Area. The Surveillance Zone is a part of the Free Area.

**Free premises**

Poultry premises that are not in an HPAI Control Area and are not Contact or Suspect premises.

**Hatchery**

A commercial establishment that produces day-old poults from hatching eggs. Commercial hatcheries receive hatching eggs from off-site breeder farms and produce day-old poults that are shipped to brooder operations.

**Incident Command System (ICS)**

A management system designed to enable effective and efficient domestic incident management by integrating a combination of facilities, equipment, personnel, procedures, and communication within a common organizational structure.

**Infected Zone**

In an outbreak of HPAI, the Infected Zone will encompass the perimeter of all presumptive or confirmed positive premises (“Infected Premises”) and include as

many “Contact Premises” as the situation requires logistically or epidemiologically. Activities in an infected zone include:

- Preventing products from birds and other susceptible animals from leaving the zone unless a risk assessment determines that such movement can be permitted.
- Preventing movement of vehicles, equipment, and non-susceptible animals out of the zone unless appropriate biosecurity procedures (as determined by a risk assessment) are followed.

**Infectious period**

The period of time that an individual bird is infectious (i.e., shedding HPAI virus at sufficient levels that transmission could result if there is adequate contact with a susceptible host).

**Latent period**

The period of time between infection of a bird and when it becomes infectious.

**Line of Separation (LOS)**

The LOS is a clearly identified boundary around or within a poultry premises to separate off-farm traffic from on farm-movements of vehicles, people, and animals. The purpose of the LOS is to prevent movement of HPAI onto or from a premises. Crossing the LOS through a controlled access point requires following appropriate biosecurity measures.

**Local area spread**

Refers to risk pathways which have an increased likelihood for disease transmission with proximity to infected flocks.

**Movement permit**

A VS Form 1-27, a State-issued permit, or a letter—customized to the applicant’s situation—generated by the Permit Team and issued at the discretion of Incident Command to allow the movement of poultry industry products from a premises or a geographic area described in a quarantine order.

**National Poultry Improvement Plan (NPIP)**

A cooperative state-industry-federal program that establishes guidelines for evaluation of poultry products and poultry production relative to disease and eligibility for interstate/international trade.

**Observation period**

The time interval between moving birds from a brooder house to the placement of new poults into the brooder house during an HPAI outbreak where the previously raised flock is observed for clinical signs of HPAI. The purpose of observation time is to gain confidence that birds previously raised in the brooder house were not infected.

**Personal Protective Equipment (PPE)**

Special clothing and equipment designed to act as a barrier between an individual and a hazard; in this case, the hazard is a highly contagious pathogen (HPAI). PPE in the event of an HPAI outbreak serves to prevent the spread of the disease

agent between animals and locations. For purposes of this report, appropriate PPE is considered protective boot covers, clothing, and gloves.

**Premises**

A geographically and epidemiologically defined location, such as a ranch, farm, plant, or other establishment.

**Secure Broiler Supply Plan (SBS Plan)**

A science-based plan that is composed of outbreak measures and protocols proposed by the broiler sector working group to mitigate the risk of HPAI spread associated with the movement of hatching eggs and day-old chicks into, within, and outside of a Control Area. The SBS Plan includes various categories of measures such as active surveillance, holding time, biosecurity, cleaning, and disinfection.

**Secure Poultry Supply (SPS) Plan**

A harmonized plan to facilitate poultry industry and state regulatory agency preparedness for product movement in an HPAI outbreak.

**Secure Turkey Supply (STS) Plan**

A set of science-based outbreak measures developed by the Turkey Sector Working Group to mitigate the risk of HPAI spread associated with the movement of turkeys, turkey eggs, and turkey semen in a Control Area.

**Standard Operating Procedure (SOP)**

Established or prescribed methods to be followed routinely for the performance of designated operations in a designated situation.

**Turkey Sector Working Group (TWG)**

A working group, which is made up of representatives from the turkey industry, academia, SAHOs, and the USDA:APHIS, to support permits for the movement of turkey hatching eggs, poults, or birds during an HPAI outbreak.

**TCID<sub>50</sub>**

50 percent tissue culture infectious dose. One TCID<sub>50</sub> unit is the amount of virus that will cause cytopathic effects in 50 percent of exposed host cells. The Madin-Darby Canine Kidney cell line is often used to estimate TCID<sub>50</sub> for HPAI viruses.

**Zoonosis**

A disease caused by an infectious agent that can be transmitted between (or shared by) animals and humans.

## 2 Executive Summary

In the event of a highly pathogenic avian influenza (HPAI) outbreak in the United States, poultry industry, local, State, and Federal authorities will implement a foreign animal disease emergency response. In these circumstances, permit requests to move poultry and poultry products must be supported by risk assessments which demonstrate that the risk of HPAI spread associated with the movement is acceptable. Performing the risk assessments prior to an HPAI outbreak can enhance emergency response and facilitate timely movement permitting decisions during an outbreak. This document assesses the risk that the movement of market-age broilers to processing (i.e., broilers to market), during an HPAI outbreak, from a premises located within the Control Area, will result in HPAI virus spread to a virus-free poultry premises.

This risk assessment is a joint effort of the Secure Broiler Supply (SBS) Working Group, which is made up of representatives from the broiler industry, academia, State Animal Health Officials (SAHOs), and the United States Department of Agriculture Animal and Plant Health Inspection Service (USDA:APHIS), to support permits for the terminal movement of broilers to processing during an HPAI outbreak. This assessment is applicable to intensively raised commercial or contract grow-out broiler premises that do not have other poultry on the premises and practice an all-in, all-out single-age growing system. These broiler facilities must participate in the USDA APHIS National Poultry Improvement Plan (NPIP) and follow the SBS Plan in the event of an HPAI outbreak. The SBS Plan contains science-based outbreak measures developed by the SBS working group to mitigate the risk of HPAI spread associated with the terminal movement of live birds to market.

This risk assessment considers applicable current industry practices and biosecurity measures (e.g., the NPIP) as well as outbreak-specific measures stipulated within the SBS Plan. The main categories of outbreak measures outlined in the SBS Plan for broiler premises that wish to move birds to slaughter from a Control Area include:

- Establishing all criteria needed to meet the definition of a Monitored Premises
- Active surveillance (e.g., rRT-PCR [real-time reverse transcription polymerase chain reaction] testing, detection of abnormally high mortality)
- Observing the greatly enhanced biosecurity measures of the Pre-Movement Isolation Period (PMIP)
- Following specific infection mitigation measures pertaining to load-out vehicles, crews, and equipment as determined by the duration of load-out process

The Pre-Movement Isolation Period (PMIP) is a critical biosecurity component that involves a defined period of greatly intensified biosecurity for an entire premises prior to permitted movement of live poultry within, into, or out of a regulatory Control Area during an HPAI outbreak. The PMIP starts a specified number of days prior to the scheduled movement date and ends when load-out begins (i.e., the hours or days of load-out are not considered part of the PMIP). The PMIP duration to move broilers to market is set at five days and includes the following stipulations:

- No live or dead poultry will be moved onto or off the premises.
- Only critical operational visits to the premises will continue.

- Manure, litter, and garbage will not be removed from the premises; the producer is responsible for managing the risks associated with any on-site movement or handling of manure, litter, and garbage that must occur.
- Enhanced biosecurity will be implemented for people, vehicles, and equipment entering the premises; no off-site equipment will be pre-staged on-site.

The length of the PMIP decided upon by the SBS working group (five days) generally provides a high probability of detection. However, a five-day PMIP is not entirely robust to allow high probabilities of detection (i.e., > 95%) of all potential HPAI virus strains and contact rates.

The emphasis in this assessment is on the risk of HPAI virus spread to a susceptible poultry premises associated with the movement of broilers from a premises located within a Control Area. We assume that movement of infected and undetected broilers to processing would pose a high likelihood of HPAI spread to susceptible poultry and have high adverse consequences, and therefore we rated the overall risk according to the likelihood of moving infected and undetected birds. The probability of detection before movement improves as the number of days after exposure increases. As HPAI moves through the flock, there is an exponential increase in mortality, which consequently increases the likelihood of including at least one infected bird in the pooled mortality sample taken for diagnostic testing or of observing total mortality above the threshold amount. Thus, the PMIP serves a dual purpose of (1) reducing the chances of exposure to HPAI close to the time of movement, and (2) allowing sufficient time for the infection to manifest itself within the flock and be detected.

To assess the overall risk of moving broilers to market, this risk assessment evaluated the possible pathways for virus transmission to broiler premises. Each pathway may consist of combinations of several activities. We have grouped these pathways into several categories: 1) components of local area spread; 2) people, vehicles, or equipment; and 3) load-out processes. Local area spread refers to risk pathways which cause an increased likelihood of disease transmission with proximity to infected poultry flocks. If, due to a lapse in PMIP biosecurity practices or other unforeseen events, broilers are moved from the finishing barn within a short time after being exposed to the HPAI virus, it is unlikely that HPAI would be detected by the time of movement. Therefore, pathways for HPAI infection of market-age broilers close to scheduled movement combined with the likelihood of detecting the infection prior to movement and the likelihood of infection during the load-out process were considered in order to evaluate the overall risk of spread associated with movement of broilers to processing. These pathways and the corresponding likelihood and risk ratings are described below. The overall finding and conclusion qualitatively integrates the results from the pathway assessments taking into account the assumed high consequence assessment.

## **2.1 Likelihood of Broiler Flock Becoming Infected with HPAI via Components of Local Area Spread Resulting in Infected but Undetected Movement to Market**

- **Insects.** The likelihood of a broiler premises becoming infected with HPAI virus via insect transmission varies with distance and with source premises infection status, where proximity to a known infected premises directly influences likelihood. Of note, for premises located closer than 1.5 km to an infected flock, there are too many variables to accurately assess the risk of becoming infected with HPAI via insect transmission. The

following is a breakdown for the likelihood of HPAI spread to a broiler flock via insect transmission:

Source Premises Type	Composite likelihood rating (Insects) Distance from source (km)		
	>1.5	>2	>3
Known infected premises	<i>Negligible to moderate</i>	<i>Negligible to low</i>	<i>Negligible</i>
Infected but undetected premises	<i>Negligible to low</i>	<i>Negligible to low</i>	<i>Negligible</i>

- Aerosols.** The likelihood of a broiler premises becoming infected with HPAI virus via bio-aerosols varies with distance and viral load at the source premises. Literature review and most previous outbreak reports indicated that aerosol transmission was not an important factor at distances more than 1.5 km from an infected flock. However, there is some evidence of aerosol transmission over shorter distances. The following is a breakdown for the likelihood of HPAI spread to a broiler flock via bio-aerosol transmission:

Source Premises Type	Composite likelihood rating (Aerosols) Distance from source (km)		
	0.5	1	>1.5
Known infected premises	<i>Moderate to high</i>	<i>Moderate</i>	<i>Low</i>
Infected but undetected premises	<i>Low to moderate</i>	<i>Low</i>	<i>Negligible to low</i>

- Wild birds.** The likelihood of HPAI virus spread to a broiler grow-out premises via wild birds depends upon the type of wild birds and exposure to the wild birds. Aquatic species and larger non-aquatic species have not been known to gain entry to poultry barns, while passerine birds may access the inside of a broiler grow-out barn. With an effective PMIP, the increased barn-to-barn biosecurity, specifically the use of house-specific footwear, may decrease HPAI infection via wild aquatic birds and via non-passerine non-aquatic birds, as their waste is unlikely to access or be tracked into a broiler grow-out barn. Given that passerine birds may access the inside of broiler grow-out barns (even during a PMIP) and have been shown to be capable of shedding the virus, the likelihood of HPAI spread to a broiler flock via each of these bird categories is described below:

Wild bird category	Composite likelihood rating (Wild birds)
Aquatic wild birds	<i>Low</i>
Non-aquatic wild birds (passerine-type)	<i>Low to moderate</i>
Non-aquatic wild birds (non-passerine)	<i>Low</i>

- Live-haul routes.** The risk of HPAI virus spread to broiler grow-out premises near poultry live-haul routes via feathers, feces, and other fomites is both distance- and source flock-dependent. Given that poultry and live-haul vehicles passing a premises within the Control Area may originate from within or outside the Control Area, the following risk ratings are provided:

Characteristics of live-haul vehicle	Risk rating at given distance (between live-haul road and poultry premises)		
	<100 meters	100-1000 meters	>1000 meters
Truck hauling birds that had no PMIP and no tests	<i>High</i>	<i>Moderate</i>	<i>Low</i>
Truck hauling birds that had less than optimum PMIP and tests (80% effective PMIP; delayed testing; or load-out >24 hours)	<i>Low</i>	<i>Very Low</i>	<i>Negligible</i>
Truck hauling birds that had a PMIP & rRT-PCR negative birds (100% effective PMIP; two tests within 24 hours of move and completion within 24 hours)	<i>Very Low</i>	<i>Negligible</i>	<i>Negligible</i>

## 2.2 Likelihood of Broiler Flock Becoming Infected with HPAI via Movements of People, Vehicles, or Equipment, Resulting in Infected but Undetected Movement to Market

- Feed and Critical Operational Visits.** Critical operational visits will be limited during PMIP; however, delivery of feed during this period is likely and the potential for emergency maintenance visits also exists. Provided the biosecurity stipulations of the PMIP are in place and strictly followed, the likelihood of a broiler flock becoming infected with HPAI via feed and critical operational visits during PMIP was assessed as follows:

Critical operation component	Composite likelihood rating (Critical Operational Visits)
Contaminated feed	<i>Negligible</i>
Feed delivery (i.e., driver and/or vehicle)	<i>Low</i>
Other critical visitors (i.e., personnel and/or vehicle)	<i>Low to moderate</i>

- Growers, Employees, and Their Vehicles.** During the PMIP, vehicle and visitor traffic to a poultry premises may include only critical visitors, employees, and growers. Provided the SBS PMIP measures for people are strictly followed (e.g., use of farm-specific clothing and barn-specific footwear), we rate the likelihood of a broiler flock



becoming infected with HPAI via people (namely growers or employees) and their vehicles during the PMIP as follows:

Person type	Composite likelihood rating (People)
Persons entering poultry barns	<i>Low</i>
Persons not entering poultry barns	<i>Very low</i>

- Shared Equipment.** Previous poultry disease outbreaks demonstrate a known risk for virus spread as a result of movement of contaminated and shared equipment. Equipment that is brought onto a poultry premises may contaminate the ground or personnel who work with poultry, or if used inside a barn, may come into direct contact with live poultry. During the PMIP, no off-site equipment will be pre-staged and only equipment associated with critical operational visits may be brought to the premises. Provided the biosecurity stipulations of the PMIP are in place and strictly followed, the likelihood of a broiler flock becoming infected with HPAI virus via shared machinery or equipment is *low*.

Pathway	Composite likelihood rating (Shared equipment)
Shared equipment	<i>Low</i>

- Dead Bird Disposal.** The risks of HPAI introduction associated with off-site dead bird disposal methods, such as rendering, are well documented, and off-site disposal of dead birds must be discontinued during PMIP. Nevertheless, the risky practice of off-site dead bird disposal may still occur in the Control Area on other premises, and on a broiler premises in the days leading up to a PMIP. Off-site dead bird disposal methods prior to a PMIP result in possible premises contamination, although the implementation of a PMIP does reduce the likelihood that such contamination will be tracked inside a grow-out barn during the PMIP. Additionally, many scavenger species can biologically or mechanically carry HPAI virus and have home ranges of adequate size to contain adjacent poultry farms. As such, access to any on-farm dead bird storage container or disposal method represents a pathway for HPAI spread. Provided the SBS PMIP measures—specifically discontinuing any off-farm mortality disposal and utilizing barn-specific footwear—are strictly followed, we rate the likelihood of a broiler flock becoming infected with HPAI via dead bird disposal as follows:

Mortality disposal practice	Composite likelihood rating (Dead bird disposal)
Likelihood of a broiler flock becoming infected via the mechanical or biological transfer of HPAI virus from on-farm dead bird disposal during PMIP	<i>Low to moderate</i>
Likelihood of a broiler flock becoming infected via the mechanical or biological transfer of HPAI virus from off-site dead bird disposal that takes place prior to the PMIP	<i>Moderate</i>

- Garbage Management.** Multiple types of potentially contaminated items have been reported to be disposed of in garbage on poultry operations, and there is potential for HPAI virus associated with garbage management to be tracked into the broiler house. During a PMIP, no off-site movement of garbage is allowed. Provided the SBS PMIP measures (specifically discontinuing any off-farm garbage disposal and utilizing barn-specific footwear) are strictly followed, we rate the likelihood of a broiler flock becoming infected with HPAI via garbage management during the PMIP as *low*.

Pathway	Composite likelihood rating (Garbage)
Garbage management	<i>Low</i>

### 2.3 Likelihood of Broiler Flock Becoming Infected with HPAI via Load-out Crews, Vehicles, or Equipment Resulting in an Infected but Undetected Movement to Market

- Load-out.** Previous outbreaks have implicated contaminated load-out crews and equipment in the spread of AI. If a flock were infected via contaminated load-out crews or equipment, shortening the duration of premises load-out to slaughter limits the time that the virus may spread within the flock. Furthermore, depopulation of the premises (i.e., no “split” or “partial load-outs” permitted) leaves no susceptible hosts on-site, as load-out results in a terminal movement. Given that PMIP enhanced biosecurity and testing measures are strictly implemented, and that additional load-out mitigation measures are in place commensurate with the duration of the premise-wide load out process, the risk of a broiler flock becoming infected with HPAI virus via load-out operations and resulting in an infected but undetected movement to market is estimated to be *low to moderate*.

Pathway	Composite risk rating (Load-out)
Load-out and transport to slaughter	<i>Low to moderate</i>

This assessment aids, but does not replace, the judgment of on-scene officials. This document is an evolving product-specific risk assessment that will be reviewed and updated as necessary before and during an outbreak to incorporate the latest scientific information and preventive measures. If the Incident Command System is activated in response to an HPAI outbreak, APHIS (and Incident Command staff) will review this risk assessment with respect to the situation in order to assess industry requests for movement of broilers to market.

### **Overall Finding and Conclusion**

The risk that movement of broilers to market into, within, and out of a Control Area during an HPAI outbreak results in the infection of susceptible poultry is *low to moderate*, provided that all applicable preventive measures from the Secure Broiler Supply Plan (SPS Plan), in particular the Pre-Movement Isolation Period, are strictly followed.

### 3 Introduction

In the event of a highly pathogenic avian influenza (HPAI) outbreak in the U.S. poultry industry, local, State, and Federal authorities will implement a foreign animal disease emergency response. This response consists of a control and eradication strategy utilizing depopulation, quarantine, and movement control measures within a Control Area to prevent further spread of HPAI virus. State and/or Federal authorities may also issue official permits to allow movement of birds and their products from premises not known to be infected within the Control Area to promote business continuity. A request for a movement permit must be supported by a risk assessment (or some scientifically based logical argument) to demonstrate that the risk of HPAI spread associated with the movement of the product in question is acceptable; ultimately, whether or not the assessed risk level is acceptable will be determined by regulatory authorities and industry.

Completing these types of risk assessments in a timely manner during an outbreak can be challenging. Integrated poultry systems precisely manage grow-out facilities to maximize carcass value and minimize cost of chicken meat production. Extended movement restrictions may result in delays to processing, increased cost of production, and loss of carcass value. Proactive risk analysis identifies areas of risk and incorporates mitigation steps in order to minimize the spread of infection. Evaluating risk before an outbreak occurs facilitates timely emergency response and movement permitting decisions and minimizes unintended disruptions to business continuity.

Previous assessments have explored the risk of HPAI infection of day-old chicks at the hatchery via horizontal transmission from breeder premises (See Broiler Hatching Egg risk assessment; risk found to be *negligible to low*), and the risk of HPAI infection of day-old chicks due to local area spread (See Day Old Chicks risk assessment). These pathways were evaluated to be *negligible to low* when the outbreak measures specified in the SPS Plan are implemented.

The purpose of this assessment is to provide regulators and industry with an objective and defensible method of assessing the disease risk associated with the movement of live broilers (meat-type chickens) to slaughter. As broilers are generally slaughtered between 35 and 40 days of age, HPAI infection early in the brood or grow period would likely be detected before movement. However, it is less likely that HPAI would be detected by the time of movement if they became infected during load-out or in the days leading up to movement, due to a delay between infection and the manifestation of clinical signs or mortality.

In order to evaluate the risk of movement of broilers in a Control Area to market, plausible pathways were identified for the spread of HPAI infection. This analysis focused on pathways for HPAI infecting a broiler flock in the days leading up to movement (entry assessment of HPAI virus introduction onto broiler farms at or before scheduled time of movement to slaughter) as well as the pathways by which this movement of broilers to market could infect another flock in the area (exposure assessment of HPAI spread as the result of moving an infected but undetected broiler flock). Each pathway may consist of combinations of several activities. These pathways have been grouped into several categories: 1) local area spread; 2) people, vehicles, or equipment; and 3) load-out.

Local area spread refers to risk pathways that pose an increased likelihood for disease transmission due to proximity to an infected premises. The components of local area spread considered in this analysis include:

- transmission of HPAI virus through insects;
- bio-aerosols generated from neighboring infected flocks;
- transmission of HPAI virus by wild birds (aquatic and nonaquatic); and
- fomite transmission from poultry live-haul roads.

Other pathways considered in this analysis include transmission through:

- feed delivery;
- vehicles associated with essential visitors;
- fomites associated with visitors or grower premises employees who may have had contact with infected poultry or poultry waste;
- shared machinery or equipment;
- mechanical or biological transmission from dead bird disposal via wildlife
- garbage collection and disposal; and
- equipment and crews used for load-out

This assessment applies only to the movement of broilers off premises located in the Control Area to slaughter either inside or outside the Control Area. This assessment considers current industry practices and biosecurity measures as well as outbreak-specific measures applicable for the movement of broilers to market in the risk evaluation. Specific biosecurity measures may vary widely by farm and geographic area. Categories of outbreak-specific measures from the SBS Plan considered here include a Pre-Movement Isolation Period (PMIP) for flocks prior to movement to market. Other measures include:

- Limiting visitors to critical operations visits
- Specific feed truck and driver biosecurity measures
- Biosecurity measures for grow-out farm personnel and other essential visitors
- Measures for persons collecting surveillance samples
- Load-out truck and crew biosecurity, including truck routing
- Following PMIP, specific downtime measures

This assessment is an evolving product-specific risk assessment that will be reviewed and updated as necessary before and during an outbreak to incorporate the latest scientific information and preventive measures. If the Incident Command System (ICS) is activated in response to an HPAI outbreak, U.S. Department of Agriculture Animal and Plant Health Inspection Service (USDA:APHIS) and Incident Command staff will review this risk assessment regarding the situation in order to assess industry requests for movement of broilers to market.

## **4 Scope**

This section describes the scope of the assessment regarding the type of movements addressed and the facilities covered.

### **4.1 Facilities Covered under this Risk Assessment**

This risk assessment is applicable to intensively raised commercial or contract grow-out facilities producing market-weight broilers that meet all of the criteria listed below:

- Are in an HPAI Control Area
- Participate in the USDA APHIS National Poultry Improvement Plan (NPIP) as stated in 9CFR145 and 9CFR147 and in conjunction with biosecurity principles approved at the 43<sup>rd</sup> NPIP Biennial Conference<sup>2,3</sup>
- Implement the SBS Plan in the event of an HPAI outbreak
- Do not have other poultry on the premises and practice an all-in all-out single-age growing system.

### **4.2 Types of Movements Addressed under this Risk Assessment**

This risk assessment will address only the pathways that may potentially affect movement of market-age broilers within the Control Area directly to commercial USDA-inspected slaughter facilities inside or out of the Control Area.

## 5 Overview of Data Analysis Approaches

This assessment follows the general qualitative risk assessment principles recommended by the World Organization for Animal Health (OIE) import risk analysis guidelines.<sup>4</sup> However, the risk assessment organization has been modified from that proposed in the OIE import risk analysis handbook as appropriate for the movement of market-weight broilers to slaughter facilities. As noted in the introduction, many of the described pathways may play a role in both entry assessment (i.e., entry of HPAI virus onto broiler farms at or before the scheduled time of movement to slaughter) and exposure assessment (i.e., spread of HPAI to a broiler flock as a result of the movement of an infected but undetected flock to slaughter). A consequence assessment was assumed to be high as the risk of moving infected but undetected broilers could have considerable adverse consequences with regard to HPAI spread.

The assessment utilizes an evaluation approach that rates the likelihood of individual pathways on a qualitative scale. The likelihood for each pathway was assessed and categorized using the descriptive scale in **Table 1**. The qualitative ratings for the pathways were determined using multiple data sources and evaluation approaches such as literature review, expert opinion, quantitative simulation model predictions, and past outbreak experiences. Quantitative simulation model results from previously completed proactive risk assessments were used to estimate the prevalence of infectious birds in potentially infected but undetected poultry flocks located near the grow-out facility. Steady-state aerosol dispersion models recommended by the U.S. Environmental Protection Agency (EPA) were used to partially inform the risk of aerosol spread from infected and undetected farms, along with other approaches. To determine the rating for pathways involving a chain of events in which all have to occur for the pathway to be completed, relatively more weight was given to events with lowest likelihood in the chain.

**Table 1:** Descriptive scale to estimate the likelihood for an event to occur.

Likelihood Rating	Description
<i>Extremely High</i>	The event is almost certain to occur
<i>High</i>	There is more than an even chance that the event will occur
<i>Moderate</i>	The event is unlikely but does occur
<i>Low</i>	It is very unlikely that the event will occur
<i>Very Low</i>	There is a remote chance that the event will occur
<i>Negligible</i>	The likelihood that the event will occur is insignificant, not worth considering

The descriptive rating scale specific to the hazard (HPAI) in this assessment is provided below.

*Negligible Risk:* HPAI spread to other susceptible poultry through the risk pathway is insignificant or not worth considering.

*Very Low Risk:* HPAI spread to other susceptible poultry through the risk pathway is remote.

*Low Risk:* HPAI spread to other susceptible poultry through the risk pathway is very unlikely.


*Moderate Risk:* HPAI spread to other susceptible poultry through the risk pathway is unlikely but does occur.

*High Risk:* There is more than an even chance that HPAI spread to other susceptible poultry through the risk pathway will occur.

*Extremely High Risk:* HPAI spread to other susceptible poultry through the risk pathway is almost certain to occur.

Uncertainty within the likelihood/risk estimations was accounted for by using a range defined by the terms in the descriptive rating scale. A risk estimate of *negligible to low* includes the true risk, which is not deterministically known, where the interval between the two ratings represents the uncertainty in the analysis. For example, a *negligible to low* rating if the premises is located 1.5 km from an infected but undetected poultry farm was used with regard to aerosol transmission where there is considerable uncertainty in the aerosol dose-response relationship in chickens and the particle size distribution of aerosols generated in poultry houses depends on the ventilation design, production type, and age of the birds. Other areas of uncertainty were handled similarly during the analysis.

The overall risk estimate for the movement of broilers to market was determined by qualitatively combining the likelihoods of the individual pathways assuming that all applicable preventive measures from the Secure Broiler Supply Plan (SPS Plan), in particular the Pre-Movement Isolation Period, are strictly followed (see Figure 1 below).



**Figure 1:** Diagrammatic representation of the overall assessed risk with the relative amount of risk increasing as the width of the figure increases (the overall risk of component parts is not to scale). The overall risk assessment is based on consideration of the steps needed to move live birds to market and the pathways that could lead to infection of a flock, the subsequent likelihood of detection of the infected flock, and potential movement of an infected but undetected flock.



## 6 Significant Assumptions Used in the Risk Assessment

This assessment is proactive in nature and cannot address the specific circumstances surrounding an outbreak in detail. Therefore, we must make some assumptions to establish context and applicability. These assumptions are that:

- An HPAI outbreak has been detected, APHIS is implementing the HPAI Response Plan, and some form of local and state planning has taken place. The APHIS HPAI Response Plan is intended to complement regional, State, and industry plans. APHIS recommends their continued development.
- Broiler farms may have undetected HPAI infection in their flocks. If there were absolute certainty that a broiler flock arrives at slaughter without HPAI, there would be no risk of HPAI spread from movement of birds from a broiler farm. On the other hand, if HPAI infection has been detected on the premises, it is assumed that Incident Command would quarantine the premises. If infection were detected, the movement of broilers to market would not be allowed (and the facility would be depopulated, cleaned, and disinfected before resuming production).
- Movement of infected and undetected broilers to processing would have a high likelihood of spreading HPAI to susceptible poultry and have high adverse consequences, and therefore we rated the risk according to the likelihood of moving infected and undetected birds.
- The movement of broilers to market in the Control Area is in accordance with the SBS Plan, and all relevant preventive measures from the SBS Plan are strictly followed. The assessment does not evaluate the risk that the preventive measures are incorrectly implemented either intentionally or unintentionally.
- Other mechanisms outside of the SBS Plan may be utilized for HPAI control at the discretion of the Incident Commander. Risks associated with movement of birds to slaughter at a date earlier than usually marketed (i.e., early marketing) in order to decrease the number of susceptible species within the Control Area falls outside the scope of this Risk Assessment.
- The assessment focuses on the risk that movement of broilers to market will result in the spread of HPAI to other susceptible poultry. Although the risks to humans or wildlife associated with the production or movement of live poultry are critical concerns that should be addressed, they are outside the scope of this assessment. The Highly Pathogenic Avian Influenza Response Plan has personnel safety measures designed to mitigate the risk to humans.
- The broiler grow-out premises is a standalone facility without other poultry on the premises. It is assumed that the premises practices an all-in, all-out grow-out schedule with birds of a single age on a given premises.
- The adverse consequences of movement of infected broilers are assumed to be high. Hence, the risk rating was determined on the basis of the likelihood of HPAI spread, and the consequences of the event were not evaluated.

- The risk assessment applies to HPAI virus strains that cause clinical infection and increased mortality in infected broiler chickens. The risk assessment may not apply to strains that do not cause clinical signs representative of HPAI infection (i.e., AI strains that are classified as highly pathogenic on a molecular basis only). For such strains, this risk assessment would have to be revised to reflect the biological characteristics of the virus.
- The disinfectants used to implement various C&D measures in the SBS Plan during an outbreak have been approved by the Incident Command and are applied according to the manufacturer's label directions or recommended procedures.
- This assessment does not evaluate the risk of transmitting poultry diseases other than HPAI. Risk management decisions for poultry diseases other than HPAI are not directly supported by this work.

## 7 Background

### 7.1 Definition of the Grow-out and Harvest Process

A broiler is a chicken raised at a broiler farm for meat. A broiler production farm receives chicks from a hatchery typically within one day after eggs are hatched.<sup>5</sup>


At the broiler production farm, the chicks go through the brooding period (the first 1 to 2 weeks) in a facility where proper temperatures are provided, because they cannot fully regulate their body temperature. Brooding is followed by the grow-out period until chickens reach the desired market live weight between 4 and 10 weeks of age. Broilers are then caught and transported to processing.<sup>6</sup>

This risk assessment specifically focuses on the movement of broilers from broiler production farms to processing plants/slaughter facilities.

### 7.2 Overview of Broiler Meat Production in the United States

The broiler industry in the United States sold USD 25 billion worth of products at the wholesale level and generated USD 40 billion in retail in 2010. During 1990 to 2001, chicken consumption per capita surpassed that of beef<sup>7</sup> and by 2010 chicken production on both a carcass and boneless basis had surpassed that of beef production.<sup>8</sup> Production of ready-to-cook chicken meat was 39,620 million pounds in 2015, a nearly 4 percent increase from 2014.<sup>9</sup>

The broiler industry structure contains two main segments: primary breeder facilities and broiler facilities (**Figure 2**). Primary breeder facilities consist of elite farms producing chicks for great grandparent farms, which in turn supply chicks to grandparent farms. Grandparent farms supply chicks to the broiler segment, which operates multiplier and broiler production farms. Multiplier farms raise parent flocks, producing hatching eggs to provide chicks to be raised at broiler grow-out farms (**Figure 3**).<sup>6</sup> Hatcheries provide the proper environment and management for hatching eggs and day-old chicks to be supplied to the next facility downstream.



**Figure 2.** Overview of primary breeder and broiler facilities. Image courtesy of USDA APHIS Poultry Industry Manual

Broiler grow-out farms receive day-old chicks from hatcheries, raise the birds to market weight, and then transport them to processing plants. The majority of large-sized broiler farms have an average house size of 24,000 birds with 50,000 to 99,999 birds at maximum capacity.<sup>10</sup>



**Figure 3.** Hatching eggs waiting to be transferred to an incubator at the hatchery. Photo courtesy of GNP Company

### 7.2.1 Vertical Integration

Most of the broiler operators in the U.S. are vertically integrated, such that a company (integrator) owns and manages two or more successive stages of the marketing system. Typically, a single company owns or controls multiplier farms, hatcheries, feed mills, broiler farms, and/or a contract grower network for all live-rearing operations, and processing plant(s).<sup>8</sup> Most large-sized broiler companies receive chicks produced by company-owned multiplier flocks and hatcheries.<sup>6</sup> The vast majority of farmers who grow chickens for slaughter are

contract farmers. The integrator provides and retains ownership of the feed and the chickens, while the farmers receive payment either per bird or per weight unit (pound).<sup>11</sup>


Integrators closely monitor factors affecting cost of production down to hundredths of a cent per pound. These factors include morbidity, mortality, rate of daily gain, feed conversion, and condemnations. Company veterinarians constantly monitor flocks for disease problems, which are quickly identified and corrected. Integrated companies are closed marketing systems and do not buy or sell birds in open livestock markets. Broilers are produced under single-age, all-in-all-out production schemes which break the chain of disease transmission from one flock to the next. Biosecurity programs limit access to poultry houses and encourage use of dedicated clothing and boots, discourage sharing of equipment, and prohibit contact with other birds including pets and other poultry.<sup>12</sup>

### **7.2.2 Service Technicians and Poultry Health Monitoring**

Service technicians are a critical part of broiler flock husbandry. Service technicians are employed by the integrator and act as a liaison between contract growers and company management. Company expectations and policy changes are presented to the growers by service technicians. Service technicians assist with scheduling chick arrivals, feed deliveries to farms, and final load-out for processing. Typically, service technicians visit each farm weekly and provide advice on best management practices for the strain of broiler chicken being grown to achieve the company's target market weight.

### **7.2.3 Broiler Distribution and Logistics**

According to the USDA National Animal Health Monitoring System (NAHMS) Poultry Survey in 2010, 95.2 percent of multiplier farms ship eggs within state. The most common method for transporting birds from hatcheries onto broiler farms, and from broiler farms to slaughter, is by company-dedicated vehicles.<sup>13</sup> U.S. broiler production is concentrated in the southeastern U.S. due to favorable climate and proximity to grains and final markets (see **Figure 4**).<sup>12</sup>



**Figure 4.** Geographic overview of broiler production in U.S. by state. Image courtesy of USDA APHIS Poultry Industry Manual



## 7.3 Overview of Major Steps in Production of Broilers during Routine Operations

### 7.3.1 Broiler Facility Operations

Generally, farms follow an “all-in, all-out” principle; that is, individual sites manage birds of a single age and load all birds out for slaughter at the same time.

The major steps in broiler production and processing during normal operations are described in the following sections.

#### 7.3.1.1 Broiler House Preparations

Downtime is the period after the old flock is removed and before the new flock arrives, and is generally recommended for at least two weeks.<sup>10</sup> Litter used for brooding successive flocks is referred to as “built up” litter. If new litter material can be acquired at a reasonable price and land is available for disposal of old litter, then most broiler houses are completely cleaned once per year or whenever a disease problem is encountered. Old litter is spread on farm land as a fertilizer and new litter is placed in the house.

After a flock has been removed from a broiler house at the end of a production cycle, litter intended for reuse is “de-caked.” “De-caking” is accomplished by removing the wettest and most compressed parts of litter (top few inches) from the chicken house and then tilling and evenly spreading the remaining litter across the floor in preparation for the next flock. “De-caking” releases trapped ammonia and allows the litter to dry.

Another procedure that is gaining favor is to windrow the used litter down the center of the house and let it compost. Composting is an excellent way of reducing pathogens in litter. After it is spread out again in the house, old litter is usually “top-dressed” by placing 1 to 2 inches of new bedding over it. It is important to maintain proper levels of litter moisture (20-30 percent). Wet litter can result in excessive ammonia production by bacteria in the litter and contributes to development of breast blisters, foot pad ulceration, and excess condemnations at processing. Wet litter can be caused by water line leaks, excess humidity, inadequate moisture removal due to poor ventilation, watery droppings associated with enteritis, high-salt or high-protein rations, poor quality fats, and improper drinker management. When proper litter conditions are maintained, growth will be closer to optimum targets, there are few to no flies and parasites, feed efficiency is improved, and feathering is better.<sup>12</sup>

Complete removal of litter from a production house is typically performed by tractor-mounted box scrapers or front-end loaders. The material can immediately be used for land application or temporarily stored. If stored, it needs to be covered by a well-secured impermeable tarpaulin.<sup>14</sup>

#### 7.3.1.2 Grow-out Period Management

##### 7.3.1.2.1 Receiving the Chicks

Upon delivery of chicks, all personnel involved must follow the farm’s biosecurity guidelines and wear clean uniforms and boots. Often drivers participate in unloading and depositing the chicks in the broiler house. Once empty, the chick boxes should be removed and returned to the hatchery for washing and disinfection, but chick boxes may be used in subsequent deliveries

before washing and disinfecting (J. Smith, personal communication, June 2016). Cardboard chick boxes should be disposed of on farm.



**Figure 5.** Chicks are loaded into boxes at the hatchery to be transported to broiler farms. Photo courtesy of GNP Company

### 7.3.1.2.2 Brooding

Newly hatched chicks are not capable of fully regulating their body temperature until sometime between 1 and 2 weeks of age.<sup>8</sup> Proper brooding temperatures and ventilation must be provided.

Two systems are commonly used to control the temperatures during the brooding period.

- *Whole-room heating system:* The recommended starting air temperature is 88°F with a gradual decrease to 65 to 70°F at week 6 and until they are transported to market. Floor temperature should remain at 83°F or warmer.<sup>8</sup>
- *Localized brooding system:* Because the heating is localized, temperatures need not be exact, as chicks can move toward or away from the heat source to seek a comfortable location. A hover (pancake) or radiant brooder is used in this system. The air temperature should be at 90°F at day one, then reduced by approximately 5°F per week, and held at 70°F until market. Floor temperature can vary from 75 to 140°F, depending on the location in the house.<sup>8</sup>

### 7.3.1.2.3 Brood and Grow Ventilation

Ventilation needs to be controlled to maintain good air quality. Because chicks are susceptible to air quality problems and drafts, maximum ammonia levels and air speed have been established. Draft speed ranges from 0.3 meters per second for 0- to 14-day-old birds to 1.75 to 3.0 meters per second for chicks 28 days and older. If circulation fans are used, they should be directed toward the ceiling to minimize the downward draft.<sup>15</sup>

It should be noted that many producers brood chicks in only a portion of the broiler house to conserve fuel used for heating, especially during cold weather. Curtains are placed to divide the brooding area from the rest of the house and then removed to allow access to the entire house when chicks are between 7 and 14 days of age, depending on the weather.<sup>8</sup>

Grow-out, the second stage of broiler growth, begins when the house dividers are removed and chickens move into the entire house. This period typically begins around 11 to 16 days of age and continues until the broilers are removed for processing. During grow-out, broilers are no longer as fragile and not quite as susceptible to disease or changes in temperature. However, care must continue in the grow-out stage to enable the broilers to reach their maximum potential.<sup>12</sup>



During the growing and finishing period, the environment has to be controlled in order to maintain the biological and social equilibrium of the chicken house. In biological equilibrium, the body weight per square meter must balance the requirements for oxygen, water, and feed. For social equilibrium, temperatures are designed to make chickens position themselves comfortably and take advantage of feed and water.<sup>10</sup> Temperature recommendations were outlined in the brooding section (Section 7.3.1.2.2 Brooding). Daily record keeping is essential to provide an early warning of potential problems. The records should be on display for each house.<sup>15</sup>



**Figure 6.** Whole-room brooding. Photo courtesy of GNP Company

### 7.3.1.3 Load-out

Prior to arrival of crews, feeders, waterers, and other equipment that could injure birds or interfere with crew activities in the barn are raised or removed.<sup>8</sup> Usually, feed is withdrawn about 6 hours before loading, while water is provided up until load-out begins. This timing allows birds to be processed between 8 and 12 hours after last feeding to minimize fecal contamination of carcasses.<sup>16</sup> On some operations, load-out crates or other equipment may be dropped off the day before scheduled load-out to expedite crew activities (D. Halvorson, personal communication, August 2015), while other systems operate on a “just in time” availability to align arrival of crews and equipment (Broiler Working Group [BWG], personal communication, July 2016).

Almost all broilers are caught by hand and loaded into coops, cages, or crates. The coops, cages, or crates are then placed on the live-haul trucks.<sup>8</sup> Catching at night is best, as it diminishes the risk of scratches and suffocation because the broilers are calmer. However, the second shift at processing plants makes it necessary to catch and transport a portion of birds during daylight hours.<sup>8,10</sup>

Temperature permitting, catching crews commonly wear coveralls and boots and consist of 7 to 10 people. (J. Smith, personal communication, June 2016). If mechanical equipment (automatic bird catcher) is used for catching, it must be operated at moderate speeds to prevent damage and stress.<sup>17</sup> The farm does not own



**Figure 7.** Broilers are caught by hand and loaded into crates. Photo courtesy of GNP Company

the automatic bird catcher; the machine travels from farm to farm when catching must be performed. Cages, crates, or coops filled with broilers are loaded onto a live-haul truck by a forklift.



**Figure 8.** Crates filled with broilers are being loaded onto a live-haul truck by a forklift. Photo courtesy of GNP Company

#### **7.3.1.4 Transportation of Broilers to Processing and Awaiting Processing**

Forklifts transfer loaded packing modules to a transport truck that usually holds 20 modules stacked two high. Approximately 6,000 broiler chickens are loaded onto each truck.<sup>12</sup> Once the truck is loaded, it travels to the processing plant without unnecessary or prolonged stops, and the transport time is managed according to the local current guidelines or legislation.<sup>8</sup> Broilers are unloaded in the holding area at the slaughterhouse, which ideally provides protection against heat and extreme cold weather. Trucks used for transporting broilers to the processing plant for slaughter are usually dedicated for this purpose only and are owned by the company.

Although cages and coops may be washed and disinfected after birds are removed at the processing plant, there is no current requirement to do so, and it is not standard practice. In the case of an HPAI outbreak, washing and disinfecting vehicles and equipment after unloading the birds may be very difficult under certain conditions, such as extreme cold, or may create wastewater handling problems (J. Smith, personal communication, June 2016).

At the processing plant, birds must be kept well-ventilated in holding areas. Air is circulated by large fans. Misting may be used if temperatures are high, and curtains may be lowered in cold weather to protect broilers from wind or precipitation. Birds remain on the trailer until ready for processing,



**Figure 9.** Loaded trucks arrive at processing plant. Photo courtesy of GNP Company



when they are removed from crates onto a belt that transports them to the hanging room. Broilers are hung by hand from their legs in shackles that are attached to a moving chain. The area is kept darkened and at some facilities, a bar contacts the breast of the birds to help keep them quiet, reduce wing flapping, minimize distress, and reduce parts hemorrhages, bruising, or broken wings.

**Figure 10.** After unloading broilers at the processing plant, the crates are cleaned and put back onto the truck. Photo courtesy of GNP Company



**Figure 11.** Carcasses move along automated processing line. Photo courtesy of GNP Company.



## 7.4 Overview of Current Disease Prevention and Biosecurity Efforts in Broiler Production

Biosecurity involves procedures that reduce the probability of disease outbreaks and includes two components: (1) bioexclusion (keeping pathogens out) prior to an outbreak, and (2) biocontainment (keeping pathogens from leaving a flock) after an outbreak occurs. Farms with poor biosecurity are vulnerable to diseases, which have the potential to ruin an entire flock. Loss of income from disease can be an enormous financial burden on growers, so the importance of biosecurity cannot be overstated.<sup>12</sup>

### 7.4.1 Current Disease Prevention and Containment Measures in Grow-out Operations during Normal (non-outbreak) Situations

The NPIP is a cooperative industry-state-federal program focused on preventing disease in poultry and promoting safety of poultry products throughout the country. Participation in NPIP provides breeders and hatcheries with standardized guidelines for poultry and egg management, as well as biosecurity practices. NPIP Provisions 9CFR145 and 9CFR147 are pertinent to hatchery and breeder facilities and contain various C&D and biosecurity measures for the production and transportation of hatching eggs. Most commercial broiler breeders and hatcheries participate in the NPIP program. Some of the typical preventive biosecurity measures practiced in the poultry industry currently include: (1) monitoring the health status of flocks, (2) C&D of reusable materials, and (3) segregation of setting, hatching, and chick-processing operations.

Participation by broiler growers in NPIP programs is also widespread. Recently, minimum biosecurity standards for growers were approved at the 43<sup>rd</sup> Biennial NPIP Conference.<sup>2</sup> According to NPIP, the biosecurity program should include a designated Line of Separation (LOS) and Perimeter Buffer Area (PBA), and provisions to address personnel biosecurity practices; control of wild birds, rodents and insects; equipment and vehicle management; mortality disposal; manure and litter management; water supplies and feed; and replacement litter management. How individual producers meet these guidelines is variable, depending on farm layout and resources.

Other biosecurity plans and standards are often set by integrators and company policy. Company veterinarians and staff are responsible for training the contracted grower on company policy; and the grower is responsible for training staff and implementing biosecurity practices on his or her facility (BWG, personal communication, June 2016). The following sections describe some of these practices and recommendations, though they may not be implemented in all facilities.

### 7.4.2 Structural Biosecurity

#### 7.4.2.1 Conceptual Biosecurity

Conceptual biosecurity includes planning grower site development in a way that limits disease transfer. Some key concepts employed in the broiler industry include<sup>12</sup>:

- Locating broiler farms so they are isolated from other premises with domestic poultry.
- Using single-age sites, which help reduce cycling of field and live vaccine viruses.
- Locating roads so as to allow clean vehicles to approach chicken houses and to divert potentially contaminated vehicles away from the houses.

To prevent disease introduction and subsequent transmission to other premises if infection occurs, sanitation and biosecurity measures are used at all farms, though to varying degrees. Some recommended sanitary and biosecurity measures are outlined below.

#### **7.4.2.2 Secured Farm Entry**

- Only essential visitors (such as veterinarians, poultry catchers, repairmen, and service personnel) are allowed in the farm areas.<sup>12,18</sup>
- Non-essential visitors (e.g., sales representatives and tours) are limited and usually allowed onto farm premises only with authorization.<sup>12</sup>
- Farms keep a record of all visitors and their previous farm visits.<sup>10,12,15,19</sup>
- All visitors should wash their hands and put on protective outer clothing, including clean boots and head gear, before working with the flocks.<sup>12,18-20</sup>
- Movements of visitors on the farm should be restricted.<sup>20</sup>
- Any visitors having birds of their own should not be allowed to be near birds at all.<sup>21</sup>
- Farm gates are not standard on broiler farms and doors should be kept locked at all times. Signs at farm entry and entrances to each house should announce that the area is a biosecure zone and unauthorized entry is strictly prohibited.<sup>19</sup>
- A PBA, an outer control boundary around the poultry houses, should be clearly delineated such that nonessential vehicles do not enter into it and personnel do not leave it in the course of their daily tasks.<sup>22</sup>
  - If personnel must enter or exit the PBA, a specified entrance is used.
  - Vehicles entering the PBA must be cleaned and decontaminated (via disinfectant or heat) before entering.
  - Personal vehicles of employees and visitors should be parked in a designated area outside the PBA.
- Facilities should establish an LOS and PBA aimed at reducing the potential for virus to enter and contaminate the production site.<sup>23</sup>

### **7.4.3 Operational Biosecurity**

#### **7.4.3.1 Farm Personnel, Service Technicians, and Visitors**

- Biosecurity training stresses the importance of not owning, and avoiding contact with, other birds not owned by the business, including wild birds, waterfowl, backyard poultry, pet birds, or show birds. Any inadvertent contact by growers or laborers requires a shower and change of clothes before entering farm premises.<sup>19</sup>
- In the case of hunting waterfowl or inadvertent contact with non-company birds, it is suggested that personnel let 72 hours pass before contact with company birds.<sup>12,19</sup>
- Personnel or any visitors entering the PBA should shower and change into clean clothes before arriving on poultry site.<sup>23</sup>

- All visitors, company, and farm personnel should employ a “no shoes touch the ground” method whereby disposable shoe covers are donned before exiting the vehicle.<sup>19</sup>
- Disposable items used during flock visits should be bagged and left on the farm.<sup>12</sup>
- Farm staff should wash hands with soap, water and a disinfectant before entering bird areas and handling birds.<sup>21</sup>
- After returning from a location where birds are present, including a feed store, all equipment, truck tires, clothing, and shoes should be cleaned and disinfected.<sup>21</sup>

#### **7.4.3.2 Feed Delivery**

- Generally, one poultry house requires feed delivery approximately three times per week. In a larger production site, with 10 houses for example, this equates to at least 20 to 30 trips per week. Therefore, distances between farms and the feed mill must be kept at a minimum, while still allowing adequate separation of production sites for disease control and manure management purposes.<sup>8</sup>
- Feed delivery trucks are generally not washed and sanitized between visits to farms (J. Smith, personal communication, June 2016). Drivers should wear clean boots (rubber or disposable) before getting out of the vehicle.<sup>12</sup>
- Any uncovered or spilled feed must be cleaned up and disposed of to avoid attracting waterfowl or pests. Spilled feed may never be fed to broilers as it may have become contaminated on the ground.<sup>19</sup>

#### **7.4.3.3 Sanitation Facilities on Farm**

- Hand-sanitizing facilities and a place to change footwear or well-maintained footbaths are normally provided at the entrance to each poultry house.<sup>19</sup>
- Clean footwear is usually provided to avoid organic material that could inactivate the disinfectant.<sup>19</sup>
- Toilets and hand-washing facilities may or may not be present (J. Smith, personal communication, June 2016).
- Showers and changing rooms are generally not provided on the farms for visitors and employees (J. Smith, personal communication, June 2016).

#### **7.4.3.4 Cleaning & Disinfection (C&D)**

##### **7.4.3.4.1 Vehicles and Drivers**

- Wheel dips or wheel spraying facilities are rarely provided at a broiler farm entrance for all vehicles coming onto the farm (J. Smith, personal communication, June 2016).
- The feed truck driver remains outside of the broiler houses.<sup>10</sup>
- Service technicians and company personnel should spray tires and underside of vehicles with an antimicrobial solution between farm visits.<sup>12</sup>

#### 7.4.3.4.2 Equipment

- Sharing of equipment or supplies with other poultry premises should be minimized. If sharing is necessary, the items should be thoroughly cleaned and free of visible dirt, feathers, manure, and other debris and then disinfected between uses (J. Smith, personal communication, June 2016).
- Equipment should be effectively sanitized between uses; sharing of equipment should be minimized.<sup>23</sup>
- Equipment should be inspected for contamination prior to entry onto farm.<sup>12</sup>
- Equipment that comes in contact with birds or their droppings must be thoroughly cleaned and disinfected before entering broiler houses<sup>19,21</sup> and before being placed back into a vehicle.<sup>12</sup>
- Organic matter such as manure, litter, debris, and feathers must be removed with soap and water before disinfecting.<sup>21</sup>

#### 7.4.3.4.3 Water Supplies

- Water should come from deep wells or sources that have been treated to eliminate any potential contamination with live virus.<sup>22,23</sup>
- If water comes from a surface source for use in cleaning, cool cells (i.e., evaporative cooling systems), or drinking, experts in water treatment should be consulted on how to continuously treat water to eliminate viable virus.<sup>19,23</sup>

#### 7.4.3.4.4 Housing Area

- After the flock is removed, unused feed is then removed from the feed system, including all bins and augers (J. Smith, personal communication, June 2016).
- When cleaning litter from each house, it should be removed by covered vehicles.<sup>15</sup>
- Doors should be secured even when houses are empty and all possible entries for wild birds should be sealed and checked frequently.<sup>12</sup>
- When manure/litter is completely removed from the house, the building must be C&D(using disinfectant or heat) before birds are reintroduced.<sup>22</sup>
- When manure/litter is reused, a modified C&D procedure may be used.<sup>22</sup>

#### 7.4.3.4.5 Load-out and Slaughter Plant

- Transport trucks and load-out equipment should be C&D prior to entry onto the farm,<sup>12</sup> although this is not common practice in some regions (J. Smith, personal communication, June 2016).
- Delivery trucks transporting broilers to slaughter, and crates or containers, are sometimes C&D before loading the broilers.<sup>10</sup>

#### 7.4.3.5 Animal, Pest, Insect Control

- Feed should be stored securely and spilled feed cleaned up immediately to limit attracting waterfowl and pests to the property.<sup>19</sup>

- Operations should have control measures to protect poultry from wild birds, their feces, and their feathers.<sup>23</sup>
- If a pond or surface water source is on the property, steps should be taken to deter waterfowl from entering the area. Surface water should not be used as drinking water, for C&D, or in cool cells on poultry houses. If surface water is used, daily monitoring must be performed to ensure the water is C&D before use.<sup>19</sup>
- Chicken houses should be bird-proof and have insect and rodent control programs in place.<sup>23</sup>
- Dogs and cats should not be allowed in broiler houses.<sup>12</sup>
- Farms should have bait stations around the perimeter of poultry houses that prevent pets and livestock from gaining access to the bait, but provide easy access for rodents. Insect control should also be in place. Rotation of bait is essential.<sup>12</sup>

#### **7.4.3.6 Dead Bird Disposal**

Disposal of dead birds is regulated by local, state, and federal governments to control the impact of carcass disposal on air quality, water quality, and the spread of disease. Disposal of mortality is a daily necessity since dead birds can harbor pathogenic microorganisms with potential transmission to other poultry. Cost of supplies, labor reliability, maximum anticipated daily mortality, and degree of biosecurity associated with each method must be assessed.<sup>12</sup>

- Composting requires construction of a solid floor and covered roof system that has adequate primary and secondary bin capacity to meet the predicted mortality level for the farm. The area should be managed so carcasses are covered to prevent access by wild animals and to maintain adequate temperatures for composting.<sup>12</sup>
- If incineration or rendering is used, stored carcasses should be protected to prevent exposure to wild animals.<sup>12,23</sup>
- Trucks moving manure or dead birds should be covered and follow a designated approved route. Trucks should be cleaned and disinfected after deliveries and before entering another farm.<sup>12</sup>
- Disposal methods should avoid the potential for cross-contamination with dead birds from other facilities.<sup>23</sup>

#### **7.4.3.7 Manure and Litter Management**

- Manure and spent litter should be removed in a manner that prevents exposure of susceptible poultry, either on or off the farm of origin, to disease agents.<sup>23</sup>
- Fresh litter should be stored and handled so it cannot be contaminated by insects, wild birds, or rodents.<sup>23</sup>



## 8 Hazard Identification: HPAI overview

Hazard identification consists of listing the pathogenic agents associated with the species from which a commodity is derived and whether the agents can be classified as hazards for further consideration in the risk assessment.<sup>24</sup> For movement of broilers to market, the pathogenic agent of concern is HPAI virus. Properties of HPAI viruses, including environmental persistence, transmission characteristics, and physical and chemical inactivation, have been extensively reviewed in comprehensive texts.<sup>25</sup> This section is a brief summary of the key properties of HPAI viruses from published scientific literature and expert opinion, with emphasis on the variability between HPAI virus strains and transmission characteristics in poultry.

### 8.1 Agent

AI viruses are negative-sense, segmented, ribonucleic acid viruses of the family *Orthomyxoviridae*. The *Orthomyxoviridae* family includes several segmented viruses including the Type A, B, and C influenza viruses. The Type A influenza viruses, which include all AI viruses, can infect a wide variety of animals including wild ducks, chickens, turkeys, pigs, horses, mink, seals, bats, and humans. The type B and C viruses primarily infect humans and occasionally pigs.<sup>25-27</sup>

Two surface glycoproteins of the influenza A virus, hemagglutinin (HA) and neuraminidase (NA), are the most important antigenic sites for the production of protective immunity in the host; however, these proteins also have the greatest variation. For AI viruses there are sixteen known different subtypes of HA (H1 to H16), nine known different subtypes of NA (N1 to N9), and 144 different HA:NA combinations (H17N10 and H18N11 were only recently isolated from bats).<sup>25,27</sup> Although relatively few of the 144 subtype combinations have been isolated from mammalian species, all subtypes, in the majority of combinations, have been isolated from avian species.

#### 8.1.1 Definition of Highly Pathogenic Notifiable Avian Influenza

For the purpose of disease control programs and international trade in domestic poultry products, HPAI is defined in the Code of Federal Regulations, Title 9, Section 53.1 as<sup>28</sup>:

- (1) Any influenza virus that kills at least 75 percent of eight 4- to 6-week-old susceptible chickens [or six out of eight birds], within ten days following intravenous inoculation with 0.2 ml of a 1:10 dilution of a bacteria-free, infectious allantoic fluid;
- (2) Any H5 or H7 virus that does not meet the criteria in paragraph 1 of this definition, but has an amino acid sequence at the hemagglutinin cleavage site that is compatible with HPAI viruses; or
- (3) Any influenza virus that is not an H5 or H7 subtype and that kills one to five [out of eight inoculated] chickens and grows in cell culture in the absence of trypsin.

The World Animal Health Organization (OIE) Terrestrial Animal Health Code Article 10.4.1 defines HPAI viruses to be AI viruses that “have an IVPI [intravenous pathogenicity index] in six-week-old chickens greater than 1.2 or, as an alternative, cause at least 75 percent mortality in four-to eight-week-old chickens infected intravenously. H5 and H7 viruses which do not have an IVPI of greater than 1.2, or cause less than 75 percent mortality in an intravenous lethality test, should be sequenced to determine whether multiple basic amino acids are present at the cleavage

site of the haemagglutinin molecule (HA0); if the amino acid motif is similar to that observed for other high pathogenicity avian influenza isolates, the isolate being tested should be considered as high pathogenicity avian influenza virus.<sup>29</sup>

All H5 or H7 virus isolates of both low and high pathogenicity, and all HPAI virus isolates regardless of subtype, are reportable to state and national veterinary authorities and to the OIE.<sup>30</sup> Although other low-pathogenic avian influenza (LPAI) viruses may cause considerable morbidity and production losses, they are not reportable diseases to the OIE (but may be reportable in some states).

### 8.1.2 Host Range

Wild waterfowl are considered the natural reservoirs of LPAI viruses, but the role of wild birds as reservoirs for most HPAI viruses responsible for high mortality in domestic birds is not fully elucidated.<sup>31</sup> Recent surveillance and phylogenetic analyses, however, suggest that migratory waterfowl are important in the maintenance, reassortment, and spread of HPAI viruses.<sup>32-34</sup> The phrase “highly pathogenic for chickens” does not indicate or imply that the AI virus strain is highly pathogenic for other bird species, especially wild ducks or geese (Anseriformes). However, if a virus is highly pathogenic for chickens, the virus will usually be highly pathogenic for other birds within the order Galliformes, family Phasianidae, such as turkeys and Japanese quail.

Most HPAI viruses for chickens are generally non-pathogenic for ducks and geese in experimental studies.<sup>26</sup> However, the lethality of HPAI viruses has changed since the re-emergence of H5N1 HPAI viruses in Hong Kong in 2002, as some strains have become highly lethal in some naturally and experimentally infected waterfowl.<sup>31</sup> The evolving H5 HPAI viruses spread throughout Asia and Europe between 2005 and 2014.<sup>35</sup> In late 2014, the Eurasian H5 clade 2.3.4.4 viruses were detected in North American wild birds,<sup>33,36,37</sup> reassorted with American AI viruses, and similar Eurasian/American HPAI H5 viruses were identified during the domestic poultry outbreak in 2015 in the United States.<sup>38</sup>

Characterization of the Eurasian/American HPAI H5 viruses found in wild birds was done by the National Wildlife Health Center and USDA National Veterinary Services Laboratory. Researchers at these agencies suggest identifying these HPAI H5 viruses as intercontinental group A (icA) to differentiate this changing subset of viruses from other Asian H5N1 HPAI.<sup>36</sup> Some wild birds—including ducks and geese—that were found to be positive for icA H5N8 and icA H5N2 exhibited morbidity/mortality at the time of sample collection.<sup>39</sup> Experimentally, both strains—H5N8 (A/GF/WA/14) and H5N2 (A/NP/WA/14)—led to some mortality in domestic geese (Chinese geese) but not in domestic ducks (Pekin) (Pantin-Jackwood, personal communication, August 2016). An icA HPAI H5N2 strain isolated from infected turkeys in Minnesota in 2015 (A/Tk/MN/12582/2015) was experimentally inoculated into mallard ducks and caused mortality in individual birds in each group at medium ( $10^4$ ) and high ( $10^6$ ) inoculation doses, with a mean death time of 9 days.<sup>40</sup> Thus, the host range affected by icA H5 viruses is broad and the clinical signs in each host are variable.

HPAI strains are known to emerge in poultry after the introduction of LPAI viruses from wild birds, and after circulation of virus for varying lengths of time in domestic poultry.<sup>41</sup> This is likely what occurred in the U.S. turkey industry in early 2016 when the first HPAI caused by an H7N8 virus (in any species), A/turkey/Indiana/2016, was detected in commercial turkeys.<sup>42</sup>

Subsequent detections of H7N8 LPAI occurred on additional turkey premises; all HPAI and LPAI viruses were found to be of North American wild bird lineage. The 2008 identification of an H5N2 virus with an HPAI genotype—with evidence of non-lethal infection in wild waterfowl and without evidence of prior extensive circulation in domestic poultry—suggests that some AI strains with potential high pathogenicity for poultry could be maintained in a wild waterfowl community prior to introduction.<sup>31</sup>

Host adaptation is a key determinant of the ability of an HPAI virus to maintain transmission within domestic poultry. Once adapted to gallinaceous birds, HPAI viruses are unlikely to circulate again among wild birds because they are adapted to poultry.<sup>43</sup> However, the emergence of Asian-origin HPAI H5 strains has led to increased uncertainty regarding the role of wild birds as reservoirs in the maintenance of HPAI viruses in nature.<sup>32,44</sup> Prior to the outbreak of HPAI H5N1 virus in Europe, Asia, and Africa starting in late 2003, HPAI viruses had only rarely been isolated from wild birds—usually associated with outbreaks in domestic poultry—with one exception: An outbreak of HPAI H5N3 (A/Tern/South Africa/1961) in South Africa in 1961 was observed in a population of terns<sup>45</sup>. Now, Eurasian HPAI H5 strains have been isolated from multiple species of wild birds, both from healthy birds and from sick, moribund, or dead birds.<sup>46-48</sup> However, despite extensive global wildlife surveillance efforts, infection with H5N1 HPAI viruses has not been detected in healthy wild birds, except for a few isolated cases.<sup>45,48</sup> The significance of wild birds as a source of infection and their influence on the epidemiology of HPAI viruses are yet to be fully established.<sup>31,47</sup>

Additional hosts also may play a role in the epidemiology of these viruses as they continue to spread and reassort. Experimental studies have shown that various LPAI and HPAI viruses can infect and replicate in multiple mammalian species (e.g., cats, ferrets, mink, pigs, rabbits, raccoons, skunks).<sup>49-52</sup> Several species of concern (e.g., wild animals that may have contact with commercial poultry premises such as rabbits, skunks, and raccoons) have been shown to be capable of shedding AI virus and, in some cases, of experimentally transmitting the virus to ducks via indirect contact (under conditions meant to simulate contact in a natural environment).<sup>51,53</sup>

## 8.2 Geographic Distribution of H5 and H7 HPAI

- The current list of all confirmed affected countries with H5 or H7 infection in animals is maintained by the OIE at <http://www.oie.int/en/animal-health-in-the-world/update-on-avian-influenza/2015/>.<sup>28</sup>
- A recent publication graphically displayed the outbreaks of HPAI virus, H5 subtype, that occurred in the United States in 2014-2015 both in relation to time and to poultry distribution and wild bird migratory patterns; the maps can be viewed at [http://wwwnc.cdc.gov/eid/article/22/1/15-1053\\_article#tnF1](http://wwwnc.cdc.gov/eid/article/22/1/15-1053_article#tnF1).<sup>54</sup>
- The Global Early Warning System for Major Animal Diseases Including Zoonosis (GLEWS)—a joint effort of the Food and Agriculture Organization of the United Nations (FAO), OIE, and the World Health Organization (WHO)—provides a regular update on global H5N1 HPAI events in the Global Animal Disease Intelligence Report, which can be viewed at <http://www.glews.net/>.<sup>55</sup>

### 8.3 Virus Shedding

HPAI viruses have been isolated from respiratory secretions, blood, feces, and feathers, as well as the eggshell surface, albumen, yolk, meat, and other tissues (e.g., spleen and lung) from infected poultry. Estimates of HPAI virus concentrations in chicken and turkey secretions, feces, feathers, and other tissues generally range between  $10^3$  and  $10^7$  EID<sub>50</sub> per gram or per milliliter, although higher concentrations have been observed in some cases.<sup>56-64</sup>

H5N2 HPAI (A/chicken/Pennsylvania/1370/1983) viruses have been isolated from the eggshell surface, yolk, and albumen of eggs laid by experimentally inoculated hens.<sup>65</sup> In these experimental studies, H5N2 HPAI viruses were not recovered from eggs laid on the first day post-inoculation of hens. This may have been due to the developing egg being protected from exposure in the shell gland (uterus) during the later stages of eggshell formation (about 15 hours), in combination with the latently infected period of at least 6 hours in individual birds in this study. In contrast, HPAI virus was recovered from the yolk and albumen of eggs forming in the oviduct of dead chickens at postmortem, 35 to 37 hours after being experimentally infected with an HPAI virus strain (Dutch East Indies) isolated from chickens.<sup>66</sup> Italian HPAI H7N1 (A/chicken/Italy/445/99) viruses have also been isolated from eggs laid by infected hens.<sup>67</sup>

In an experimental study, the concentration of H5N2 HPAI (A/chicken/Pennsylvania/1370/1983) virus ranged from 0.97 to  $10^{5.9}$  EID<sub>50</sub>/eggshell; from 0.97 to  $10^{6.1}$  EID<sub>50</sub>/ml in albumen; and from 0.93 to  $10^{4.8}$  EID<sub>50</sub>/ml in yolk of eggs laid by infected hens.<sup>65</sup>

As compared to chickens, AI viruses in turkeys demonstrate a relatively high degree of affinity for oviduct tissue, relative to respiratory and digestive tissue.<sup>68</sup> A predilection for replication within these tissues may explain the precipitous drops in egg production reported in turkey breeder hen flocks during natural outbreaks.<sup>69-72</sup> Narayan et al. (1969) recovered AA 5-turkey/Ontario 7732/66 HPAI virus from the yolks of each of three eggs laid by 30-week-old turkey hens that were infected through contact with a hen experimentally infected with an HPAI virus.<sup>73</sup> In turkey breeder hens experimentally inoculated with swine-origin LPAI H3N2 (A/turkey/Ohio/313053/04), virus was recovered from eggshells and egg contents.<sup>68</sup> In this study, the percentage of viral detection on shell surfaces was significantly higher ( $P < 0.005$ ) than in albumen, when shell-less eggs were excluded from the analysis.

### 8.4 Chemical and Physical Inactivation

AI viruses are inactivated by physical factors such as heat, extremes of pH, hyper-isotonic conditions, and dryness; however, their infectivity can be maintained for several weeks under moist, low-temperature conditions.

Due to their lipid envelope, AI viruses are relatively sensitive to disinfection agents and inactivation by lipid solvents such as detergents. The EPA maintains a list of disinfectants with label claims for AI viruses. These products include halogens, aldehydes, quaternary ammoniums, phenols, alcohols, peroxides, and some detergents.<sup>74-76</sup> To ensure effective disinfection, appropriate operational conditions as recommended by the manufacturer have to be maintained. Operational conditions such as disinfectant concentration, temperature, contact time, pH, and organic load may impact the degree of virus inactivation.

## 8.5 Persistence of HPAI Virus in Manure and Other Media

Persistence of AI viruses at various humidity levels and temperatures and on various substrates is summarized in Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates. The HPAI virus shed by infected birds may be protected environmentally by accompanying organic material that shields the virus particles from physical and chemical inactivation. Specific environmental conditions such as cool and moist conditions increase survival times in organic media and on surfaces. For example, H5N2 virus (A/chicken/Pennsylvania/1370/1983) remained viable in wet poultry manure in a barn up to 105 days following bird depopulation in the Pennsylvania 1983-1984 outbreak (presumably in winter under freezing conditions). Experimentally, an HPAI strain from this outbreak survived for at least 35 days under moist conditions, but only 9 to 21 days under dry conditions at 4° C (39 °F).<sup>77,78</sup> H5N1 HPAI virus was viable in allantoic fluid for ten days at 25 to 32 °C (77 to 90 °F) when kept out of direct sunlight, but was killed within 30 minutes of placing the sample in sunlight (32 to 35 °C; 90 to 95 °F).<sup>79</sup>

## 8.6 Transmission

Contact with migratory waterfowl, water birds, or shore birds is a risk factor for introduction of AI virus into domestic poultry populations.<sup>80</sup> Because AI virus can be isolated in large quantities from feces and respiratory secretions of infected birds, an important mode of transmission is the mechanical transfer of infective feces.<sup>25</sup> Once introduced into a flock, AI virus can spread directly from flock to flock by movement of infected birds and indirectly via contaminated equipment, egg flats, feed trucks, off-site mortality disposal, garbage trucks, service crews, or other means. Windborne transmission may occur when farms are closely situated and appropriate air movement exists.<sup>81,82</sup> Wild animals such as raccoons and foxes have also been implicated in local area spread; some wild animals, specifically skunks and cottontail rabbits, have been shown experimentally capable of transmitting virus to birds via indirect contact through shared environments.<sup>53,83</sup> Other mechanisms of transmission are outlined below.

### 8.6.1 Vertical transmission in chickens and turkeys

Evidence of vertical transmission of AI virus from infected hens to day-old chicks or turkey poults has been lacking thus far, as most strains are lethal to embryos.<sup>84-87</sup> Groups of turkey hens in egg production, with no clinical evidence of influenza A virus infection, were inoculated intravenously, or intratracheally, or were inseminated with semen contaminated with two LPAI viruses (T/Calif/meleagrium/64, T/Calif/5142/66), and virus was not recovered from poults hatched from eggs laid by exposed turkey hens.<sup>88</sup> Chicks hatched from eggs produced by two broiler breeder flocks infected with HPAI H7N3 (A/Chicken/Canada/AVFV2/04) tested negative for AI during an outbreak in British Columbia in 2004. The outbreak report of the Canadian Food Inspection Agency states, “Because avian influenza does not survive long at incubator temperatures, day-old chicks are not a likely source of infection for broiler growers.”<sup>89</sup> In the 1983 Pennsylvania HPAI H5N2 (A/chicken/Pennsylvania/1370/1983) outbreak, eggs from four severely infected layer breeder flocks were incubated and assayed for AI virus. None of the dead embryos yielded HPAI virus in this study.<sup>90</sup> Also, the 214 chicks hatched from these eggs showed no sign of AI disease and had not developed AI antibodies.<sup>90</sup>

Transmission of HPAI or LPAI viruses from infected breeder flocks to day-old poults via hatchery dissemination has not been observed in previous outbreaks. Turkey industry



veterinarians and AI experts have stated that although there have been several LPAI outbreaks in the United States, vertical transmission or hatchery transmission has not been observed.<sup>91</sup> In a small-scale survey conducted by the University of Minnesota, turkey industry representatives provided reports of 26 flocks that had undergone avian and other influenza A virus infections and where eggs from those flocks were set and not removed from incubation.<sup>91</sup> There was no evidence of horizontal or vertical transmission of AI within the hatchery to day-old poults in any of these instances.

### **8.6.2 Transmission via artificial insemination in turkeys**

As compared to chickens, there is an additional risk in turkeys of viral transmission via the artificial insemination process. It is not anatomically or practically possible to collect semen without the collection device touching the cloaca; semen could also be contaminated during the semen preparation process in the laboratory. Contaminated fomites, such as hands or equipment of insemination crews and contaminated turkey semen, have been implicated in the spread of AI viruses between commercial turkey breeder operations and to commercial turkeys from humans.<sup>92-95</sup> Although semen was implicated in the spread of AI in field outbreak investigations, isolation of AI virus from tom turkey semen was not reported in these studies.

AI virus has previously been isolated from tom turkey semen, but titer levels were not reported.<sup>96</sup> It was unclear whether this virus came from the semen *per se*, or from the cloaca contaminated by fecal material.<sup>97</sup> Other studies have demonstrated that AI viruses can be transmitted to turkey breeder hens through artificial insemination with semen experimentally contaminated with AI virus on the day of collection.<sup>98</sup> Pantin-Jackwood et al. (2010) transmitted pandemic H1N1 (A/Chile/3536/2009) virus to hens by intracloacal or intrauterine inoculation, demonstrating that transmission is possible through contamination of these mucosal surfaces by semen or fomites.<sup>99</sup>

H5N1 virus antigen has been observed in testes, suggesting that virus could be present in semen.<sup>100</sup> In a 2013 study, tom turkeys were inoculated intranasally with  $10^6$  TCID<sub>50</sub>/0.5ml of triple-reassortant H3N2 influenza A virus of swine (IAV-S) A/Turkey/OH/313053/2004.<sup>101</sup> Low viral titers were detected in the reproductive tract (testicles, epididymis, vas deferens, and phallus) and semen by real-time reverse transcription polymerase chain reaction (rRT-PCR), but virus isolation was unsuccessful. The authors suspect that the low virus titers and/or the seminal environment may have adversely affected virus isolation. Nonetheless, based on the presence of viral RNA in the reproductive tract and semen, there remains a potential for venereal transmission of influenza virus in turkeys.

## 8.7 Dose Response

### 8.7.1 Dose Response in Turkeys

Both intraocular and intranasal inoculation were used in an experimental study of infectious and lethal doses of two HPAI strains in turkeys.<sup>102</sup> In this study, turkeys were inoculated with H5N1 (A/turkey/Turkey/1/05) and H7N1 (A/ostrich/Italy/984/00) strains, and all birds shown to be infected died. The ID<sub>50</sub> and LD<sub>50</sub> were thus equal; the median was 10<sup>1</sup> EID<sub>50</sub> (or less) for H5N1 and 10<sup>2.2</sup> EID<sub>50</sub> for H7N1. Turkeys were found to be more susceptible than chickens by over 200-fold for both H5N1 and H7N1.

In another study, turkeys were inoculated with different doses of A/ostrich/Italy/984/2000 H7N1 HPAI by a combined intranasal/intraocular route.<sup>103</sup> Although ID<sub>50</sub> and LD<sub>50</sub> were not explicitly measured, the latter can be extrapolated from their data and was shown to be both dose- and time-dependent. There was no mortality with 10<sup>1</sup> EID<sub>50</sub> by 7 days post-inoculation (PI), while there was greater than 50 percent (4/5) mortality with 10<sup>6</sup> EID<sub>50</sub> at 48 hours PI. At 72 hours PI, the LD<sub>50</sub> was 10<sup>3</sup> EID<sub>50</sub>, and it was 10<sup>2</sup> EID<sub>50</sub> by 96 hours PI.

In their studies using a highly poultry-adapted LPAI strain (A/turkey/Ohio/313053/04), Pillai et al. (2010) demonstrated a markedly lower ID<sub>50</sub> for turkeys (10<sup>1.4</sup> EID<sub>50</sub>) than for chickens (10<sup>2.6</sup> EID<sub>50</sub>).<sup>68</sup> They cautioned that virus strain as well as genetic make-up of the study birds may affect the minimum infectious dose, such that it may not be possible to generalize results from a few isolates in a certain breed of turkey.

As stated above, the infectious dose for turkeys through intranasal inoculation for HPAI viruses (H5N1 and H7N1) has been found to be 2 to 3 logs lower than that for chickens.<sup>102</sup> Given a 50 percent chicken infectious dose of 5 to 6 log EID<sub>50</sub> for aerosol transmission from the dose-response models, it is possible that the turkey infectious dose is between 3 and 4 log EID<sub>50</sub>. Transmission of LPAI (A/turkey/Wisconsin/1966) to turkeys has been demonstrated via an estimated aerosol dose between 3 and 4 log EID<sub>50</sub>.<sup>104</sup> Data from this experimental study suggests that the 50 percent aerosol infectious dose is close to or less than 3 to 4 log EID<sub>50</sub>.

HPAI infection via the gastric route is not well-documented in turkeys. In one small study, 50-day-old turkeys were inoculated by the direct esophageal route with A/turkey/Italy/4580/1999 HPAI H7N1 in a dose of 2 grams of 10<sup>3.6</sup> EID<sub>50</sub>/0.1g infective meat homogenate (for a total dose of 10<sup>4.9</sup> EID<sub>50</sub>).<sup>62</sup> Tracheal and cloacal swabs collected up to day 7 remained negative, as did serum samples up to day 21, and no clinical signs were observed. These results imply that the infective dose for HPAI via esophageal inoculation is likely more than 20 times 10<sup>3.6</sup> EID<sub>50</sub>. However, since the choanal cleft was bypassed, no inference can be made as to the infective dose with exposure that may occur through natural feeding.

Although transmission of HPAI via artificial insemination is strongly suspected in turkeys, data on dose response to such exposure are lacking.

### 8.7.2 Dose Response in Chickens

Most experimental studies in chickens used intranasal inoculation as an entry point. For the intranasal route, in one study, the 50 percent chicken infectious dose (CID<sub>50</sub>) for 11 H5 and H7 HPAI strains (of chicken and turkey origin) varied between 10<sup>1.2</sup> and 10<sup>4.7</sup> EID<sub>50</sub> with a geometric mean of 10<sup>2.9</sup> EID<sub>50</sub>.<sup>105</sup> All but one strain (A/chicken/Rostock/1934 HPAI H7N1, which was endemic in Europe in the early 1900s) in this study had a mean CID<sub>50</sub> above 10<sup>2</sup>

EID<sub>50</sub> with strains less adapted to chickens having the higher CID<sub>50</sub> values. Other studies have also found similar estimates for the CID<sub>50</sub> through the intranasal route, with higher CID<sub>50</sub> values indicating a lack of adaptation for infection in chickens.<sup>64,105,106</sup> The initial cases in wild birds in the U.S. with Eurasian HPAI H5N8 (A/GF/WA/14) and reassortant H5N2 (A/NP/WA/14) viruses had high CID<sub>50</sub> values (i.e., near or above 10<sup>4.7</sup>) and thus were likely poorly adapted to chickens, possibly explaining why poultry outbreaks were limited in the Pacific flyway during the 2014-2015 outbreak.<sup>64</sup>

Single-hit dose-response models (e.g., exponential) have been used for HPAI virus in chickens and mammals.<sup>107,108</sup> These models assume that each virion has the capacity to independently act and cause infection in the host. Dose-response models enable us to estimate the probability of infection when a bird is exposed to a dose different from the 50 percent infectious dose. For example, given a CID<sub>50</sub> less than 10<sup>2.82</sup> EID<sub>50</sub>, a chicken exposed to 10 EID<sub>50</sub> would have a 1 percent chance of infection according to the single-hit exponential dose-response model.

Given limited data, there is greater uncertainty regarding the infectious dose for other routes such as oral consumption of infected material. Kwon and Swayne (2010) found a substantially higher 50 percent infectious dose for HPAI H5N1 (A/Whooper Swan/Mongolia/244/) via oral consumption of chicken meat (10<sup>7</sup> EID<sub>50</sub>) or drinking of contaminated water (10<sup>6.7</sup> EID<sub>50</sub>).<sup>109</sup> However, in this study, a group of three to five chickens were fed contaminated meat with a single virus concentration, and details regarding the uncertainty in the estimates were not provided. The study also found higher infectious doses for the intragastric inoculation route by gavage (10<sup>6.2</sup> EID<sub>50</sub> for liquid and 10<sup>7.4</sup> EID<sub>50</sub> for meat) compared with the intranasal route.

In Swayne and Beck (2005), feeding of finely chopped meat from chickens infected with H5N1 HPAI viruses at higher doses (10<sup>7.8</sup> EID<sub>50</sub>/bird) resulted in transmission of H5N1 HPAI (A/chicken/Korea/ES/2003) virus.<sup>110</sup> However, feeding of HPAI H5N2- (A/chicken/Pennsylvania/1370/1983) infected chicken breast or thigh meat to Specific Pathogen Free (SPF) chickens at lower doses (10<sup>3.5-3.6</sup> EID<sub>50</sub>/bird) did not produce infection. The authors reasoned that lack of direct exposure of the respiratory tract (i.e., minced meat likely did not pass through the choanal cleft and contact nasal surfaces) could explain the lack of infection in H5N2 trials with lower doses. Moreover, a reference is made to a feeding trial by Purchase et al. (1931), in which 0.5g of blood fed to chickens resulted in HPAI transmission whereas feeding 5 g of meat did not, suggesting that transmission is more likely if a feedstuff is conducive to passage into the nasal cavity.<sup>111</sup> However, in the Purchase et al. study, the HPAI concentration in blood was not estimated, and it may have been sufficient to cause infection via the intragastric route.

Sergeev et al. (2013) found a CID<sub>50</sub> for H5N1 HPAI (A/Chicken/Suzdalka/Nov-11/2005) virus of 10<sup>3.9</sup> EID<sub>50</sub> for oral inoculation and 10<sup>5.2</sup> EID<sub>50</sub> for intragastric inoculation via gavage tube.<sup>112</sup> The authors suggested contamination of the nasal mucosal membranes from the oral cavity via the choanal slit as a possible internal mechanism for transmission via the fecal-oral route.

There is considerable uncertainty regarding the infectious dose via the aerosol route. Direct aerosol data from Spekrijse et al. (2013) suggest very low transmission rates, even after 24 hours of exposure to H5N1 HPAI (A/turkey/Turkey/1/2005) virus in a concentration of more than 10<sup>3</sup> EID<sub>50</sub>/m<sup>3</sup> in air coming from a room housing infectious chickens.<sup>113</sup> When we fit exponential and logistic dose-response models to data from Spekrijse et al. (2013), maximum likelihood estimation suggested a CID<sub>50</sub> for the aerosol route between 5 and 6 log EID<sub>50</sub>.<sup>113</sup> An



estimate of 5 to 6 log EID<sub>50</sub> is more consistent with the lower transmission rates for AI observed between chickens housed in adjacent cages in most studies.<sup>114</sup>

Sergeev et al. (2013) found considerably lower CID<sub>50</sub> estimates (approximately 1 log EID<sub>50</sub>) for various HPAI H5N1 strains when susceptible chickens were exposed to 0.5- to 2- $\mu$ m diameter aerosols generated from liquid contents of HPAI-infected embryonating eggs.<sup>112</sup> The results from this paper are not consistent with other studies that indicate lower aerosol transmission between infected and susceptible chickens housed in adjacent cages, and are also not consistent with data published in Spekrijse et al. (2013).<sup>113</sup> A possible explanation for the differences between this study and Spekrijse et al. (2013) is that the characteristics of 0.5- to 2- $\mu$ m diameter contaminated aerosols generated by nebulizing embryonating egg contents are different from naturally contaminated aerosols emanating from a chamber with infectious chickens. In addition, Spekrijse et al. (2013) allow that the viral titer determined by RT-qPCR includes inactivated virus, such that the titer of viable virus in the air sample was actually lower.<sup>113</sup>

### 8.7.3 Route of Entry and 50 Percent Infectious Dose Estimate Used in this Assessment

In poultry, the choanal cleft (palatine fissure)—located on the roof of the mouth—is a papillae-lined, narrow slit that connects the oral and nasal cavities. During mastication or drinking, contents of the oral cavity may pass through this slit and contact the mucosal surfaces lining the nasal cavity.

Because of the variability in the susceptibility of different tissues to infection with HPAI virus (intranasal vs. intragastric) observed in laboratory inoculation and experimental feeding trials, there is considerable uncertainty as to the infectious dose that is appropriate for natural exposure via feeding of contaminated materials. The route of entry impacts the dose-response parameters in the exposure assessment.

We had obtained expert opinion regarding the appropriate route of entry and associated infectious dose (intranasal or intragastric) that best represents oral exposure in chickens, given the limited data on this aspect.<sup>115</sup> Experts stated that it is reasonable to assume that transmission may occur if contaminated food or water were to pass through the choanal cleft into the nasal cavity. Therefore, due to the limited studies on exposure via natural feeding of contaminated materials and the associated uncertainty, we conservatively assumed that transmission of HPAI viruses through consumption of contaminated materials might occur with exposure to doses infectious for the intranasal route, in turkeys as well as in chickens.

## 8.8 Latently Infected and Infectious Periods

In individual birds, the incubation period is dependent on the dose, route of exposure, and individual host susceptibility. At the flock level, detection is highly dependent on the level of clinical signs and the ability of the grower to detect them.<sup>116</sup> For trade purposes, the OIE defines the flock incubation period as 21 days.

### 8.8.1 Latently Infected and Infectious Periods in Turkeys

The latently infected and infectious periods may vary considerably with HPAI strain and turkey breed. Saenz et al. (2012) estimated the mean infectious period for HPAI H7N1 (A/ostrich/Italy/984/00) in turkeys to be 1.47 days (95% CI [confidence interval], 1.3 to 1.7)

from experimental transmission studies.<sup>29,117</sup> The data from this study also suggested that the latent period for HPAI H7N1 in turkeys is likely less than 16 hours.<sup>117</sup>

Aldous (2010) evaluated the virus shedding patterns and mortality in turkeys inoculated with various doses of HPAI H5N1 (A/turkey/Turkey/1/05) virus.<sup>102</sup> Analysis of these data indicated a mean latent period of 1.27 days (std.dev. 0.40 days) and a mean infectious period of 1.28 days (std.dev. 1.17 days).<sup>102,118</sup> Further details on the estimation of these parameters are provided in Appendix 2 of Turkey Hatching Eggs Risk Assessment.<sup>119</sup>

## 8.8.2 Latently Infected and Infectious Periods in Chickens

Latent and infectious periods have been documented for multiple HPAI virus strains, and periods may vary depending on virus strain and chicken type used in experimental conditions. Using 6-week-old SPF white Leghorn chickens, Van der Goot et al (2005) determined an infectious period 6.3 days (95% CI 3.9-8.7 days) when birds were inoculated with HPAI H7N7 (A/Chicken/Netherlands/621557/03).<sup>120</sup> In another experiment also using 6-week-old SPF white Leghorn chickens, this time inoculated with HPAI H5N2 (A/Chicken/Pennsylvania/1370/83), the mean infectious period was 6.8 days (95% CI 4.91-8.69 days) and a modeled latent period was 1-2 days.<sup>121</sup>

Mean time to death (which includes both latent and infectious periods) was observed in 2- to 4-week-old SPF white Leghorn chickens using multiple strains of HPAI H5N1. Death was observed in 100 percent of birds in less than 36 hours when inoculated with one of the four following strains: DK/Vietnam/201/05, DK/Vietnam/206/05, DK/Vietnam/207/05, Muscovy DK/Vietnam/213/05. Mean time to death was estimated at less than 48 hours for DK/Vietnam/218/05 and at 48 hours for DK/Vietnam/203/05.<sup>122</sup>

In a study using 4-week-old SPF chickens of a layer breed inoculated with HPAI H5N1 (A/Chicken/Legok/2003), researchers reported a mean latent period of 0.24 days (95% CI 0.099-0.48 days) and a mean infectious period of 1.6 days (95% CI 0.90-2.5 days).<sup>123</sup>

In 5- and 8-week-old broiler chickens inoculated with 2015 EA/AM HPAI H5N2 (Tk/MN/2015), a mean time to death of 4.8 and 3.2 days was observed, respectively.<sup>124</sup>

## 8.9 Clinical Signs

The presence and severity of clinical signs of HPAI infection depend on the virus strain and bird species affected.<sup>43</sup> Infected wild and domestic ducks may be asymptomatic, whereas clinical signs in gallinaceous poultry are usually severe, resulting in high mortality.<sup>125</sup> In chickens and turkeys, the clinical signs associated with HPAI infection include marked lethargy with ruffled feathers, lack of appetite, excessive thirst, neurological signs (e.g., tremors, torticollis, opisthotonos, etc.), decreased egg production, soft-shelled or misshapen eggs, respiratory signs (coughing and sneezing), watery diarrhea, or sudden, unexpected death.<sup>91,125</sup> Mature chickens frequently have swollen, cyanotic combs and wattles, and edema surrounding the eyes.<sup>125</sup> In turkeys, a cessation in flock vocalization ("cathedral syndrome") often accompanies infection.<sup>92</sup> Progressive somnolence, reduction of normal vocalization, swollen sinuses, oculonasal discharge, edema of the face, and hemorrhages on the shanks are other clinical signs observed in turkeys.<sup>116,118,126</sup>

The mortality rate in an infected flock can reach 100 percent.<sup>127</sup> In mature birds, gross lesions on necropsy may consist of subcutaneous edema of the head and neck; fluid in the nares, oral cavity, and trachea; congested conjunctivae and kidneys; and petechial hemorrhages which cover the abdominal fat, serosal surfaces, peritoneum, and surface under the keel.<sup>91,125</sup> In layers, the ovary may be hemorrhagic or degenerated and necrotic.<sup>128</sup> Ruptured ova have been reported in layers and broiler and turkey breeders; the peritoneal cavity may be filled with yolk from ruptured ova, causing severe peritonitis in birds that survive long enough.<sup>125</sup> In addition, most HPAI viruses can cause necrosis of the pancreas;<sup>91</sup> all species of birds affected in the 1999-2001 H7N1 HPAI outbreak in Italy had lesions at necropsy of pancreatitis, but this was most pronounced in turkeys and chickens.<sup>129</sup>

## 8.10 Diagnosis

HPAI is a differential diagnosis to be considered in any flock in which marked lethargy, inappetence, or a drastic decline in egg production are followed by sudden deaths. While a confirmed diagnosis is dependent on the isolation and identification of the virus, it is typically advantageous (for rapid control and eradication purposes) to respond to a presumptive positive H5 or H7 result by polymerase chain reaction (PCR) in accordance with any case definition.<sup>130</sup> In the United States, confirmation of an HPAI outbreak is made by the National Veterinary Services Laboratories in Ames, IA (NVSL). After positive confirmation of HPAI, subsequent samples from premises inside the established CA may be sent to approved laboratories that are part of the National Animal Health Laboratory Network (NAHLN).<sup>130</sup>

The reference standard for diagnosis of viable AI virus is virus isolation—an accurate method of confirming the presence of a virus that could infect other birds.<sup>131</sup> In the laboratory, 9- to 11-day-old embryonated chicken eggs are inoculated with swab or tissue specimens. Additional tests on fluids from the egg are required to confirm the presence of AI virus and determine its HA and NA subtype.<sup>25</sup>

The application of molecular methods for detection of viral nucleic acid and genetic sequencing for viral genes have become important tools in recent years. The rRT-PCR has advantages for outbreak surveillance such as speed, scalability for high throughput, high sensitivity, and high specificity.<sup>25</sup>

Antigen detection immunoassay kits have also been used in prior outbreaks and have advantages of speed (15-20 minutes), simplicity, and good specificity. While the low analytical sensitivity (detection limit greater than  $10^4$  EID<sub>50</sub>) is a limiting factor, birds with clinical signs of AI, or that died of AI infection, generally shed adequate virus antigen for detection with these kits. In contrast, the assays are not recommended for screening of apparently healthy poultry, due to the lower level of shedding before the disease is clinical.<sup>25</sup>

## 8.11 Differential Diagnosis


HPAI can resemble several other avian diseases, including velogenic viscerotropic Newcastle disease, infectious bronchitis, infectious laryngotracheitis, mycoplasmosis, infectious coryza, fowl cholera, aspergillosis, and *Escherichia coli* infection. It also must be differentiated from heat exhaustion, toxicities, and severe water deprivation.

## 9 Risk Evaluation

### 9.1 Pathways for a Broiler Flock Becoming Infected with HPAI via Local Area Spread Components other than those Involving Movements of People, Vehicles, and Equipment

#### 9.1.1 Role of Local Spread Components in Previous AI Outbreaks

Local area spread refers to mechanisms whereby the transmission likelihood increases with proximity to infected farms. The implementation of a Control Area (e.g., minimum 3-km infected zone plus 7-km buffer zone) is based on potential for local spread. A review of past outbreak experiences indicates that the majority of local area spread of AI virus between farms can be attributed to the movement of people and equipment. We evaluated the likelihood of local spread occurring via wild birds, aerosols, and insects in this chapter.



**Figure 12.** Relationship between the daily likelihood of exposure and distance from infected premises estimated from past HPAI outbreak data (also called a “transmission kernel”). Note that all these transmission kernels are not “mechanism-specific” and, hence, include movement of people, vehicles, and equipment between farms as possible transmission mechanisms.<sup>132-134</sup>

Several HPAI outbreak studies have evaluated proximity as a risk factor in general without differentiating between component mechanisms. Spatial and risk-factor analysis from HPAI outbreaks in the Netherlands and Italy indicates a considerable decrease in the chances of infection with distance from infected premises. For example, in Busani et al. (2009), farms within 1.5 km of an infected premises had a 4 to 5 times greater chance of infection relative to farms located more than 4.5 km away.<sup>135</sup>

**Figure 12** above shows the relationship between the daily likelihood of infection and distance from infected premises based on transmission equations estimated from different HPAI outbreaks. The predicted likelihood of exposure steadily decreases with distance in all curves. The specific mechanisms by which the transmission likelihood increases with proximity is

ambiguous based on these studies (see Appendix 2: Literature Review on the Role of Local Area Spread in Previous Outbreaks for a summary of past outbreak studies on proximity). Nevertheless, the transmission likelihood estimates from these studies can be considered as a conservative (an upper bound or maximum) estimate of the spread that occurs due to mechanisms not associated with movement of people, vehicles, and equipment.

Apart from the above spatial analyses, most other AI outbreak observations indicate limited spread of AI among poultry premises by local spread mechanisms such as via insects, aerosols, and wildlife. For example, in a 2008 HPAI outbreak in the United Kingdom, there was no spread to 78 other farms within 3 km of an infected farm.<sup>136</sup> There are several instances where spread did not occur to other houses even on the same premises. (See Appendix 2: Literature Review on the Role of Local Area Spread in Previous Outbreaks for a summary of past outbreak studies on proximity).

### 9.1.2 Role of Insects in the Transmission of HPAI Virus

Insect or fly transmission of AI virus has been suspected in previous HPAI outbreaks based on anecdotal reports.<sup>137,138</sup> However, there are no quantitative epidemiological studies establishing transmission via flies. Houseflies (*Muscidae*) and blowflies (*Calliphoridae*) are reservoirs and vectors of a wide variety of pathogenic organisms affecting poultry.<sup>139</sup> The housefly is usually the most abundant and pestiferous fly species in poultry houses.<sup>139</sup> Most blowflies result from improper disposal of dead birds in a poultry operation, with very few associated with manure.<sup>139</sup> Some biosecurity plans and guidelines for AI control recommend fly control to minimize the spread of AI because of the existing uncertainty about fly transmission of HPAI.<sup>140,141</sup> A majority of broiler integrators specify some sort of fly control program within their biosecurity plans (BWG, personal communication, June 2016).

Below is a summary of the literature from previous outbreaks implicating insects in transmission of HPAI, survivability of AI viruses in flies, dispersion likelihood, and transmission of HPAI to a poultry flock in the two weeks prior to marketing. For a more in-depth discussion on transmission via flies, please see the Day Old Chicks Risk Assessment.<sup>142</sup>

#### 9.1.2.1 Literature Review

- Blowflies were considered as a potential transmission route in the 2004 HPAI H5N1 outbreak in Japan.<sup>143,144</sup> In this outbreak, the prevalence of H5 virus genes was highest in blowflies collected 600 to 700 meters from the infected farm (20 to 30 percent), and HPAI virus gene-positive flies (10 percent) could be detected up to two kilometers from the infected premises. The authors estimated that prevalence of viable virus at 5 percent in flies around the epidemic area.<sup>144</sup>
- Experimental studies indicate that flies can ingest AI virus and that there is a steady decrease in the viable virus titer over time.<sup>145-147</sup> Sawabe et al. (2009)<sup>147</sup> evaluated the survivability of H5N1 virus in blowflies after experimental exposure. Viable virus was recovered in the crop and intestine up to 24 hours post-exposure. However, there was a steady decrease in viral titers from the gut contents over time. Most of the flies had viral titers below the level of detection for the assay (0.50 log TCID<sub>50</sub>/0.05 ml of fly homogenate) at 24 hours. All of the flies had viral titers below the level of detection at 48 hours post-exposure.

- Wanaratana et al. (2013)<sup>148</sup> evaluated the potential of the housefly to serve as a mechanical vector of the H5N1 virus. H5N1 virus could survive within the body of the housefly and remain infective for up to 72 hours post-exposure. The viral titers in housefly homogenate varied between 10<sup>5.43</sup> EID<sub>50</sub>/ml at 6 hours post-exposure to 10<sup>2</sup> EID<sub>50</sub>/ml at 72 hours post-exposure. In this study, the potential for virus transmission via virus on the fly body was also investigated. At 24 hours post-exposure, the virus concentration was 1.90 log ELD<sub>50</sub>/ml (the concentration at time 0 was 4.70 log ELD<sub>50</sub>/ml), whereas virus could not be recovered by 48 hours post-exposure.
- Wanaratana et al. (2013), demonstrated transmission to chickens fed fly homogenate via oral drop with a pipette one day after exposure to 10<sup>8.5</sup> ELD<sub>50</sub>.<sup>148</sup> Based on the timing of virus shedding,<sup>a</sup> between 1 and 3 chickens out of 10 appeared to have been directly exposed from the fly homogenate in this study.<sup>148</sup>
- Tsuda et al. (2009)<sup>149</sup> proposed a mechanism of transmission whereby poultry directly feed on HPAI-infected blowflies. It has been shown that a chicken can eat 31 blow flies placed in its cage in just 7 minutes.<sup>147</sup> However, feeding dead flies (*C. nigribarbis*) contaminated with H5N1 virus did not result in transmission (unpublished data).<sup>150</sup> The frozen dead flies were not attractive to chickens, and only small numbers of flies were consumed by the chickens in this experiment.<sup>150</sup>
- Fly dispersal behavior varies by species and environmental conditions. Houseflies tend to remain close to their breeding site as long as they find suitable food, breeding sites, and shelter. Also of note, the dispersal rate of flies decreases at temperatures below 53°F and increases during premises cleanout or spreading of litter.<sup>151</sup> A summary of fly dispersal rates appears in **Table 2**.

**Table 2.** Reported dispersal rates for types of flies implicated in the mechanical transmission of H5N1 HPAI.

Common name	Reported dispersal rates	Reference
Housefly	1-3 km/day	James & Harwood <sup>152</sup>
Housefly	Generally range less than 2 miles (3.2 km); range in a radius of 328-1,640 feet from breeding site if suitable food available; only 8-30% disperse beyond a poultry facility	Stafford <sup>151</sup>
Blowfly	Estimated 1,250-1,789 meters/day on average	Tsuda et al. <sup>149</sup>
Blowfly	2-3 km in 24 hours	Sawabe et al. <sup>143</sup>

- Beetles have also been implicated as a possible vector for transmitting AI viruses in a few studies.<sup>153-155</sup> However, there are minimal data on the experimental transmission of AI via

<sup>a</sup> Only 3 birds out of 10 were shedding by day 2 postinoculation. In experimental studies in the literature, most HPAI strains had a mean latent infection period of less than 1.5 days.




beetles. In the 1983 HPAI H5N2 outbreak in Pennsylvania, testing of 144 pools of beetles (*Coleoptera*) yielded only two positive pools. One of the positive pools consisted of darkling beetles, and the second of hide beetles.<sup>138</sup>


- Given that 10 to 60 insects were pooled together in each sample, Bayesian analysis indicates that the actual prevalence among beetles would be between 0.01 and 0.15 percent, which is quite small (see Appendix 3: Estimating an Approximate Posterior Distribution for the Prevalence Among Insects).

### 9.1.2.2 Expert Opinion

We obtained expert opinion on insects as a risk factor from veterinarians in the broiler and turkey sector working groups who had field experience in managing AI outbreaks. Overall, eight experts, including industry and regulatory veterinarians, rated this risk factor on a categorical scale ranging from negligible to extremely high (see Appendix 4: Expert Polling on Insect Transmission Routes, for details of the questionnaire). A majority of experts (six out of eight) rated the likelihood of insect transmission from known infected premises as low or negligible at 1.5 km, and negligible at 2 km or farther (**Figure 13**). The ratings for the likelihood of insect transmission from infected but undetected premises were similar; however, slightly more concern was noted at 1.5 km, where three of eight experts rated risk of infection as moderate or high (**Figure 14**).



**Figure 13.** Ratings for likelihood of insect transmission from a known HPAI-infected premises by veterinarians with experience managing AI outbreaks.



**Figure 14.** Ratings for likelihood of insect transmission from an **HPAI-infected but undetected premises** by veterinarians with experience managing AI outbreaks.

### 9.1.2.3 Qualitative Analysis

We considered the following factors in evaluating this pathway:

- In the period leading up to load-out, the inside of a broiler house likely contains a large amount of manure and other environmental conditions that may attract flies. Winpisinger et al. found the number of house flies was significantly higher near (within 3.2 km) large (>2 million) caged layer operations, compared with background fly levels in rural areas.<sup>156</sup> However, dispersal may depend on outdoor environmental and other factors. The number of flies caught at a distance of 0.8 km (3 to 22 percent of the mean value at layer farm) and 1.6 km (2 to 8 percent of the mean value at layer farm) was much lower than the number of flies trapped at the layer facilities.
- While fly transmission has been proposed as a possible mechanism for spread of HPAI, there has not been any epidemiological analysis evaluating flies as a risk factor for spread. Furthermore, local area spread components (other than mechanisms involving movement of people, vehicles, and equipment) have historically played a minimal role in most AI outbreaks. (See section 9.1.1, Role of Local Spread Components in Previous AI Outbreaks, for more detail.)
- Chickens have been shown to ingest live and actively flying houseflies,<sup>147</sup> but there has been no experimental evidence of chickens or turkeys becoming infected with AI virus through feeding on contaminated whole flies in previous outbreaks. Infection was achieved experimentally in chickens using fly homogenate administered via pipette, which likely approximates the oral or possibly nasal/choanal route of infection.<sup>148</sup> We hypothesized that HPAI transmission via feeding of whole flies as opposed to homogenate would have a low likelihood for the following reasons:
  - For HPAI virus encapsulated in the fly body (i.e., virus ingested by a fly), the most likely inoculation route to the chicken is intragastric. As chickens do not grind or masticate their food within the oral cavity, the likelihood that fly gut contents would



contact the choanal cleft during ingestion is decreased. Intra-gastric infectious dose (CID<sub>50</sub>) estimates are quite high at 10<sup>5.2</sup> EID<sub>50</sub> to 10<sup>6.2</sup> EID<sub>50</sub> based on two studies.<sup>109,157</sup>

- Wanaratana et al. (2013) have found a considerable decrease in the external HPAI virus concentration on an exposed fly within 24 hours.<sup>148</sup> While HPAI virus is inactivated at a slower rate in fly gut content, the likelihood of infection due to the virus encapsulated in the fly gut would be reduced because of the higher infectious dose needed for the intra-gastric route.
- Contamination of fly perching surfaces with virus from the fly body, vomit, or feces is a possibility. However, available experimental studies indicated that there would be a considerable reduction in the virus concentration in fly body, vomit, or feces by 6 to 24 hours post-exposure of the fly to virus. (See section 9.1.2.1, Summary of Literature on Insect Transmission.) The relatively rapid inactivation of virus present externally on flies would result in reduced likelihood of transmission at greater distances.
  - In addition, the oral infectious dose for HPAI virus in chickens is also relatively high compared with intranasal (or choanal) exposure (estimates range from 10<sup>3.9</sup> to 10<sup>6.7</sup> for HPAI H5N1 and 10<sup>8</sup> for LPAI H9N2).<sup>109,157,158</sup>
- The proportion of flies around an infected premise that could contain viable virus is likely low. Literature estimates report between 2 and 5 percent of flies may contain virus. Dispersal behavior may vary depending on environmental conditions and fly species, and dispersal is hypothesized to increase during outbreak activities such as premises depopulation.

**9.1.2.4 Likelihood Rating and Conclusion**

We rated the likelihood of a broiler premises becoming infected with HPAI virus via insect transmission to vary with distance as described in **Table 3**. Of note, at premises located closer than 1.5 km to an infected flock, there are too many variables to accurately assess the risk of becoming infected with HPAI via insect transmission.

**Table 3.** Likelihood of a broiler premises becoming infected with HPAI virus via insect transmission based on qualitative analysis and expert opinion.

Source premises type	Composite likelihood rating			
	Distance from source (km)			
	1.5	2	3	5
Known infected premises	<i>Negligible to moderate</i>	<i>Negligible to low</i>	<i>Negligible</i>	<i>Negligible</i>
Infected but undetected premises	<i>Negligible to low</i>	<i>Negligible to low</i>	<i>Negligible</i>	<i>Negligible</i>

### 9.1.3 Role of Aerosol Transmission of HPAI Virus

Aerosol spread of AI virus between premises has been implicated in some outbreaks, although most considered it to have played a limited role.<sup>81,159</sup> Aerosol transmission of AI is an active research area with considerable data gaps. We used a combination of approaches including literature review of past outbreak experiences and experimental studies, exploratory dispersion models, and expert opinion to evaluate the role of aerosol transmission.

#### 9.1.3.1 Aerosol Transmission of AI Virus in Past Outbreaks

- The limited role of local area spread through all mechanisms not involving movements of people and equipment in most previous AI outbreaks indicates a limited role for aerosol spread as well. Aerosol spread has been implicated in very few HPAI outbreaks.
  - In several AI outbreaks, such as the LPAI H7N2 outbreak in Virginia, the geographic distribution of affected farms did not show a specific pattern, suggesting that aerosols were not a primary mode of transmission.<sup>72</sup> In an HPAI H5N1 outbreak in the United Kingdom, there was no transmission to 78 other farms within 3 km of an infected turkey farm. The authors concluded that there was no evidence of local area spread beyond 1 km.<sup>136</sup> Appendix 2: Literature Review on the Role of Local Area Spread in Previous Outbreaks summarizes the literature on the role of local spread in previous outbreaks.
  - Ypma et al. (2012) estimated the contribution of a possible wind-mediated mechanism to the total amount of spread during the 2003 HPAI H7N7 outbreak in the Netherlands to be around 18 percent.<sup>82</sup> This estimate was based on the observed correlation between the wind direction and the direction of the spread of disease, estimated through phylogenetic and epidemiological data. The possibility of the direction of spread coinciding with the wind direction by chance was also accounted for in their statistical analysis. We note that this outbreak occurred in a region of very high poultry density (~4 farms per km<sup>2</sup>), which may increase the likelihood of spread over short distances.
  - Aerosol transmission between poultry barns that were in close proximity was suspected as a possible means of spread in the 2004 HPAI H7N7 outbreak in British Columbia. In this outbreak, there were anecdotal reports of some of the infected farms being in close proximity and downwind of other infected flocks. Some of these anecdotal reports were associated with depopulation methods used early in the outbreak, such as grinding carcasses outside the barn or bringing birds outside the barn to depopulate. Although it was suspected, there is no conclusive evidence that aerosol transmission played a major role in this outbreak.<sup>160</sup>
  - In a case-control study of infected layer facilities in Iowa and Nebraska in the 2014-2015 HPAI outbreak, the authors were not able to determine if aerosol transmission was responsible for infection at a facility.<sup>38</sup>
  - A plume analysis model of infected farms in the 2014-2015 HPAI outbreak in Minnesota found that farms located 7 to 15 km from an infected farm were at low to moderate risk of infection via aerosol transmission; however, wind speed and direction may impact the distance at which transmission can occur. Farms located

- within 5 km of an infected premises were at increased risk regardless of wind conditions.<sup>38</sup>
- Activities that can generate AI virus-contaminated dust or aerosols very close to susceptible poultry have been implicated as a transmission mechanism.
    - Live haul trucking of birds actively infected with AI virus within 200 meters of a susceptible flock can pose a risk for aerosol transmission (D. Halvorson, personal communication, July 2016).<sup>137</sup>
    - Depopulation activities up to 400 yards (366 meters) upwind from a susceptible flock can present a risk for aerosol transmission.<sup>161</sup> In an LPAI H7N2 outbreak in Pennsylvania, aerosols generated by stirring up organic materials during depopulation were considered a potential mechanism of spread to farms within 1 to 1.25 miles.<sup>162</sup> Depopulation methods used early in the 2004 HPAI outbreak in Canada, such as grinding carcasses outside the barn or bringing birds outside the barn to depopulate, were implicated in spread of HPAI.<sup>160</sup>
    - Spreading of non-composted contaminated litter on adjacent fields was suspected as a transmission mechanism during the 1983 HPAI H5N2 AI outbreak (D. Halvorson, personal communication, March 2016)<sup>137</sup>. Spread of non-composted manure from infected farms approximately 1.25 miles from susceptible poultry was suspected to have resulted in transmission in one instance during an LPAI H7N2 outbreak in Pennsylvania.<sup>162</sup>
    - A 2015 survey of HPAI-infected turkey farms in the Midwest highlighted anecdotal evidence of aerosol spread related to recent nearby bird transport, blowing sawdust, and depopulation of nearby farms.<sup>38</sup>
  - Only a couple of studies have reported air-sampling results from or around HPAI-infected houses during previous outbreaks. These studies demonstrate the effect of dilution on aerosol concentration with increasing distance from the generating source.
    - High-volume air sampling was conducted in and near an infected layer flock that had high mortality during the HPAI H7N7 outbreak in Canada.<sup>163</sup> Inside the barn, a viral titer of 292 TCID<sub>50</sub>/m<sup>3</sup> was detected in air samples.<sup>b</sup> Air sampling at a command post outside the barn showed a much lower viral load of 12.5 TCID<sub>50</sub>/m<sup>3</sup> based on quantitative PCR. However, no viable virus was recovered. Low concentration and inactivation of virus by sunlight was hypothesized as a possible explanation for the apparent absence of viable virus in these samples.
    - In the 1983 H5N2 HPAI outbreak in Pennsylvania, 5 of 6 samples taken 3 to 6 meters downwind of affected flocks on six farms were positive by virus isolation, whereas only 1 of 12 samples taken 45 to 85 meters downwind of affected flocks on 8 farms was virus-positive; the positive sample was taken 45 meters downwind.<sup>137</sup>
    - The 2015 USDA epidemiology investigation report describes the results of air and environmental sampling of three turkey flocks in Minnesota and three layer flocks in Iowa and Nebraska. Air samples were collected inside and immediately outside (5

---

<sup>b</sup>TCID<sub>50</sub> refers to the 50% tissue culture infectious dose. The MDCK cell line was used for the tissue culture.

meters) of affected barns, and at extended distances ranging from approximately 70 to 1,000 meters downwind from the barns. Five of the six flocks had at least one air sample test positive.<sup>164</sup>

### 9.1.3.2 Experimental Studies of Aerosol Transmission of AI Virus

Several experimental studies indicate that airborne transmission of HPAI infection between turkeys and chickens in adjacent pens or cages is possible but inefficient. These studies also suggest that aerosols may not be a primary route of transmission within a flock.

- In several studies, aerosol transmission of AI was not observed between groups of inoculated and susceptible chickens housed in adjacent cages or chambers with direct airflow.<sup>26,57,165,166</sup>
- A few studies have shown inefficient transmission or low transmission of AI between groups of inoculated and susceptible chickens housed in adjacent cages or chambers with direct airflow.
  - LPAI (Turkey/Wis/66) virus was transmitted via aerosols between groups of 400 turkeys in different compartments of a building. In this experiment, AI virus was transmitted to one out of three exposed groups of turkeys in different compartments. Infection was detected based on serology and hemagglutination inhibition (HI) titer, and no virus was recovered from tracheal swabs.<sup>167</sup>
  - Two out of six strains of LPAI H9N2 were transmitted via aerosol from a cage with four infected chickens to chickens in an adjacent cage 100 cm away.<sup>168</sup>
  - For chickens housed in cages 10 cm apart, airborne transmission of HPAI H5N1 occurred inefficiently when 1 to 2 chickens were infected, but efficiently when 4 to 8 chickens were infected.<sup>169</sup> With likely similar distances, Yee et al. (2009) found the aerosol route to be an important mode of AI virus transmission among chickens in a simulated live bird market setting (i.e., stacked cages) using LPAI H6N2 (A/chicken/California/1772/02).<sup>114</sup>
  - For H5N1 (A/turkey/Turkey/1/2005), Spekrijse et al.<sup>113,170</sup> estimated a transmission rate of 0.10 new infections per infectious bird per day for chickens housed 1 meter away.
- Experimental studies indicate that variability between strains can impact transmissibility via aerosols. For example, Zhong et al. (2014) found different strains of LPAI H9N2 to have markedly different aerosol transmissibility between chickens.<sup>171</sup> The study proposed that the influenza virus genes HA and PA are important in determining aerosol transmissibility.
- Several studies have indicated efficient transmission of HPAI H5N1 and LPAI H9N2 virus to chickens by aerosols that were mechanically generated by nebulizing virus containing stock fluid to very small particle sizes (2-5 µm).<sup>112,158</sup>
- Several studies have found that influenza A viruses show decreased survivability in aerosols at higher temperature and higher relative humidity.<sup>172,173</sup>

AERMOD plume models (see Appendix 5: Live Broiler Movement Aerosol Modeling for model parameters and scenarios) were utilized in the context of this risk assessment. The measure of interest was HPAI virus concentration:

- These models estimate the risk of transmission to a house of near market-weight broilers holding 20,000 birds.
- In a scenario in which a house of 25,000 broilers was infected, aerosol concentration was predicted to be highest downwind from the infected flock; concentration of virus is predicted to fall sharply as distance increases. In this model, infectious dose was estimated at  $10^{5.44}$  EID<sub>50</sub>/m<sup>3</sup>, meteorological parameters and particle size were accounted for, and the predicted concentration of aerosolized virus farther than 2.5 km from the infected premise was considered to be low (Scenario A, see Appendix 5: Live Broiler Movement Aerosol Modeling).
  - When the infectious dose was lowered to  $10^4$  EID<sub>50</sub>/m<sup>3</sup>, the AERMOD model predicted that transmission likelihoods are much higher at longer distances (Scenario C, see Appendix 5: Live Broiler Movement Aerosol Modeling).
- In an alternate scenario involving multiple different variables (the source of infection was a somewhat smaller turkey flock and weather conditions were from a different geographic area), the predicted HPAI virus concentration at a given distance from the infected source was greater than when broilers were the source flock, and transmission likelihoods increased somewhat as well (Scenario B, see Appendix 5: Live Broiler Movement Aerosol Modeling).

#### 9.1.3.3 Expert Opinion

- We obtained expert opinion on aerosol spread as a risk factor from veterinarians in the broiler and turkey sector working groups with field experience managing AI outbreaks. Overall, eight experts, made up of industry and regulatory veterinarians, rated this risk factor on a categorical scale ranging from negligible to extremely high (see Appendix 6 for details of the questionnaire). In a scenario in which depopulation activities were not taking place, a majority of experts (7 out of 8) rated the likelihood of aerosol transmission from known infected premises as low or negligible at 1.5 km and negligible at 2 km or farther. The ratings for the likelihood of aerosol transmission from infected premises where depopulation activities were taking place were similar, but more concern was noted at 1.5 km, where 5 of 8 experts rated risk of infection as moderate or high (see Figures 1-2, Appendix 6: Expert Polling on Aerosol Transmission Route ).

#### 9.1.3.4 Qualitative analysis

We considered the following factors in evaluating this pathway:

- The housing and ventilation systems utilized in commercial broiler operations likely represent at least a partial barrier to local area spread when compared with alternative housing systems (free range or pasture-raised), which are not within the scope of this assessment.
- Transmission via the aerosol pathway involves many constantly changing variables.

- Virus viability may change with temperature, humidity, and UV exposure, as increased temperature, humidity, and UV exposure may or may not cause virus inactivation.<sup>172-174</sup>
- Weather conditions (temperature, humidity, wind speed and direction) vary widely by season and geography. Dispersion of particulate matter and virus from an infected premises may not be consistent over time.
- To date, all exploratory models have assumed the source to be a static premises (i.e., infected poultry house). Other sources of infection, such as proximity to trucking routes or road traffic, have not been investigated.

### **9.1.3.5 Likelihood Rating and Conclusion**

#### **9.1.3.5.1 Likelihood of HPAI Spread to a Broiler Flock in a Control Area via Aerosol Transmission from a Known HPAI-Infected Flock**

Given the higher predicted prevalence of infectious birds in known infected flocks, both the expert opinion ratings and exploratory dispersion modeling results indicated higher potential risk for this category. Literature review and most previous outbreak reports indicated that local area spread and aerosol transmission were not an important factor at distances more than 1.5 km from an infected flock. However, there is some evidence of aerosol transmission over shorter distances.

We provided the following risk ratings, considering the above factors.

- *Moderate to high* if the broiler flock is located 0.5 km from a known infected poultry farm.
- *Moderate* if the broiler flock is located 1 km from a known infected poultry farm.
- *Low* if the broiler flock is located 1.5 km from a known infected poultry farm.

#### **9.1.3.5.2 Likelihood of HPAI Spread to a Broiler Flock in a Control Area via Aerosol Transmission from an Infected but Undetected Flock**

In this case, the expert opinion ratings and dispersion modeling results indicated lower risks. We rated the risk of broilers becoming infected with HPAI via aerosols from an infected but undetected poultry flock at a specific distance from the infected premises as follows:

- *Low to moderate* if the broiler flock is located 0.5 km from an infected but undetected poultry farm.
- *Low* if the broiler flock is located 1 km from an infected but undetected poultry farm.
- *Negligible to low* if the broiler flock is located 1.5 km from an infected but undetected poultry farm.

#### **9.1.3.5.3 Conclusion**

The risk of exposure of a broiler flock from bioaerosols ranges from *negligible to high*, depending on the distance from, and prevalence of virus in, the source flock. The assessed risk is highest for flocks located within 0.5 km from a known infected poultry farm. We estimate the risks of exposure of a broiler flock to be *negligible to low* if the premises is located 1.5 km from

an infected but undetected poultry farm, and *low* if the premises is located 1.5 km from a known infected poultry farm.


#### 9.1.4 Role of HPAI Spread to a Broiler Flock in a Control Area via Wild Aquatic Birds in the Farm Vicinity

Wild aquatic birds are the main reservoir of influenza A viruses in nature. They harbor all 16 (H1-H16) HA and all 9 (N1-N9) NA subtypes of AI in their population. Most of the isolates from aquatic birds have been LPAI, which generally does not cause disease in the wild population. It is understood that the virus circulates continuously in the wild population, but often at low levels.<sup>175</sup>

Various species of wild aquatic birds are implicated in the maintenance of AI viruses:

- Wild waterfowl are considered to be the primary source of new H5 or H7 LPAI outbreaks in poultry, particularly in poultry reared in semi-intensive or extensive (free-range) conditions.<sup>176</sup> Wild ducks have been found to carry a higher prevalence of virus during their southern migration in the fall (22.2 percent) than during their spring northerly migration (0.3 percent). This difference may be due to the increased number of susceptible young birds during the fall migration.<sup>175</sup>
- Anecdotally, during the 2014-2015 HPAI outbreak in the Midwest, snow geese were observed in the proximity of poultry houses that later became infected with H5N2.
- Shorebirds have also been found to carry influenza viruses in a higher percentage than that in ducks during the spring migration.<sup>175</sup>
- Gulls are susceptible to HPAI viruses<sup>177</sup> and are a known reservoir of AIVs.<sup>178</sup> Gulls are suspected to have been the source of a 2002 outbreak in the Chilean poultry industry. In this instance, the HPAI virus likely mutated from an LPAI strain.<sup>179</sup> The role of gulls in the transmission of AI is likely twofold because of their susceptibility to infection and their opportunistic nature when they scavenge for food. Gulls are susceptible to AI and thus can contract but transmission from gulls to other species is less clear.<sup>177</sup> Because they are opportunists, gulls are likely to be present near poultry barns and may come into contact with dead birds. In this case, gulls may act as fomites in the dispersal of AIVs (more in-depth analysis of the role of scavengers can be found in the “Dead bird disposal” chapter of this Risk Assessment).

Influenza viruses have been shown to affect all types of domestic birds, and the primary infection depends on the degree of contact with wild birds. As mentioned in section 9.1.1., Role of Local Spread Components in Previous AI Outbreaks, secondary spread usually results from human activities that transfer infective feces to susceptible birds.<sup>180</sup> Potential pathways of HPAI virus transmission through wild aquatic birds in the farm vicinity are illustrated below.



**Figure 15.** Pathway of HPAI virus transmission through wild aquatic birds

#### 9.1.4.1 Literature Review

Historically, HPAI viruses rarely have been isolated from wild birds. Where HPAI viruses were identified, they were usually from isolates obtained from dead wild birds found in the vicinity of HPAI-infected poultry farms<sup>181</sup> or from aquatic bird population surveillance sampling.<sup>175</sup>

Recent studies have shown that HPAI viruses, in particular Eurasian H5N1 and H5N8, are present in populations of different wild aquatic bird species covering wide geographical areas globally.

- In a survey conducted in China from 2004 to 2007, 14,472 wild bird samples (cloacal swabs, organ tissues, or fresh excrement) were collected from 10 bird orders. The samples from Anseriformes had the highest prevalence of H5N1 virus. The positive samples were collected from nine species of ducks, geese, and swans.<sup>182</sup>
- HPAI outbreaks in migratory water birds from 2005 to 2011 in Mongolia, a country with very few domestic poultry (fewer than 100,000 birds), provided strong evidence that wild birds can carry HPAI virus over at least moderate distances, but may not be competent as indefinite reservoirs.<sup>183</sup>
- A large-scale surveillance program detected HPAI H5N2 in healthy birds of two wild waterfowl species sampled in Nigeria and genetically-related LPAI H5N2 in Eurasian domestic poultry.<sup>31</sup>
- HPAI H5N8 was identified in poultry in South Korea in January 2014, and closely related strains subsequently appeared in Japan, China, and Europe. Several reassortant H5 HPAI viruses recently isolated in North America show 99 percent similarity to the Korean H5 strains.<sup>184,185</sup>



- Wild bird sampling activities in the Netherlands between November 2014 and February 2015, following H5N8 virus outbreaks in poultry, detected HPAI H5N8 virus in two samples (out of 4,018 birds sampled) from ducks of the Eurasian wigeon species.<sup>186</sup>
- Between December 2014 and February 2015, Eurasian/North American reassortant HPAI H5N1, H5N2, and H5N8 were found in several species of wild ducks, as well as wild raptors, in the states of Washington, Oregon, California, Utah, Idaho and Nevada. After February 2015, new H5N2 cases in wild aquatic birds and raptors were also detected in Minnesota, Wisconsin, Michigan, Wyoming, Kansas, and Kentucky (**Table 4**).<sup>39</sup>

**Table 4.** H5N2 cases in U.S. aquatic birds, December 2014 to June 2015 <sup>38</sup>

Bird Species	Number	State	Cause of death
Canada goose <i>Branta canadensis</i>	5	Michigan	Morbidity/mortality
	1	Wyoming	
	1	Kansas	
	1	Washington	
Lesser snow goose <i>Anser caerulescens caerulescens</i>	1	Kentucky	Morbidity/mortality
	2	Montana	
Ring-necked duck <i>Aythya collaris</i>	1	Kentucky	Morbidity/mortality
American green-winged teal <i>Anas crecca</i>	1	Idaho	Hunter harvest
	1	Oregon	
Mallard <i>Anas platyhrynchos</i>	2	Idaho	Hunter harvest
	5	Washington	
	3	Oregon	
Northern pintail <i>Anas acuta</i>	2	Oregon	Hunter harvest
	1	Washington	
Northern shoveler <i>Anas clypeata</i>	3	Oregon	Hunter harvest
Wood duck <i>Aix sponsa</i>	3	Oregon	Hunter harvest

During the 2014-2015 H5N2 outbreak in the midwestern U.S., sampling of wildlife took place on five infected and five uninfected farms. Out of 419 individual birds sampled, killdeers were the only aquatic birds collected, and none tested positive for HPAI. It should be noted, however, that the samples were collected 2 to 4 weeks after clinical signs of HPAI were observed in the poultry flocks, and while depopulation was complete at some infected farms, it was ongoing at others.<sup>38</sup>

Interestingly, the role of wild aquatic birds in perpetuating HPAI viruses remains unresolved. AI researchers have examined current and historical aquatic bird influenza A virus surveillance and outbreaks of highly pathogenic H5 viruses in poultry in the U.S. and Canada dating back 43 years prior to the 2014-2015 outbreak.<sup>187</sup> This analysis failed to detect HPAI viruses in wild aquatic birds before or after the resolution of that outbreak, suggesting that there are yet undetermined mechanisms preventing wild aquatic birds from perpetuating HPAI viruses.<sup>187</sup>

Experimental studies suggest that while most aquatic bird species show minor or no clinical signs after being infected with HPAI viruses, some can efficiently transmit the viruses to their contacts. **Table 5** summarizes the results of several studies on HPAI virus in wild and domesticated aquatic birds.

**Table 5.** Summary of experimental studies of HPAI virus in wild and domesticated aquatic birds.

HPAI virus	Bird species	Inoculation	Findings	Reference
<b>H7N3</b> (A/chicken/Chile/184240-1/02)	Chiloe wigeon and cinnamon teal	10 <sup>6</sup> EID <sub>50</sub> (intranasal)	No ducks developed disease or died.  Oral and/or cloacal shedding in all virus-inoculated cinnamon teals and oral shedding in 2/8 chiloe wigeons at day 2 post-inoculation  Virus efficiently transmitted to cinnamon teal contacts, not to chiloe wigeon contacts	Sá e Silva et al., 2011 <sup>188</sup>
<b>H5N1</b> (A/chicken/Scotland/59) <b>H5N2</b> (A/chicken/Pennsylvania/1370/83) <b>H5N2</b> (A/chicken/Pennsylvania/1/83) <b>H5N9</b> (A/turkey/Ontario/7732/66)	Khaki-Campbell duck	0.1 ml of diluted infectious allantoic fluid (intramuscular and intranasal routes, and contact with inoculated ducks)	No infection and no shedding established.	Alexander et al., 1986 <sup>189</sup>
<b>H5N8</b> (A/turkey/Ireland/83) <b>H5N8</b> (A/duck/Ireland/113/84)	Khaki-Campbell duck	0.1 ml of diluted infectious allantoic fluid (intramuscular and intranasal routes and contact with inoculated ducks)	Virus shedding in cloaca and trachea and transmission to in-contact ducks  No clinical signs or deaths	Alexander et al., 1986 <sup>189</sup>
<b>H7N7</b> (A/Chicken/Netherlands/621557/03)	Ringed teal	0.2 ml of tenfold diluted allantoic fluid (intravenous)	All unvaccinated ringed teals became infected and rapidly transmitted to all contact teals.  Shedding through cloaca and trachea in all animals  2/10 developed conjunctivitis; no clinical signs in others.	van der Goot, 2005 <sup>120</sup>

**Table 5. Summary of experimental studies of HPAI virus in wild and domesticated aquatic birds, cont.**

<b>HPAI virus, cont.</b>	<b>Bird species</b>	<b>Inoculation</b>	<b>Findings</b>	<b>Reference</b>
<b>H5N2</b> (A/chicken/Pennsylvania/1/83)	Ring-billed gull	10 <sup>8</sup> EID <sub>50</sub> (intranasal/intraocular)	Virus detected in the intestine, lung, and spleen  No transmission to in-contact birds	Wood et al., 1985 <sup>190</sup>
<b>H5N1</b> (A/Whooper Swan/ Mongolia/244/05) <b>H5N1</b> (A/Duck Meat/ Anyang/01)	Mallard, northern pintail, blue-winged teal, redhead, wood duck, and nestling laughing gulls.	0.1 ml of diluted allantoic fluid from inoculated eggs diluted in brain-heart infusion (intranasal)	Wood ducks were the only species of duck to exhibit illness or death after inoculation with either of the HPAI viruses. Severe clinical signs appeared in all of the inoculated gulls. In both species virus was isolated from internal organs. Viral titers were higher in oropharyngeal swabs than in cloacal swabs.	Brown et al., 2006 <sup>177</sup>
<b>H5N8</b> (A/Gyrfalcon/Washington/41088/2014) <b>H5N2</b> (A/Northern Pintail/Washington/40964/2014)	(1) White Chinese Goose (2) Pekin duck (3) Mallards	10 <sup>6</sup> EID <sub>50</sub>	<b>Geese:</b> few clinical signs, some mortality <b>Pekin duck:</b> no mortality <b>Mallards:</b> no mortality or clinical signs, but lower body weight and elevated body temperature	Mary Pantin-Jackwood, personal communication, August 2016

A recent study of several H5 and H7 HPAI virus strains in mallard ducks further illustrates the variability in shedding and transmission to contacts, depending on the virus strain.<sup>191</sup> These findings are summarized in **Table 6**.

**Table 6.** Shedding and transmission results of experimental infection of mallard ducks with H5 and H7 HPAI virus at  $10^6$  EID<sub>50</sub> intranasally.<sup>191</sup>

Virus Strain	Shedding (days)	OP vs. CL	Trans. to contacts	> Chicken BID <sub>50</sub>
H7N3 A/chicken/Chile/184240-1/2002	14	CL	3/3	na
H7N3 A/chicken/Canada/314514-2/2005	14	CL	3/3	na
H7N3 A/chicken/Jalisco/CPA1/2012	14	CL	3/3	na
H7N7 A/chicken/Victoria/1985	11	CL	3/3	>2.9
H7N7 A/chicken/North Korea/7916/2005	11	CL	3/3	na
H7N7 A/chicken/Netherlands/1/2003	11	=	3/3	na
H7N1 A/turkey/Italy/4580/1999	11	=	3/3	>2
H5N2 A/chicken/Pennsylvania/1370/1983	14	=	3/3	>3
H5N2 A/chicken/Queretaro/14588/1995	4	OP	1/3	>3
H5N8 A/turkey/Ireland/1378/1983	11	OP	2/3	<4.7
H5N3 A/tern/South Africa/1961	14	=	1/3	>3.4

OP: primarily oropharyngeal shedding; CL: primarily cloacal shedding; =: equal OP and CL shedding.

BID<sub>50</sub>: 50 percent bird infectious dose. One BID<sub>50</sub> unit is the amount of virus that will infect 50 percent of inoculated birds.

The evidence that connects wild birds to infected farms is divergent. In a case-control study of layer and pullet premises in Iowa and Nebraska in the 2015 HPAI outbreak, no consistent association was observed between infected or control farm status and wild bird sighting.<sup>192</sup> In other cases, evidence has been found linking wild birds to infected premises.

- Observations of wild bird activity in two provinces in Canada showed seven species of wild aquatic birds—Canada goose, mallard, ring-billed gull, glaucous-winged gull, mew gull, killdeer, and trumpeter swan—were seen in the immediate barn area at least twice.<sup>193</sup> They were most frequently observed near feed silos. No wild aquatic birds were observed entering the poultry houses.

Some recent events have provided additional evidence for outbreaks resulting from possible introduction of HPAI virus into domestic birds from wild aquatic birds.

- A North American outbreak of HPAI with H5 of Eurasian lineage began on December 1, 2014, and H5N2 was detected in 11 commercial broiler breeder, table egg layer, and turkey farms in British Columbia by December 17, 2014.<sup>194</sup> In addition, the Canadian Food Inspection Agency confirmed HPAI H5N1 on a noncommercial poultry farm on February 7, 2015.<sup>194</sup> Influenza viruses had been previously isolated from wild and domestic ducks in British Columbia.<sup>195</sup>

- Eurasian H5N8 was confirmed in a backyard mixed poultry flock in Oregon on December 19, 2014, followed by Eurasian/North American reassortant H5N2 outbreaks in backyard flocks in Washington, Oregon, and Idaho in January and February 2015.<sup>196,197</sup> Various positive aquatic birds were found during the outbreak, as shown in **Table 7**, cementing the possibility of introduction from wild aquatic birds.

**Table 7.** Hunter-harvested wild bird surveillance for HPAI virus H5 intercontinental A (icA) results for AI matrix gene, Pacific Flyway, December 2014 through February 1, 2015, as reported in Bevins et al.<sup>197</sup>

Species	n	HPAI virus icA positive
Mallard, <i>Anas platyrhynchos</i>	1,410	15
Northern shoveler, <i>Anas clypeata</i>	555	3
Green-winged teal, <i>Anas crecca</i>	724	4
American wigeon, <i>Anas americana</i>	777	31
Northern pintail, <i>Anas acuta</i>	460	5
Cinnamon teal, <i>Anas cyanoptera</i>	67	0
Wood duck, <i>Aix sponsa</i>	27	3
Gadwall, <i>Anas strepera</i>	185	1
Canvasback, <i>Aythya valisineria</i>	68	0
Ruddy duck, <i>Oxyura jamaicensis</i>	46	0
Bufflehead, <i>Bucephala albeola</i>	35	0
Canada goose, <i>Branta canadensis</i>	148	1
Cackling goose, <i>Branta hutchinsii</i>	33	0
Lesser scaup, <i>Aythya affinis</i>	14	0
Ring-necked duck, <i>Aythya collaris</i>	65	0
Common goldeneye, <i>Bucephala clangula</i>	39	0
All other species sampled	76	0

- Commercial turkey flocks in Stanislaus County, California, were infected with a novel Eurasian HPAI H5N8 in January 2015, and the outbreak is considered related to the HPAI events in wild birds. This novel AI virus of Eurasian origin (EA-H5N8 clade 2.3.4.4) spread rapidly along wild bird migratory pathways during 2014.<sup>198</sup> On February 12, 2015, Eurasian H5N8 was also confirmed in a commercial chicken flock in Kings County, California.<sup>196</sup>
- Between March and June of 2015, an outbreak of H5N2 was observed in the Midwest; turkey barns were the most impacted in Minnesota and chickens were more involved in Iowa.<sup>38</sup> Although 3,139 waterfowl fecal samples were tested during this outbreak, HPAI virus was not isolated from any aquatic bird fecal sample.<sup>199</sup>

#### 9.1.4.2 Qualitative Analysis

We considered the following factors in evaluating this pathway:

- Generally, total confinement and biosecurity measures practiced on commercial poultry operations effectively prevent wild aquatic birds and other pests from entering broiler barns.
  - The spread of the viruses via migratory waterfowl routes is far less likely to occur in poultry farms with bird-proof confinement.<sup>200</sup>
  - Feed bins are maintained so that wild birds do not frequent or access broiler premises, and spilled feed should be promptly cleaned up to avoid attracting wild birds.<sup>201-203</sup>
- Experimental studies suggest that the possibility of HPAI infection in wild aquatic birds varies considerably, and their ability to transmit viruses depends on the combination of virus strain and host.
  - A study in 1986 by Alexander et al. challenged ducks with eight different HPAI and LPAI viruses via three different routes (intranasal, intramuscular, and contact with inoculated ducks).<sup>180</sup> Ducks became infected with only four strains of viruses, and the infection rate for each strain varied with the route of infection. In addition, tracheal and cloacal viral shedding were inconsistent.
  - As illustrated in **Table 6**, mallard ducks experimentally infected intranasally with four strains of H5 HPAI viruses at  $10^6$  EID<sub>50</sub> responded differently from those infected with strains of H7 HPAI viruses at the same EID<sub>50</sub>.<sup>191</sup>
- Despite the possibility of wild aquatic birds introducing HPAI to susceptible farmed poultry, there is no known evidence of secondary spread from total confinement farms within control zones via wild birds. Anecdotally, there have been reports of suspected movement of LPAI virus between flocks of free-range turkeys (Mahesh Kumar, personal communication, November 1995), but these free-range flocks fall outside the scope of this assessment. Once the viruses move from wild birds to poultry, it is assumed that human activities—especially movement of personnel and equipment from farm to farm—are responsible for transferring infective materials from infected to susceptible birds.<sup>200</sup> Secondary spread caused by wild birds between poultry premises should be considered possible in rare instances.
  - Wood et al. (1985) demonstrated little to no fecal shedding of HPAI H5N2 in wild ring-billed gulls (and domestic Pekin ducks), suggesting these birds were unlikely to transmit virus from farm to farm in the 1983 Pennsylvania outbreak.<sup>190</sup>
  - None of the HPAI-infected wild ducks (H5N2, H5N1, and H5N8) found in the 2014-2015 U.S. outbreak have been implicated in transferring the virus from one poultry farm to another.<sup>38,184,185,204</sup>
  - In the above-mentioned HPAI H5N8 outbreak in commercial California turkeys, other houses on the premises remained negative, and spread of the disease within the Control Area did not occur.<sup>198</sup>



### **9.1.4.3 Likelihood Rating and Conclusion**


While wild aquatic birds are natural reservoirs for influenza A viruses (including several strains of HPAI virus) and could potentially cause a spillover of disease to domestic poultry, primary infection in broilers or other domestic poultry depends upon the degree of contact with wild birds. In addition, there is no historical evidence of secondary spread from total confinement barns within Control Areas via wild aquatic birds. Modern poultry management systems, in combination with stringent biosecurity measures, make the contact between wild and domestic birds—and resulting secondary spread of HPAI virus among domestic poultry via wild aquatic birds—unlikely within the Control Area. Therefore, we conclude that the likelihood of HPAI infection in poultry via wild aquatic birds in the farm vicinity is *low*.

### **9.1.5 Role of HPAI Virus Spread to a Broiler Flock via Wild Non-Aquatic Birds in Farm Vicinity**

An AI virus was first identified in wild birds in 1961 when HPAI H5N3 was isolated from common terns (*Sterna hirundo*) in South Africa.<sup>205</sup> A compilation of more recent surveys of wild birds describes an overall AI virus prevalence of 15.2 percent in Anseriformes (waterfowl), 2.9 percent in Passeriformes (perching birds), and 2.2 percent in Charadriiformes (waders, gulls, and auks).<sup>200</sup> Influenza viruses are primarily spread from wild birds to domestic poultry through the mechanical transfer of infective feces, usually via human activity.<sup>200</sup> For a thorough review of pathways associated specifically with aquatic bird species, please see section 9.1.4, Role of HPAI Spread to Broiler Flock in a Control Area via Wild Aquatic Birds in the Farm Vicinity.

#### **9.1.5.1 Likelihood of Infection via Passerine or Columbiforme Birds in Farm Vicinity**

Since its appearance, HPAI H5N1 has demonstrated the unique ability among HPAI viruses to infect a wide variety of species, including wild birds. Small perching birds of the order Passeriformes (passerines) commonly frequent poultry farm areas and thus have the potential to serve as biological or mechanical vectors of H5N1, or as so-called bridge species in its transmission.<sup>206</sup> This group includes commonly encountered species such as sparrows, swallows, and starlings. Other potential bridge species include the Columbiforme birds, which include pigeons and doves.<sup>207</sup> The potential pathways for HPAI transmission via passerine or Columbiforme birds include infection or contamination of the wild bird on an infected poultry farm or premises contaminated with infected wild bird feces, with subsequent primary or secondary transmission into an uninfected broiler house. The flying distances that some wild bird species travel depend on food supply and weather. For example, starlings and blackbirds disperse as far as 15 to 25 miles on average, with some individuals traveling up to 50 miles daily from their roost to their feeding grounds.<sup>208</sup> Any of these movements provide an individual bird the opportunity to contact and disseminate AI viruses.



**Figure 16.** Pathway for exposure of a broiler farm via wild passerine or columbiforme birds

#### 9.1.5.1.1 Literature Review

Small species of wild birds have been observed entering poultry barns.

- European starlings and house sparrows are frequently located near poultry houses.<sup>209</sup> During a field survey to estimate the incidence of bacterial pathogens in passerines near broiler houses, starlings were seen trying to gain entrance to all chicken houses on one farm, and a nest with young starlings was seen in the eaves of one house.<sup>209</sup> Numerous droppings on the sides of the houses on another farm indicated that sparrows and starlings were attracted to the house and possibly trying to gain entrance.
- In a survey of table-egg layer operations in California regarding pest management practices, producers ranked wild birds (passerines) as being somewhat more pestiferous on southern ranches<sup>210</sup> than on northern ones<sup>206</sup> when asked to rank pests in order of perceived importance.<sup>211</sup>
- Craven et al. note that starlings have the ability to peck through plastic wire mesh on the sides of chicken houses.<sup>209</sup> Burns et al. counted wild birds in the vicinity of poultry farms in Ontario and British Columbia<sup>193</sup> and found:
  - Barn swallows (*Hirundo rustica*), rock doves (*Columba livia*), and European starlings (*Sturnus vulgaris*) were all observed entering poultry barns, which included broiler, broiler breeder, layer, and turkey production.
  - Rock doves were observed entering barns the most frequently.
- In a survey of infected turkey farms during the 2014-2015 outbreak in Iowa, Minnesota, North Dakota, South Dakota, and Wisconsin (n=81), 35 percent of farms reported seeing wild birds (unspecified species) in poultry barns daily to occasionally; most reported that small perching birds are present on farm year-round.<sup>38</sup>

- Some species or populations of passerines could be termed synanthropic, as they occupy a distinct ecological niche in and around human agricultural activities. The behavioral characteristics of passerines that may contribute to their ability to play a role in the transmission of AI to domestic poultry are summarized in **Table 8**.

**Table 8.** Behavioral characteristics of several members of the order Passeriformes that impact the potential role of transmission of HPAI virus in environments on the farm around poultry houses.

Common name (species)	Migration	Habitat	Nesting behavior	Food
Common Grackle ( <i>Quiscalus quiscula</i> )	Resident or short-distance migrant	Agricultural fields, feedlots, woodland, forest edges, marshes	Nearly always in scattered trees, rarely in barns	Omnivorous; seeds (agricultural grains)
House Sparrow ( <i>Passer domesticus</i> )	Resident	Closely associated with people and their buildings	Prefers structures; eaves or walls of buildings	Grains and seeds (livestock feed)
European Starling ( <i>Sturnus vulgaris</i> )	Resident or short-distance migrant	Countryside near human settlements; feed in fields	Trees, buildings, structures	Focus on insects and invertebrates; also fruits, berries, grains (livestock feed)
House Finch ( <i>Haemorhous mexicanus</i> )	Resident or short-distance migrant	Farms, parks, urban centers, backyards	In or near buildings; trees	Plant materials almost exclusively; millet, milo, etc.

Table from USDA-APHIS Poultry Feed Risk Assessment.<sup>212</sup>

Wild birds may also be attracted to poultry feed. For more information on specific risks of feed contamination if passerines breach biosecurity at feed mills or on farms, see USDA-APHIS Poultry Feed Risk Assessment.<sup>212</sup>

While passerine birds have not been directly implicated in the spread of HPAI in previous outbreaks, such birds have tested positive for AI in the vicinity of poultry outbreaks.

- In a 1985 H7N7 HPAI outbreak in chickens in Australia, an antigenically closely related strain was isolated from starlings on the affected farm, and serologic evidence of H7N7 infection was found in sparrows as well.<sup>213</sup>
- In a 1995 survey to establish disease freedom for poultry operations during an outbreak of HPAI H5N2 virus in Mexico, serologic evidence of infection of three passerine birds (species not specified) to an H5N2 serotype was reported.<sup>214</sup> However, an LPAI H5N2 virus had been circulating in poultry in 11 Mexican states prior to the outbreak; it is ambiguous as to which virus resulted in the exposure.
- In Pakistan in 2007, four wild crows were found to be H5N1-positive following outbreaks in backyard poultry and zoo birds.<sup>215</sup>

- In Hong Kong in 2009, among 22 birds found dead, including chickens, one large-billed crow (*Corvus macrorhynchos*) was infected with H5N1.<sup>215</sup>
- In Jalisco, Mexico, in 2012, 81,000 general surveillance samples in an H7N3 outbreak region yielded one positive common grackle (*Quiscalus quiscula*) and one positive barn swallow.<sup>216</sup>
- A chickadee recovered in metropolitan Ramsey County, Minnesota, and delivered on June 10, 2015, to a wildlife rehabilitation center later tested positive for AI by immunohistochemical stains of fixed brain (Minnesota Department of Natural Resources 2015).<sup>217</sup> No virus was isolated, but the chickadee tissues were positive by the H5 intercontinental A (icA) molecular assay, which targets the Eurasian H5 clade 2.3.4.4 viruses. However, hemagglutinin gene sequencing attempts were negative. Where the bird may have become exposed to icA H5 is unknown since complete information about submission circumstances was unavailable.
- As part of a case-control study of layer flocks in northwest Iowa in 2015, wild birds and mammals around the flocks were sampled.<sup>38</sup>
  - Of over 1,600 wild bird samples collected—caught using a mist-net around a nest built in a walkway between two poultry barns on an infected premises—a single sample from lung tissue of a juvenile European starling was positive for Eurasian H5 (icA).
  - Additional serologic evidence of positives for icA H5 were found in a sparrow (*Passer domesticus*), another European starling (*Sternus vulgaris*), and two American robins (*Turdus migratorius*) sampled around the same positive farm.

With one exception, surveillance of passerines for AI virus has demonstrated a zero to low prevalence in the wild population.

- In a summary of three studies from 1979 to 1980, in which a total of 11 passerine species were tested, AI virus isolation was reported from 17 out of 586 birds.<sup>38,218</sup>
- No influenza virus was isolated from 83 cloacal swabs collected from four adult and 79 juvenile reed warblers (*Acrocephalus scirpaceus*) in 1993, despite proximity to aquatic habitats of known AI reservoir species.<sup>176</sup>
- In a survey of passerine birds in the state of Georgia from 1999 to 2009, zero of 234 birds of 25 different species tested positive for AI antibodies.<sup>219</sup>
- On Helgoland Island in the North Sea in 2001, 543 migrating passerine birds of different species all tested negative for AI virus subtypes H5 and H7.<sup>220</sup>
- In China, from 2004 to 2007, RT-PCR on 7,320 cloacal, tissue, or fecal samples from 33 Passeriforme species identified 0.36 percent to be H5N1-positive; 1.09 percent of tree sparrows were positive.<sup>182</sup>
- During active surveillance of Passeriformes for HPAI H5N1 in Mongolia from 2005 to 2011, zero of 80 live-bird, fecal, and sick-bird samples were positive.<sup>183</sup>
  - No passerine birds were sampled during five H5N1 wild bird outbreak investigations.

- In 2006, out of 8,961 Passeriformes sampled via RT-PCR in Europe, one (0.01 percent) was H5N1 positive and eight (0.09 percent) were LPAI positive.<sup>206</sup>
- From a total of 670 cloacal swabs from 37 different species of migratory passerine birds in Slovenia from 2004 to 2006, there was one positive rRT-PCR in the only common starling (*Sturnus vulgaris*) tested, but virus isolation was unsuccessful.<sup>221</sup>
- In a 2007 study in Slovakia, 30 percent of 155 passerine birds of 12 species were AI virus positive via RT-PCR on cloacal and/or oropharyngeal samples, including three of six swallows (*Hirundo rustica*).<sup>222</sup> The authors speculate that the higher than typically reported prevalence may be due to the increased sensitivity of *nested* RT-PCR used in this study.
- Rectal samples from 1,300 tree sparrows (*Passer montanus*) in China in 2011 yielded no AI virus, while 94 of 800 were serologically positive for H5N1, and zero of 800 were seropositive for H7.<sup>223</sup>

Experimental susceptibility of passerine birds to HPAI H5N1 depends on the species of bird and strain of virus.

- In sparrows inoculated with four different H5N1 strains, mortality was 66 to 100 percent, oropharyngeal and cloacal titers were as high as 4.7 and 4.1 log<sub>10</sub> EID<sub>50</sub>/ml, respectively, at 4 days post-inoculation (DPI), and there was no same-species contact transmission.<sup>224</sup> Mortality was 0 percent in European starlings, maximum cloacal titer was 3.8 log<sub>10</sub> EID<sub>50</sub>/ml at 2 DPI, and there was only one unduplicated instance of contact transmission. Oropharyngeal and cloacal titers were very low in pigeons (*Columba* spp.), and their mortality was 0 percent.
  - The authors deduce that sparrows may act as intermediate hosts for transmission to both poultry and mammals, but the lack of contact transmission and high mortality preclude them from serving as a reservoir species for H5N1.
  - While starlings may also act as intermediate hosts, the authors conclude the low contact transmission rate likely indicates they could not serve as an H5N1 reservoir.
  - Pigeons were determined to be likely to play a minor role in the ecology of H5N1.
- Brown et al. (2009) found similar mortality rates (60-100 percent at 10<sup>2</sup> to 10<sup>6</sup> EID<sub>50</sub> inoculum/bird) and maximum oropharyngeal titers (4.2 log<sub>10</sub> TCID<sub>50</sub>/ml) in house sparrows (*Passer domesticus*) inoculated with A/whooper swan/Mongolia/244/05 HPAI H5N1, but maximum cloacal titers were significantly (P=0.002) lower than oropharyngeal. While 40 percent of pigeons (*Columba livia*) inoculated with the highest dose of H5N1 died, they and the survivors shed virus only briefly and at low titers. All pigeons in the low- and medium-dose groups survived and remained AI virus-free.<sup>225</sup>
  - These authors conclude that sparrows could play a role in AI virus transmission in an outbreak, though more likely via contamination of the environment and feed, due to their predominantly oropharyngeal shedding, or via chickens scavenging infected sparrow carcasses.
- Two studies with the HPAI H5N1 strain A/chicken/Hong Kong/220/97 resulted in no mortality and infrequent histopathology lesions in house sparrows and European

starlings.<sup>226,227</sup> While mortality among house finches (*Carpodacus mexicanus*) averaged 44 percent, histopathology lesions were absent to mild and viral antigen rare in the nasal cavity and gastrointestinal tract. The authors were not able to draw any definitive conclusions regarding the role of these species as biological vectors.

- In another study, house sparrows experimentally infected with A/duck/Laos/25/06 H5N1-shed virus for several days and rapidly contaminated their drinking water.<sup>228</sup> On the other hand, inoculated chickens shed undetectable levels of virus into their water troughs, despite high oropharyngeal and cloacal shedding; the authors surmise that this was due to rapid disease progression in the chickens.
  - These authors concluded that sparrows may be unlikely to become infected by chickens under normal field conditions in an H5N1 outbreak.
  - They also inferred that the behavior of infected sparrows may be a determining factor in their potential as intermediate H5N1 hosts via viral shedding.
- In tree sparrows inoculated with A/chicken/Miyazaki/K11/2007 and A/chicken/Shimane/1/2010 H5N1, mortality was 100 percent within 11 days (mean >6 days), with oral swabs positive from 1 to 8 DPI and maximum titers of 10<sup>6.5</sup> to 10<sup>7.3</sup> EID<sub>50</sub>/ml.<sup>229</sup> While there was no intraspecies transmission among sparrows, 10 of 16 (62.5 percent) contact chickens died when housed with infected sparrows.
  - Due to the prolonged viral shedding observed here, the authors concluded that tree sparrows have the potential to serve as biological vectors of H5N1.
- Nestorowicz et al. infected house sparrows and starlings with 10<sup>5</sup> log EID<sub>50</sub> of an isolate of an HPAI H7N7 virus from chickens (A/Chicken/Victoria/1/85) via the oral/tracheal and nasal cleft route.<sup>213</sup> Uninfected birds were placed in contact with infected birds of the same species. Transmission to starlings was observed. More details from the experiment are provided in **Table 9** below.

**Table 9.** Summary of the experimental transmission of H7N7 HPAI virus in house sparrows and starlings by Nestorowicz et al. (1987)<sup>213</sup>

Common name	Mortality	Virus isolation	Transmission
Starlings	100%; All inoculated birds died within 48 hr. post-inoculation	Not reported	Contact birds died within 4 days of being placed with infected birds
Sparrows	30% mortality rate	Isolated from all tissues from birds that died within 2 days post-inoculation	Not transmitted to uninfected contact birds

- 23 of 23 stonechats (*Saxicola torquata*) inoculated with A/Cygnus cygnus/Germany/R65/2006 H5N1 died within three to seven days, most with no clinical signs.<sup>210</sup> Oropharyngeal shedding peaked at 10<sup>3</sup> to 10<sup>4</sup> TCID<sub>50</sub>/ml on 3 to 6 DPI.

### 9.1.5.1.2 Qualitative Analysis

We considered the following qualitative factors for evaluating this pathway:

- A majority of the studies cited above examine strains of HPAI H5N1.
  - To date, HPAI H5N1 has proven to be unique in its ability to infect a variety of species, and more ubiquitous in its prevalence than any other HPAI virus.
- The risk of AI transmission is much lower from a single infected bird than from a population of birds in which infection is established.
  - As noted in the literature review, surveys of passerine birds have demonstrated a low prevalence of AI virus, including the more pervasive H5N1.
  - Several experimental studies have resulted in no intraspecies transmission in passerine species.<sup>210,224,229</sup>
- Given the preponderance of passerine birds, more disease spread out of Control Areas in previous outbreaks would be expected to have occurred if these birds played an important role in the transmission of HPAI.
  - As discussed in section 9.1.1, Role of Local Spread Components in Previous AI Outbreaks, most studies indicate limited spread of AI between poultry premises via mechanisms that do not involve the movement of people, vehicles, or equipment.
- The  $CID_{50}$  for HPAI H5N1 infection via aerosol was estimated to be  $10^5$  to  $10^6$   $EID_{50}$ , with higher doses deemed necessary for infection via the oral route (see section 8.7.2, Dose Response in Chickens).
  - The studies cited above demonstrate variability in oropharyngeal and cloacal HPAI virus titers in passerines, depending on the bird species and the H5N1 strain, but when shed titers were measured, most studies indicate they would be adequate to infect chickens via the aerosol route.
- Biosecurity guidelines dictate measures to prevent wild bird access to broiler barns and maintenance of feed bins such that wild birds are neither frequenting nor accessing broiler premises (section 7.4.3.5, Animal, Pest, Insect Control).<sup>201,203</sup>
  - Nonetheless, Burns et al. observed wild birds frequenting and entering poultry barns on premises where the producers were “highly involved in poultry industry management” and, the authors note, may have thus been practicing more stringent biosecurity than other producers.<sup>193</sup>
  - Similarly, while it is industry standard to discourage wild birds from accessing poultry barns as part of a company biosecurity plan, assuring that no birds are entering broiler barns is not considered feasible (BWG, personal communication, June 2016).
  - For more information on potential for contamination of finished feed products by passerine birds, see USDA-APHIS Poultry Feed Risk Assessment.<sup>212</sup>
- Secondary transmission of HPAI from a passerine bird outside the broiler house is unlikely.

- As potential biological vectors, passerine birds shed lower cloacal viral titers, and their fecal volume is small.
- There also is the possibility of mechanical transmission of HPAI virus if plumage or feet were to become contaminated. Preliminary results from a survey of 419 passerine birds<sup>c</sup> on five farms infected with HPAI H5N2 virus and five uninfected farms in Iowa indicates that mechanical transmission through contamination of the external surface of passerine birds is a possibility, although the likelihood is very low (only one external surface swab was positive by matrix gene rRT-PCR and submitted for further testing).<sup>38</sup> As potential mechanical vectors, Passeriformes, due to their small size, can transfer a very small volume of contaminated feces from an infected broiler premises.

#### 9.1.5.1.3 Likelihood Rating and Conclusion

While passerine and columbiforme birds have proven unlikely to play an important role in the transmission of HPAI virus in poultry outbreaks, uncertainty remains as to their potential as vectors, and they have demonstrated the ability to enter secure broiler houses. Thus, the likelihood of HPAI infection via passerine birds in the farm vicinity is *low to moderate*.

#### 9.1.5.2 Likelihood of Infection via Other Non-Aquatic Bird Species in Farm Vicinity

Other non-aquatic avian species such as birds of prey or scavenger species vary greatly in number and behavior around poultry farms. Unlike passerines or columbiforme species, these birds are unlikely to be able to enter poultry barns, resulting in only secondary transmission pathways. Several studies have clearly shown that flying birds transport viruses such as HPAI H5N1.<sup>44</sup> These birds might have contact with manure stored outside the poultry house or manure that is land-applied. Although the quantity of manure wild birds can carry is unknown, as well as the host adaptability of other HPAI virus strains to different wild bird species, it was conservatively (and hypothetically) assumed that wild birds will carry HPAI-contaminated manure if they have contact with it. Additionally, a predatory bird or scavenger may become contaminated with feathers or body fluids of an infected prey item, or potentially pick up and move that prey item to a location closer to a broiler premises.

Common predator and scavenging wild birds undertake a variety of short- and long-distance trips to search for food and cover. These include daily movements to and from hunting/feeding and roosting areas, post-fledging dispersal, and seasonal movements.<sup>44</sup> Scavenger species may be attracted to premises with improperly secured daily mortality carcasses. Species known to scavenge avian carcasses in the U.S. considered in this assessment include vultures, some hawks and eagles, crows, ravens, and magpies. Some gull species known to scavenge are covered separately in section 9.1.4, Role of HPAI Spread to Broiler Flock in a Control Area via Wild Aquatic Birds in the Farm Vicinity.


Finally, wild galliformes also may be attracted to poultry operations.

---

<sup>c</sup> 220 individual birds across 18 species on infected farms, 199 individual birds across 16 species on uninfected farms



- In a survey of infected turkey farms during the 2014-2015 outbreak in Iowa, Minnesota, North Dakota, South Dakota, and Wisconsin (n=81), 26 percent reported seeing wild turkeys, pheasants, and quail around their poultry barns.<sup>38</sup>



**Figure 17.** Pathway for exposure of a broiler farm via scavenging birds or raptor species. A similar process could be demonstrated for wild gallinaceous birds

#### 9.1.5.2.1 Literature Review

Non-passerine non-aquatic birds have not been directly implicated in the spread of HPAI in previous outbreaks, and few such birds have tested positive for AI in the vicinity of outbreaks in poultry or wild waterfowl.

- Turkey vultures (*Cathartes aura*) may visit poultry farms to feed on dead birds. Turkey and black vultures (*Coragyps atratus*) both belong to the order Accipitriformes, family Cathartidae.
  - During the 1983-1984 HPAI H5N2 outbreak in Pennsylvania, Virginia, and Maryland, eight turkey vultures and 22 black vultures from the quarantine zones were tested for H5N2 and none were positive.<sup>230</sup>
- Experimentally infected wild pheasants (order Galliformes) shed the virus in their feces for up to 15 days, demonstrating the potential to transmit HPAI H5N2 (A/Chicken/Penn./1370/83). However, surveillance of wild pheasants in the quarantine area did not support this.<sup>190</sup>
- Raptors found dead during an H5N1 outbreak in wild water birds in Germany in 2006 revealed evidence of H5N1 infection in common buzzards and peregrine falcons.<sup>231</sup> The authors hypothesize that in this H5N1 outbreak in wild water birds, raptor exposure and mortality likely occurred more often in species that hunted or scavenged sick or dead medium-sized prey birds.
  - The highest concentration of H5N1 was found in brain tissue and air sacs, with marked encephalitis as a common finding on histopathology.

- The suspected main cause of death in H5N1-positive raptors is encephalitis.
- No infection was found in other species tested (Eurasian sparrow hawk, common kestrel, white-tailed sea eagle, undetermined species buzzard, undetermined species raptor, red kite, rough-legged buzzard, western marsh-harrier, and goshawk).
- Other birds of prey in the order Accipitriformes, such as the common buzzard (*Buteo buteo*), have become infected in previous HPAI H5N1 outbreaks.
  - 10.5 percent of wild birds testing positive during the 2006 HPAI H5N1 outbreak in Germany were birds of prey, including common buzzards, peregrine falcons, kestrels, and European eagle owls.<sup>232</sup>
    - The buzzards reportedly displayed severe central nervous system infection without systemic virus distribution (unpublished data).
  - An HPAI H5N1-positive common buzzard carcass found in Bulgaria in 2010 contained no gross pathological lesions, suggesting the bird died shortly after infection and would not have served as a reservoir of infection.<sup>233</sup>
- The U.S. Interagency Steering Committee on Avian Influenza in Wild Birds has compiled all U.S. wild bird cases of HPAI H5 from December 2014 to June 2015.<sup>39</sup> Out of 100 positive birds, only seven were from non-passerine non-aquatic species (**Table 10**).

**Table 10:** HPAI H5-positive samples from non-passerine non-aquatic species collected from December 2014 to June 2015 in the U.S.<sup>39</sup>

Date	Species	Lineage	Sampling type (location)
4/14/15	Cooper's hawk	EA/AM H5N2	Mortality (MN)
4/13/15	Snowy owl	EA/AM H5N2	Mortality (WI)
1/20/15	Bald eagle	EA H5N8	Mortality (ID)
1/9/15	Red-tailed hawk	EA/AM H5N2	Mortality (WA)
12/31/14	Red-tailed hawk	EA/AM H5N2	Mortality (WA)
12/29/14	Cooper's hawk	EA/AM H5N2	Mortality (WA)
12/29/14	Peregrine falcon	EA H5N8	Mortality (WA)

- There also were cases of HPAI confirmed in captive wild birds (**Table 11**).<sup>234</sup>

**Table 11.** HPAI-positive samples from captive wild birds in the U.S.<sup>234</sup>

Date	Species	Lineage	Sample location
3/27/15	Captive gyrfalcon	EA/AM H5N2	MT
3/27/15	Captive falcon (hybrid)	EA/AM H5N2	MO
1/29/15	Captive gyrfalcon (2)	EA H5N8	ID
1/16/15	Captive falcons, great horned owl	EA/AM H5N2	ID
12/14/14	Captive gyrfalcon	EA H5N8	WA

- Similarly, cases of HPAI in captive falconry birds in Dubai suggests that these raptors were likely infected through consumption of infected farmed or wild prey items.<sup>235</sup>
- An outbreak of H5N1 clade 2.3.2.1c in captive falconry birds in Dubai and avian prey species at a breeding facility included mortality in gyrfalcons and hybrid gyr/peregrine falcons.<sup>235</sup>

Surveillance of non-passerine non-aquatic birds for AI virus has demonstrated zero to low prevalence in the wild.

- A 2010 survey of antibodies to AI in wild birds revealed zero positives out of 184 black vultures sampled in Mississippi.<sup>102</sup> The authors note that nearly all species of terrestrial birds tested were negative for AI antibodies.
- A survey of antibodies to influenza A in 616 raptors admitted to two U.S. wildlife rehabilitation centers, and 472 peregrine falcons caught at a migratory banding station, found relatively low prevalence of antibodies (subtype not described) in a variety of species. Results of the survey are summarized in the table below (**Table 12**).<sup>236</sup>
  - Antibodies to influenza A (subtyping not possible due to low HI ratio in sera) were found in bald eagles, peregrine falcons, great horned owls, and Cooper’s hawks.
  - No influenza A was found in turkey vultures or black vultures.

**Table 12.** Serologic evidence of influenza A in raptors admitted to two U.S. wildlife rehabilitation centers<sup>236</sup>

Species	Number tested	Number positive	Percent positive
Bald eagle	406	22	5.1
Peregrine falcon	472	1	0.2
Great horned owl	81	1	1.2
Cooper's hawk	100	1	1.0
Turkey vulture	21	0	0
Black vulture	8	0	0

- Gunnarson et al. (2010) sampled nestling white-tailed sea eagles (n=181) and peregrine falcons (n=168) in Sweden for influenza A infection.<sup>237</sup>
  - No evidence of infection or antibodies was detected in any samples.
  - Authors acknowledge that maternal antibodies last less than three weeks in nestlings, and sampling older nestlings that haven't fledged may be a less sensitive screening population than adult birds.
- Peterson et al. (2002) found a 0 percent prevalence of AI virus in wild turkeys (*Melleagris gallopavo*) in a survey of 70 turkeys in Texas.<sup>238</sup>
- Another study of wild captured or hunter-harvested wild bobwhite quail (*Colinus virginianus*) in Texas found prevalence of 1.4 percent using rRT-PCR; however, no virus could be isolated from culture.<sup>239</sup>

Experimental susceptibility of non-passerine birds to HPAI is relatively unstudied.

- Experimentally infected American kestrels (*Falco sparverius*) with H5N1 (A/whooperswan/Mongolia/244/05) demonstrated 100 percent mortality within seven days of inoculation.<sup>240</sup>
  - The American kestrels demonstrated oral viral RNA shedding and infectious virus and, to a lesser extent, cloacal shedding. Infectious viral particles as detected by embryonated egg inoculation were not detected in cloacal samples.
  - Seroconversion occurred by DPI 4 to 5.
  - The most consistent histopathological lesions occurred in brain and pancreas; all infected birds had some evidence of meningitis and encephalitis.
- In commercial Japanese quail, Chukar partridges, ring-necked pheasants, and Guinea fowl experimentally infected with H5N8 (A/GF/WA/14) and H5N2 (A/NP/WA/14), there was 100 percent mortality, with a mean time to death of 2.5 to 3 days (M. Pantin-Jackwood, personal communication, Nov 2015).
  - Clinical signs included listlessness and ruffled feathers.
  - Histopathological signs included necrotic pancreas, mottled spleen, petechial

hemorrhages on the myocardium, and pulmonary hemorrhage.

- Conspecific birds placed in contact with infected partridges or pheasants also showed high mortality from both H5N8 and H5N2 infections.

#### 9.1.5.2.2 Qualitative Analysis

We considered the following qualitative factors in evaluating this pathway:

- To date, HPAI H5N1 has proven to be unique in its ability to infect a variety of species, and more ubiquitous in its prevalence than any other HPAI virus.
- The risk of AI transmission is much lower from a single infected bird than from a population of birds in which infection is established. Additionally, low circulating antibodies, solitary living patterns, and apparent rapid mortality in raptors make risk of spread within these predatory species less likely.
  - As noted in the literature review, surveys of birds of prey and scavenging birds have demonstrated a low prevalence of AI virus, including the more pervasive H5N1.
- Given that some scavenger and other non-passerine species may have relatively large home ranges,<sup>241-243</sup> spread beyond the Control Areas in previous outbreaks would have been expected if these birds played an important role in the transmission of HPAI.
  - Further discussion of avian scavenger species, home ranges, and factors for likelihood of transmission can be found in the dead bird disposal section (see section 9.2.4, Role of HPAI Virus Spread to a Broiler Flock via Dead Bird Disposal).
  - As discussed in section 9.1.1 (Role of Local Spread Components in Previous AI Outbreaks), most studies indicate limited spread of AI between poultry premises via mechanisms that do not involve the movement of people, vehicles, or equipment.
- Biosecurity guidelines dictate measures to prevent wild bird access to broiler barns, manage dead bird disposal, and maintain feed bins such that wild birds are neither frequenting nor accessing broiler premises (section 7.4.3.5, Animal, Pest, Insect Control).<sup>201-203</sup>
  - Given proper disposal of dead birds on broiler premises and the fact that non-passerine and non-columbiforme species have not been observed inside broiler houses, direct contact with poultry is extremely unlikely with standard biosecurity measures in place in the broiler industry (BWG, personal communication, June 2016).

#### 9.1.5.2.3 Likelihood Rating and Conclusion

Other (not passerine or columbiforme) non-aquatic bird species have the potential to contract HPAI virus and have home ranges of adequate size to contain adjacent broiler farms where they potentially may access contaminated carcasses, manure, or other material at an infected broiler premises. However, they are unlikely to have direct contact with poultry flocks if standard biosecurity measures are in place, and their ability to shed virus has not been studied in many species. For the above reasons, the likelihood of HPAI infection via non-passerine non-aquatic birds in the farm vicinity was rated as *low*.


### 9.1.6 Role of HPAI Virus Spread to Broiler Grow-Out Premises near Poultry Live-Haul Routes via Feathers, Feces, and Other Fomites

The evaluation of the risk of HPAI virus transmission to broiler grow-out premises in a Control Area near poultry live-haul routes assumes the release of potentially HPAI-contaminated material from live-haul trailers along roadways and transportation routes in close proximity to a broiler grow-out premises. The birds in transit may originate from premises inside or outside the Control Area. This evaluation is written specifically for broiler grow-out premises. However, since multiple poultry commodities utilize live-bird movements, including broilers to market, turkeys to market, spent layer hens to market, layer pullets to egg production, turkeys from brood to grow-out, and breeder movements, the concepts here can be translated across these other live-bird movements.

As a requirement of the Secure Poultry Supply Plans, the Pre-Movement Isolation Period (PMIP) decreases the likelihood of permitted movement of infected but undetected flocks within a Control Area. Additionally, broiler grow-out premises in a Control Area that wish to request permitted movement must adhere to the greatly intensified biosecurity of the PMIP, which minimizes the likelihood of exposure to virus in the days leading up to movement (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period).

#### 9.1.6.1 Risk of HPAI Virus Transmission to Broiler Grow-out Premises near Poultry Live-haul Routes

The transport of an infected but undetected flock near a broiler grow-out facility represents a potential pathway for local area spread. HPAI virus transfer to premises near the live-haul route could occur via HPAI-contaminated feathers, feces, and other fomites, which may contaminate broiler premises close to the route and may subsequently be tracked into broiler barns. The two specific pathways identified are: (1) HPAI-contaminated fomites from a live-haul truck blow into or are tracked onto a broiler premises and introduce virus to the broiler flock, and (2) a contaminated live-haul road results in a vehicle bringing virus onto broiler premises and subsequent virus transfer into a barn. **Figure 18** diagrams the exposure pathway.



**Figure 18.** Pathway for exposure of a broiler grow-out premises via fomites originating from nearby live-haul route.

### 9.1.6.2 Literature Review

- If infected poultry are transported to processing, the extent of virus contamination available to infect a flock near the live-haul route is affected by virus shedding, virus persistence, and quantity of virus transferred between transfer steps.
  - Estimates of HPAI virus concentrations in feathers, feces, and blood from HPAI-infected poultry generally range between  $10^3$  and  $10^7$  EID<sub>50</sub> per gram or per milliliter of tested substrate, although higher concentrations have been observed in some cases. Various units of measure are used.
    - Immature feathers: In chicken feathers, the median viral titers for three HPAI H5N1 virus strains (A/duck/Sleman/BBVW-1003-34368/2007, A/duck/Sleman/BBVW-598-32226/2007, and A/Muscovy duck/Vietnam/453/2004) tested were  $\sim 10^5$ ,  $\sim 10^6$ , and  $\sim 10^{5.7}$  TCID<sub>50</sub>/0.1 mL for immature pectorosternal feathers, immature flight feathers, and immature tail feathers, respectively, in inoculated broilers after feather samples were ground with a mortar and pestle.<sup>244</sup> From chicks inoculated with an HPAI H7N1 strain (A/Chicken/Italy/5093/99) at 15 days old, viral RNA load was higher in feather pulp than in oropharyngeal and cloacal swabs for most days tested post-inoculation.<sup>245</sup> Feather pulp was obtained by squeezing the calamus (i.e., the feather quill).<sup>245</sup> In detached feather quills from ducks, HPAI viral titers were  $10^{5.5}$  EID<sub>50</sub>/mL and  $10^{6.3}$  EID<sub>50</sub>/mL at day 10 at 4°C (39.2°F) for the two H5N1 virus strains (A/chicken/Miyazaki/ K11/2007 and A/whooper swan/Akita/1/2008) tested, respectively, when four-week-old ducks were inoculated with  $10^7$  EID<sub>50</sub>.<sup>246</sup>
    - Mature feathers: In chickens, viral antigen was detected in feather stromal cells and feather epidermal cells in inoculated (Ck/Miya/K 11/07, Ws/Akita/1/08) seven- and eight-week-old chickens.<sup>247</sup> In ducks, 3.8 percent of mature pectorosternal feather samples were positive post-challenge and, of the virus-positive feathers, viral titers ranged from  $\sim 10^{0.6}$  to  $\sim 10^{4.5}$  TCID<sub>50</sub>/0.1 mL.<sup>244</sup> From inoculated (A/duck/Nigeria/1071-23/2007) 24-week-old Pekin ducks, 31.25 percent of breast and tail feather calami and 37.5 percent of wing feather calami were positive by rRT-PCR at 3 days post-inoculation.<sup>248</sup>
    - Feces: In chicken feces, HPAI viral titers were greater than  $10^9$  ELD<sub>50</sub>/g when chickens were inoculated with 1983 Pennsylvania H5N2 (SEPRL-PA isolate).<sup>78</sup> In turkey feces, HPAI viral titers were estimated to be between  $10^3$  and  $10^5$  EID<sub>50</sub>/mL with 2015 HPAI H5N2 viruses (A/turkey/MN/12528/2015 and A/chicken/IA/13388/2015), interpolated from cloacal swab data (E. Spackman, personal communication, May 2016).<sup>249</sup>
    - Blood: In turkey blood, HPAI viral titers ranged from  $10^1$  to  $10^{5.8}$  EID<sub>50</sub>/0.1 mL at 1-3 days post-inoculation when inoculated with  $10^6$  EID<sub>50</sub> of an H7N1 virus strain (A/chicken/Italy/1067/1999).<sup>62</sup>
  - Once virus is shed, it remains viable for a varying amount of time depending on viral strain and environmental conditions, such as humidity and temperature. Virus persistence is generally longer at cooler temperatures and in more humid conditions.

- For virus persistence data in a range of conditions and on substrates relevant to this pathway, such as feathers, feces, and water, see Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates.
  - This transmission pathway is likely multi-step. The available literature suggests virus concentration decreases when transferred between surfaces. In an experimental setting, mechanical transmission of an enveloped virus after multiple contact steps have occurred.<sup>250</sup>
    - Virus transfer between surfaces for non-AI viruses ranges from undetectable to 46 percent transferred.<sup>251</sup>
    - Mechanical transmission via a multiple-step pathway was documented using porcine reproductive and respiratory syndrome virus (PRRSV) in 1 of 10 replicates by virus isolation and in 8 of 10 replicates by RT-PCR at less than 0°C (32°F) in a swine industry-like setting.<sup>250</sup>
      - Similar to HPAI virus, PRRSV is an enveloped virus shed in feces, urine, semen, aerosolized respiratory secretions, and other bodily fluids.
      - Experimental design simulated a four-step transmission pathway: *PRRSV-inoculated (field strain MN 30-100) carrier attached to undercarriage of vehicle and driven 50 km*→ *Contact between PRRSV-inoculated carrier and driver's boots*→ *Driver re-entered vehicle and drove 50 km*→ *Driver's boots entered farm anteroom*→ *Contact between farm anteroom floor and containers of four surfaces (cardboard, Styrofoam, metal, and plastic)*.
      - PRRSV RNA was detected by PCR in 8 of 10 replicates on three container surfaces (Styrofoam, metal, and plastic) and 7 of 10 replicates on a cardboard container after the final transmission step at less than 0°C (32°F).<sup>250</sup>
      - At 10-16°C (50-60.8°F), infectious PRRSV RNA was detected by PCR in 2 of 10 replicates on the farm anteroom floor.<sup>252</sup>
- Findings from previous disease outbreaks suggest virus transmission to a poultry premises near a live-haul route is possible.
  - In a review of infectious laryngotracheitis (ILT) outbreaks on U.S. broiler operations, some experts have implicated live-haul trucks transporting infectious birds as a probable means of indirect spread to nearby susceptible flocks along the route.<sup>253,254</sup>
  - In the 2002-2003 outbreak of ILT on Mississippi broiler farms, mean distance of the nearest live-haul road to case farms was 0.40 miles, while distance of the nearest live-haul road to control farms was 1 mile (distance to nearest live-haul road [miles]: Odds Ratio=0.54; P-value=0.0392; univariate analysis).<sup>255</sup>
  - In the 1995 outbreak of LPAI H9N2 in Minnesota, spatial observations suggested exposure to the live-haul route used to transport a known infected turkey flock that was sent to slaughter was a risk for premises infection (eight of nine premises within 250 meters of live-haul route became infected) (D. Halvorson, personal communication, June 2016).



- Close proximity to an infected premises has been associated with an increased risk of infection.<sup>133,256,257</sup> As a function of distance, the pathway of infection is not clear. For a detailed examination of the literature on local area spread in AI outbreaks, see Appendix 2: Literature Review on the Role of Local Area Spread in Previous Outbreaks.
- If virus is transferred into a barn, the likelihood of infection is dependent on infectious dose. Mean infectious dose varies with poultry species and virus strain.<sup>105</sup>
  - In chickens, the infectious doses post-challenge with 11 HPAI-viruses of chicken and turkey origin ranged from  $10^{1.2}$  to  $10^{4.7}$  BID<sub>50</sub>.<sup>105</sup> In this study, the infectious doses for most strains tested were between  $10^2$  and  $10^3$  BID<sub>50</sub> and were lower for isolates of chicken origin.<sup>105</sup> With four 2015 H5N2 poultry isolates (A/turkey/Minnesota/12582/2015, A/turkey/South Dakota/12511/2015, A/chicken/Iowa/13388/2015, A/turkey/Arkansas/7791/2015), infectious doses in chickens ranged from  $10^{3.2}$  to  $10^{5.1}$  BID<sub>50</sub>.<sup>40</sup>

### 9.1.6.3 Qualitative Analysis

We considered the following factors in evaluating this pathway:

- While this risk assessment is limited to evaluating risk of HPAI infection on premises located within the Control Area, poultry transport on routes passing through the Control Area may include flocks originating inside or outside the Control Area, which have different movement requirements.
  - Permitted terminal and transfer movements of live poultry originating from within a Control Area likely will require movement from a Monitored Premises, adherence to a PMIP, and rRT-PCR testing in the days preceding movement. The duration of PMIP may vary by sector and type of movement but is determined in part to provide a 95 percent probability or greater of detection in flocks exposed to HPAI virus before the PMIP begins, given a 100 percent effective PMIP [See Appendix 7: Cross-Commodity Pre-Movement Isolation Period). As an example of movement originating from inside the Control Area, **Table 13** shows simulation results of the detection probability for broilers with SPS pre-movement testing and PMIP. This modeling assumed a 100 percent effective PMIP, which prevents flock exposure to virus during the PMIP. For modeling with a PMIP that is not 100 percent effective, see Appendix 10: Supplementary Testing Protocols.
  - Premises located outside a Control Area may not be subject to permitted movement. There may be variation in pre-movement testing as States or Incident Command may require testing for poultry movements from premises in the Free Area,<sup>130</sup> but if not, these premises may not be subject to pre-movement testing requirements beyond routine NPIP surveillance for LPAI. There is also likely variation of biosecurity practices in the Free Area. Biosecurity measures may be heightened in an outbreak scenario, but implementation may differ markedly between premises. For this analysis, the pre-movement surveillance modeled as the method to detect infection prior to movement from outside the Control Area consists of rRT-PCR testing of 2 pools of 11 swabs and a mortality trigger of 3 birds per 1000. Simulation results for broilers are shown in **Table 13**. When detection by a mortality trigger is obtained

depends on transmission parameters and virus characteristics.<sup>258</sup> In the models for movement originating outside a Control Area, the flock could be exposed 1 to 10 days prior to movement as a PMIP is not implemented. Introduction close to movement is more likely to go undetected, and, if infection is not detected, there may be fewer infected undetected birds at movement.

- During the 2014-2015 HPAI outbreak in the United States, approximately one third (36/103) of the positive commercial premises in Minnesota were located outside a Control Area at the time of detection (P. Bonney, personal communication, September 2016).

**Table 13.** Detection probabilities for broilers using three biosecurity and surveillance protocol scenarios. \*

Biosecurity and Surveillance Protocol	Detection Probability
<i>Scenario A</i>	
○ rRT-PCR testing of a pooled sample of 11 swabs each on 2 consecutive days with a 5-day 100% effective PMIP. Second test within 24 hours of movement.	0.98
<i>Scenario B</i>	
○ rRT-PCR testing of a pooled sample of 11 swabs each on 2 consecutive days. Second test within 24 hours of movement. No PMIP implemented.	0.74
<i>Scenario C</i>	
○ Detection under mortality trigger of 3 birds per 1,000 only. No PMIP implemented.	0.54

\*Probabilities estimated from 6,000 simulation iterations using EA/AM HPAI H5N2 strain characteristics and considering virus exposure within 10 days of movement.

- If infected poultry are transported to processing, the initial contamination for this pathway is dependent on HPAI-contaminated material falling from the live-haul trailer. Feathers, feces, and other potential fomites fall from live-haul trailers because they are not enclosed, as shown in **Figures 19-21** (D. Halvorson, personal communication, July 2016). Day-old chicks and poults are transferred in different vehicles and are totally enclosed.
  - Netting systems to contain feathers in the live-haul trailer typically are not used because they are ineffective and create an additional biosecurity issue as nets are difficult to clean. Thus, nets were not used on live-haul trucks during the 2014-2015 or 2016 U.S. HPAI outbreaks (Turkey Working Group [TWG], personal communication, July 2016).

- **Figures 19-21** show the openings in crates used for live-haul in the broiler and turkey industry.



**Figure 19.** Crates filled with broilers to be loaded onto a live-haul truck (Photo courtesy of GNP Company).



**Figure 20.** Live-haul trailer of turkeys after load-out (Photo: Anonymous)



**Figure 21.** Live-haul trailer of turkeys (Photo: Jill Nezworski.)

- The likelihood of this contamination reaching the premises and infecting the flock may depend on the distance of the premises from the live-haul road, weather conditions, natural barriers/landscape, and virus transfer steps.
  - Close proximity to a road seems quite likely for the majority of poultry premises, regardless of species.
    - In a survey of growers of broiler and breeder-layer premises in Georgia, 68 percent of growers were within 440 yards (0.25 miles) of a public road.<sup>259</sup>
    - Among turkey premises, a case-control study following the 2016 H7N8 outbreak in Indiana reported similar distances between a majority of farms and the nearest public gravel or dirt road (55 percent of case farms were within 499 yards of a public road; 60 percent of control farms were within 499 yards of a public road).<sup>260</sup>
  - During disease outbreaks, the distance between the live-haul road and poultry premises may be efficiently maximized by routing, when possible or based on company requests. Poultry live-haul routes are determined by individual companies based on transit time and welfare of the birds on the truck, in addition to consideration of any poultry premises in the area (BWG, personal communication, July 2016; TWG, personal communication, August 2016).
    - Poultry companies near outbreaks have communicated frequently and shared locations of premises; although knowledge of the locations of other poultry

premises by a particular company or veterinarian varies (BWG, personal communication, July 2016; TWG, personal communication, August 2016).

- In geographic areas with many poultry production premises, routing may take on increased importance due to the density of susceptible birds near a route.
- During the 2014-2015 HPAI outbreaks in the U.S., live-haul routing requirements were not needed for approval of permitted movement in Minnesota (Minnesota Board of Animal Health, personal communication, October 2016) and were not mandated by Incident Command in 2016 in Indiana (TWG, personal communication, July 2016).
  - However, for permitted movement from premises in an HPAI Control Area, both the Secure Broiler Supply (SBS) Plan and Secure Turkey Supply (STS) Plan recommend live-haul route approval from the Incident Command team or routes selected in consultation with a poultry veterinarian or production manager.<sup>261,262</sup>
- In the management of ILT outbreaks, geographic information system (GIS)-assisted live-haul route planning has been used to minimize the number of farms within a specified distance along the route to processing from a broiler premises in a Biosecurity Zone.<sup>254</sup>
- The transmission steps of this pathway could be affected by weather conditions, natural barriers/landscape, and C&D.
  - Since feathers are lightweight, transmission to the premises via feathers over short distances might be a possibility. Weather conditions such as wind and precipitation as well as natural barriers/landscape between the live-haul route and broiler premises may affect whether virus arrives on-farm. As most broiler barns are fully enclosed, feathers are not likely to blow directly into a poultry house.
- Virus transmission from a live-haul trailer to a premises close to the road represents a multi-step transmission pathway. With each virus transfer step, virus concentration is likely to decrease. Among the potential pathways identified, blowing of HPAI-contaminated fomites from a live-haul trailer to a broiler premises, with subsequent transfer into the barn, involves fewer transfer steps compared to a vehicle bringing virus to a broiler premises from a contaminated live-haul road, followed by transfer into the barn.
- The minimum biosecurity guidelines for poultry premises participating in the NPIP and the greatly intensified biosecurity of the PMIP for premises in a Control Area that wish to request movement during an outbreak are designed to reduce the likelihood that contamination which reaches the premises would subsequently infect the flock.
  - Standardized biosecurity in the poultry industry, such as rules about entering the perimeter buffer area, crossing the line of separation, and managing vehicle access, are intended to prevent flock exposure to diseases.<sup>2</sup>
  - For broiler grow-out premises in a Control Area that wish to request permission to move live birds, the enhanced biosecurity of the PMIP minimizes the chances of a flock being exposed to HPAI for the five-day period prior to movement. The PMIP

reduces the likelihood of a vehicle contaminated from a live-haul road bringing virus to a broiler operation, as all vehicles will be cleaned and disinfected before entering the premises. A requirement to use barn-specific footwear to enter the poultry house during the PMIP minimizes introduction of virus via tracking into the barn on the boots of personnel. The pertinent biosecurity guidelines of the PMIP are:

- Limiting visits to the premises to critical operational visits
- Requiring specific biosecurity for those critical visits (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period)
- Vehicles and equipment arriving on a broiler premises may be difficult to disinfect thoroughly, especially during harsh winter conditions. Thus, virus may remain on vehicles contaminated from the live-haul route, despite C&D steps.
- Previously, ten experienced poultry veterinarians evaluated the risk of infecting susceptible poultry flocks via the microbial load from two truckloads of turkeys shedding a generic pathogen at varying distances (results shown in **Table 14**).<sup>263</sup>

**Table 14.** Perceived qualitative risk posed by two truckloads of turkeys at varying distances from susceptible poultry based on expert opinion, as reported in Halvorson and Heuston 2006.<sup>263</sup>

Distance to susceptible poultry	10 m	100 m	1,000 m	10,000 m
Risk rating*	Intolerable	Intolerable	Low	Negligible

\*Risk rating scale of negligible, low, moderate, high, and intolerable.

- The results of the veterinarian survey were strongly correlated ( $P < 0.01$ ) with the values calculated with an exposure risk index, which took into account mass of contaminant, percentage available for transmission, initial titer of the pathogen, age of contaminant/half-life of virus, and distance to susceptible poultry.<sup>263</sup>

#### 9.1.6.4 Risk Rating and Conclusion

##### 9.1.6.4.1 Risk of HPAI Transmission to Broiler Grow-Out Premises in a Control Area near Route of Live-Haul Trailer

Literature review and expert opinion indicate a potential for increased risk when a poultry premises is located close to live-haul routes for transporting infectious birds. The requirements for permitted movement of live birds in the SPS Plans, specifically implementing an effective PMIP, increase the likelihood of detection prior to scheduled movements that originate in a Control Area. Vehicles transporting live poultry from a Monitored Premises that has met the requirements of the SPS plan (PMIP, PCR testing) are less likely to represent an infected but undetected movement than if the PMIP and testing were not in place. As presented in Appendix 10: Supplementary Testing Protocols, it is also unlikely that flocks moved after a PMIP and testing would contain large numbers of clinically infected birds.

During the 2014-2015 HPAI outbreak in the U.S., infected premises were identified both inside and outside Control Areas at the time of detection. It is expected that biosecurity may be heightened during an outbreak scenario; however, there may be variation in biosecurity and pre-

movement testing from the Free Area. With the use of a mortality trigger alone or pre-movement testing without implementing a PMIP, the likelihood of detecting HPAI virus in a flock before movement is estimated to be substantially lower than the detection probability with a PMIP in place.

**9.1.6.4.2 Conclusion**

Considering the above factors, assuming that the preventive measures specified in the SPS Plans are strictly followed when moving live poultry and given that live-haul vehicles passing a premises within the Control Area may originate from within or outside the Control Area, the following risk ratings are provided:

The likelihood of HPAI infection at a broiler grow-out Monitored Premises located within a Control Area due to HPAI-infected poultry or contaminated live-haul vehicles passing on a nearby road is rated:

Characteristics of live-haul vehicle	Likelihood rating at given distance (between live-haul road and poultry premises)		
	<100 meters	100-1000 meters	>1000 meters
Truck hauling birds that had no PMIP and no tests	<i>High</i>	<i>Moderate</i>	<i>Low</i>
Truck hauling birds that had less than optimum PMIP and tests (80% effective PMIP; delayed testing; or load-out >24 hours)	<i>Low</i>	<i>Very Low</i>	<i>Negligible</i>
Truck hauling birds that had a PMIP & rRT-PCR negative birds (100% effective PMIP; two tests within 24 hours of move and completion within 24 hours)	<i>Very Low</i>	<i>Negligible</i>	<i>Negligible</i>

**9.2 Pathways for a Broiler Flock Becoming Infected with HPAI via Movements of People, Vehicles, or Equipment**

**9.2.1 Role of HPAI Virus Spread to a Broiler Flock via Critical Operational Visits during PMIP**

Routine operational visits to a farm include feed delivery, gas delivery, veterinarians, shavings suppliers, meter readers, repairmen, service personnel, managers, vaccination crews, clean-out services, de-caking services, pit inspectors, and other visitors. The SBS Plan requires most of the operational visits to be scheduled to occur outside of the PMIP before moving broilers. However, some critical operational visits, such as feed delivery or emergency repair services, would need to continue during the PMIP. A survey of broiler growers (convenience sample) indicated an approximate mean frequency of feed delivery of once every two days (range one to five days).<sup>264</sup> The frequency of outside repair or maintenance services varied between two and five times a year in studies in the literature.<sup>264,265</sup>

**9.2.1.1 Likelihood of Infection via Feeds**

During the 2015 HPAI outbreak in Minnesota and Iowa, risk managers were concerned about biosecurity practices related to storage of feed ingredients and finished feed. Specifically, the observation of corn piles stored on the ground and contaminated with wild bird feces raised concerns about the possibility that contaminated corn might be a pathway for HPAI virus introduction and spread. Additional concerns include the chance that finished feed could become contaminated by wild birds through breaches in biosecurity at the feed mill or feed storage bins on a farm.<sup>212</sup> Feedback from the BWG revealed that feed spilled on the farm (outside the barn) would not be fed to poultry without first being reprocessed at a feed mill to limit exposure to contaminants or infected material (BWG, personal communication, June 2016).

On most broiler operations, birds over three weeks of age have transitioned to a finisher feed. Some operations may use a withdrawal-compliant feed in the days leading up to slaughter to adhere to any drug residue withdrawal periods.<sup>10,15,17</sup> Finisher formulations may be pellets, crumbles (broken pellets), or mash. The Feed Risk Assessment assessed the risk of HPAI transmission to poultry fed contaminated feed in a variety of scenarios listed in **Table 15**. Further information can be found in the Feed Risk Assessment.<sup>212</sup>

**Table 15.** Risk ratings for various types of poultry feed products<sup>212</sup>

<b>Pathway</b>	<b>Risk</b>
Potential that corn stored on ground is contaminated with feces from wild migratory birds	<i>Low to very low*</i>
Potential that pelleted feed made with contaminated corn transmits HPAI to poultry flock	<i>Negligible</i>
Potential that untreated mash feed made with contaminated corn transmits HPAI to poultry flock	<i>Low to very low</i>
Potential that formaldehyde-treated mash feed made with contaminated corn transmits HPAI to poultry flock	<i>Negligible</i>
Potential that finished feed contaminated by perching birds at feed mill or storage bins on farm transmits HPAI to poultry flock	<i>Low to very low</i>


\*Under fall and spring seasonal conditions

**9.2.1.2 Likelihood of Infection via Feed Delivery or Other Critical Operations Visits**

Under normal operations, feed vehicles may deliver to multiple farms the same day (a range of 0 to 5 deliveries per day was used in Dorea et al. [2010]).<sup>265</sup> The possible pathways for transmission via feed delivery involve contamination of the vehicle or driver at an infected but undetected farm, and subsequent cross-contamination of a virus-free broiler premises. During the PMIP, only the following critical operational visits to the premises are allowed:

- Feed delivery in a dedicated truck directly from a stand-alone feed mill
- Emergency repair of critical mechanical equipment
- Service visits to address changes in bird health

It has been suggested that company feed delivery should involve dedicated trucks for use in the quarantine zone. Additionally, trucks within the Control Area should be limited to a single premises delivery per load rather than delivering feed to several farms on a route.<sup>266</sup> In addition to feed delivery, other critical operations visits (emergency maintenance, service visits, etc.) are assumed to offer a similar potential pathway to that of feed trucks.



**Figure 22.** Pathway for exposure of a broiler farm via a feed delivery or critical visitor vehicle

### 9.2.1.2.1 Literature Review

- Feed truck deliveries were the most commonly reported visits at 84 percent of Georgia poultry farms surveyed.<sup>259</sup>
- In a survey of contract broiler growers in the U.S., feed delivery averaged once every two days (range one to five days). In this analysis, feed delivery accounted for 74 percent of total point estimates of risk for farms using the same integrator as index farm.<sup>264</sup> Of note, this model considered all vehicle/visitor traffic to a farm, even activities that would not be allowed under PMIP, and did not account for differences in magnitude of virus contamination in different types of visitor contacts.
- Similarly, a stochastic model by Dorea et al (2010) predicted that off-farm spread of HPAI by visitors is most frequently associated with feed trucks and company personnel.<sup>265</sup>
  - Of the reviewed HPAI and LPAI outbreaks in the U.S., feed delivery or contaminated feed was implicated in only the 1983-1984 Pennsylvania outbreak (mixed LPAI/HPAI).<sup>267</sup>
- In a model of risk for ILT infection during an outbreak, farms with more visits per month by feed trucks were associated with higher risk for ILT (OR=1.18; P=0.0099).<sup>255</sup>



### 9.2.1.2.2 Qualitative Analysis

We considered the following factors in evaluating this pathway:

- Feed truck visits and feed delivery are likely to occur on most, if not all, broiler operations during the PMIP.
- While feed truck visits will be the most frequent type of contact during PMIP, they are subject to specific biosecurity guidelines outlined in the SBS plan (see SBS PMIP recommendations for a full list of biosecurity requirements relevant to feed trucks and drivers).<sup>261</sup>
  - Within the Infected Zone, the outside of the vehicle must be C&D at or near the broiler premises both before entering and after leaving the broiler premises.
  - A dedicated feed truck will be used for the Infected Zone.
  - Feed truck drivers may not enter the poultry house and must put on disposable boots and gloves before exiting the truck cab.
  - Feed truck drivers will sanitize or wash hands before leaving and upon re-entering the cab, and will spray the cab interior floors, pedals, and bottoms of feet after every stop.
- The SBS plan also outlines biosecurity practices for other critical or emergency operations visits (such as those by poultry house repair persons).<sup>261</sup>
  - Personnel who have contact with poultry on other premises must shower and change clothes before entering the premises and also wear necessary protective clothing and footwear as described in appropriate biosecurity protocols.
  - All vehicles and equipment will be C&D prior to entering premises.
- Critical visitors other than those associated with feed delivery may be required to enter a poultry house to complete their necessary tasks. Visitors who enter poultry houses during PMIP may contact birds directly, thus decreasing the number of steps in the potential pathway to infection diagrammed above.
  - Relative to feed deliveries, however, other critical operations visits likely happen less frequently, and visitors may or may not have visited other poultry farms recently.

### 9.2.1.2.3 Likelihood Rating and Conclusion

Critical operations visits will be limited during PMIP; however, delivery of feed during this period is likely and the potential for emergency maintenance visits also exists. Assuming all requirements for biosecurity during PMIP are followed, the likelihood of introducing HPAI virus to a broiler flock by feed, feed delivery, and critical operations visits during PMIP is as follows:


Pathway	Likelihood
Contaminated feed	<i>Negligible</i>
Feed delivery (driver and/or vehicle)	<i>Low</i>
Other critical ops visitor (personnel and/or vehicle)	<i>Low to Moderate</i>

## 9.2.2 Role of HPAI Virus Spread to a Broiler Flock via Growers or Employees and their Vehicles Entering the Premises

Off-site movements of poultry growers, their families, and their employees have been implicated as risk factors for disease transmission in previous outbreaks.<sup>83,255</sup> During the PMIP, growers will not be permitted to visit other poultry farms during the five days leading up to load-out. Off-site social contacts with other growers, however, may still occur. In addition, farm employees will not be allowed to enter the poultry house during the PMIP if they have contact with other live poultry.<sup>261</sup> Other non-critical visitors are prohibited on the premises during the PMIP. For a discussion on critical operations visitors and their vehicles, see section 9.2.1, Role of HPAI Virus Spread to a Broiler Flock via Critical Operational Visits During PMIP. For a discussion of any critical tools or equipment brought onto poultry premises during these visits, please see section 9.2.3, Role of HPAI Virus Spread to a Broiler Flock via Machinery or Equipment Shared between Multiple Premises.

### 9.2.2.1 Likelihood of Infection via Movement of Growers

In a national survey of broiler poultry growers, Leibler et al. (2010) reported an average frequency of social visits between growers of once per month, with a range of 10 times a month to no visits.<sup>264</sup> In two Georgia counties, however, 49 percent of growers reported personal interactions with another grower in the previous week.<sup>259</sup> While 64 percent of these growers reported interacting one or two times during that week, 20 percent interacted with another grower five or more times during that week. The possible pathways for transmission via social contacts between growers involve contamination of the grower’s clothes, shoes, hands, or vehicle at a meeting place with a grower from an infected but undetected farm, and subsequent cross-contamination of a virus-free poultry premises. These pathways are shown below in **Figure 23**.



**Figure 23.** Pathway for exposure of a poultry premises due to virus introduction by grower. Similar pathway likely exists for employees working on poultry premises.

### 9.2.2.1.1 Literature Review

- Social contacts between growers have been evaluated as a risk in disease transmission.
  - In a study combining a national survey of broiler poultry growers and stochastic disease modeling, Leibler et al. (2010) determined that broiler grower social contacts contribute less than 1 percent to the AI disease transmission risk attributable to contacts between farms.<sup>264</sup> Their estimated frequency of social contacts of once monthly is, however, far lower than that determined by another survey,<sup>259</sup> which the authors allow may lead to an underestimated risk. Nonetheless, the social contacts in Leibler et al. (2010) were “at a given visitor-receiving farm,” and the PMIP measures prohibit such on-farm visits.<sup>264</sup>
  - In addition to other factors, Dunn et al. (2003) noted that there were significant social ties between growers in the H7N2 LPAI in the 2001-2002 Pennsylvania outbreak in broiler chickens. Five infected flocks had growers who all belonged to the same church group, two flocks were managed by brothers, and two others were managed by individuals with close social and business ties.<sup>268</sup>
  - Analytical studies on disease transmission resulting from off-farm social contact between growers are lacking.
- HPAI virus has the potential to be transmitted via feces-caked shoes or vehicle tires, depending on ambient temperature, humidity, and elapsed time. For additional information on virus survival on various surfaces and under various conditions, see Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates.
  - At low ambient temperatures of 4.0-6.7°C (39-44°F) and low to moderate relative humidity (15.2 to 46.3 percent), HPAI H5N1 (A/Vietnam/1203/2004) in chicken feces remained viable until day 13.<sup>269</sup>
    - However, at temperatures closer to summer conditions in the United States (72.3-74.6°F and 89.1-91.2 percent relative humidity), the same HPAI H5N1 virus strain in chicken feces was inactivated at day 4.<sup>269</sup>
  - On two rubber surfaces (gumboot and tire) at an unspecified room temperature, LPAI H13N7 was below the detectable limit at day 6.<sup>270</sup>
- Glanville et al. (2010) used modeling to predict the average probability of HPAI H5N1 virus transmission via contaminated shoes from a house *in which an infection is beginning* into the house on another farm (without cleaning and disinfecting shoes) to be P= 0.039 to 0.15 per transfer event.<sup>271</sup>
  - The model was based on a small-scale broiler farm in Indonesia, and model parameters were estimated from survey data, literature review, and expert opinion.
  - Variables affecting the risk estimation include viral concentration on shoes after arriving at the second broiler farm, as well as the proportion of fecal matter (and virus) transferred from the shoes.

- In the same study, imposing a mandatory 24-hour downtime between farms decreased the predicted probability of transmission to P=0.0016 in this exploratory model.
- Respiratory viruses can be transmitted via human hands, though studies with HPAI virus are lacking.
  - As detailed in Appendix 5 of the Risk Assessment of the Movement of Broiler Hatching Eggs During an HPAI Outbreak, several studies have determined the transfer rate for various non-AI viruses between different surfaces, including from fingerpad to fingerpad.<sup>251</sup> Depending on the virus, percentage transferred via fingerpads ranged from undetectable to 23 percent.
  - Ansari et al. (1991) demonstrated that 20 minutes after deposition on donor fingertips, 0.7 percent of human rhinovirus transferred to recipient fingertips.<sup>272</sup> On the other hand, transfer of human parainfluenza virus was undetectable at 20 minutes post-deposition. Both parainfluenza and rhinovirus are single-stranded RNA viruses similar to influenza.
  - Assuming a virus transmission efficiency of 0-20 percent, and based on data extrapolation from other viruses (including the above study), modeling by Glanville et al. (2010) demonstrated an average 5 percent chance of a bird being infected with HPAI H5N1 virus via hand contact with someone who had *directly* handled an infected bird at another farm.<sup>271</sup> This estimate applies only to the first bird handled and incorporates the effect of estimated travel time—specific to the study locale in Indonesia—on virus decay.

#### 9.2.2.1.2 Qualitative Analysis

We considered the following qualitative factors for evaluating this pathway:

- There is the potential for growers or members of their households to have regular social or other contacts with other growers. During the PMIP, however, these contacts will likely occur off the poultry premises.<sup>261</sup>
- Growers or household members who may potentially become contaminated via social contacts should, however, change clothes and shoes before coming into contact with birds on their premises.
  - Cross-commodity PMIP measures state that during PMIP, growers must wear clothing dedicated to the farm and shoes dedicated to the barn before entering poultry houses. See Appendix 7: Cross-Commodity Pre-Movement Isolation Period.
- The level of contamination on the person a grower is meeting, however, may be variable.
  - Other growers whose premises are operating under heightened PMIP biosecurity may represent a lower risk as they will have taken measures to remove any potential virus contamination before departing the premises.
    - SBS measures state that growers should shower and change to clean clothes before leaving the farm during PMIP.<sup>261</sup>


- As detailed in Appendix 6 of Broiler Hatching Eggs Risk Assessment,<sup>5</sup> several studies have demonstrated the effectiveness of showering and changing clothes in preventing the transmission of infectious diseases.
    - There are no cleaning or disinfection stipulations for growers who are not observing a PMIP. While it is reasonable to assume that biosecurity may be heightened in the face of an HPAI outbreak, the practices utilized on individual commercial or noncommercial poultry premises will likely vary.
  - Additionally, there are no required mitigation steps for personal vehicles moving on or off the premises prior to or during PMIP.
    - Virus could contaminate the exterior of a truck that has been used close to a poultry barn.
      - As outlined above, virus may survive days to weeks, depending on weather conditions and type of contaminated surface.
    - A grower with contaminated boots, hands, or clothing may drive on his or her premises (for example, from working in a poultry barn to residence on the same premises) without any C&D step. This contamination may remain in the cab of a vehicle, thus re-contaminating an individual who uses that vehicle to drive off-site to meet with another grower.
  - The potential pathways involve multiple virus transfer steps between contact surfaces. In general, the chances of the pathway resulting in virus transmission decrease with the number of contact steps that need to occur. Furthermore, even if the transfer steps occur, there would likely be a substantial reduction in the virus concentration transferred with each contact step. This is because only a fraction of the virus (6 to 27 percent) on a donor surface is transferred to the recipient surface in each direct contact event.<sup>5</sup>
    - Viral contamination on the exterior of the vehicle on an infected and undetected farm, already reduced by dilution outside the house, would undergo multiple transfer steps with viral load reduction (*e.g., vehicle tires → travel to social meeting place → ground → tires of vehicle from uninfected farm → travel to uninfected farm → ground → grower's boots → uninfected barn*).
    - If, however, the social contact were directly contaminated and the grower contaminated the interior of the vehicle, which is not cleaned or disinfected before use on farm, fewer contact steps are needed (*e.g., contaminated grower colleague → grower → vehicle → re-contamination of grower hands/clothes → uninfected barn*).
      - In this scenario, contamination in the interior of a vehicle serves as a point of re-contamination even if a grower were to change clothes and boots before working with poultry.
    - In the period before the PMIP begins, growers may visit other poultry premises, thus decreasing the number of transfer steps needed to bring virus onto the premises, where it may be tracked into the barn during PMIP.
  - Biosecurity measures such as PPE, dedicated work clothing, barn-dedicated footwear, showers, and hand hygiene further reduce the likelihood of virus transmission. In an

outbreak situation, it is expected that biosecurity measures may be heightened on many premises in addition to those undergoing the PMIP (BWG, personal communication, May 2016)

- Appendix 6 of the Broiler Hatching Eggs Risk Assessment details the effectiveness of PPE and hand hygiene in mitigating the transmission of infectious diseases.<sup>5</sup>

### 9.2.2.2 Likelihood of Infection via Farm Employees

For broiler premises whose employees work exclusively on one poultry premises and do not have contact with other commercial or noncommercial flocks, risks are likely similar to those for poultry growers. However, in a small national survey of broiler poultry growers, Leibler et al. (2010) found that 12.5 percent of respondents employed part-time workers.<sup>264</sup> The authors assumed that part-time workers move among farms that employ day laborers. As stated in the PMIP measures, laborers who are employed on other poultry premises or have contact with other poultry are prohibited from entering poultry houses during the PMIP (SBS Plan), but continue to work on the premises. Additionally, contact (such as through hunting) with waterfowl species known to harbor HPAI virus is not addressed in the SBS Plan guidance. The potential pathway is described below:



**Figure 24.** Pathway for exposure of a poultry premises due to virus introduction by an employee who has contact with birds outside the premises of interest. While these employees are not allowed inside barns during PMIP, they may be permitted to continue working on a broiler premises.

#### 9.2.2.2.1 Literature Review

- Part-time employees have been implicated in the spread of poultry viruses in previous outbreaks.
  - In their study of risk factors in the spread of LPAI H7N2 in the 2002 Virginia outbreak, McQuiston et al. (2005) found the likelihood of infection to be significantly

- higher among farms using nonfamily caretakers to work in the poultry houses, with an odds ratio of 2.1 in the multivariate analysis.<sup>83</sup>
- In the 1999-2000 H7N1 outbreaks in Italy, which included LPAI and HPAI outbreaks in many poultry types, it has been suggested that temporary staff on larger farms may have contributed to the identification of larger farm sizes as a risk factor for infection.<sup>135</sup>
  - In the 1979 LPAI H7N3 outbreaks in four Texas turkey flocks, two of the flocks with AI shared personnel and an insemination crew.<sup>273</sup>
  - Although this does not apply exclusively to part-time workers, Alexander stated that the dominant route of secondary spread in domestic poultry has occurred via people and that farm owners and caretaker staff have been implicated in the spread of AI.<sup>45</sup>
  - As detailed in Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates, HPAI virus has the potential to survive for prolonged periods in cool, moist conditions.

#### 9.2.2.2.2 Qualitative Analysis

In addition to the factors outlined above for growers, we considered the following qualitative factors for evaluating this pathway:

- PMIP measures reduce the risk of HPAI virus transmission via part-time employees.
  - Farm laborers will not be allowed to enter the house during the PMIP if they work on another commercial operation with live poultry or have contact with a noncommercial flock. This adds an additional viral transfer step, e.g., *contaminated hands/shoes/tires of part-time employee* → *travel from infected and undetected farm to uninfected farm* → *ground/surfaces at uninfected farm outside of poultry house* → *hands/shoes of full-time personnel* → *poultry house*.
  - Farm laborers will also be required to follow strict biosecurity measures during the PMIP.
    - If a full-time employee were to become cross-contaminated via contact with a part-time employee outside the poultry house, farm-specific clothing required during PMIP may still permit virus to enter the barn; however, barn-specific footwear may decrease the amount of contamination on boots or shoes worn into the poultry house.
- According to industry representatives, many company biosecurity plans dictate that employees shower and change clothes after contact with wild waterfowl (for example, when hunting) before entering a broiler premises. It is uncommon, however, for companies to prohibit waterfowl hunting by growers or employees working with broilers (BWG, personal communication, July 2016).
  - A detailed assessment of the risks of HPAI contamination or infection due to wild birds can be found in section 9.1.4, Role of HPAI Spread to Broiler Flock in a Control Area via Wild Aquatic Birds in the Farm Vicinity, and section 9.1.5, Role of HPAI Virus Spread to Broiler Flock via Wild Non-aquatic Birds in Farm Vicinity.



- Contact with potentially infected or contaminated wild bird carcasses or contamination of a vehicle by an employee who does not have contact with other domestic poultry species may be one type of contact with potentially infectious material in which the employee is not expressly prohibited from entering a poultry barn during PMIP.

**9.2.2.3 Likelihood Rating and Conclusion**

Although some contact may be unavoidable, it is recommended that growers minimize unnecessary contact with other growers during the PMIP and limit travel to other poultry premises during the entire grow period. Still, social and other non-business contacts have been documented between growers and members of their families. Similarly, while task-specific crews (such as vaccination or turkey insemination workers) and day-laborers have been documented in the literature, many companies suggest that employees should limit contact with other commercial or noncommercial poultry flocks outside their job duties. During the PMIP, vehicle and visitor traffic to a poultry premises is likely to be decreased to include only critical visitors, employees, and growers. The prevention of HPAI virus transmission by growers and employees during the PMIP is dependent on close adherence to the biosecurity measures outlined in the PMIP.

Provided the SBS PMIP measures for growers and employees are strictly followed, the likelihood of HPAI transmission during the PMIP is as follows:

<b>Person type</b>	<b>Likelihood Rating</b>
Critical operations visitors and vehicles	See section 9.2.1.1 Likelihood of infection via feeds
Growers and employees entering poultry barns during PMIP	<i>Low</i>
Employees who may contact other birds (not entering barns during PMIP)	<i>Very low</i>


**9.2.3 Role of HPAI Virus Spread to a Broiler Flock via Machinery or Equipment Shared between Multiple Premises**

Many types of equipment are used to maintain a broiler flock and premises, including specialized types of equipment that may be shared between premises for activities such as house maintenance, removal of manure, and preparing an empty house for the next flock of chicks. Shared equipment may be owned and managed by an integrator, a grower, or a visitor who utilizes an item on multiple poultry operations. During PMIP, no equipment may be brought onto the premises, except in the case of tools or items required to complete a critical operations visit. Such equipment should be cleaned and disinfected before entering a poultry barn during PMIP.<sup>261</sup> There are, however, no stipulations about the types and disinfection of equipment which may be brought onto the premises in the days leading up to PMIP.



### 9.2.3.1 Likelihood of Infection via Shared Machinery or Equipment

Shared equipment between premises may be returned to its owner (if grower-owned) or staged on a broiler premises (if owned by an integrator or a third party) at any point in the days leading up to PMIP. While birds may not directly contact this equipment, it may arrive contaminated with HPAI virus, potentially contaminating the ground around a barn and/or personnel who may then track virus into a broiler barn. Additionally, some shared equipment may enter a poultry barn in the final grow-out period during routine tasks, disease investigation, or building maintenance. **Figure 25** illustrates the transmission pathway.



**Figure 25.** Pathway for exposure of a broiler flock via shared machinery or equipment during final grow-out period.

### 9.2.3.2 Literature Review

- Movement of contaminated equipment and, more specifically, shared equipment has been implicated as a potential transmission pathway in previous AI outbreaks in the U.S. poultry industry.
  - In the 2015 U.S. HPAI outbreak on turkey premises in the Upper Midwest (IA, MN, ND, SD, and WI), a descriptive analysis of epidemiologic findings based on narrative responses from infected turkey farms identified highly likely transmission routes for some of them. Of those routes, two involved sharing equipment, with the onset of clinical signs occurring 10 to 11 days after potential exposure to the shared equipment.<sup>38</sup>
  - On layer premises in the 2014-2015 HPAI outbreak, preliminary univariate analysis of a case-control study considered sharing of egg racks or pallets and egg flats as potential infection risk factors for entry into the farm-level multivariate model (egg racks or pallets shared in previous 14 days: 29% of case farms, 11% of control farms,  $P=0.08$ ; egg flats shared in previous 14 days: 30% of case farms, 14% of control farms,  $P=0.17$ ). Because of model instability due to sparse data, equipment sharing

- was not included in the multivariate model, but the authors concluded that this could be a risk factor for infection.<sup>192</sup>
- In the 2002 LPAI H7N2 outbreak in Virginia the pattern of virus spread indicated likely movement by fomites, people, and equipment contaminated with virus.<sup>72</sup> However, in a case-control study of infected premises during the same outbreak, there was no statistically significant difference in the occurrence of equipment sharing on infected premises versus non-infected premises (farm equipment borrowed or loaned: case farms, 17.3%, control farms, 16.7%; P=0.88).<sup>83</sup>
  - In the 1996-1998 LPAI H7N2 outbreak in Pennsylvania, the source of virus exposure for one of 25 positive premises was suspected to be equipment contact.<sup>162</sup>
  - Virus spread was attributed in part to equipment in the 1983-1984 HPAI H5N2 outbreak in Pennsylvania and in the 2000-2002 LPAI H6N2 outbreak in California.<sup>267</sup>
    - During the 1983-1984 HPAI H5N2 outbreak in Pennsylvania, Utterback identified movement of products, materials, equipment, tools, and supplies that may contact birds as a potential mode of AI transmission between flocks. In the analysis, the potential high-risk equipment included egg flats; equipment used for manure, dead bird disposal, and feed; and equipment moved from farm to farm.<sup>274</sup>
  - Sharing of equipment has been associated with disease spread in outbreaks of other viral pathogens of poultry. In a multivariate model with matched controls, removing litter using shared equipment increased the odds of infectious laryngotracheitis (ILT) infection on Mississippi broiler farms in a 2002-2003 outbreak (caked and/or total litter removed from house and/or farm using shared equipment: OR=5.39; P=0.0378).<sup>255</sup>
  - Based on the available data in the literature, equipment sharing occurs in the broiler, turkey, and layer industries during both outbreak and non-outbreak situations. The types and frequency of equipment sharing vary between operations, sectors, and regions.
    - Regional biosecurity surveys of Georgia broiler and breeder-layer growers reported that 8 to 25.8 percent of respondents share equipment.<sup>259,275</sup> Findings by Dorea et al. indicate shared equipment is most commonly used by one other grower, although this ranged from 1 to 20 growers utilizing the same piece of equipment.<sup>275</sup>
  - During outbreak scenarios (LPAI H7N2 in Virginia in 2002 and HPAI/LPAI H7N8 in Indiana in 2016), equipment was shared at 17.3 percent (26/150) and 11.1 percent (1/9) of surveyed case premises, respectively.<sup>83,260</sup> Among case premises in Indiana in 2016, the only equipment reported to be shared was a live-haul loader.<sup>42</sup>
    - Data on sharing of specific types of equipment, as published in a case-control study of infected Iowa layer flocks and a case series of infected turkey flocks in the upper Midwest during the 2015 HPAI outbreak, are shown in **Table 16**.

**Table 16.** Percentage of layer and turkey premises reporting equipment sharing (i.e., not farm-specific equipment) during the 2015 HPAI outbreak.<sup>38,164</sup>

Equipment	Layer premises, 2015 HPAI outbreak		Turkey premises 2015 HPAI outbreak <sup>a</sup>
	Case farms (n=26)	Control farms (n=33)	Case farms (n=67-80) <sup>b</sup>
	Percent sharing equipment		
Gates/panels	-- <sup>c</sup>	--	9%
Lawn mowers	20%	28%	37%
Pressure sprayer/washer	8%	9%	43%
Skid-steer loader	12%	0%	39%
Tillers	--	--	13%
Manure handling	12%	3%	--

<sup>a</sup>Investigation questionnaire to turkey premises in 2015 case series asked if equipment used on the premises was farm-specific. In this analysis, non-farm-specific equipment was interpreted to be equivalent to shared equipment.

<sup>b</sup>The number of respondents varied for each equipment type listed in the survey.

<sup>c</sup>Dashes indicate data on that specific type of equipment was not reported.

- Historically, disinfecting shared equipment is a described mitigation measure in the event of an infectious disease outbreak in poultry. Little is documented, however, on the specific procedures for C&D during an outbreak.
  - All layer case and control premises (n=59) surveyed in a case-control study in the 2015 HPAI outbreak reported that shared pressure sprayers/washers, skid-steer loaders, and manure-handling equipment were disinfected. Other shared equipment was disinfected at variable rates (lawn mowers, egg flats, racks, or pallets). Shared lawn mowers were disinfected by 65 percent of surveyed farms.<sup>164</sup>
  - For case turkey premises in the Upper Midwest 2015 HPAI outbreak, the majority of farms responded that pre-loaders were cleaned and disinfected by first power washing (to remove organic material, manure, and feathers) and then applying disinfectant.<sup>38</sup>
  - During an outbreak of ILT in northern Georgia, the most frequently reported cleaning procedures for equipment were power washing (64.4 percent), water washing only (33.8 percent), disinfectant use (20.0 percent), and disassembly and cleaning of parts (3.8 percent); 3.5 percent of equipment was not cleaned.
    - In southern Georgia, where there was no ongoing ILT outbreak, the methods used to clean equipment were not statistically different; however, relatively fewer farms utilized a disinfectant in this region (9.4% versus 20% of farms, P=0.056).<sup>275</sup>

### 9.2.3.3 Qualitative Analysis

We considered the following qualitative factors in evaluating this pathway:

- Sharing equipment between poultry operations represents a known risk to biosecurity and an opportunity for disease introduction. Although poultry industry biosecurity guidelines recommend not sharing equipment when possible,<sup>2,19</sup> this may not be economically or logistically feasible for all types of equipment and scenarios.<sup>38</sup>
  - Equipment sharing between poultry premises is documented in the literature, and industry representatives report that sharing of equipment such as that used for tilling, de-caking barns, and litter spreading is difficult to avoid in some production systems (BWG, personal communication, June 2016).
    - Feedback from industry is that frequently the shared large equipment is delivered just in time and is not stored on a poultry premises before use. (BWG, personal communication, June 2016)
  - During PMIP, sharing of equipment is prohibited. The exception is for equipment needed to complete a critical operational visit, such as emergency repair of critical mechanical equipment or service visits to address changes in bird health (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period).
  - However, any equipment brought to the premises during the final grow-out period prior to implementation of a PMIP may continue to be used during PMIP.
    - Recent HPAI outbreak experience suggests this scenario may still present a risk of infection, as there were epidemiological links during the 2015 HPAI outbreak for turkey premises that used shared equipment 10 to 11 days prior to the onset of clinical signs.<sup>38</sup>
    - Arrival of contaminated shared equipment well in advance of movement to processing could provide more opportunities for personnel or equipment to track virus into the barn, especially under certain environmental conditions (e.g., cool temperatures and high humidity) in which viable virus could persist for days to months (see Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates).
- On shared equipment, the potential level of virus contamination depends on the infectious material, the contaminated substrate, and the survival characteristics of the virus.
  - Machinery or equipment that enters a poultry barn, has poultry contact, or comes in contact with poultry feces is at high risk for contamination at an infected but undetected premises. Such equipment includes de-caking or tilling equipment, skid steers/front loaders, hand tools, and poultry monitoring equipment.
  - Estimates of HPAI virus concentrations in infected poultry carcasses, feces, and feathers from infected poultry generally range between  $10^3$  and  $10^7$  EID<sub>50</sub> per gram or milliliter of tested substrate, although higher concentrations have been observed in some cases.
    - For a detailed summary of the literature on virus titers in feces, feathers, blood, and poultry carcasses from infected poultry, see section 9.2.4, Role of HPAI Virus

Spread to a Broiler Flock via Dead Bird Disposal, and on virus titers in immature and mature feathers from infected poultry and ducks, see section 9.1.6, Role of HPAI Virus Spread to Broiler Grow-Out Premises Near Poultry Live-Haul Routes via Feathers, Feces, and Other Fomites.

- Virus survival is generally longer in cooler temperatures and moist conditions. In chicken feces, virus can remain infectious between two and seven weeks at cooler temperatures (39 to 46°F), similar to winter conditions in many regions of the United States.<sup>269,276,277</sup> Persistence in chicken feces is reduced to less than five days in warmer temperatures (71 to 77°F).<sup>269,276</sup>
  - For virus persistence data on feathers, feces, and equipment surfaces, see Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates.
- In a non-outbreak setting, premises participating in the NPIP will follow the site-specific biosecurity guidelines outlined in 9 CFR,<sup>2</sup> which include C&D procedures for equipment when applicable.
  - As described in the literature above, disinfecting shared equipment is the best practice in non-outbreak situations, and regional broiler industry representatives reported this likely occurs for a high proportion of equipment types (BWG, personal communication, Nov 2016). During an HPAI outbreak, attention to disinfecting most equipment is even more likely.
    - While C&D protocols should be described in most company biosecurity plans, strict compliance with plans cannot be assumed, especially when growers or third-party contractors are not directly observed (BWG, personal communication, Oct 2016).
    - C&D procedures for some equipment types may vary depending on if a flock is present on premises or if equipment is used during downtime between flocks (BWG, personal communication, Nov 2016).
  - There may be limitations to C&D of some equipment because of environmental concerns (e.g., excessive waste water from cleaning large equipment) or concerns about damaging the functionality of mechanical or electrical equipment that cannot be heated or wet (i.e., laptops, cell phones, or other monitoring devices). Weather conditions, such as harsh winter weather, may make thorough C&D of equipment more difficult.
- For shared equipment that is not used inside barns during the final grow-out period, the pathway to infect a broiler flock requires multiple steps to introduce HPAI virus into a barn.
  - A potential pathway in this scenario involves 4 contact steps: *shared contaminated equipment* → *ground area on uninfected broiler premises* → *farm personnel's boots or clothing* → *broiler house*. However, if personnel handle the equipment directly and then enter a broiler barn (e.g., to move equipment to a new location on the premises before working with the flock), fewer steps are needed in this pathway: *shared*

*contaminated equipment* → *farm personnel's boots, clothing, or hands* → *broiler house*.

- With each transfer step, there would likely be a reduction in the virus concentration transferred to the recipient surface. This is because only a fraction of the virus (6 to 27 percent) on the donor surface is transferred to the recipient surface in each direct contact.<sup>5</sup> However, depending on the initial viral load and infectious dose in broilers for that strain, the potential level of virus concentration tracked into the barn may still be adequate for infection.
- If the equipment remains outside the PBA, there is a decreased likelihood of cross-contamination to the ground near the poultry house.
  - NPIP stipulations state that all biosecurity plans for poultry premises utilize a PBA.<sup>2</sup>
- During a PMIP, individuals who have direct contact with poultry must wear premises-specific clothing and barn-specific footwear (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period).
  - For more information on risks and mitigation measures related to farm personnel introducing virus into a barn, see section 9.2.2, Role of HPAI Virus Spread to a Broiler Flock via Growers or Employees and their Vehicles Entering the Premises.
- Ssematimba et al. assessed the risk of people and/or equipment that access only the poultry premises or storage rooms on the premises on broiler and layer farms in the Netherlands. They proposed the exposure risk classification for a majority of these contacts to be medium. In the analysis, the risks identified for such contacts are the potential of serving as fomites, expanding the farm network, and lack of or non-adherence to biosecurity protocols, such as non-thorough C&D.<sup>86</sup>
- Equipment that enters the poultry house may be a greater risk due to closer proximity to or potential direct contact with poultry.
  - In the final grow-out period, shared equipment may be placed inside the poultry house. If so, site-specific biosecurity plans should have provisions regarding crossing the line of separation, as required by the NPIP standardized biosecurity principles.
    - Regional industry feedback indicated most broiler companies likely require C&D of shared equipment that will be used inside barns, although some do not require such protocols between houses on the same premises. (BWG, personal communication, Nov 2016)
  - During the PMIP, equipment used in the house may include shared equipment that arrived before the onset of the PMIP and equipment used for a critical operational visit during a PMIP.
    - Any equipment brought on farms for critical operations visits during a PMIP must be cleaned and disinfected, if possible, before use in a poultry barn.<sup>261</sup>
  - Critical visits during the PMIP, such as emergency repair of mechanical equipment or service visits to address changes in bird health, are assumed to occur at a situation-

specific frequency. The available information from studies conducted in non-outbreak scenarios suggests a maintenance visit may be required during a five-day PMIP.

- A survey of poultry growers in Georgia found that the total number of high-risk visits (classified as people who enter the poultry houses), during a seven day period, did not differ significantly between low and high poultry density regions. However, one type of high risk visit (repair personnel entering the poultry house) was significantly more likely among broiler farms in the low-density region (approximately 18 percent) compared with farms in the high-density region (0 percent).<sup>259</sup>
- Among eight premises observed for four days in Ontario, Canada, there was one maintenance visit to service barn ventilation equipment, and this contact had visited three other premises over a three-day period.<sup>278</sup>
- During a PMIP, it is plausible that required repair or service persons performing critical maintenance or evaluating changes in bird health may use equipment that has been used on multiple poultry premises.
  - For further information on the risks associated with the person or vehicle involved in a critical operational visit during PMIP, see section 9.2.1, Role of HPAI Virus Spread to a Broiler Flock via Critical Operational Visits During PMIP.
- In a comparable evaluation of broiler and layer premises in the Netherlands, the exposure-risk classification was proposed to be high for people and equipment that access the poultry house largely because of the frequency of the human contacts. Such contacts (e.g., veterinarians) may visit up to 100 farms per year, and their equipment may not always be thoroughly cleaned and disinfected between farm visits.<sup>86</sup> The same evaluation proposed an overall exposure-risk classification of medium for people and equipment that did not access the poultry house.<sup>86</sup>

#### 9.2.3.4 Likelihood Rating and Conclusion

Previous poultry disease outbreaks demonstrate a known risk for virus spread as a result of movement of contaminated and shared equipment. Equipment that is brought onto a poultry premises may contaminate the ground or personnel who work with poultry, or if contaminated equipment is used inside a barn, live poultry may directly contact it. In the U.S., C&D of shared equipment should be addressed in the biosecurity protocols for all NPIP-approved grower premises. However, adherence to protocols may be limited by feasibility, consistency, and logistics. In the absence of a PMIP, we rate the likelihood of a broiler flock becoming infected with HPAI virus via shared machinery or equipment to be *moderate*.

During the PMIP, in addition to standard biosecurity measures, no off-site equipment will be pre-staged and only critical operational visits may continue, such as emergency repair of critical mechanical systems or service visits to address changes in bird health. Provided the SPS Plans, including the greatly intensified biosecurity of the PMIP, are strictly followed, we rate the likelihood of a broiler flock becoming infected with HPAI virus via shared machinery or equipment as *low*.

## 9.2.4 Role of HPAI Virus Spread to a Broiler Flock via Dead Bird Disposal

The process of dead bird disposal in the Control Area addressed in this risk evaluation relates to normal mortality on broiler premises, as opposed to mortality from known infected premises, including depopulation. Processes described are recommended within the SBS Plan<sup>261</sup> and the cross-commodity PMIP working document (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period).

Dead broilers must be collected and removed in a biosecure manner from houses each day and moved to an on-site location that is as far away from the barns as possible; containers (dumpsters) for dead broilers should never leave the farm. It is not advised for multiple poultry farms to share initial dead bird collection containers or disposal sites located on poultry premises. In normal operations, broiler premises may employ a variety of methods to dispose of daily mortality, both on- and off-site. But the SBS Plan recommendations restrict off-site transportation of carcasses in the days leading up to movement of live birds to processing, i.e., for the duration of the PMIP.

### 9.2.4.1 Dead Bird Disposal during PMIP

Due to the potential spread of HPAI via carcass disposal, the pre-movement isolation measures restrict off-site carcass transportation for disposal during the PMIP. Dead bird disposal is limited to secure on-site storage or disposal during the PMIP, as outlined in the SBS Plan.<sup>261</sup> Secure on-site storage or disposal options include refrigerator/freezer storage, composting, pit burial, incineration, individual burial, and carcass fermentation. As individual burial and carcass fermentation are not widely practiced, this risk evaluation will focus on the more common on-site practices of pit burial, composting, incineration, and refrigerator/freezer storage.

#### Pit Burial

In large poultry operations, pit burial is the most common means of carcass disposal, although it is restricted in certain states, such as Virginia.<sup>279</sup> Environmental (specifically groundwater) contamination and carcass scavenging by wildlife are concerns that can lead to restrictions on pit burial sites. Areas with nonporous soil and a deep water table are the most amenable to pit burial, while sandy soil requires reinforcement of the sides of the pit. To prevent access by animal and insect pests, the top of the pit must be solid and have a tight-fitting lid. Due to groundwater concerns, the placement of burial pits is usually closely regulated by State Departments of Agriculture.<sup>280</sup>

#### Composting

In states that have banned burial pits, composting (controlled decomposition under thermophilic and aerobic conditions) is now the most widely used method of carcass disposal.<sup>1</sup> Under conditions of routine mortality, carcasses are composted together in piles or bins to which a supplemental carbon source, such as litter or sawdust, has been added. Under good composting practices, the carcasses are positioned and layered within the carbon source in a manner optimal for complete and odor-free composting. The resulting product is humus-like, with only feathers and small bone fragments remaining, and the process is generally able to deactivate many pathogens due to the high temperatures (130-150°F) achieved. Composted chicken carcasses may be used as fertilizer, soil amendment, or as a source of organic material for composting additional material.<sup>281</sup>



Poultry composters are typically constructed on a concrete slab to prevent nutrient leaching and vermin entrance (Figure 26). They typically are three-sided and have an overhead roof.<sup>1</sup> If mass depopulation is needed, in-house composting may be used. Multiple peridomestic species have been shown to access poultry carcass compost piles (Figure 27), including raccoon (*Procyon lotor*), opossum (*Didelphis virginiana*), striped skunk (*Mephitis mephitis*), and domestic cats (*Felis catus*).<sup>282</sup>




Figure 26. Mortality composter profile<sup>1</sup>



Figure 27. Wild mammals accessing poultry mortality compost piles. Photos courtesy of USGS

Incineration

Incineration is a commonly used method for poultry carcass disposal and one of the most biosecure methods. Complete carcass combustion occurs within the incinerator unit and the resultant residue does not attract animal or insect pests.<sup>283</sup>

Refrigeration/Freezer storage

Carcasses may be stored in a vermin-proof refrigerator/freezer until off-site carcass movement for rendering or another disposal method is permitted. In large broiler operations, daily mortality usually requires the use of a high-capacity unit.<sup>283</sup> The freezer typically contains leak-proof carcass storage boxes, which are also used to transport the carcasses to the rendering plant<sup>284</sup> or other destination.


Use of freezers in daily operation on broiler farms may vary based on geography and integrator. In some states, freezer carcasses may be picked up from broiler farms for rendering. They are kept in an on-farm freezer if picked up less frequently than daily. In Alabama, for example, this is regulated by the State Department of Agriculture, such that carcass freezers must be placed a minimum of 150 feet from the nearest poultry house, the pick-up truck must not pass between

any poultry houses or within 150 feet of any poultry house to reach the freezer, the driver must disinfect the truck tires before entering the road to the poultry farm and again as the truck leaves the farm, and the grower must minimize contact around the freezer area and with carcasses.<sup>285</sup>

In areas where refrigeration/freezing is not commonly practiced during normal operations, short-term on-site refrigeration/freezing of carcasses may provide a viable choice for dead bird management during PMIP (e.g., using refrigerated trucks or “reefers”) until other on-site or off-site disposal can be coordinated or permitted.

#### 9.2.4.2 Likelihood of a Broiler Flock Becoming Infected via On-farm Dead Bird Disposal and Scavengers during PMIP

Carcass disposal on the farm presents an opportunity for vermin and scavengers to access infected wildlife or poultry carcasses and transmit the HPAI virus to a neighboring uninfected broiler house or mortality disposal site, either mechanically or via virus shedding. The virus could subsequently be transmitted into the broiler house via farm personnel or other mechanisms.



**Figure 28.** Pathway for exposure of a broiler farm via dead bird disposal on-site

##### 9.2.4.2.1 Literature Review

- Several studies have evaluated the impact of composting on HPAI virus:
  - Using a small-scale duplicate of a typical on-farm compost bin (depicted above, **Figure 26**), Senne et al. (1994) composted HPAI H5N2-infected chicken carcasses for 20 days at 22°C (72°F) ambient temperature, with compost turning at day 10.<sup>281</sup>
    - Peak composting temperatures were 57.3 and 58.3°C (135 and 137°F) during the first and second phases of composting, respectively, for the upper layer of carcasses, and 41.5 and 42.8°C (107 and 109°F), respectively, for the lower layer.
    - Despite the lower temperatures at the lower carcass layer, no HPAI virus was detected from any of the carcasses at 10 and 20 days, including from carcasses placed at the periphery of the bin, within 15 to 20 cm (6 to 8 inches) of the walls.
  - Elving et al. (2012) composted HPAI H7N1, a strain with known prolonged survival in manure at 5 to 22°C.<sup>286</sup> In laboratory-scale reactors at 35, 45 and 55°C (95, 113 and

131°F), they found a 12-log viral load reduction within 6.4, 1.7 and 0.5 hours, respectively, in a manure/straw mixture, and within 7.6, 9.8 and 0.5 hours, respectively, in a manure/straw/embryonated egg mixture.

- They recommend:
  - No turning of compost pile during the first phase of composting, to avoid aerosolization of HPAI virus
  - An insulating top layer on the compost to maintain adequate temperature
  - Monitoring of the surface temperature as a parameter for HPAI virus inactivation
- Ahmed et al. could no longer isolate an H5N1 virus strain by day 15 from a closed composter used to dispose of infected birds and their wastes, with temperatures reaching 60°C (140°F).<sup>287</sup>
- Using a static pile passive aeration composting system, Guan et al. (2009) demonstrated inactivation of H6N2 virus in chicken tissue samples and embryonated eggs by day 10 at 61.5°C (143°F) at the top and 50.3°C (123°F) at the bottom of the bin.<sup>288</sup> While still detectable at day 10, viral RNA was degraded in all samples by day 21.
- In the 2004 LPAI H7N2 outbreak on the Delmarva Peninsula in Delaware, in-house windrow composting was the method of carcass disposal.<sup>289</sup> AI virus was undetectable in all samples from the compost and house environment upon compost turning at days 14 to 19 and again upon compost removal at four to five weeks.
  - In this case, as an additional measure, the houses were heated to 37.8°C (100°F) for three consecutive days after windrow formation and again after compost turning.
  - The outbreak was contained to three farms in a dense poultry production area, which the authors attribute largely to on-site composting, as opposed to off-site disposal, for carcass disposition.
- The observation of scavengers near poultry houses has been identified as a risk factor for AI transmission.
  - In the 2002 LPAI H7N2 outbreak in Virginia, multivariate analysis determined that the presence of foxes, raccoons, and opossums posed an approximately two-fold increase in risk of infection.<sup>83</sup>
- Multiple studies have demonstrated the susceptibility of mammals, including scavenger species, to HPAI.
  - Ferrets and foxes fed HPAI H5N1-infected chicken meat developed respiratory and/or digestive infections, demonstrating mammalian potential to shed HPAI virus after consuming HPAI virus-tainted meat.<sup>290,291</sup> Lipatov et al. (2009) measured presence of viral antigen in ferret tissue, not actual viral shedding. Reperant (2008), however, demonstrated pharyngeal shedding in foxes for three to seven days, peaking at 10<sup>3.5</sup> to 10<sup>5.2</sup> TCID<sub>50</sub>/ml following intratracheal inoculation. Pharyngeal shedding

- peaked at  $10^{4.2}$  to  $10^{4.5}$  TCID<sub>50</sub>/ml and lasted for three to five days after feeding infected carcasses. Rectal shedding was detected in one of three foxes inoculated intratracheally at approximately  $10^2$  TCID<sub>50</sub>/ml, only at two days post-inoculation(DPI), and in one of three foxes fed infected meat, at approximately  $10^1$  TCID<sub>50</sub>/ml, on 1 DPI only. All foxes were euthanized at 7 DPI, and virus isolation was negative from all organs sampled from foxes fed infected carcasses.
- Following experimental gastrointestinal HPAI H5N1 infection, cats became systemically infected and viral shedding was detected (via RT-PCR) in pharyngeal and rectal swabs.<sup>292</sup> Pharyngeal shedding occurred in both cats with gastrointestinal exposure, beginning 2 DPI. Rectal shedding was observed in only one of these cats, and only 2 DPI.
  - Songserm et al. (2006) describe a fatal HPAI H5N1 infection in a dog following ingestion of infected duck carcasses.<sup>293</sup>
  - When they were experimentally fed carcasses of LPAI H4N6-inoculated mallards or H4N6-spiked and coated chicken eggs, raccoons failed to subsequently shed AI virus RNA. While this study does not support scavengers as a source of LPAI biological transmission, the authors propose that HPAI virus may be more likely to be shed by scavengers because of its ability to cause more disseminated infection.<sup>294</sup>
    - More recently, experimentally infected striped skunks (as well as cottontail rabbits) have been shown capable of transmitting LPAI H4N6 to birds (mallards) through contact with shared resources (i.e., through contaminating the environment).<sup>295</sup>
  - Both striped skunks and raccoons have been shown to shed LPAI H4N8 and H4N6, respectively, following experimental nasal inoculation with those strains.<sup>51,296</sup> For most of the skunks, nasal shedding of H4N8 peaked at 8 DPI at an average  $10^{5.65}$  PCR EID<sub>50</sub><sup>d</sup> equivalents/ml, and oral shedding at 7 DPI at an average  $10^{4.82}$  PCR EID<sub>50</sub> equivalents/ml. Nasal shedding of H4N6 in the raccoons varied from 1 to 6 days of shedding and between  $10^{0.02}$  and  $10^{1.1}$  EID<sub>50</sub> equivalents/ml. Both species (plus cottontail rabbits) also have been shown to shed novel avian-origin H7N9 (A/Anhui/1/2013) influenza virus at more than  $10^5$  PFU/ml nasal flush.<sup>297</sup>
  - Turkey vultures (*Cathartes aura*) may visit poultry farms to feed on dead birds. Turkey and black vultures (*Coragyps atratus*) both belong to the order Accipitriformes, family Cathartidae. While a review of the literature revealed a paucity of studies of AI in turkey vultures and other Cathartidae, other birds of prey in the order Accipitriformes, such as the common buzzard (*Buteo buteo*), have become infected in previous HPAI H5N1 outbreaks.<sup>233</sup>
    - For a detailed assessment of susceptibility and pathogenicity in avian scavenger species, please see section 9.1.5, Role of HPAI Virus Spread to Broiler Flock via Wild Non-aquatic Birds in Farm Vicinity.

<sup>d</sup> PCR EID<sub>50</sub> equivalent is a measure based on comparing the viral load in the experimental samples with the viral load in samples with known virus titers, as measured by rRT-PCR

#### 9.2.4.2.2 Qualitative Analysis

We considered the following qualitative factors for evaluating this pathway:

- The pathway may involve one or more virus transfer steps between contact surfaces. For example, if a wild animal is acting as a mechanical vector, the pathway *infected undetected carcass*→*scavenger*→*ground area on uninfected premises*→*farm personnel's boots*→*broiler house* involves four contact steps. In general, the chances of the pathway resulting in virus transmission decreases with the number of contact steps that need to occur. Furthermore, even if the transfer steps occur, the virus concentration transferred would likely decrease substantially with each contact step. This is because only a fraction of the virus (6 to 27 percent) on a donor surface is transferred to the recipient surface in each direct contact.<sup>5</sup> The ground traveled by the scavenger between the carcass and the uninfected broiler premises would further lessen the amount of virus present on the scavenger for transmission once at the premises.
  - If a wild animal were contaminated by an infected carcass, we would expect virus may be transferred via feces, bodily fluids, or feathers of that carcass. One gram of organic matter from a poultry carcass may contain  $10^6$  EID<sub>50</sub>/g.<sup>251</sup>
  - For perspective, using a mid-range viral transfer concentration, if 15 percent of virus is transferred at each contact step described above, enough virus particles still remain after four steps to infect five birds (assuming an infectious dose of  $10^2$  EID<sub>50</sub>) if only a single gram of feathers, fluid, or feces is present at the first step of the pathway.
- If, however, the scavenger becomes infected with and subsequently sheds HPAI virus on the grounds outside the uninfected broiler house, there are only two contact steps: from the contaminated grounds to the personnel's boots, and from the boots to the broiler house floor. The likelihood of a scavenger actively shedding HPAI virus following ingestion of an infected carcass is, thus, a critical consideration.
  - The studies cited above demonstrate that mammalian and avian scavengers can become infected with HPAI virus following ingestion of infected chicken, both naturally and experimentally.
    - In the studies in which rectal shedding following consumption of HPAI-infected meat was studied, it was short-lived and occurred inconsistently.<sup>291,292</sup>
  - Additionally, HPAI H5N1 strains that replicate mostly in the lower respiratory tract may not be readily excreted via the upper respiratory system of mammals.<sup>298</sup> The role of other excretory systems, such as the gastrointestinal and urinary tracts, as portals of viral exit is unknown at this time.
- Other plausible pathways where fewer contact steps are involved include those where the grower or other poultry farm personnel directly contact an infected or contaminated scavenger species:
  - An infected or contaminated scavenger species is trapped and killed on an uninfected farm. The grower or employee disposes of the scavenger and then enters a broiler house, introducing virus to the flock.

- A domesticated scavenger (e.g., dog or cat) is infected or contaminated on an infected neighboring farm. The grower or employee touches the pet and then enters a broiler house, introducing virus into the flock.
  - In a study of commercial poultry farms in Virginia, over half of all farms had cats on the premises, and over two-thirds of farms had dogs on premises.<sup>83</sup>
  - The number was slightly lower in a case-control study of turkey flocks affected by the 2016 HPAI outbreak in Indiana; between 30 and 44 percent of flocks had dogs and/or cats on the premises.<sup>260</sup>
  - 25 to 30 percent of HPAI-positive turkey farms in the 2015 outbreak in the northern Midwest had dogs and/or cats.<sup>38</sup>
- The distance between poultry farms (i.e., the distance a scavenger must travel between a carcass and an uninfected farm), also impacts the likelihood of HPAI transmission by the scavenger. The infected carcass and the uninfected farm must be within the likely range of the scavenger for transmission to potentially occur and, based on knowledge of scavenger ranges, this is very likely.
  - The home range of red foxes (*Vulpes vulpes*) is generally up to 8 km (5 miles) in diameter, being largest in the winter.<sup>299</sup>
  - Raccoons (*Procyon lotor*) normally have a home range diameter of 1.8 to 3 km (1.1 to 1.95 miles).<sup>300</sup>
  - The diameter of the opossum's (*Didelphis virginianis*) home range is between 1.3 and 2 km (0.8 to 1.2 miles).<sup>301</sup>
  - The striped skunk (*Mephitis mephitis*) has a home range 2.2 to 2.5 km (1.4 to 1.6 miles) in diameter.<sup>302</sup>
  - Turkey vultures can travel up to 225 km (140 miles) per day.<sup>303</sup>
- The enhanced biosecurity required during the PMIP applies only to farms located in a Control Area that wish to move birds off the premises. While it is assumed that biosecurity practices may be elevated in an outbreak situation, other farms in the Control Area are not subject to any particular stipulations on dead bird disposal or other movements on and off the farm. It is assumed that there may be marked variation in the practices on farms within the Control Area that are not currently adhering to a PMIP.
- In a case-control study of commercial poultry farms in Virginia in 2002, a 1.9-fold increased risk of LPAI H7N2 infection was associated with the sighting of wildlife near poultry houses.<sup>83</sup> Scavengers must, however, gain access to the infected carcass at the source farm in order to contact and transmit HPAI virus.
  - As described above, it may be unlikely for scavengers to access carcasses in burial pits, incinerators, or storage freezers, as all are designed to prevent animal entrance.
    - However, we assume some industry variation in frequency of mortality collection and type of storage container used to gather carcasses from the time they are removed from the poultry house to the point when they are placed in the pit, incinerator, or storage freezer.

- These intermediate transport or storage containers should also prevent access by scavengers on premises observing PMIP.<sup>304</sup>
- While most often constructed on a concrete slab, in part to prevent vermin access, compost bins typically are not completely enclosed. The top layer of litter or sawdust, however, is of a depth (10 to 12 inches) designed to prevent odor production that would attract scavengers.<sup>1</sup>
  - When the carcass is surrounded by a sufficient carbon source and the proper moisture level is maintained, odorous gases enter an aerobic zone and are degraded to CO<sub>2</sub> and water.<sup>305</sup>
  - Reports vary on the prevalence of vermin and scavengers with a properly managed composter.<sup>1,282</sup>
  - In their univariate analysis, McQuiston et al. (2005) found that uninfected farms were significantly more likely to dispose of dead birds via composting than infected farms (77.9 % versus 63.9%, P=0.008).<sup>83</sup>

#### 9.2.4.2.3 Likelihood Rating and Conclusion

The risks for HPAI introduction associated with off-site disposal methods such as rendering are well-documented in the literature (see section 9.2.4.1, Dead Bird Disposal during PMIP, above), and off-site disposal of mortality must be discontinued during PMIP. Best practices for on-site carcass disposal and SBS Plan biosecurity measures should decrease the likelihood of attracting scavenger species to poultry mortality on a broiler farm in the days leading up to marketing. Other poultry farms in the Control Area including an infected farm, however, may not be subject to the intensified biosecurity practices required by PMIP. Mammalian and avian scavengers have the potential to biologically or mechanically carry HPAI virus. Given that many scavenger species have home ranges of adequate size to contain adjacent poultry farms, the likelihood of HPAI introduction to a broiler farm during the PMIP via scavengers within the Control Area is *low to moderate*.

#### 9.2.4.3 Likelihood of a Broiler Flock Becoming Infected with HPAI via Dead Bird Disposal that Takes Place before PMIP

Broiler operations are free to utilize their preferred disposal method for daily mortality in the days leading up to PMIP. For facilities that exclusively use an on-farm disposal method listed above, refer to protocols and procedures listed in section 9.2.4.2., Likelihood of Broiler Flock Becoming Infected with HPAI via Dead Bird Disposal during PMIP, for an evaluation of risk.

Off-site disposal methods include rendering, landfill (garbage), or sourcing animal byproduct as feed for other farmed carnivorous species (such as fox and mink for fur production). When dead birds are moved off a farm, trucks should be covered (to prevent dissemination of potentially contaminated feathers) and should follow a designated approved route. Trucks that carry dead birds must be C&D, using an appropriate protocol, after delivery of the carcasses and before returning to the broiler farm.<sup>266</sup>

#### Rendering


The typical mortality collection process for rendering was described during in communications (D. Halvorson and M. Smeltzer, personal communication, June 2016). During non-outbreak

operations, dead birds are hand collected inside each house throughout the day and deposited in one or more designated areas (sometimes a bin, sometimes the ground) on one or both ends of the house, either inside or outside. Once or twice per day, an employee collects the carcasses or the bin containing the carcasses from each house and loads them into a pickup truck, utility vehicle, or small front-loader (such as a skid-steer or mini-track loader). The carcasses are transported to a common collection bin at some distance from the houses. The distance between the houses and the collection bin varies from farm to farm. The bins typically are dumpster-type containers and may or may not be covered. On large farms, roll-off containers may be used.

Several times per week a company-owned or contracted rendering truck collects the contents of the collection bins: lift arms on the truck engage the dumpster and raise it over the bed of the truck, dumping the carcasses into the truck. While the truck bed is sealed against leakage, it may or may not be covered. Depending on the company ownership, the rendering truck may or may not visit other premises on the same day for carcass pick-up.

Ideally, provisions should be in place to limit contact with dead birds from other farms. These may include prohibiting growers or farm employees from using company or personal vehicles to transport carcasses to a rendering site, utilizing a neutral off-site area for pickup by the rendering company, and ensuring all containers used in transporting mortality from the poultry house to the final destination are secure against wildlife, leaks, and spills.<sup>306</sup>

The transfer of dead birds via onsite equipment to collection bins from which carcasses are then collected by a rendering truck presents an opportunity for truck and ground contamination with HPAI virus and subsequent virus transfer to other premises and other houses on the same premises. The rendering truck may directly enter other premises, and/or transfer virus to other premises along the route if the truck is uncovered. The virus could subsequently be transmitted inside other poultry houses via farm personnel. For further analysis of infection of premises near the route, please see section 9.1.6, Role of HPAI Virus Spread to Broiler Grow-out Premises near Poultry Live-Haul Routes via Feathers, Feces, and other Fomites.



**Figure 29.** Pathway for exposure of a broiler farm via rendering. Other off-site disposal methods are assumed to share a similar pathway.



Other off-site dead bird disposal

- *Landfill* – Less commonly, broiler farm mortality may be transported to a landfill with other garbage products from the operation. A survey of broiler industry veterinarians revealed that two out of eight respondents knew of growers who would “maybe” dispose of broiler carcasses in their trash in the event of an HPAI outbreak (see Appendix 9: Poultry Industry Survey on Garbage Management Practices). Further information about this pathway can be found in section 9.2.5, Role of HPAI Virus Spread to a Broiler Flock due to Garbage Management.
- *Transportation for use as feed for other carnivore-raising operations* – Use of poultry carcasses and byproducts from the poultry industry has been described domestically and internationally in the feeding practices for fur-bearing animals (such as mink and fox),<sup>298,307,308</sup> other exotic species (e.g., alligator farms,<sup>309</sup> captive wildlife and zoos<sup>310</sup>) and anecdotally in some commercial dog breeding operations. Carcasses may be transported directly to local feeding operations (Jill Nezworski, personal communication, July 2016),<sup>308</sup> or may be consolidated at regional collection centers for distribution.<sup>311</sup>

**9.2.4.3.1 Literature Review**

- Rendering has been implicated in the spread of AI virus in previous outbreaks.
  - Following the 2015 HPAI H5N2 outbreak in Minnesota, a case control study of 43 case and 40 control turkey farms found that the rendering of dead birds was a risk factor for H5N2 infection, with an odds ratio of 9.8 (i.e., farms that used rendering were 9.8 times more likely to be infected than farms that did not use rendering).<sup>312</sup>
  - Also in the 2015 HPAI H5N2 outbreak, a case-control study of Iowa layer farms and barns found that 39 percent of case farms versus 13 percent of control farms reported that the renderer came onto the farm. Rendering trucks came near the barns in 29 percent of case farms, compared with 3 percent of control farms.
    - The adjusted odds ratio (OR) for rendering trucks coming near the barns was 22.3 (P < 0.001).<sup>192</sup>
  - In a case series of 81 infected turkey farms in Iowa, Minnesota, North Dakota, South Dakota, and Wisconsin in the 2015 HPAI H5N2 outbreak, 47 percent of case farms used off-site carcass disposal (renderer, landfill, other) in the 14 days prior to disease detection.<sup>38</sup>
  - Data from the 2003 HPAI H7N7 outbreak in the Netherlands were used to estimate H7N7 transmission probabilities to susceptible farms by individual contact types (e.g., feed delivery, egg transport, etc).<sup>81</sup>
    - The analysis determined that, per-contact, rendering visits posed a 25 percent chance of transmission.
  - A case-control study of the 2002 LPAI H7N2 outbreak in Virginia found dead bird disposal by rendering to be the most significant risk factor for AI infection on turkey and chicken farms.
    - The odds ratio was 7.3 (P < 0.001) in a multivariate analysis.<sup>83</sup>

- Rendering pick-ups of dead birds likely played a part in the spread of AI virus in the California LPAI H6N2 outbreak from 2000 to 2002.<sup>141,313</sup>
- Among carnivorous species that may be fed poultry carcasses or byproducts from the poultry industry, HPAI infection has been documented in a variety of species that may be used in the commercial or exotic pet trade, in zoos, or in the fur industry.
  - In reviews of HPAI H5N1 infection in carnivorous species, tigers, leopards, other exotic felids, domestic cats, domestic dogs, civets, and ferrets were identified as potential host species. Within-population transmission was documented in various cat species and in ferrets.<sup>298,314</sup>
    - It was hypothesized in all cases, except that of infection in a colony of civets, that carnivores were infected by consuming or scavenging infected bird carcasses.<sup>298</sup>
  - In captive large felids, onset of clinical signs has been correlated with feeding poultry carcasses in areas with ongoing HPAI outbreaks.<sup>310</sup>
- In captive mink, clinical signs when inoculated with H9N2 (A/Chicken/Hebei/4/2008) were relatively mild, including lethargy and dry nose. No mortality was observed, but pulmonary edema and inflammatory infiltrates were noted on histopathology of lung tissue. Upper respiratory shedding of virus was evident up to 15 days post-inoculation.<sup>307</sup>
- In non-mammalian carnivores, there is limited evidence for AI virus infection. One study noted antibodies against an unknown subtype of AI virus in captive crocodilians of three species in Florida, and PCR testing revealed over 99 percent identity with the NS1 gene of duck AI virus isolates in four crocodilian species tested.<sup>315</sup>

#### 9.2.4.3.2 Qualitative Analysis

The following qualitative factors were considered for evaluating off-site dead bird disposal:

- Collection dumpsters may not be consistently or securely covered, allowing potential access to scavengers.
  - As discussed in other sections of this risk assessment (concerning visitors/people, wild non-aquatic birds, and on-farm disposal during PMIP), inconsistently covering collection dumpsters presents the opportunity for mechanical or biological transfer of HPAI virus via scavengers from infected and undetected carcasses onto the surrounding grounds. This practice could potentially result in cross-contamination of the rendering truck or other mortality transport truck tires and personnel boots, with subsequent contamination of other premises and broiler houses.
- The transfer of infected and undetected poultry carcasses from the collection dumpster into the rendering truck or mortality transport truck bed can result in feathers and bodily fluids contaminating the truck's lift arms, the outside of the truck bed, and the ground surrounding the truck. When the same truck collects a load on another premises, the lift arms could contaminate the dumpster there, and the truck tires could contaminate the ground near the dumpster.
  - Many studies have demonstrated high titers and the persistence of HPAI virus in various poultry tissues and fluids:

- When turkeys were experimentally infected oro-nasally with 100 µl of 10<sup>6</sup> EID<sub>50</sub> of HPAI H7N1, virus persisted for >1 day at >10<sup>4</sup> EID<sub>50</sub>/g of muscle tissue at 4°C (39°F).<sup>59</sup>
- Toffan et al. (2008) experimentally infected turkeys with A/turkey/Italy HPAI H7N1 and measured titers up to 10<sup>6.8</sup> EID<sub>50</sub>/ml in their blood.<sup>62</sup>
- In experimental infections of chickens with EA/AM HPAI H5N2, viral titers were 10<sup>7</sup> to 10<sup>8</sup> EID<sub>50</sub>/g in spleen and lung samples.<sup>124</sup>
- Chicken thigh meat contained up to 10<sup>7.5</sup> EID<sub>50</sub>/g of HPAI H5N1 virus at 24 hours after experimental infection.<sup>63</sup>
- EA/AM HPAI H5N2 viral titers of 10<sup>3</sup> to 10<sup>5</sup> EID<sub>50</sub>/mL of turkey feces were interpolated from cloacal swab data (E. Spackman, personal communication, May 2016).<sup>249</sup>
- Experimental infections with the 1983 Pennsylvania HPAI H5N2 strain resulted in ~10<sup>9</sup> ELD<sub>50</sub>/g of chicken feces.<sup>90</sup>
- Turkey feather tip pools from experimentally infected birds contained 10<sup>4.168</sup> to 10<sup>5.79</sup> EID<sub>50</sub>/ml of HPAI H5N1 virus. (M. Slomka, personal communication, January 2014)
- Indiana HPAI H7N8 viral titers were 10<sup>5.9</sup> EID<sub>50</sub>/ml in feather root samples. (M. Pantin-Jackwood and E. Spackman, personal communication, May 2016)
- A conservative compilation of these results indicates that 1.0 g of tissue or 1.0 ml of feather pulp could contain a minimum 10<sup>4</sup> EID<sub>50</sub> of HPAI virus.
  - Assuming a relatively low infectious dose of 10<sup>2</sup> viral particles, based on findings discussed in Section 8.7.2, only 1.5 ounces (~44 ml) of carcass fluid contains enough viral particles to infect approximately 4,400 birds.
  - In a study of 1-week-old chicks that had been infected intratracheally with 2.5 × 10<sup>4</sup> TCID<sub>50</sub> of HPAI virus (H5N1), homogenates of liver, lung, kidney, and brain from those infected chicks contained 10<sup>6.3</sup> to >10<sup>9.3</sup> TCID<sub>50</sub>/g tissue. On the basis of the relative weight of the lungs, liver, kidneys, and brain of 1-week-old chicks weighing 50 to 55 g, the volume of two chick carcasses represents a minimum of 10<sup>10</sup> TCID<sub>50</sub>.<sup>291</sup>
- If the rendering truck or other mortality transport truck bed is not covered, feathers and other material may escape and result in contamination along the truck's route, with the potential for subsequent transfer into other poultry houses along the route.
  - Transport trucks may be owned by the integrator or by a third-party contractor. Trucks operated by contractors may or may not be covered, and it may be difficult to require use of covered trucks in these situations (BWG, personal communication, April 2016). Additionally, even if a truck were covered, feathers or other material may still escape at driving speeds.
  - In addition to the literature cited above, another study found that feathers harvested from ducks experimentally infected with HPAI H5N1 maintained viral titers of 10<sup>5.5</sup>

- EID<sub>50</sub>/ml for 3 to 6 days at 4°C (39 °F) and 10<sup>4.0</sup>EID<sub>50</sub>/ml for 3 to 6 days at 20°C (68°F).<sup>246</sup>
- The authors point out that while fecal material containing high viral loads may be quickly diluted in the environment, contaminated feathers may persist as solid materials in the field.
  - In the 1983-1984 LPAI and HPAI H5N2 outbreak in Pennsylvania, contaminated transport trucks and coops, and movement of dead (and live) birds, were some of the factors implicated in spread of the virus.<sup>313</sup>
  - Some companies require rendering trucks to be C&D between farms.
    - This is more likely to be effective at eliminating virus contamination in warmer and drier conditions than in colder and wetter conditions.
      - Appendices 4-6 in the Nest Run Shell Eggs risk assessment discuss the effectiveness of disinfectants and C&D under different environmental conditions.<sup>316</sup>
    - If C&D is inadequate to remove viral contamination or not performed between farms, HPAI virus survival is not likely to be affected by the relatively short time period between farms, especially in the protected, moist environment provided by poultry carcasses.
      - Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates details HPAI virus survival on various substrates under different environmental conditions.
  - Similar to rendering, collection of mortality from broiler grower operations for use in the feeding of captive carnivores or fur-industry commercial mink and fox operations represents a risk of premises contamination.
    - Growers may individually contract with a local mink operation or may source mortality to a centralized feeding operation that collects byproducts from many agricultural industries, including egg, dairy, swine, beef, fish, and slaughter facilities.<sup>311</sup>
    - Protocols for truck and driver biosecurity of such operations, in addition to C&D of equipment used to store and transport mortality, are not known and are suspected to vary by premises.
    - Feed for many mink and some farmed fox is prepared on-site and may include ground raw poultry carcasses combined with other agricultural products that the grower may source and store on-site. This practice occurs both in the U.S. and in other major fur-producing countries such as Canada and China.<sup>307,317,318</sup>
  - Ranches and farms in the U.S. that raise fur-bearing animals are geographically limited; they are mostly located in states with cooler winter climates, and the number of registered premises is likely under 400.<sup>319</sup>
  - Both mink and foxes have been shown experimentally to be susceptible to multiple HPAI and LPAI viruses,<sup>291,320</sup> and mink have been shown to transmit avian H3N8, H11N4, H7N7, H5N3, and H9N2 by contact with other mink.<sup>307,320</sup>

- In a Chinese study of farmed mink, there was serological evidence of exposure to H5N1 strains (RE-5 and RE-7) and H9N2 strains (A/Chicken/Hebei/4/2008 and A/chicken/Shanghai/10/01) in both juvenile and breeding adult mink.<sup>307</sup> Of note, researchers could not find a mink farm where poultry carcasses or byproducts were not fed to mink as part of their regular rations.
- This may represent a means of perpetuating virus and a potential source of contamination for broiler premises that contract with mink farms for mortality disposal.
- Biosecurity measures such as using footwear and clothing that are specific to each poultry house (e.g., barn-specific footwear) have been proposed as a means to minimize introduction of virus into the poultry house on the boots/clothes of personnel (PMIP document).
  - During PMIP, growers and employees must wear clothing dedicated to the farm and barn-specific boots before entering poultry houses (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period).
    - If virus from off-site dead bird disposal contaminated the ground before the onset of PMIP, premise-specific clothing and footwear could still become contaminated before entering the poultry house.

#### **9.2.4.3.3 Likelihood Rating and Conclusion**


While off-site disposal methods are prohibited during the PMIP leading up to load-out because of the associated high risk of virus transmission, there remains the potential for contamination of the ground around a poultry house that could lead to virus being tracked into the house on personnel clothing or equipment. The PMIP is designed to increase the probability of detection if virus were introduced six or more days before load-out (since PMIP duration for broilers to market is five days), but virus could be tracked into the barn any time during PMIP, potentially allowing an infected but undetected flock to move to market. Given that off-site dead bird disposal occurs in the Control Area before a PMIP, the likelihood of a broiler flock becoming infected as a result of HPAI virus introduction to the flock (before or during the PMIP) via off-site dead bird disposal that takes place prior to the PMIP is *moderate*, provided that best on-site carcass disposal practices and the cross-commodity and SBS Plan PMIP measures are followed.

### 9.2.5 Role of HPAI Virus Spread to a Broiler Flock due to Garbage Management

Garbage is typically removed from poultry premises by contracted garbage management services (see Appendix 9: Poultry Industry Survey on Garbage Management Practices). Garbage trucks coming near the barns were a significant risk factor in a case-control study of egg layer flocks in the 2015 U.S. HPAI outbreak.<sup>192</sup> This evaluation considers the possible ways a broiler flock could become infected with HPAI-virus by garbage management before movement to processing.

#### 9.2.5.1 Likelihood of HPAI-Virus Infection via Garbage Management

Garbage management represents a potential pathway for HPAI-virus infection of a broiler flock, as multiple poultry premises may share a common disposal site (e.g., landfill), trash collection provider, or trash collection site (i.e., shared dumpster for multiple premises). HPAI-virus may enter a broiler premises via contaminated garbage trucks or drivers. **Figure 30** diagrams the transmission pathway.



**Figure 30.** Pathway of HPAI virus infection of broiler flock via garbage management.

#### 9.2.5.2 Literature Review

- In the 2014-2015 HPAI outbreak, garbage management was identified as a novel risk factor for disease spread.<sup>192</sup>
  - In the 2014-2015 outbreak of HPAI H5N2 in the U.S., a case-control study with multivariable analysis of infected egg layer flocks in Nebraska and Iowa identified garbage trucks coming near the barns as a risk for infection at the farm level

- (OR=14.7; P <.001). This practice occurred at 61 percent of case farms and 23 percent of control farms.<sup>192</sup>
- The univariate analyses (of factors considered for the farm-level multivariable model) showed that 39 percent of control farms had garbage trucks come to the perimeter of the premises; this did not occur at case farms (P=0.003). The frequency of garbage trucks entering the farm but not nearing barns was reported to be comparable among case and control farms (case farms, 21%; control farms, 26%).<sup>192</sup>
  - The frequency with which garbage trucks visited the farms in this study is not known.
  - Prior to 2015, epidemiologic trace-back questionnaires in AI outbreaks did not specifically identify garbage management services as a risk factor. However, previous studies have assessed the risk related to non-company visitors that, similar to garbage collectors, do not typically need to access the poultry house and may visit or contract with multiple poultry premises in an area.
    - In the 2002-2003 outbreak of ILT virus on Mississippi broiler farms, each gas supplier visit to the farm per month increased the likelihood of infection (gas suppliers per month: OR=6.89; P=0.0132; multivariate model, matched controls).<sup>255</sup>
      - The authors suggest gas suppliers may have contributed to viral spread by transporting contaminated material between farms.
    - Based on a stochastic model predicting the spread of HPAI virus between Georgia broiler farms in low- and high-poultry-density regions, gas delivery and utility management visitors contributed minimally (approximately 2 to 4 percent) to off-farm transmission.<sup>265</sup>
      - The models estimated the percent contribution to off-farm transmission. Visitor activities in the high-poultry-density region (1.45 farms/5 miles<sup>2</sup>) and low-poultry-density region (0.49 farms/5 miles<sup>2</sup>) were calculated separately.
  - In many areas, noncommercial poultry operations (i.e., live poultry markets and backyard flocks) may utilize the same garbage management contractors as commercial poultry farms. On noncommercial poultry operations, disposal of mortality in garbage has been identified as a risk factor for AI.
    - In an evaluation of risk factors for live bird markets in New York, New Jersey, Pennsylvania, and New England, markets that disposed of dead birds and offal in the trash were 2.4 times more likely to have a repeated presence of LPAI H5 and H7 viruses (OR, 2.4; 95% CI, 1.8-3.4).<sup>321</sup>
    - In an analysis of risk factors associated with H5N1 in backyard poultry in Egypt from 2010-2012, disposing of mortality and poultry feces in garbage piles outside was significantly correlated in the regression model (F=15.7; P<0.0001).<sup>322</sup>
  - Landfills may serve as a potential site for cross-contamination as garbage management services for poultry premises may transport garbage to the same landfill. This risk likely increases if landfills are used as an off-site disposal method for positive depopulated flocks, which has been reported in previous LPAI outbreaks.<sup>72,268</sup>

- In the 2002 LPAI H7N2 outbreak in Virginia, disposal of depopulated flocks mainly occurred at “mega-landfills” by sealed, leak-proof trucks that were cleaned and disinfected on-farm and at the landfill.<sup>72</sup>
- During the 2001-2002 Pennsylvania H7N2 LPAI outbreak, some euthanized case flocks were disposed of at landfills after being transported in closed containers.<sup>268</sup>

### 9.2.5.3 Qualitative Analysis

We considered the following qualitative factors in evaluating this pathway:

- The types of potentially infectious or contaminated material disposed of in garbage vary by sector of the poultry industry. However many potentially contaminated or infectious materials have been reported to be routinely disposed of in the trash, according to survey responses from representatives of the broiler, turkey, and layer sectors, as listed in **Table 17**.
- The survey results referenced throughout this qualitative analysis were obtained from a small convenience sample of individuals with knowledge of garbage industry practices in various poultry sectors and with a low response rate. Statistical analyses were not conducted for these data. Absence of an affirmative response cannot be assumed to indicate a high-risk activity is not occurring, and this has been taken into account in the assessment. Despite these limitations, the data are informative for the purpose of the risk assessment. For a summary of the survey results, see Appendix 9: Poultry Industry Survey on Garbage Management Practices.



**Table 17.** Survey results concerning material disposed of in garbage on premises in the broiler, turkey, and layer industries.<sup>a</sup>

<b>Item</b>	<b>Broiler sector (n=8 respondents)</b>	<b>Turkey sector (n=15 respondents)</b>	<b>Layer sector (n=39 respondents)</b>
Dead wildlife/wild birds	Yes (1/8)	Yes (5/15)	Yes (1/39)
Rodents	Yes (3/8)	Yes (5/15)	Yes (10/39)
Mortality or poultry carcasses	No (0/8)	Yes (1/15)	Yes (9/39)
Eggs or egg products <sup>b</sup>	Yes (1/8)	Yes (1/15)	Yes (8/39)
Manure	No (0/8)	No (0/15)	Yes (1/39)
Spilled feed	Yes (2/8)	Yes (8/15)	Yes (7/39)
Disposable chick transport boxes <sup>b</sup>	Yes (4/8)	Yes (4/15)	Yes (24/39)
Used needles/syringes/diagnostic supplies that have contacted birds <sup>b</sup>	Yes (1/8)	Yes (5/15)	Yes (14/39)
PPE (boot covers, gloves, coveralls, etc.)	Yes (8/8)	Yes (14/15)	Yes (36/39)
Feathers	No (0/8)	Yes (2/15)	Yes (4/39)
Offal	No (0/8)	No (0/15)	No (0/39)
Equipment or supplies from inside barns <sup>c</sup>	Yes	Yes	Yes (22/39)
Household garbage from farm manager or any other residence <sup>3</sup>	--	Yes	Yes (20/39)
Trash associated with waterfowl hunting <sup>c</sup>	--	--	No (0/39)
Garbage from processing operation <sup>c</sup>	--	--	Yes (23/39)
Lunch room and restroom garbage <sup>c</sup>	--	--	Yes (37/39)

<sup>a</sup>Yes indicates materials disposed of in the garbage by one or more survey respondents within each industry. In parenthesis, numerator indicates number of survey respondents reporting disposal of item and denominator indicates total number of respondents.

<sup>b</sup>Language of selection choice modified in survey distributed to representatives of layer industry.

<sup>c</sup>Item explicitly asked only in survey distributed to representatives of layer industry. Yes in the broiler and turkey industries for these items indicates at least one respondent manually reported disposing of that item in the garbage.

- Additional items reported to be disposed of in the garbage on broiler premises were boxes, buckets, jugs from disinfectants, litter treatments, and disposable chick feeder lids. On turkey premises, additional items included empty medication containers, poult box papers, and supply containers.
- Of potential HPAI-contaminated or infectious material reported to be disposed of in the garbage on poultry premises (i.e. dead wildlife, poultry carcasses, egg shells, and potentially contaminated materials that have contacted poultry), the hypothetical expected virus concentration on each type of item varies.

- HPAI virus has been recovered in many tissues of poultry carcasses, such as muscle, liver, kidney, brain, spleen, and blood. For detailed information on virus concentration in various tissues, feces, and feathers, see section 9.2.4, Role of HPAI Virus Spread to a Broiler Flock HPAI via Dead Bird Disposal.
  - If garbage is contaminated with infectious poultry carcasses, the risk of infection via garbage management is likely similar to the risk via off-site dead bird disposal. Off-site dead bird disposal has been implicated in previous AI outbreaks. For a detailed literature review, see section 9.2.4.2.1, Literature Review.
- There are reports of disposing of dead wildlife in trash on commercial poultry premises.
  - Evidence of AI virus infection of multiple mammalian species, such as ferrets, foxes, cats, dogs, skunks, raccoons, and mink, has been demonstrated by virus isolation, antigen detection, and PCR. For a detailed description on mammalian susceptibility, see section 9.2.4, Role of HPAI Virus Spread to a Broiler Flock via Dead Bird Disposal.
  - Wild and domesticated bird species can be infected with HPAI virus. For a detailed description of experimental studies in wild and domesticated aquatic birds, see section 9.1.4, Role of HPAI Spread to Broiler Flock in a Control Area via Wild Aquatic Birds in the Farm Vicinity. For a detailed review of HPAI detections, prevalence, and susceptibility of passerine birds and non-passerine non-aquatic birds, see section 9.1.5, Role of HPAI Virus Spread to Broiler Flock via Wild Non-Aquatic Birds in Farm Vicinity.
- Eggs from infected hens have tested positive for HPAI virus, including shells, albumen, and yolk. Measured concentrations have varied. See the Secure Egg Supply Egg Shell Risk Assessment for more details.<sup>115</sup>
- Influenza virus survival varies depending on strain and environmental conditions, such as humidity and temperature. Virus persistence is generally longer at cooler temperatures and in more humid conditions. For virus persistence data on materials that may be disposed of in the garbage, such as poultry carcasses, feathers, egg shells, egg trays, wood, steel, glass, and PPE, see Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates.
- In a survey sent to representatives of the broiler, turkey, and layer sectors, it was reported that garbage is likely to be transported to a landfill by a contracted service provider for the majority of commercial poultry operations.
  - Transport trucks may become contaminated at municipal landfills; it has been noted that upon arrival at landfills, garbage management vehicles may drive over previously deposited garbage (D. Halvorson, personal communication, June 2016).
    - The CFR provides standards for design and operation at landfills.<sup>323</sup> For municipal solid waste landfills, these include 6 inches of covering on disposed solid waste each day or as necessary, disease vector control, and access requirements.<sup>323</sup>

- Garbage management contractors used by some turkey and broiler premises have been reported to visit multiple poultry premises on one route before depositing a load at the landfill; thus, HPAI-virus-contaminated garbage from an undetected premises may be present on the truck when it arrives at a broiler farm.
  - The types of potentially contaminated trash from other types of poultry operations (e.g., backyard poultry, processing facilities, live bird markets, etc.) are not known, but are assumed to include materials similar to those reported in garbage from commercial poultry operations.
    - In the Netherlands, poor management practices pertaining to liquid waste (e.g., waste water) and solid waste have been identified as potentially increasing the risk of AI transmission in the neighborhood of infected farms (A. Ssematimba, personal communication, August 2016).<sup>86</sup>
    - A shared dumpster or common trash collection point for multiple poultry premises, while not commonly used in the poultry industry, represents an additional site of potential cross-contamination between commercial poultry operations.
- Garbage trucks and drivers typically do not contact live poultry while completing contracted duties on a poultry premises. Biosecurity recommendations and site-specific biosecurity plans may not stipulate specific measures for garbage management drivers, but it is recommended that visitors follow procedures to cross the PBA and LOS.<sup>2</sup>
  - In a qualitative evaluation of potential AI transmission pathways on broiler and layer premises in the Netherlands, Ssematimba et al. proposed an exposure risk classification of "medium" for the majority of contacts assessed that access only the premises and have no contact with live poultry.<sup>86</sup> The analysis considered contact frequency, biosecurity practices, and risk category.
- Virus introduction into poultry houses via garbage management may involve one or more virus transfer steps. Although there would likely be reduction in the virus concentration (6 to 27 percent) between a donor surface and recipient surface in each direct contact,<sup>5</sup> the virus concentration potentially tracked into the barn may still be above the infectious dose. This depends on the initial viral load and infectious dose of that virus strain in broilers.
  - It is assumed that the ground traveled by the vehicle between the time of contact with infected garbage and the uninfected broiler premises may lessen the amount of virus present for transmission once at the premises. However, mechanical transmission of a similar type virus (PRRSV) has been demonstrated experimentally in a swine industry-like setting.<sup>250</sup>
  - Alternatively, if an infected load of garbage is in the truck at the time of arrival on an uninfected broiler premise, fewer transfer steps are required.
  - The enhanced biosecurity required during a PMIP applies only to farms located in a Control Area that wish to move birds off the premises. It is assumed that there may be marked variation in the biosecurity and garbage practices on farms within the Control

Area that are not currently adhering to a PMIP, despite a likely elevation of biosecurity during an outbreak.

- If garbage management activities and visits occur outside of the PBA, there is a decreased likelihood of cross-contamination between contaminated garbage management and personnel, equipment, or other potential fomites that may access the poultry house.
  - Based on survey responses from representatives of the broiler industry, it is common practice for the dumpster or trash collection point to be located at the entrance or perimeter of the farm. This equates to a distance of 100 to 150 feet from the nearest poultry barn for the majority of respondents, but this distance varies.
- As is true with other third-party contractors, poultry growers or integrators may find it difficult to control or influence certain practices by contract garbage haulers, including C&D of garbage trucks, pickup routing, and landfill practices.
- On broiler premises, the frequency of garbage pickup is most often weekly or every other week, based on survey responses from representatives of the broiler industry.

#### **9.2.5.4 Likelihood Rating and Conclusion**

##### **9.2.5.4.1 Likelihood of a Broiler Flock Becoming Infected with HPAI-Virus due to Garbage Management**

Garbage management was identified as a novel risk factor for HPAI virus introduction in the 2014-2015 outbreak in the U.S. Epidemiological studies of past outbreaks have not specifically investigated garbage as a potential route for HPAI virus entry onto a poultry premises, but a recent survey identified a number of items disposed of in trash across poultry industry sectors that could be potentially infectious or contaminated by HPAI virus. Given that there is potential for HPAI virus associated with garbage management to be tracked into the poultry house, the likelihood of a broiler flock becoming infected with HPAI virus due to garbage management without a PMIP is *moderate to high*.


##### **9.2.5.4.2 Likelihood of a Broiler Flock Becoming Infected with HPAI Virus due to Garbage Management when a PMIP is Implemented**

During the PMIP, garbage will not be removed from the premises, given the risk it presents, and the producer is responsible for managing the risks associated with any on-site garbage movement that must occur. The greatly intensified biosecurity measures of the PMIP, such as using footwear specific to each poultry house (e.g., barn-specific footwear), should decrease the likelihood that virus is tracked into barns during the final days before load-out (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period). Provided the on-farm biosecurity measures are strictly followed during a PMIP, the likelihood of a broiler flock becoming infected with HPAI-virus due to garbage management during PMIP is *low*.

### 9.3 Pathways for a Broiler Flock Becoming Infected with HPAIV via Load-Out Operations

Movements of load-out equipment and crews have been implicated in AI transmission in previous outbreaks. According to Poss et al., load-out crews, which may load out more than one flock within 12 hours, have been associated with the spread of AI.<sup>324</sup> Several large LPAI outbreaks in turkeys in Minnesota, such as the 1978 and 1995-1996 LPAI outbreaks, were attributed in part to potentially contaminated load-out crews and equipment or processing trucks coming into close contact with birds that remained on the farms after partial flock removal.<sup>267,325</sup> During the 1986 LPAI H5N2 outbreak in Pennsylvania, restricting farm access to only sanitized load-out trucks and crates interrupted infection transmission.<sup>326</sup> Currently, a majority of commercial broiler grower premises practice an “all-in, all-out” management system, in which a single-age flock is placed on a premises and then the farm is completely depopulated at load-out.<sup>266</sup>

In this chapter we are assessing the likelihood that a broiler flock becomes infected during the load-out process, resulting in movement of infected but undetected birds to market. Pathways considered include contaminated load-out equipment, vehicles, and/or crews from an infected and undetected farm, followed by introduction of HPAI virus into a flock at the next load-out premises.<sup>e</sup>



**Figure 31.** Pathway for exposure of a broiler house via load-out operations

<sup>e</sup> Premises contamination with HPAI virus by load-out crews or processes may also represent a pathway by which day-old chicks could become infected upon placement of the next flock in the same poultry house if virus were not inactivated during downtime. These pathways leading to infection of the next flock are outside the scope of this risk assessment.

### 9.3.1 PMIP Measures for Moving Broilers to Slaughter

For premises within a Control Area that wish to move broilers to slaughter, a Pre-Movement Isolation Period (PMIP) is defined that limits non-critical visits and personnel on the farm, while biosecurity and flock disease surveillance is increased (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period).<sup>261</sup> Adherence to enhanced biosecurity principles during this isolation period prior to scheduled movement minimizes the likelihood that the flock will become exposed to HPAI via contact with people, vehicles, or equipment that may be contaminated with HPAI in the final days leading up to load-out. Similarly, decreasing the likelihood of late introduction of virus to a flock will increase the sensitivity of surveillance and sampling performed during the PMIP. For further information on the likelihood of detecting infection close to movement, see section 9.4, Likelihood of Detecting HPAI in an Infected Broiler House.

#### 9.3.1.1 *Additional Load-out Mitigation Measures for Permitted Movement of Poultry to Market from a Control Area*

Load out begins as the first crew, vehicle, or equipment arrives on the premises and ends when the last load of birds departs the premises. Pre-staging of equipment during PMIP is prohibited (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period).

The biosecurity and sampling stipulations pertinent to load-out of broilers become more stringent as the duration of load-out increases. If birds are infected because of contaminated load out equipment entering the premises, they have the potential to shed virus up until the time of slaughter. Viral contamination may be tracked into occupied poultry houses which are still awaiting load-out. This extends the period for HPAI virus to replicate and spread through the flock, and includes any time remaining in the poultry house until load-out, in addition to transit time, and any hold time at the plant before slaughter. Load-outs of longer duration pose an increased risk of transporting a considerable number of infected but undetected birds to market.

To meet the permit guidance criteria for movement from a Control Area, all broiler premises (regardless of load-out time) should adhere to mitigation measures after completion of load-out, such as load-out crew stipulations and live-haul routing requirements, and sanitation procedures for live-haul equipment when moving it within the Control Area from the Infected Zone to the Buffer Zone.<sup>261</sup>

A greater emphasis is placed on decreasing the likelihood of HPAI-contaminated load-out equipment being used for permitted movement within the Control Area, and on diligent biosecurity between barns to minimize spread between poultry houses in the event of a virus introduction during load-out. These suggested measures become more stringent as the amount of time needed to completely load-out a premises increases (i.e. premises that take 25 hours or more to completely depopulate). Specifically, load-out equipment must be adequately cleaned and disinfected or meet other specifications so that that it remains free or nearly free of contamination. These specifications can be met through use only on Monitored Premises during an outbreak. For the relatively low proportion of broiler premises where load-out takes more than 48 hours, PCR testing is required of barns after the initial 48 hours of load-out in addition to the PCR testing in the final day(s) of the PMIP (See Appendix 11: Load-out Mitigation Methods).

Further detail on load-out mitigations recommended for broiler premises where the time to complete the load-out, transport to the processing plant, and yard time at the processing plant for

the premises is brief (i.e., 24 hours or less) or prolonged (i.e., 25-48 hours or more than 48 hours) are in Appendix 11: Load-Out Mitigation Measures, and on the SBS Plan website. Results of modeling simulations to support the increased biosecurity and sampling requirements for premises as this duration increases can be found in Appendix 10: Supplementary Testing Protocols.

### 9.3.2 Literature Review

In past LPAI outbreaks in the U.S. poultry industry, load-out equipment and crews have been implicated as a means of virus spread between farms, specifically involving partial flock removal and movement of load-out crews between premises.

- In the 1995-1996 LPAI H9N2 outbreak in Minnesota, likely transmission pathways between commercial turkey premises included exposure from contaminated processing trucks during partial flock load-outs and contaminated load-out personnel and equipment from an infected flock.<sup>325</sup>
- In a 1978 outbreak of LPAI (H6N1, H4N8, H6N2) in Minnesota turkey flocks, the management practice of marketing turkey hens while leaving growing toms on the farm allowed potentially contaminated load-out equipment and crews to contact birds (toms) that would remain on the farm.<sup>267</sup>
- In an overview of the Minnesota Cooperative Control Program, Poss et al. identified orderly marketing as a procedure to prevent AI virus spread, as there is potential for heavy contamination to personnel and equipment involved in the transport of an infected flock. Previously, load-out crews, which may load out more than one flock within 12 hours, have been associated with the spread of AI.<sup>324</sup>
- The use of clean load-out vehicles and equipment has been protective in a past AI outbreak. During the 1986 LPAI H5N2 outbreak in Pennsylvania, which likely spread in part through movement of contaminated crates, transmission was interrupted when premises access was restricted to sanitary crates and clean trucks.<sup>326</sup>

During previous poultry disease outbreaks (LPAI, HPAI, and ILT), movement of contaminated transport vehicles, transport equipment, and infected poultry likely contributed to virus spread between farms.

- A 2001-2002 outbreak of LPAI H6N2 in Northern California is suspected to have spread from flock to flock in part through moving live birds to slaughter.<sup>141,313</sup>
- In the 1983-1984 H5N2 LPAI and HPAI Pennsylvania outbreak, contaminated transport trucks and coops and movement of live birds were among the factors that contributed to spread of infection.<sup>267</sup>
- Findings from previous disease outbreaks suggest that virus transmission to poultry premises near live-haul routes is possible. For a review of literature on infection of premises near the route in past outbreaks, see section 9.1.6, Role of HPAI Virus Spread to Broiler Grow-Out Premises near Poultry Live-Haul Routes Via Feathers, Feces, and Other Fomites.

The load-out process inherently places crews, vehicles, and equipment in close contact with live poultry, poultry feces, and poultry feathers.

- Estimates of HPAI virus concentrations in chicken secretions, feces, feathers, and other tissues generally range between  $10^3$  and  $10^7$  EID<sub>50</sub> per gram or per milliliter, although higher concentrations have been observed in some cases.<sup>57,63,64</sup>
- For further information on viral load on substrates related to live-bird movement, see section 9.1.6, Role of HPAI Virus Spread to Broiler Grow-Out Premises near Poultry Live-Haul Routes via Feathers, Feces, and Other Fomites, and section 9.2.4, Role of HPAI Virus Spread to a Broiler Flock via Dead Bird Disposal.

Viral persistence depends on substrate, temperature, and humidity, among other factors. Virus may persist for days to weeks or longer in a climate like that of the continental U.S..

- Kurmi et al., Beard et al., and Wood et al. reported that HPAI virus strains were inactivated in chicken feces in less than five days in warm temperatures (71 to 77°F) and nearly two to eight weeks in cooler temperatures (39.2 to 46.04°F).<sup>90,269,277</sup> In these experimental studies, when temperature was constant, time to virus inactivation in feces usually increased as moisture level increased.<sup>90,269</sup> On substrates that may be found on vehicles or poultry transport crates, an LPAI virus strain (A/Herring gull/Delaware 471/86 [H13N7]) was below detectable limit at day 6 on tires, steel, and plastic, and at hour 72 on wood.<sup>270</sup> On glass and soil in cool temperatures (39.2-46.0 °F), an HPAI H5N1 strain (A/Vietnam/1203/2004 [H5N1 clade 1]) was recovered at day 13 in low relative humidity and day 9 in high relative humidity.<sup>269</sup>
- For further data on viral persistence on different substrates and in varying environments, see Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates.

### 9.3.3 Qualitative Analysis

We considered the following qualitative factors for evaluating this pathway:

- The time required to load out and completely depopulate a broiler premises depends on size of operation, crew and equipment logistics, and capacity of the slaughter facility.
  - Industry representatives report that most individual barns can complete the cumulative process of load-out, transit, and yard time in under 12 hours. This timeline is optimized to minimize transit mortality and protect carcass value and likely cannot be shortened further (BWG, personal communication, Sept 2016).
  - Broiler farm size and slaughter plant capacity vary with geography and logistics. Industry representatives from the BWG estimate that for an entire premises, the proportions of U.S. commercial broiler farms that would be able to complete cumulative load-out, transit, and yard time within certain time limits are shown in **Table 18**.



**Table 18.** Estimated percentage of U.S. commercial broiler farms capable of completing the cumulative load-out process, transport to slaughter, and yard time at processing plant for entire premises in certain time intervals during an outbreak.

Cumulative load-out, transit, and yard time for an entire premises	Percent of premises
Less than or equal to <u>12 hours</u>	40 to 50
Less than or equal to <u>24 hours</u>	80 to 90
Less than or equal to <u>48 hours</u>	90

- As discussed in section 9.4, Likelihood of Detecting HPAI in an Infected Broiler House, the likelihood of a broiler house becoming infected with HPAI in the days leading up to movement is lower when PMIP enhanced biosecurity measures are implemented and the premises is located far enough from infected premises. Increased biosecurity and greater distance help reduce the chances of moving birds that are infectious because of exposure to HPAI during PMIP.
- The enhanced biosecurity measures in place during PMIP are not required to continue into the load-out period (see Appendix 7: Cross-Commodity Pre-Movement Isolation Period); thus it is possible that contaminated equipment or crews brought to the premises for load-out could lead to cross-contamination of houses that have yet to be depopulated via people or equipment entering poultry houses.
  - In the event of a single point-source infection, **Table 19** shows the estimated number of birds in a given barn which may be infected, depending on duration of time between infection and slaughter (i.e., load-out, transit time, and yard time at plant).

**Table 19.** Estimated number of infectious birds in a single poultry house if exposed at the time of load-out from a single-point source infection.<sup>a</sup>

Initial number of infected birds	Mean number of infectious birds post-exposure to HPAI					
	12 hr	24 hr	48 hr	72 hr	96 hr	120 hr
1 bird	1	3	32	346	2,769	9,487

<sup>a</sup>Transmission model estimations are from 6,000 iterations using PA HPAI H5N2 strain characteristics, a mean flock size of 24,482 birds, and a mean contact rate of 5.1 per day.

- If, however, equipment were heavily contaminated or virus strain characteristics caused initial infection in multiple birds, the mean number of infectious birds in a barn at a given time-point is much higher (**Table 20**).

**Table 20.** Estimated number of infectious birds in a single poultry house if exposed at the time of load-out from an initial infection in multiple birds.<sup>a</sup>

Initial number of infected birds	Mean number of infectious birds post-exposure to HPAI					
	12 hr	24 hr	48 hr	72 hr	96 hr	120 hr
10 birds	10	32	320	2,900	11,456	18,106
100 birds	102	318	2,839	12,257	19,024	18,351 <sup>b</sup>

<sup>a</sup>Transmission model estimations are from 6,000 iterations using PA HPAI H5N2 strain characteristics, a mean flock size of 24,095 birds, and a mean contact rate of 5.1 per day.

<sup>b</sup>In the model, a large proportion of birds died before 120 hr, leading to smaller mean numbers of infectious birds at this time.

- If birds are infected by load out equipment, they have the potential to shed virus up until the time of slaughter. This includes load-out, transit time, and any hold time at the plant before slaughter. Load-outs of longer duration thus pose an at increased risk of transporting a considerable number of infected but undetected birds to market.
- In the absence of a disease emergency, live-haul equipment is not routinely cleaned and disinfected between terminal movements in the broiler industry. Feces, feathers and other contaminants may remain on surfaces that will contact a subsequent flock.
  - There are some commercial operations in which load-out crates and equipment are cleaned and disinfected at the processing plant before moving to another premise. However, operationalizing C&D of all vehicles and equipment used in load-out during an outbreak is generally considered not feasible on a national scale (BWG personal communication, Aug-Sept, 2016).
  - The SBS downtime measures state that when moving live-haul equipment from the Infected Zone to the Buffer Zone, equipment must be C&D (power washed) and have a down period of at least 24 hours.<sup>261</sup> The plan does not stipulate any type of downtime or C&D for crews or their personal vehicles.
- As pre-staging of load-out crews or equipment is not allowed during PMIP, potential viral introduction via contaminated crews or load-out equipment would occur after the conclusion of flock sampling for HPAI.
  - The latent period for an individual bird varies with virus strain and infectious dose, but has been estimated to be less than one day.<sup>123,327</sup> Thus, considering both latent period and contact rate in the event of exposure to HPAI virus, the number of infectious birds shedding virus in a flock at the end of the 48-hour load-out period would be low (**Table 19** above).
    - Greater variation in infectious period and mean time to death has been reported. All chickens inoculated with one of six 2005 HPAI H5N1 Vietnam strains died in 48 hours or less,<sup>122</sup> while inoculation with 2015 EA/AM HPAI H5N2 (Tk/MN/2015) resulted in a mean time to death of 4.8 and 3.2 days in 5- and 8-week-old broiler chickens, respectively.<sup>124</sup>

- For a more detailed review of experimental studies of latency period, infectious period, and mean time to death from AI infections in chickens, see Section 8, Hazard Identification: HPAI Overview.
- Very large flocks may require more than 48 hours to load out. In the transmission model shown in **Table 19**, extending load-out to 72 hours results in more than a 10-fold increase in the number of infectious birds compared with load-outs of 48 hours or less.
  - In a scenario in which contaminated load-out crews or equipment infect more than one bird initially, the potential number of infectious birds at the point of slaughter could be much higher. For example, if there are 10 birds initially infected, the model estimates 2,900 infectious birds 72 hours after exposure to HPAI (**Table 20**).
- Transport time from farm to slaughter plant represents additional time for potential viral shedding. The average transportation time for commercial broiler systems in the U.S. is generally less than two hours.
- More stringent load-out biosecurity for premises with longer load-out times should decrease the likelihood of heavily contaminated people or equipment bringing HPAI virus to a premises during the load-out process.
  - Premises with a cumulative load-out, transport, and yard time of 25 to 48 hours are required to adequately clean and disinfect load-out equipment or use equipment that has been used only on Monitored Premises during an outbreak.
  - For premises with a cumulative load-out, transport, and yard time of greater than 48 hours, this risk is further mitigated by requiring load-out equipment used previously only at Monitored Premises with PCR testing within 24 hours of movement and premises depopulation completed in 24 hours or less, as an alternative to cleaning and disinfection. At these premises, the load-out crew should access the premises only after showering and wearing clean clothes to decrease the amount of virus contaminating their clothing or bodies.
  - For additional details on mitigation measures, see Appendix 11: Load-out Mitigation Measures.
- Most companies expect that load-out crews enter only the barn where they are currently working. However, growers and employees may still need to enter multiple barns on the premises in the course of caring for birds and preparing barns for downtime (shutting off water, heat, etc.)(BWG, personal communication, 2016).
  - At premises with prolonged load-out durations, barn-to-barn biosecurity measures should be implemented to limit likelihood of contaminating barns still occupied by poultry during the load-out, such as no pre-staging of equipment in barn before loadout, utilizing barn-specific footwear and farm-specific clothing, and handwashing (see Appendix 11: Load-out Mitigation Measures).
- Flocks which are infected via load-out equipment may not be detected by clinical signs or mortality trigger alone.

- PCR testing of barns during load-out on a premises should increase the probability of detecting infections that occurred because of the load-out process, especially as the duration of load-out increases.
- For further information on load-out testing and surveillance protocols and sensitivity analysis of such protocols, see Appendix 10: Supplementary Testing Protocols.
- Premises outside the Control Area are less likely to be infected, as the role of local area spread diminishes with distance. However, if HPAI-contaminated load-out crews, vehicles, or equipment were used outside the Control Area, there are fewer safeguards in place to decrease likelihood of heavy contamination of a poultry flock during load-out that could lead to movement of an infected but undetected flock.
  - While some companies may choose to implement it where feasible, the current SBS Plan does not stipulate any C&D for load-out equipment, vehicles, or crews exiting the Control Area.
  - Premises outside the Control Area may not be subject to surveillance and pre-movement testing requirements beyond routine NPIP surveillance for LPAI.
  - Premises outside the Control Area have no limitations on load-out duration or pre-staging of load-out equipment.
  - Load out vehicles and crews outside the Control Area may have less stringent biosecurity requirements.
  - However, industry representatives have said that they track the use of load-out equipment and crews in routine operations, and hypothetically could dedicate “lower risk” crews in the event that dedicated equipment/crews for the CA were not logistically feasible (BWG, personal communication, Nov 2016).
- Load-out crews have the potential to carry virus off the premises on clothing, boots, and vehicles, and may pose additional risks in transit to/from job sites, by living in homes with backyard poultry or shared with individuals who work on other poultry operations.
  - SBS Plan measures require a change of clothes and shower for load-out crews before going on to work with other poultry.<sup>261</sup>
  - Interaction between load-out crews and other poultry industry employees or other poultry is addressed in section 9.2, Likelihood of Broiler Flock Becoming Infected with HPAI via Movements of People, Vehicles, or Equipment.

### 9.3.4 Risk Rating and Conclusion

Previous outbreaks have implicated contaminated load-out crews and equipment in the spread of AI. In the U.S. commercial broiler industry, C&D of vehicles and equipment associated with terminal movements is not routine practice. If a flock were infected via contaminated load-out crews or equipment, shortening the time from load-out to slaughter for the complete premises limits how long the virus may spread within the flock. However, the time required to load out and completely depopulate a broiler premises depends on size of operation, crew and equipment logistics, and capacity of the slaughter facility.

Given that PMIP enhanced biosecurity and testing measures are utilized, and that additional load-out mitigation measures are in place that are commensurate with the duration of the premise-wide load-out process, we estimate the likelihood of a broiler flock becoming infected with HPAIV via load-out operations and resulting in an infected but undetected movement to market to be *low to moderate*.

## 9.4 Likelihood of Detecting HPAI in an Infected Broiler House

### 9.4.1 HPAI Surveillance Measures

#### 9.4.1.1 Current Measures

Current routine influenza surveillance measures involve testing of meat-type chicken flocks for H5/H7 subtypes of AI for birds processed at slaughter plants participating in the U.S. H5/H7 Avian Influenza Monitored program of the NPIP (see 9 CFR part 146.33 for further information).

#### 9.4.1.2 Outbreak Measures

##### Active Surveillance by rRT-PCR Testing

The active surveillance protocol option outlined in the SBS Plan involves testing two pooled samples via rRT-PCR at National Animal Health Laboratory Network (NAHLN) labs. One pooled sample with swabs from 11 dead birds must be tested by rRT-PCR for every 50 dead birds from each house on the premises for two consecutive days prior to the start of load-out for broilers. Samples for the second (later) rRT-PCR test must be collected within 24 hours before the start of load-out.<sup>261</sup>

Current USDA:APHIS HPAI emergency response plans assume same-day turnaround for submitted rRT-PCR samples. For example, the results of samples collected and submitted to NAHLN labs for rRT-PCR testing in the morning are assumed to be available to the Incident Command at the end of the same business day. However, this may not always be feasible, in which case earlier sample collection times for rRT-PCR tests may be needed on a case-by-case basis. An active surveillance protocol involving one pooled sample of 11 swabs on two consecutive days in which the second (later) test is collected within 48 hours prior to the start of load-out is evaluated in Appendix 10: Supplementary Testing Protocols for those Premises that Anticipate the Turnaround Time for rRT-PCR Results to Take Longer than 12 Hours. Collecting rRT-PCR samples earlier reduces the likelihood of detecting HPAI prior to the load-out start; thus, it is important to note that this alternate testing protocol is outside the scope of preferred testing protocols as outlined in the SPS Plan.

##### Detection through Trigger for High Mortality

If daily mortality is abnormally high (more than 3/1,000 birds in a house, excluding culls) immediately prior to a scheduled movement, broilers shall not move until diagnostic steps have been initiated and HPAI has been ruled out as the cause of elevated mortality.<sup>261</sup>

## 9.4.2 Quantitative Methods for the Estimated Likelihood of Detection prior to the Start of Load-out on a Premises

The likelihood of detecting HPAI in a broiler house prior to the start of load-out is estimated via simulation. The approach consists of a stochastic disease transmission model, which simulates the spread of HPAI within a house, and an active surveillance model, which uses the output from the disease transmission model to simulate the probability of detection under a given active surveillance protocol. A technical description of the simulation model algorithms can be found in Weaver et al.<sup>328</sup> These simulation models from Weaver et al. have been reparametrized for broilers for use in the current analysis.<sup>328</sup> A summary of the input parameters is given in **Table 21**, and details on their estimation are given in Appendix 8: Modeling Technical Details. A brief overview of the disease transmission and active surveillance models is given below.

### 9.4.2.1 Overview of Disease Transmission and Active Surveillance Models

The likelihood of detecting HPAI depends on the following factors:

- The HPAI spread dynamics in a house, which impact the rate of mortality and morbidity, rise over time. The HPAI spread dynamics depend on parameters such as the length of latently infected and infectious periods in individual birds and the “contact rate” between infectious and susceptible chickens.
- The variability in the steps of the detection process, given an active surveillance protocol option. Factors such as the normal mortality (mortality not related to HPAI) and HPAI mortality rates impact the chances of including a virus-positive swab in the test sample. The chances of detecting a virus-positive sample depend on the diagnostic sensitivity of the test.

The HPAI spread dynamics in a house are simulated by the disease transmission model. Disease states included in the model are susceptible (S), latently infected (L), infectious (I), and removed (R). The number of broilers in each disease state is updated at 0.1-day intervals. Transitions from the latent to the infectious state and the infectious to removed state are determined by latent and infectious period distributions estimated for various HPAI strains based on data from experimental studies. Once a bird is in the removed state, it is considered to be deceased and remains in that state for the remainder of the simulation. The transition from the susceptible to the latently infected state is determined by the adequate contact rate and number of infectious birds in the current time period. The adequate contact rate ( $\beta$ ) is defined as the mean number of birds each bird comes in contact with per unit time such that the contact is adequate to transmit infection. Higher adequate contact rates result in a higher likelihood of infection. Similarly, as the number of infectious birds increases, the likelihood of infection increases.

The variability in the detection process is simulated by the active surveillance model. Detection of HPAI in the surveillance model occurs through either diagnostic testing or heightened mortality. Samples for diagnostic tests are randomly selected from the normal and disease mortality available on the test day. The normal mortality is simulated based on industry-provided daily and weekly mortality, while the disease mortality is drawn from the transmission model output. Provided at least one infected bird is present in the test sample, detection occurs according to a Bernoulli trial with probability equal to the test sensitivity. Detection via heightened mortality occurs if the total mortality exceeds the trigger level on one of the days prior to the start of load-out.

#### 9.4.2.2 Model Scenarios

The likelihood of detecting HPAI in a broiler house prior to movement is evaluated under four scenarios:

- Pennsylvania HPAI H5N2 with a mean time to death of 4.1 days and baseline contact rate
- EA/AM HPAI H5N2 with a mean time to death of 3.2 days and baseline contact rate
- EA/AM HPAI H5N2 with a mean time to death of 4.8 days and baseline contact rate
- EA/AM HPAI H5N2 with a mean time to death of 4.8 days and slow contact rate

The length of latent and infectious period distributions can impact the time to detection: for example, HPAI strains with a long mean time to death—the combined length of the latent and infectious periods—will generally take longer to detect via active surveillance due to the slower rise in mortality. As the latent and infectious periods are strain-specific and can vary considerably, evaluating results based on multiple strains is critical for developing robust risk management strategies. In the current analysis, the likelihood of detection is estimated for latent and infectious period distributions based on Pennsylvania HPAI H5N2 and EA/AM HPAI H5N2 strains. Two further scenarios were evaluated for EA/AM HPAI H5N2 based on two different mean times to death of 3.2 days observed in 8-week-old broilers and adult breeders, and 4.8 days observed in 5-week-old broilers (D. Swayne, personal communication, Oct 2016).

We evaluated two scenarios for the contact rate, given high uncertainty in the parameter estimates identified in a review of the available literature. The baseline contact rate distribution was determined in light of this literature review with an emphasis values estimated from outbreak data. However, since estimates from experimental data were generally lower, a slow-contact-rate scenario is considered to account for this highly plausible situation. The contact rate affects how quickly the infection moves within the house, with slower contact rates resulting in slower spread. As slow spread generally increases the time for high levels of mortality to accumulate, leading to longer detection times, the slow-contact-rate scenario represents a more conservative approach. The EA/AM HPAI H5N2 strain with a 4.8-day mean time to death is paired with the slow contact rate due to its relevance in light of the 2015 outbreak.

**Table 21.** Parameter estimates for the HPAI transmission model for broiler houses.

Parameter name	Parameter description	Distribution/Value
Contact rate (transmission parameter)	The number of direct or indirect contacts a bird has that are sufficient to transmit infection per unit time	Baseline: Beta-PERT distribution (min=2.50, mode=4.77, max=9.00, scale=4.00) of contacts per day Slow: Uniform distribution of contacts per day (minimum 1, maximum 3)
Latent period distribution	Length of the latent period	Pennsylvania HPAI H5N2: Gamma distributed (shape=0.8248, scale=0.4446; mean 0.37 days) EA/AM HPAI HPAI H5N2 (MTD <sup>a</sup> 3.2 & 4.8 days): Gamma distributed (shape=0.8926, scale=0.7145; mean 0.64 days)
Infectious period distribution	Length of the infectious period	Pennsylvania HPAI H5N2: Weibull distributed (shape=1.9651, scale=4.2376; mean 3.76 days) EA/AM HPAI H5N2 (MTD of 3.1 days): Weibull distributed (shape=1.9651, scale=2.90; mean 2.57 days) EA/AM HPAI H5N2 (MTD of 4.8 days): Weibull distributed (shape=1.9651, scale=4.70; mean 4.17 days)
Number of chickens/house	Distribution of the number of hens per house	Log normal distribution truncated at 13,000 and 50,000. Mean 24,257 birds, std.dev. 9,787 birds per house. Estimated from industry data.

<sup>a</sup>Mean Time to Death, the combined lengths of the latent and infectious periods.

**9.4.2.3 Estimated Likelihood of Detection under a Pre Movement Isolation Period (PMIP)**

As discussed previously, a PMIP involves the implementation of heightened biosecurity to minimize the chances of a house becoming exposed to HPAI close to the start of load-out. **Table 22** gives the detection probabilities for a house one to ten days following exposure to HPAI under the active surveillance protocol of one rRT-PCR test of 11 swabs taken the day of and day before load-out begins on the premises.

Under the PA HPAI H5N2 baseline contact rate scenario, for example, if a house were exposed to HPAI three days prior to the start of load-out, the estimated probability of detection is 73%. Given that the house was exposed for three days prior to load-out, testing would occur on the second and third days of HPAI's presence in the house. In this example, the probability of detection improves as the number of days post-exposure increases. This is due to the exponential



growth in mortality that occurs as HPAI moves through the house, which increases the likelihood of including at least one bird dead from HPAI in the pooled sample taken for diagnostic testing or observing total mortality above the threshold amount. Thus, by reducing the chances of exposure to HPAI close to the start of load-out, the PMIP decreases the risk of releasing infected but undetected birds by allowing sufficient time for the infection to spread within the house.

**Table 22** can be used to inform the length of the PMIP under an assumption that the PMIP is 100% effective in preventing exposure to the pathogen. In these scenarios, it is conservatively assumed that the house is infected immediately prior to implementation of the heightened biosecurity of PMIP. For example, under a four-day PMIP, a house is assumed to have been infected four days before the start of load-out, just prior to the start of the PMIP. The detection probability in this case, considering the PA HPAI H5N2 baseline contact rate scenario, is estimated to be 94%. Similarly, the scenario under a five-day PMIP is estimated to result in a 99% likelihood of detection. The length of the PMIP decided on by the SBS Workgroup is five days, which generally achieves high probabilities of detection. However, under the EA/AM HPAI H5N2 slow contact rate scenario, detection probabilities as low as 74% are estimated to be possible, which suggests a five-day PMIP is not entirely robust for all potential strains and contact rates.

**Table 22.** Simulation model results showing the predicted probability of HPAI detection for a broiler house infected a given number of days prior to the start of load-out.<sup>a</sup>

Scenario	Number of days prior to movement when exposure to HPAI occurs								
	1	2	3	4	5	6	7	8	9
PA HPAI H5N2	0.05	0.28	0.73	0.94	0.99	1.00	1.00	1.00	1.00
EA/AM HPAI H5N2: MTD 3.2 days	0.05	0.30	0.68	0.89	0.97	0.99	1.00	1.00	1.00
EA/AM HPAI H5N2: MTD 4.8 days; baseline contact rate	0.04	0.16	0.50	0.81	0.94	0.99	1.00	1.00	1.00
EA/AM HPAI H5N2: MTD 4.8 days; slow contact rate	0.04	0.11	0.30	0.54	0.74	0.87	0.94	0.97	0.99

<sup>a</sup> The detection probabilities are estimated from 6,000 simulation iterations for four different scenarios varying by the strain or contact rate. The active surveillance protocol consists of 1 sample of 11 swabs taken for rRT-PCR testing the day of and day before the start of load-out.

**Table 23** compares the probability of detection under three different active surveillance and PMIP strategies. Under the scenarios with no PMIP, exposure is assumed to occur sometime between one and ten days prior to the start of load-out. Under the scenario with a five-day, 100% effective PMIP, meaning the PMIP guarantees the house is not infected during its implementation, exposure is assumed to occur sometime between five and ten days prior to the start of load-out. Exposures occurring earlier than ten days prior to load-out are not considered since the infection is almost certain to be detected via diagnostic testing and monitoring of mortality, so the risk of moving infected but undetected broilers would be minimal in such cases.

The results in **Table 23** indicate that performing active surveillance using only a mortality trigger without implementing a PMIP is insufficient for detecting HPAI in a broiler house. Including diagnostic testing consisting of one pooled sample of 11 swabs taken for rRT-PCR testing the day of and day before load-out begins substantially improves the likelihood of detecting HPAI in the house prior to movement. However, when a PMIP is not implemented, exposures occurring within ten days of load-out still fail to generate high levels of detection. The results in **Table 23** suggest that HPAI could go undetected in roughly one third to one fifth of these cases. When exposure close to the time of movement is prevented through a five-day PMIP, on the other hand, HPAI is generally detected in the house with a high degree of confidence, although under the EA/AM HPAI H5N2 (MTD=4.8 days) slow-contact rate-scenario there is an estimated 10% chance of not detecting the infection.

Also included in **Table 23** is the mean number of infectious birds at the start of load-out in the houses that go undetected, along with the 5<sup>th</sup> and 95<sup>th</sup> percentile. The mortality-trigger-only active surveillance protocol with no PMIP is estimated to result in large numbers of infectious birds present at the start of load-out in undetected houses. As it is likely that many infectious birds will go undetected, given the low detection probabilities, surveillance via a mortality trigger alone with no PMIP represents an extremely risky practice. The mean number of infectious birds at the start of load-out in houses that went undetected is higher under the scenario of diagnostic testing with a five-day PMIP, because the infection is present in the house for at least five days, which leads to more birds becoming infected. Diagnostic testing with no PMIP, on the other hand, allows for infections to occur within five days of the start of load-out, which provides less time for large numbers of infectious birds to accumulate. The amount of mortality due to HPAI will also be lower when infections occur within five days of the start of load-out.

As HPAI is less likely to be detected when mortality is low, exposures close to the time of load-out have a higher probability of going undetected; therefore they represent a greater proportion of the cases with infectious but undetected birds and lead to the lower mean dead birds. While the mean number of infectious birds in undetected houses is higher under the scenario using both diagnostic testing and PMIP, the likelihood of detecting the infection is relatively high. Thus, this scenario poses the lowest risk for HPAI spread.

**Table 23.** Likelihood of detecting HPAI in a broiler house prior to the start of load-out under three surveillance scenarios, followed by the mean number of infectious broilers in undetected houses at the time of movement.

Strain/contact rate scenario	Active surveillance and PMIP scenario <sup>a</sup>		
	Mortality trigger only, no PMIP	rRT-PCR testing and mortality trigger, no PMIP <sup>b</sup>	rRT-PCR testing and mortality trigger, 100% effective 5-day PMIP <sup>c</sup>
PA HPAI H5N2	0.62 496 (1, 2,741)	0.80 57 (1, 207)	0.99 176 (0, 1,157)
EA/AM HPAI H5N2: MTD 3.2	0.58 193 (0, 991)	0.79 22 (0, 73)	0.99 79 (0, 383)
EA/AM HPAI H5N2: MTD 4.8; Baseline contact rate	0.54 412 (0, 2,295)	0.74 43 (0, 170)	0.98 266 (0, 1,260)
EA/AM HPAI H5N2: MTD 4.8; Slow contact rate	0.26 168 (0, 883)	0.63 15 (0, 56)	0.90 51 (0, 199)

<sup>a</sup>Parentheses indicate the 5<sup>th</sup> and 95<sup>th</sup> percentiles estimated from 6,000 iterations for different active surveillance and PMIP scenarios.

<sup>b</sup>Houses are assumed to be infected sometime within 1 to 10 days of the start of load-out with no PMIP.

<sup>c</sup>Flocks are assumed to be infected sometime within 5 to 10 days of the start of load-out with PMIP.

#### 9.4.2.4 Overall Likelihood of not Detecting HPAI in a Broiler House prior to the Start of Load-out on the Premises

The overall probability of not detecting HPAI in an infected broiler house by the start of load-out considers two events: the probability a susceptible house becomes infected provided it is some given distance from an infectious premises, and the probability that the infection is not detected in the house prior to the start of load-out. The probability that a susceptible premises located a given distance from an infectious premises also becomes infected is estimated via a spatial transmission kernel, which is discussed in section 9.4.2.4.1 below. The probability that infectious birds are not detected by the start of load-out, given that the house has been infected, is estimated using the transmission and active surveillance simulation models discussed in the previous sections. The two probabilities are combined into an overall likelihood using a method described in Weaver et al. that considers the ten days prior to the start of load-out.<sup>328</sup>

##### 9.4.2.4.1 Estimation of the Probability of Infection via a Spatial Transmission Kernel

A spatial transmission kernel uses outbreak data to estimate the hazard rate, or infection risk, posed by an infectious premises a given distance away from a susceptible premises. The spatial transmission kernel theoretically averages the risk overall transmission pathways at the given inter-premises distance, therefore providing a summary view of outbreak spread. The current analysis considers two different spatial transmission kernels: a transmission kernel estimated from the 2003 HPAI H7N7 outbreak in the Netherlands by Boender et al. (2007) and a transmission kernel estimated from the 2015 HPAI H5N2 outbreak in Minnesota.<sup>133</sup> The

Minnesota transmission kernel was estimated using the maximum likelihood method from Boender et al. (2007) with an additional parameter added to the force of infection, which is the cumulative hazard rate faced by a susceptible premises on a given day.<sup>133</sup> The force of infection on susceptible premises  $i$  on day  $t$ ,  $\lambda_i(t)$ , is given in Boender et al. (2007) as

$$\lambda_i(t) = \sum_{j \neq i} h(d_{ij}) 1\{j \text{ is infectious}\}$$

where  $h(d_{ij})$  represents the spatial transmission kernel as a function of the distance between susceptible premises  $i$  and infectious premises  $j$ .<sup>133</sup>

The force of infection as defined above assumes all spread to be lateral, dependent only on the number of infectious premises on day  $t$ . Due to phylogenetic evidence of primary introductions occurring concurrently with lateral spread in the Minnesota outbreak, an additional parameter,  $k$ , was added to the force of infection equation used to estimate the spatial transmission kernel for Minnesota, giving the following expression<sup>38</sup>:

$$\lambda_i(t) = \left[ \sum_{j \neq i} h(d_{ij}) 1\{j \text{ is infectious}\} \right] + k$$


The additional parameter represents a constant, distance-independent hazard primarily expressing the infection risk posed by distance-independent environmental factors—note that  $k$  does not depend on the number of infectious premises—such as wild birds. For more details on the estimation of the spatial transmission kernel for the Minnesota HPAI H5N2 outbreak, see Appendix 8: Modeling Technical Details.

The force of infection is used to estimate the probability that susceptible farm  $i$  is infected on day  $t$ , called  $q_i(t)$ . The expression for  $q_i(t)$  is defined below:

$$q_i(t) = 1 - e^{-\lambda_i(t)}$$

As the force of infection increases, the probability of infection increases. **Figure 32** is a comparison of the Netherlands HPAI H7N7 and Minnesota HPAI H5N2 transmission kernels under the mean maximum likelihood estimates. Both transmission kernels indicate that infection risk was primarily distance-dependent during their respective outbreaks. The Netherlands HPAI H7N7 transmission kernel suggests that local transmission pathways to infectious premises, such as wild animals, aerosols, or equipment sharing, were the primary forces behind outbreak spread, while the Minnesota HPAI H5N2 transmission kernel suggests that transmission pathways involving moderate distances, such as garbage or rendering truck visits, played a significant role in outbreak spread in addition to the local pathways.

As the mean hazard rate for the Minnesota outbreak is higher and persists over longer distances relative to the Netherlands outbreak, the probability of infection will also be higher and remain elevated at larger distances under the Minnesota transmission kernel. As the overall probability of not detecting HPAI in a house prior to the start of load-out is derived using the transmission-kernel-based probability of infection, it is expected to exhibit similar behavior.



**Figure 32.** Spatial transmission kernels estimated from the 2003 HPAI H7N7 outbreak in the Netherlands by Boender et al. (2007)<sup>133</sup> and the 2015 HPAI H5N2 outbreak in Minnesota.

#### 9.4.2.4.2 Estimated Overall Likelihood of not Detecting HPAI in a Broiler House Prior to the Start of Load-out

Estimates for the overall likelihood of not detecting HPAI in a broiler house prior to the start of load-out are given in **Table 24**. The overall likelihood is the combined probability of a house first being exposed to HPAI and then HPAI going undetected in the house prior to load-out following exposure. The probability a susceptible premises is infected with HPAI by an infectious premises located a specific distance away is estimated using the Minnesota HPAI H5N2 and Netherlands HPAI H7N7 spatial transmission kernels. The overall likelihood under the Netherlands transmission kernel is given by **Table 24** in parentheses. The probability the infection goes undetected in the house is estimated using the active surveillance simulation model under a diagnostic testing protocol of one pooled sample of 11 swabs taken for rRT-PCR the day of and day before the start of load-out. The overall likelihood is estimated for the 10 days prior to the start of load-out under the EA/AM HPAI H5N2 (MTD=4.8 days) baseline-contact rate-scenario in order to obtain relatively conservative estimates.

The overall likelihood is estimated under three scenarios varying by the effectiveness of the PMIP at preventing exposure during the five days prior to the start of load-out. Premises did not institute a PMIP during the Minnesota or Netherlands HPAI outbreaks. Since the heightened biosecurity during the PMIP should result in lower likelihoods of exposure, the spatial transmission kernels estimated from these outbreaks likely overestimate the infection risk during this time. The baseline scenario in **Table 24** assumes the daily probability of exposure does not change during the PMIP, which would be expected if no additional biosecurity measures were implemented. The second scenario assumes the PMIP is 80% effective at preventing exposure, which means the daily probability of infection during the PMIP is reduced to one fifth of the probability prior to the PMIP. The last scenario considers a 100% effective PMIP, which means the daily probability of exposure during PMIP is zero.

The estimates given in **Table 24** provide evidence that limiting exposure close to the time of movement through a PMIP reduces the overall likelihood of infection; even a partially effective PMIP leads to a considerable reduction. The overall likelihood decreases as distance from the infectious premises increases, due to the distance dependence exhibited by the spatial transmission kernels. In addition, the higher mean hazard rate estimated from the Minnesota HPAI H5N2 outbreak results in higher estimates for the overall likelihood. Biosecurity and distance from an infectious premises both play a critical role in preventing exposure to HPAI and thereby limiting the risk of not detecting the infection in a house prior to the start of load-out.

This risk can be further reduced by implementing a sound active surveillance protocol. **Table 24** indicates that the heightened biosecurity during the PMIP combined with an active surveillance protocol of one pooled sample of 11 swabs taken for rRT-PCR testing the day of and day before the start of load-out is a viable strategy for reducing the overall likelihood, yielding low likelihoods of moving infected and undetected birds even at relatively short distances from infectious premises and under the higher hazard rates of the Minnesota transmission kernel.

**Table 24.** Predicted percent likelihood of a broiler house being: (1) exposed to HPAI from an infected premises at a specific distance and (2) undetected prior to the start of load-out following exposure under three PMIP scenarios varying by biosecurity effectiveness.<sup>a</sup>

Distance from an infected premises (km)	Scenario for the daily likelihood of exposure during 5-day PMIP varying by status and effectiveness <sup>b</sup>		
	Baseline: no PMIP	80% effective PMIP	100% effective PMIP
	Predicted likelihood		
1.5	1.53 (0.32)%	0.35 (0.07)%	0.048 (0.010)%
2	1.50 (0.24)%	0.34 (0.05)%	0.047 (0.007)%
3	1.41 (0.14)%	0.32 (0.03)%	0.044 (0.004)%
5	1.13 (0.06)%	0.26 (0.01)%	0.035 (0.002)%

<sup>a</sup> The first likelihood estimate in each cell is based on the 2015 Minnesota HPAI H5N2 outbreak transmission kernel, and the estimate in parenthesis is based on the 2007 Netherlands HPAI H7N7 outbreak transmission kernel estimated by Boender et al (2007).<sup>132</sup>

<sup>b</sup> In all scenarios, an active surveillance protocol of one pooled sample of 11 swabs taken for rRT-PCR testing on the day of and day before the start of load-out on the premises was implemented under the EA/AM HPAI H5N2 (MTD=4.8 days) baseline contact rate strain.

### 9.4.3 Likelihood of Moving Infectious but Undetected Broilers Following Exposure during Load-out

Contaminated load-out crews and equipment entering a poultry premises pose an infection risk that is especially relevant during extended (i.e., multiple-day) load-outs. As discussed in Section 9.3.3, the number of infectious birds can increase rapidly in houses infected early in the load-out process, which could pose significant consequences if these birds were to be transported from the premises. Additional diagnostic testing during the load-out period can decrease the likelihood of

moving large numbers of infectious birds following exposure to HPAI during the load-out process. The estimated likelihood of detection for a single house two to six days following exposure to HPAI under an active surveillance protocol of one rRT-PCR sample of 11 swabs taken daily is given in **Table 25**.

The protocol is evaluated under three scenarios varying by the number of birds assumed to be initially infected, which represents increasing levels of contamination on the load-out crews and equipment. This model uses the Pennsylvania HPAI H5N2 strain. The testing protocol decided upon by the SBS Workgroup involves daily rRT-PCR testing of 11 swabs for all houses scheduled to be loaded out beginning 48 hours after the arrival of load-out equipment onto the premises. On large operations, it is possible that multiple poultry houses will be awaiting load out at the 48-hour mark after premises-wide load-out has begun. Since more than one house could be tested, the premises-wide likelihood of detection would be at least as high as and generally higher than the estimates given in **Table 25**, which are for surveillance in a single house only.

As expected, the likelihood of detection increases as the number of days since exposure increases. Similarly, the likelihood of detection increases as the number of initially infected birds increases, since more infectious birds results in faster growth of the infection within the house. The results suggest that the infection in a house exposed early in a four-, five-, or six-day load-out would likely be detected independent of the level of contamination. When the initial number of infected birds is 10, the probability of infection exceeds the 95% threshold three days post-exposure. When the initial number of infected birds is 100, the 95% threshold is estimated to be exceeded as soon as two days post-exposure. The low detection probabilities for houses exposed close to the time of movement can be improved through the use of supplementary AC testing. The likelihood of detection with supplementary antigen capture (AC) testing is evaluated in Appendix 10: Supplementary Testing Protocols.

**Table 25.** The likelihood of detecting HPAI in a house prior to the transportation of broilers to processing for different numbers of days post-exposure and different numbers of initially infected birds, meant to represent the contamination level of the load-out crew and equipment. <sup>a</sup>

Initial no. Of birds infected	Days post-exposure				
	2	3	4	5	6
	<b>Predicted detection probability<sup>a</sup></b>				
1	0.27	0.73	0.95	0.99	1.00
10	0.84	0.99	1.00	1.00	1.00
100	0.99	1.00	1.00	1.00	1.00

<sup>a</sup>The active surveillance protocol consists of a daily rRT-PCR sample of 11 swabs. The likelihood of detection was estimated from 6,000 simulation iterations using the Pennsylvania HPAI H5N2 strain.

As infections occurring two to three days prior to the testing day have a low likelihood of detection, the exposure mitigation and biosecurity measures implemented during a multiple-day load-out are especially important in reducing the risk of transporting infectious but undetected birds to processing. Despite these low probabilities of detection, the likelihood of sending large

numbers of infectious but undetected broilers to processing is expected to be quite low. Given in **Table 26** is the predicted percent probability of not detecting HPAI in a house where the number of infectious but undetected broilers exceeds 300 birds, given exposure occurred during load-out, some number of days prior to movement. The percent probabilities are estimated from the Pennsylvania HPAI H5N2 strain under the active surveillance protocol of daily samples of 11 swabs taken for rRT-PCR testing. Similar results were determined for numbers of infectious but undetected broilers exceeding 100, 500, and 1,000 birds at the time of movement and are given in Appendix 10: Supplementary Testing Protocols.

The results in **Table 26** suggest the risk of sending infectious but undetected broilers to processing in numbers of 300 or more is generally quite low. However, a 9.3% probability of moving at least 300 infectious but undetected broilers where 10 birds were initially infected 2 days prior to movement is not trivial. The nontrivial likelihood of such an event underscores the importance of using relevant biosecurity to prevent heavy contamination from accumulating on load-out equipment and prevent contamination of other barns yet to be loaded out.

Current recommended practices for multiple-day load-outs include either using load-out equipment used only on Monitored Premises or cleaning and disinfecting the load-out equipment prior to use on the premises. Both of these measures prevent the equipment from being highly contaminated, making the scenario where only one bird is initially infected more likely than having many infected birds initially infected. In addition, heightened barn-to-barn biosecurity, such as barn-specific footwear, is recommended, which limits the likelihood of HPAI entering a populated barn before load-out begins in that barn. This may keep HPAI virus from infecting a house for multiple days. Considering these recommended exposure mitigation measures for multiple-day load-outs, the likelihood of sending at least 300 infectious but undetected broilers to processing is expected to be low.

**Table 26.** The estimated percent probability of not detecting HPAI in a house following exposure during load-out where the number of infectious but undetected broilers at the time of movement exceeds 300 birds.

Initial number of birds infected <sup>a</sup>	Days post-exposure				
	2	3	4	5	6
	<b>Predicted percent probability of at least 300 infectious but undetected broilers at the time of movement<sup>b</sup></b>				
1	0.22%	4.72%	2.00%	0.35%	0.00%
10	9.34%	0.98%	0.00%	0.00%	0.00%
100	0.55%	0.00%	0.00%	0.00%	0.00%

<sup>a</sup>The initial number of birds infected represents the level of contamination present on the load-out crew and equipment.

<sup>b</sup>Percent probabilities are estimated from 6,000 simulations based on the Pennsylvania HPAI H5N2 strain and an active surveillance protocol of one sample of 11 swabs taken daily for rRT-PCR testing.

### 9.4.4 Conclusions

An effective PMIP increases the probability of detection by preventing exposure close to the time of load-out, which allows a longer time for HPAI to spread within the house. This leads to



higher levels of disease mortality and increases the likelihood that the total mortality exceeds the trigger level or that a swab from an HPAI-infected dead bird is included in the diagnostic test sample. A five-day PMIP generally yields high probabilities of detection, though it may not be entirely robust for all HPAI strains and within-house spread scenarios. Exposure of a broiler house to HPAI during a multiple-day load-out may be difficult to detect, since the infection occurs close to the time of movement. However, given the load-out biosecurity and active surveillance measures in place, if an infected but undetected movement were to take place, a movement containing large numbers of infectious birds would be unlikely.


Assuming that an effective PMIP is implemented, and that both mechanisms for active surveillance outlined in the SBS Plan (trigger for elevated mortality and rRT-PCR mortality testing) are utilized as described, and that load-out biosecurity measures are implemented, the likelihood of HPAI in an infected broiler house going undetected is rated as follows:

- The overall likelihood of HPAI-infected but undetected broilers in a house at the conclusion of PMIP and prior to the start of load-out on the premises is estimated to be *low* at a distance of 1.5 km or more from an infected premises.
- The likelihood of HPAI-infected but undetected broilers in a house at the conclusion of load-out, resulting in movement of large numbers of infectious birds ( $\geq 300$ ) to slaughter, is estimated to be *low*.

## 10 Overall Conclusion

The objective of this assessment was to estimate the risk that the movement of market-age broiler chickens to processing (i.e., broilers to market), from a premises located within a Control Area during an HPAI outbreak in the poultry industry in the U.S., would result in the introduction of HPAI infection onto another poultry premises (e.g., another poultry farm).

The assessment considered relevant current industry practices and current biosecurity measures as well as outbreak-specific measures from the SBS Plan, in particular the PMIP. The assessment focused on the risk pathways for HPAI infection of market-age broilers on grow-out premises located within an HPAI Control Area via components of local area spread, people and vehicles, and load-out processes. Many of these pathways do not involve the movement of live birds, and rather relate to the likelihood of infection of live birds that will move and potential missed detection prior to movement. Qualitatively compiling the assessed risks and likelihoods of the pathways analyzed yields the overall risk of HPAI spread to susceptible poultry due to the movement of broilers to market (**Figure 33**).



**Figure 33:** Diagrammatic representation of the overall assessed risk. The overall risk assessment is based on consideration of the steps needed to move live birds to market and the pathways that could lead to infection of a flock, the subsequent likelihood of detection of the infected flock, and potential movement of an infected but undetected flock.

The evaluation of the major risk pathways identified resulted in the following conclusions:

### Local Area Spread Pathways

- Insects.** The likelihood of a broiler premises becoming infected with HPAI virus via insect transmission varies with distance and with source premises infection status. The estimated likelihood ratings range from *negligible to moderate*, with a higher likelihood of infection closer to a known infected premises. For premises located closer than 1.5 km

to an infected flock, there are too many variables to accurately assess the risk of becoming infected with HPAI via insect transmission.

- **Aerosols.** The likelihood of a broiler premises becoming infected with HPAI virus via bioaerosol transmission varies with distance and with viral load at the source premises. Literature review and most previous outbreak reports indicated that aerosol transmission was not an important factor at distances more than 1.5 km from an infected flock. However, there is some evidence of aerosol transmission over shorter distances. Thus the likelihood of a broiler premises becoming infected via bioaerosol transmission is rated as follows:
  - *Negligible to low* if >1.5 km from an infected but undetected premises
  - *Moderate to high* if <0.5 km from a known infected premises
- **Wild Birds.** The likelihood of HPAI virus spread to a broiler grow-out premises via wild birds depends upon the type of wild birds and exposure to the wild birds. With an effective PMIP, the likelihood of HPAI infection via wild aquatic birds and via non-passerine non-aquatic birds is *low*, as these birds and their waste are unlikely to access or be tracked into a broiler grow-out barn. Given that passerine birds may access the inside of broiler grow-out barns (even during a PMIP) and have been shown to be capable of shedding the virus, the likelihood of HPAI infection via passerine birds in the farm vicinity was assessed as *low to moderate*.

#### People, Vehicles, and Equipment Pathways

- **Live-haul Routes.** The risk of HPAI virus spread to broiler grow-out premises near poultry live-haul routes via feathers, feces, and other fomites depends on both distance and source flock. For trucks hauling birds that had an effective PMIP and negative rRT-PCR test results, the risk is estimated to be *negligible to low* no matter the distance. In contrast, for trucks hauling birds that had no PMIP and no diagnostic tests (e.g., from premises outside the Control Area), the risk ranges from *low to high*, with premises within 100 meters of the live-haul route at highest risk.
- **Feed and Critical Operational Visits.** Critical operations visits will be limited during PMIP; however, delivery of feed during this period is likely, and the potential for emergency maintenance visits also exists. The likelihood of a broiler flock becoming infected with HPAI via critical operational visits during PMIP was assessed as *negligible to moderate*, as follows:
  - *Negligible* via contaminated feed
  - *Low* via feed delivery (i.e., contaminated driver and/or vehicle)
  - *Low to moderate* via other critical operational visits (i.e., personnel or vehicle)
- **Growers, Employees, and their Vehicles.** Provided PMIP measures for people are strictly followed and people wear farm-specific clothing and barn-specific footwear, we rate the likelihood of a broiler flock becoming infected with HPAI via people and their vehicles entering the premises during the PMIP as *low* for people entering the poultry barns and *very low* for people who do not enter the poultry barns.

- **Shared Equipment.** Previous outbreaks have demonstrated that shared equipment poses a disease transmission risk; however, during the PMIP, no off-site equipment will be pre-staged and only critical operational visits may continue. Thus, we rated the likelihood of a broiler flock becoming infected with HPAI virus via shared equipment as *low* during PMIP, and *moderate* prior to PMIP.
- **Dead Bird Disposal.** The risks of HPAI introduction associated with off-site dead bird disposal methods, such as rendering, are well documented, and off-site disposal of mortality must be discontinued during PMIP. However, the risky practice of off-site dead bird disposal may still occur outside of a PMIP.
  - For on-farm dead bird disposal, given that many scavenger species can biologically or mechanically carry HPAI virus and have home ranges large enough to contain adjacent poultry farms, we assessed the likelihood of HPAI introduction to a broiler farm during the PMIP as *low to moderate*.
  - Off-site dead bird disposal methods prior to a PMIP may possibly result in premises contamination. However, the implementation of a PMIP does reduce the likelihood that such contamination will be tracked inside a grow-out barn during the PMIP. We thus assessed the likelihood of a broiler flock becoming infected as a result of HPAI virus introduction to the flock via off-site dead bird disposal that takes place prior to the PMIP as *moderate*.
- **Garbage Management.** There is potential for HPAI virus associated with garbage management to be tracked into a poultry house, and thus we assessed the likelihood of a broiler flock becoming infected with HPAI virus due to garbage management without a PMIP to be *moderate to high*. During a PMIP, no off-site movement of garbage is allowed, and thus we assessed the likelihood of a broiler flock becoming infected with HPAI virus due to garbage management during a PMIP as *low*.

### Load-out Pathways

- **Load-out.** Assuming PMIP enhanced biosecurity and testing measures are strictly implemented, and that additional load-out mitigation measures are in place and commensurate with the duration of the premises-wide load-out process, the risk that a broiler flock will become infected with HPAI virus via load-out operations and that this will result in an infected but undetected movement to market is estimated to be *low to moderate*.

## Overall Risk

It is concluded that the overall risk of HPAI spread to susceptible poultry associated with the movement of broilers to market into, within, and outside of a Control Area is *low to moderate*, provided that all applicable preventive measures from the SBS Plan, in particular the PMIP, are strictly followed.

In using the results of this risk assessment, it should be remembered that:

- This assessment is based on current (December 2016) information and will need to be reviewed and revised as circumstances warrant.
- The assessment does not replace the judgment of on-scene officials with first-hand knowledge of the outbreak situation and the premises in question.

## Appendix 1: AI Virus Survival at Various Humidity Levels, at Various Temperatures, and on Various Substrates.

**Appendix 1 Tables 1-6** summarize the results of studies documenting survival and persistence of AI viruses at various humidity levels, at various temperatures, and on various substrates. The trend in persistence and survival time in the environment for AI viruses appears to be decreased survival in conditions of lower moisture and higher temperature. Virus survival and persistence in the environment has also been reported to be longer near neutral pH, in lower salinity, and without UV exposure.<sup>79,269,329-331</sup>

These tables are compiled to describe virus survival and persistence across a range of conditions. Of note, there are multiple methodologies to determine virus survival or persistence; readers should consult the studies listed to evaluate all parameters and methods utilized in experimental studies, as definitions of these terms are not uniformly applied. In compiling data from the literature for these tables, studies where HPAI virus was utilized were given preference over LPAI studies. Where information on AI virus was not available, data on other influenza viruses are included as indicated. Virus inactivation was prioritized as a time point in the summary tables below. In studies in which virus remained viable for all time points measured, the last reported time when virus was measured (and detected) is included in the tables for comparison.

These summary tables focus on conditions that may be similar to those encountered on commercial poultry operations and climatic parameters similar to those of the continental United States. Further summaries of virus inactivation times in eggs and egg products can be found in the OIE Terrestrial Animal Health Code (Article 10.4.25),<sup>332</sup> and inactivation times at high temperatures have been summarized by USDA documents on parameters to inactivate HPAI virus using heat treatment.<sup>333</sup>

**Appendix 1 Table 1.** Summary of experimental studies on survival of AI viruses in feces and manure by increasing temperature.

Substrate	Temperature	Humidity (as described by study authors)	Sub- type	Strain	Last time point detected (if viable for all contact times)	Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)	Reference
Duck feces	0°C (32°F)	Moist germ carrier; feces in closed 50- ml plastic tubes	LPAI H5N1	A/Teal/Wv632/ Germany/05	-	T <sub>90</sub> <sup>f</sup> value of 75 days	Nazir et al., 2011 <sup>334</sup>
Wet Chicken feces	4°C (39.2°F)	Closed vial	HPAI H5N2	#1370 isolate	Viable virus through 35 days (last time point tested)	-	Beard et al., 1984 <sup>78</sup>

<sup>f</sup> T<sub>90</sub> value: time required for 90% loss of virus infectivity

**Appendix 1 Table 1.** Summary of experimental studies on survival of AI viruses in feces and manure by increasing temperature, cont.

Substrate, <i>cont.</i>	Temperature	Humidity (as described by study authors)	Sub-type	Strain	Last time point detected (if viable for all contact times)	Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)	Reference
Commercial chicken manure (field house)	4°C (39.2°F)	Manure-virus mixture in a 50-ml sterile tube	LPAI H7N2	A/chicken/PA/3779-2/97AIV	Remained activated at 20 days	-	Lu et al., 2003 <sup>335</sup>
Wet chicken feces	4°C (39.2°F)	Capped vials	H5N1	A/Ck/Sikkim/15146/6/2008	-	0% infectivity at week 7	Kurmi et al., 2013 <sup>277</sup>
Dry chicken feces	4°C (39.2°F)	Capped vials	H5N1	A/Ck/Sikkim/15146/6/2008	-	0% infectivity at week 8	Kurmi et al., 2013 <sup>277</sup>
Chicken feces	4.0-6.7°C (39.2-44.06°F)	15.2-46.3% relative humidity	H5N1	A/Vietnam/1203/2004v	-	Virus not detected at day 13	Wood et al., 2010 <sup>269</sup>
Chicken feces	6.7-7.8°C (44.06-46.04°F)	79.0-96.9% relative humidity	H5N1	A/Vietnam/1203/2004	Day 13 (last time point tested)	-	Wood et al., 2010 <sup>269</sup>
Duck feces	10°C (50°F)	Moist germ carrier; feces in closed 50-ml plastic tubes	LPAI H6N8	A/Mute Swan/Germany/R29/27/07	-	T <sub>90</sub> value of 14 days	Nazir et al., 2011 <sup>334</sup>
Commercial chicken manure (field house)	15-20°C (59-68°F)	Manure-virus mixture in a 50-ml sterile tube	LPAI H7N2	A/chicken/PA/3779-2/97AIV	Remained activated at 2 days	-	Lu et al., 2003 <sup>335</sup>
Duck feces	20°C (68°F)	Moist germ carrier; feces in closed 50-ml plastic tubes	LPAI H4N6	A/Mallard/Wv1732-34/03	-	T <sub>90</sub> value of 4 days	Nazir et al., 2011 <sup>334</sup>
Fecal material	22°C (71.6°F)	Capped glass vials	H3N6	A/Duck/Memphis/546/74	-	Infectious virus not detected at day 13	Webster et al., 1978 <sup>336</sup>
Chicken feces	22.0-22.7°C (71.6-72.86°F)	30-42% relative humidity	H5N1	A/Vietnam/1203/2004	-	Virus not detected at day 2	Wood et al., 2010 <sup>269</sup>

**Appendix 1 Table 1.** Summary of experimental studies on survival of AI viruses in feces and manure by increasing temperature, cont.

<b>Substrate, cont.</b>	<b>Temperature</b>	<b>Humidity (as described by study authors)</b>	<b>Sub-type</b>	<b>Strain</b>	<b>Last time point detected (if viable for all contact times)</b>	<b>Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)</b>	<b>Reference</b>
Chicken feces	22.4-23.7°C (72.32-74.66°F)	89.1-91.2% relative humidity	HPAI H5N1	A/Vietnam/1203/ 2004	-	Virus not detected at day 4	Wood et al., 2010 <sup>269</sup>
Wet chicken feces	25°C (77°F)	Closed vial	HPAI H5N2	#1370 isolate	-	No viable virus at day 3	Beard et al., 1984 <sup>78</sup>
Field commercial chicken manure	28-30°C (82.4-86°F)	Manure-virus mixture in a 50-ml sterile tube	LP AI H7N2	A/chicken/PA/3779- 2/97AIV	-	Inactivated at hour 12	Lu et al., 2003 <sup>335</sup>
Duck feces	30°C (86°F)	Moist germ carrier; feces in closed 50-ml plastic tubes	LP AI H4N6	A/Mallard/Wv1732- 34/03	-	T <sub>90</sub> value of 2 days	Nazir et al., 2011 <sup>334</sup>
Dry chicken feces	37°C (98.6°F)	Capped vials	HPAI H5N1	A\Ck\Sikkim\15146 6\2008	-	0% infectivity at hour 30	Kurmi et al., 2013 <sup>277</sup>
Wet chicken feces	37°C (98.6°F)	Capped vials	HPAI H5N1	A\Ck\Sikkim\15146 6\2008	-	0% infectivity at hour 30	Kurmi et al., 2013 <sup>277</sup>
Field commercial chicken manure	37°C (98.6°F)	Manure-virus mixture in a 50-mL sterile tube	LP AI H7N2	A/chicken/PA/3779- 2/97AIV	-	Inactivated at hour 24	Lu et al., 2003 <sup>335</sup>
Dry chicken feces	42°C (107.6°F)	Capped vials	HPAI H5N1	A\Ck\Sikkim\15146 6\2008	-	0% infectivity at hour 24	Kurmi et al., 2013 <sup>277</sup>
Wet chicken feces	42°C (107.6°F)	Capped vials	HPAI H5N1	A\Ck\Sikkim\15146 6\2008	-	0% infectivity at hour 24	Kurmi et al., 2013 <sup>277</sup>
Field commercial chicken manure	56°C (132.8°F)	Manure-virus mixture in a 50-mL sterile tube	LP AI H7N2	A/chicken/PA/3779- 2/97AIV	-	Inactivated at minute 15	Lu et al., 2003 <sup>335</sup>



**Appendix 1 Table 2.** Summary of experimental studies on survival of AI viruses in compost by increasing temperature.

Substrate	Temperature	Humidity (as described by study authors)	Subtype	Strain	Last time point detected (if viable for all contact times)	Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)	Reference
Cage layer manure in middle of compost	Peak recorded 46 °C (114.8°F)	Nylon mesh bag; 65% moisture content of compost	LPAI H6N2	A/Tky/Mass/3740/ 65	-	Virus below detectable limit at day 3 (1 <sup>st</sup> time point tested)	Guan et al., 2009 <sup>288</sup>
Used litter in middle of compost	Peak recorded 46°C (114.8°F)	Nylon mesh bag; 65% moisture content of compost	LPAI H6N2	A/Tky/Mass/3740/ 65	-	Virus below detectable limit at day 3 (1 <sup>st</sup> time point tested)	Guan et al., 2009 <sup>288</sup>
Breast muscle in abdominal cavity of chicken carcass at <u>bottom</u> of compost	Peak recorded 50.3°C (122.54°F)	Plastic netting; 65% moisture content of compost	LPAI H6N2	A/Tky/Mass/3740/ 65	-	Virus below detectable limit at day 10	Guan et al., 2009 <sup>288</sup>
Embryonated chicken eggs at <u>bottom</u> of compost	Peak recorded 50.3°C (122.54°F)	Plastic mesh baskets; 65% moisture content of compost	LPAI H6N2	A/Tky/Mass/3740/ 65	-	Virus below detectable limit at day 10	Guan et al., 2009 <sup>288</sup>

**Appendix 1 Table 3.** Summary of experimental studies on survival of AI viruses in water by increasing temperature.

Substrate	Temperature	Humidity (as described by study authors)	Subtype	Strain	Last time point detected (if viable for all contact times)	Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)	Reference
Surface water (Lake Constance)	-10°C (14°F)	-	LPAI H6N8	A/mute swan/ Germany/R2927/07	-	T <sub>90</sub> value of 395 days	Nazir et al., 2010 <sup>337</sup>
Surface water (Lake Constance)	0°C (32°F)	-	LPAI H5N1	A/teal/Germany/Wv 632/05	-	T <sub>90</sub> value of 208 days	Nazir et al., 2010 <sup>337</sup>
Contaminated fecal material in river water	4°C (39.2°F)	-	LPAI H3N6	A/Duck/Memphis/5 46/74	Viable for all contact times (32 days)		Webster et al., 1978 <sup>336</sup>
Sea water (Black Sea)	5-6°C (41-42.8°F)	-	LPAI H6N2	Not specified	-	No infective virus detected at day 7	Zarkov, 2006 <sup>338</sup>
Sea water (Black Sea)	5-6°C (41-42.8°F)	-	LPAI H11N6	A/duck/England/ 56	-	No infective virus detected at day 9	Zarkov, 2006 <sup>338</sup>
Surface water (Koprinka dam)	5-6°C (41-42.8°F)	-	LPAI H6N2	Not specified	-	No infective virus detected at day 16	Zarkov, 2006 <sup>338</sup>
Surface water (Koprinka dam)	5-6°C (41-42.8°F)	-	LPAI H11N6	A/duck/England/ 56	-	No infective virus detected at day 18	Zarkov, 2006 <sup>338</sup>
Surface water (Lake Constance)	10°C (50°F)	-	LPAI H4N6	A/mallard/Germany/ Wv1732-34/03	-	T <sub>90</sub> value of 85 days	Nazir et al., 2010 <sup>337</sup>
Surface water (Ovcharitsa dam)	10-12°C (50-53.6°F)	-	LPAI H6N2	Not specified	-	No infective virus detected at day 1	Zarkov, 2006 <sup>338</sup>
Surface water (Ovcharitsa dam)	10-12°C (50-53.6°F)	-	LPAI H11N6	A/duck/England/ 56	-	No infective virus detected at day 1	Zarkov, 2006 <sup>338</sup>
Distilled water	17°C (62.6°F)	-	HPAI H5N1	A/WhooperSwan/M ongolia/244/05	-	Predicted persistence of 158 days	Brown et al., 2007 <sup>331</sup>

**Appendix 1 Table 3.** Summary of experimental studies on survival of AI viruses in water by increasing temperature, cont.

<b>Substrate, cont.</b>	<b>Temperature</b>	<b>Humidity (as described by study authors)</b>	<b>Subtype</b>	<b>Strain</b>	<b>Last time point detected (if viable for all contact times)</b>	<b>Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)</b>	<b>Reference</b>
Surface water (Lake Constance)	20°C (68°F)	-	LPAI H4N6	A/mallard/Germany/ Wv1732-34/03	-	T <sub>90</sub> value of 23 days	Nazir et al., 2010 <sup>337</sup>
Contaminated fecal material in river water	22°C (71.6°F)	-	LPAI H3N6	A/Duck/Memphis/5 46/74	-	Virus not detected at day 7	Webster et al., 1978 <sup>336</sup>
Distilled water	28°C (82.4°F)	-	H5N1	A/DuckMeat/ Anyang/01	-	Predicted persistence of 30 days	Brown et al., 2007 <sup>331</sup>
Surface water (Lake Constance)	30°C (86°F)	-	LPAI H4N6	A/mallard/Germany/ Wv1732-34/03	-	T <sub>90</sub> value of 14 days	Nazir et al., 2010 <sup>337</sup>

**Appendix 1 Table 4.** Summary of experimental studies on survival of AI viruses in poultry carcass (meat, liver, muscle, feather) by increasing temperature.

Substrate	Temperature	Humidity (as described by study authors)	Sub- type	Strain	Last time point detected (if viable for all contact times)	Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)	Reference
Duck feathers	4°C (39.2°F)	Placed in incubator	H5N1	A/chicken/Miyaza ki/K11/2007 A/WhooperSwan/ Akita/1/2008	-	Negative for virus isolation at day 200	Yamamoto et al., 2010 <sup>246</sup>
Breast muscle in abdominal cavity of chicken carcass	3.9-7.9°C (39-46.2°F)	Plastic netting outside compost bin	LPAI H6N2	A/Tky/Mass/3740/ 65	Virus detected at all times tested (21 days)	-	Guan et al., 2009 <sup>288</sup>
Liver in abdominal cavity of chicken carcass	4.0-7.9°C (39.2-46.2°F)	Plastic netting outside compost bin	LPAI H6N2	A/Tky/Mass/3740/ 65	-	Virus not detected at day 7	Guan et al., 2009 <sup>288</sup>
Duck feathers	20°C (68°F)	Placed in incubator	H5N1	A/WhooperSwan/ Akita/1/2008	-	Negative for virus isolation at day 20	Yamamoto et al., 2010 <sup>246</sup>
Chicken meat	57.8°C (136.04°F)	PCR tubes in 163hermos- cycler heating block	H5N1	A/chicken/Korea/ ES/2003	-	Predicted 11-log EID <sub>50</sub> reduction at 39.6 minutes	Thomas et al., 2007 <sup>339</sup>

**Appendix 1 Table 5.** Summary of experimental studies on survival of AI viruses in allantoic fluid and embryonated chicken eggs by increasing temperature.

Substrate	Temperature	Humidity (as described by study authors)	Subtype	Strain	Last time point detected (if viable for all contact times)	Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)	Reference
Embryonated chicken eggs	3.9-7.9°C (39-46.2°F)	Plastic mesh baskets outside compost bin	LPAI H6N2	A/Tky/Mass/3740/65	Virus detected at all times tested (21 days)	-	Guan et al., 2009 <sup>288</sup>
Allantoic fluid	55°C (131°F)	Capped centrifuge tubes	H5N1	A/chicken/Chonburi/ Thailand/CU-7/04, A/chicken/Nakorn Patom/Thailand/CU- K2/2004, A/chicken/Ratchaburi/ Thailand/CU-68/04	Infective at all contact times (60 minutes)	-	Wanaratana et al., 2010 <sup>340</sup>
Allantoic fluid	56°C (132.8°F)	Thermocycler tubes in heating block	LPAI H7N9	A/Anhui/1/2013, A/Shanghai/1/2013	-	Virus not infective at minute 30	Zou et al., 2013 <sup>341</sup>
Allantoic fluid	60°C (140°F)	Capped centrifuge tubes	H5N1	A/chicken/Chonburi/ Thailand/CU-7/04, A/chicken/Nakorn Patom/Thailand/CU- K2/2004, A/chicken/Ratchaburi/ Thailand/CU-68/04	-	Virus not infective at minute 60	Wanaratana et al., 2010 <sup>340</sup>

**Appendix 1 Table 6.** Summary of experimental studies on survival of influenza viruses on additional substrates by increasing temperature.

Substrate	Temperature	Humidity (as described by study authors)	Subtype	Strain	Last time point detected	Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)	Reference
Galvanized metal, glass, soil	4.0-6.7°C (39.2-44.06°F)	15.2-46.3% relative humidity	H5N1	A/Vietnam/1203/ 2004	Virus detected at all times tested (13 days)	-	Wood et al., 2010 <sup>269</sup>
Galvanized metal	6.7-7.8°C (44.06-46.04°F)	89.5-96.9% relative humidity	H5N1	A/Vietnam/1203/ 2004	-	Virus below detectable limit at day 9	Wood et al., 2010 <sup>269</sup>
Glass, soil	6.7-7.8°C (44.06-46.04°F)	79.0-96.9% relative humidity	H5N1	A/Vietnam/1203/ 2004	-	Virus below detectable limit at day 13	Wood et al., 2010 <sup>269</sup>
Window glass, unvarnished oak	17-21°C (62.6-69.8°F)	23-24% humidity	H1N1	A/PuertoRico/8/34 (PR8)	-	Virus not detected at hour 4	Greatorex et al., 2011 <sup>342</sup>
Stainless steel, plastic control	17-21°C (62.6-69.8°F)	23-24% humidity	H1N1	A/PuertoRico/8/34 (PR8)	-	Virus not detected at hour 24	Greatorex et al., 2011 <sup>342</sup>
Steel, tile, gumboot, tire, egg shell, plastic	Unspecified room temperature	In 14-ml round- bottom tubes and stored in a drawer	LPAI H13N7	A/Herringgull/ Delaware 471/86	-	Virus below detectable limit at day 6	Tiwari et al., 2006 <sup>270</sup>
Latex, feather	Unspecified room temperature	In 14-ml round- bottom tubes and stored in a drawer	LPAI H13N7	A/Herringgull/ Delaware 471/86	Virus detected at day 6 (last time point tested)	-	Tiwari et al., 2006 <sup>270</sup>
Wood	Unspecified room temperature	In 14-ml round- bottom tubes and stored in a drawer	LPAI H13N7	A/Herringgull/ Delaware 471/86	-	Virus below detectable limit at hour 72	Tiwari et al., 2006 <sup>270</sup>
Egg tray, polyester fabric	Unspecified room temperature	In 14-ml round- bottom tubes and stored in a drawer	LPAI H13N7	A/Herringgull/ Delaware 471/86	-	Virus below detectable limit at hour 24	Tiwari et al., 2006 <sup>270</sup>
Cotton fabric	Unspecified room temperature	In 14-ml round- bottom tubes and stored in a drawer	LPAI H13N7	A/Herringgull/ Delaware 471/86	-	Virus below detectable limit at hour 48	Tiwari et al., 2006 <sup>270</sup>

**Appendix 1 Table 6.** Summary of experimental studies on survival of influenza viruses on additional substrates by increasing temperature, cont.

<b>Substrate, cont.</b>	<b>Temperature</b>	<b>Humidity (as described by study authors)</b>	<b>Subtype</b>	<b>Strain</b>	<b>Last time point detected</b>	<b>Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)</b>	<b>Reference</b>
Stainless steel	22°C (71.6°F)	50-60% relative humidity	H1N1	A/PR/8/34	Viable virus at hour 24 (last time examined)	-	Noyce et al., 2007 <sup>343</sup>
Galvanized metal, glass	22.7-23.4°C (72.86-74.12°F)	32-38% relative humidity	H5N1	A/Vietnam/1203/2004	-	Virus below detectable limit at day 1	Wood et al., 2010 <sup>269</sup>
Soil	22.0-23.4°C (71.6-74.12°F)	30-42% relative humidity	H5N1	A/Vietnam/1203/2004	-	Virus below detectable limit at day 2	Wood et al., 2010 <sup>269</sup>
Galvanized metal, glass	22.4°C (72.32°F)	89.1% relative humidity	H5N1	A/Vietnam/1203/2004	-	Virus below detectable limit at day 1	Wood et al., 2010 <sup>269</sup>
Soil	22.4-23.4°C (72.32-74.12°F)	89.1-90.4% relative humidity	H5N1	A/Vietnam/1203/2004	-	Virus below detectable limit at day 2	Wood et al., 2010 <sup>269</sup>
Rubber glove, N95 particulate respirator, surgical mask (non-woven fabric), gown (Dupont Tyvek), coated wooden desk, stainless steel	25.2°C (77.36°F)	55% relative humidity	H1N1	A/PR/8/34	Virus detected at hour 24 (last time point tested)	-	Sakaguchi et al., 2010 <sup>344</sup>
Plastic	27.8-28.3°C (82.0-82.9°F)	35-40% relative humidity	H1N1	A/Brazil/11/78-like	Virus detected at ~10 <sup>1</sup> TCID <sub>50</sub> /0.1 ml at hour 48 (last time point tested)	-	Bean et al., 1982 <sup>56</sup>
Stainless steel	27.8- 28.3°C (82.0- 82.9°F)	35-40% relative humidity	H1N1	A/Brazil/11/78-like	-	Virus below detectable limit at hour 72	Bean et al., 1982 <sup>56</sup>

**Appendix 1 Table 6.** Summary of experimental studies on survival of influenza viruses on additional substrates by increasing temperature, cont.

<b>Substrate, cont.</b>	<b>Temperature</b>	<b>Humidity (as described by study authors)</b>	<b>Subtype</b>	<b>Strain</b>	<b>Last time point detected</b>	<b>Time to virus inactivation (experimental, estimated, or predicted based on regression analysis)</b>	<b>Reference</b>
Stainless steel	55°C (131°F)	50% relative humidity	H1N1	A/PR/8/34	Minute 60 (last time point tested)	-	McDevitt et al., 2010 <sup>345</sup>
Stainless steel	60°C (140°F)	50% relative humidity	H1N1	A/PR/8/34	-	Virus below detectable limit at minute 30	McDevitt et al., 2010 <sup>345</sup>
Stainless steel	65°C (149°F)	50% relative humidity	H1N1	A/PR/8/34	-	Virus below detectable limit at minute 15 (1 <sup>st</sup> time point tested)	McDevitt et al., 2010 <sup>345</sup>



## Appendix 2: Literature Review on the Role of Local Area Spread in Previous Outbreaks

Appendix 2 Table 1 below summarizes the results from studies (to include modeling) on the influence of local area spread in AI transmission during previous outbreaks.

**Appendix 2 Table 1.** Previous AI outbreak investigations and results associated with local area spread.

AI strain (Location)	Year of outbreak (species involved)	Study approach	Key findings	Source
HPAI/LPAI H7N8 (Indiana)	2016 (turkeys)	Geospatial analysis; case-control (9 cases, 30 controls)	The geospatial analysis showed a likely association between infected premises and a common driving route. The case-control study identified risk factors more common on case farms and in case barns than on control farms and in control barns as: shorter distance to dead bird disposal and litter compost area, presence of wild mammals, and visitors entering barns.	USDA Epi Report Indiana, March 18, 2016 <sup>260</sup>
HPAI H5N2 (Iowa)	2015 (layer chickens)	Case-control (28 cases, 31 control) with multivariate logistic regression	Farm-level analysis indicated that location in an existing control zone (10 km perimeter beyond the closest infected premises) was highly associated with infection status.	Garber et al., 2016 <sup>192</sup>
LPAI H5N2 (Japan)	2005 (layer chickens)	Case-control (37 cases, 36 controls) with multivariate logistic regression. Biosecurity factors controlled for.	Distance up to 1.5 km from infected premises identified as a risk factor for egg layer farms in Japan. Equipment sharing and visitor biosecurity were also identified as risk factors.	Nishiguchi et al., 2007 <sup>256</sup>
HPAI H5N2 (Texas)	2004 (broiler chickens)	Outbreak observation	No area spread. Samples were collected from 368 premises (39 in the 8-km affected zone, 167 in the surveillance zone [16 km], and 162 in the buffer zone [50 km]).	Pelzel et al., 2006 <sup>346</sup>
HPAI H7N7 (Netherlands)	2003 (multiple poultry species)	Spatial transmission model with distance and infectious period at premises level as factors	Exposure increased with proximity to infectious farm. Farms $\leq 1$ km from an infected premises were at least 8 times more likely to become infected than farms $\geq 5$ km.	Boender et al., 2007 <sup>133</sup>

**Appendix 2 Table 1.** Previous AI outbreak investigations and results associated with local area spread, cont.

AI strain (Location)	Year of outbreak (species involved)	Study approach	Key findings	Source
LPAI H7N2 (Virginia, West Virginia, North Carolina)	2002 (chickens and turkeys)	Outbreak observation	Spread mainly by people and fomites, including equipment; rendering especially high risk. Very little evidence for airborne spread.	Senne et al., 2005 <sup>347</sup>
LPAI H7N2 (Pennsyl- vania)	2001-2002 (broiler breeders and broiler chickens)	Outbreak observation	Local spread within 1 mile. Likely mechanisms were family ties, business connection, social contact, etc.	Dunn et al., 2003 <sup>268</sup>
HPAI H7N1 (Italy)	1999-2000 (turkeys [meat and breeder], chickens [breeders, layers, and broilers], geese, quail, ostriches, guinea fowl, pheasants)	Multivariable Cox regression; people and equipment flow not controlled for in model.	Flocks $\leq 1.5$ km from an infected premises were estimated to have a Hazard ratio of 7.9.  Poultry species and farm size also were identified as risk factors.	Mannelli et al., 2006 <sup>348</sup>
		Multivariable Cox regression; people and equipment flow not controlled for in model.	Flocks $\geq 4.5$ km from infected premises had lower risk. Flocks $\leq 1.5$ km from infected premises had highest risk (hazard ratio 4.6 in comparison to flocks $> 4.5$ km from an infected premises).  Poultry species, type of production, and farm size also were identified as risk factors.	Busani et al., 2009 <sup>135</sup>
		Spatial transmission model with distance and infectious period at premises level as factors	Proximity to infectious farms increased the risk of infection, e.g., probability of infection estimated to be 2.5 times higher for susceptible farms 1 km from an infectious farm than for farms 3 km away.  Control measures such as culling of infected farms and ban on restocking were identified through simulation to reduce infection spread.	Dorigatti et al, 2010 <sup>132</sup>

**Appendix 2 Table 1.** Previous AI outbreak investigations and results associated with local area spread, cont.

<b>AI strain (Location)</b>	<b>Year of outbreak (species involved)</b>	<b>Study approach</b>	<b>Key findings</b>	<b>Source</b>
HPAI H5N1 (England)	2007 (turkeys)	Outbreak observation, spatial simulation model	Spread to 3 houses on the same premises. No transmission to 78 other farms within a 3-km protection zone or 70 farms within a 10-km surveillance zone.  Simulation showed no evidence of local transmission above 1 km.	Sharkey et al., 2008 <sup>136</sup> ; Irvine et al., 2007 <sup>126</sup>
LPAI H5N3 (California)	1984 (turkeys)	Outbreak observation	Spread associated with insemination at 5 breeder premises across 110 miles, linked to one company and insemination crew. No spread to 193 other turkey premises or >800 chicken premises in the state.	McCapes et al., 1986 <sup>92</sup>
LPAI H6N1 (Minnesota)	1978 (layer chickens)	Outbreak observation	No spread to 1 of 4 houses on the same layer premises; the unaffected house was across a road from the 3 affected and interconnected houses. No spread to epidemiologically linked layer farms or neighboring premises.	Halvorson et al., 1980 <sup>349</sup>
LPAI A/T/Minn./67 (Minnesota)	1967 (turkeys)	Outbreak observation	Spread between houses on same premises and between premises. Spread between premises appeared associated with insemination; some houses on severely infected premises were not infected.	Kleven et al., 1970 <sup>95</sup>

### Appendix 3: Estimating an Approximate Posterior Distribution for the Prevalence among Insects

A Bayesian approach with Monte Carlo simulation was used to estimate the posterior distribution of the prevalence among insects given the observed data from testing pools of insects.<sup>138</sup> In the observed data, the pool size varied between 10 and 60 insects, and 2 of the 144 pools tested were positive for AI. The steps in the simulation iteration were as follows.

For each of the 144 pooled samples, the pool size  $N_j$  was simulated as an integer Uniform (10, 60) distribution. The prevalence among individual insects  $P_r$  was simulated using a uniform (0,1) prior (uninformed). The probability of a pooled sample being positive  $P_{pool}(j)$  was calculated according to the equation below,

$$P_{pool}(j) = 1 - (1 - P_r)^{N_j}$$

An indicator variable for whether the pooled sample  $j$  is positive,  $X(j)$  was simulated as a Bernoulli trial with the probability equal to  $P_{pool}(j)$ .

The prevalence  $P_r$  in an iteration was accepted if the sum  $\sum_j X_j = 2$ , as only 2 out of 144 pools were positive in the data. The simulation was run for 2,000,000 iterations to estimate the approximate posterior distribution of  $P_r$ . An approximate two-sided 95% credibility interval was (0.01%, 0.15%) based on 391 values.

## Appendix 4: Expert Polling on Insect Transmission Routes

A panel of eight experts in the turkey and broiler industries with field experience managing AI was surveyed between November 2013 and January 2014 on risk of HPAI transmission via multiple routes of infection. Surveys were administered through the online polling service SurveyMonkey.<sup>§</sup> Experts were asked to provide their opinion, based on previous experience, of perceived risk for given scenarios. Qualitative risk rating definitions were provided and match those used in this risk assessment (with the exception that the survey did not include a “very low risk” option) (see section 5, Overview of Data Analysis Approaches, for risk rating definitions). Below is the subset of questions that pertain to spread by insects with associated expert responses shown in **Appendix 4 Tables 1-2** and **Appendix 4 Figures 1-2**.


*Q1. Please qualitatively rate the likelihood of AI transmission from a known infected flock to an uninfected turkey flock via insects located at distances specified in the table. Please complete the following table, selecting a risk rating for each scenario as negligible, low, moderate, high, or extremely high, based on your expert opinion.*

*Q2. Please qualitatively rate the likelihood of AI transmission from an infected but undetected flock (lower prevalence) to an uninfected turkey flock via insects located at distances specified in the table. Please complete the following table, selecting a risk rating for each scenario as negligible, low, moderate, high, or extremely high, based on your expert opinion.*

**Appendix 4 Table 1.** Expert responses (n=8) to the question of likelihood of AI transmission from a **known infected** flock to an uninfected turkey flock via insects at specified distances (Question 1).

Distance from source flock	Likelihood rating				
	Negligible	Low	Moderate	High	Extremely high
1 km	1	3	2	2	0
1.5 km	3	3	1	1	0
2 km	6	1	1	0	0
3 km	7	1	0	0	0


<sup>§</sup> SurveyMonkey, Inc., Palo Alto, CA, [www.surveymonkey.com](http://www.surveymonkey.com)



**Appendix 4 Figure 1.** Expert responses (n=8) to the question of likelihood of AI transmission from a **known infected** flock to an uninfected turkey flock via insects at specified distances (Question 1).

**Appendix 4 Table 2.** Expert responses (n=8) to the question of likelihood of AI transmission from an **infected but undetected** (lower prevalence) flock to an uninfected turkey flock via insects at specified distances (Question 2).

Distance from source flock	Likelihood rating				
	Negligible	Low	Moderate	High	Extremely high
1 km	2	1	3	2	0
1.5 km	3	2	2	1	0
2 km	5	2	1	0	0
3 km	6	2	0	0	0



**Appendix 4 Figure 2.** Expert responses (n=8) to the question of likelihood of AI transmission from an **infected but undetected** (lower prevalence) flock to an uninfected turkey flock via insects at specified distances (Question 2).

## Appendix 5: Live Broiler Movement Aerosol Modeling

### **Introduction**

AERMOD<sup>350,351</sup> is a regulatory model used by, among others, the U.S. EPA for air quality assessment. The model inputs include information on source location and parameters, receptor locations, and meteorological variables (such as wind speed, wind direction, and turbulence parameters), and outputs include concentrations at various receptor locations over selected time periods. The model can output printed summaries of, for example, high values by receptor, overall maximum values for each averaging period, and tables of concurrent values summarized by receptor. These values can then be used to generate outputs such as contour plots that depict the concentrations at the various receptor locations. The model can also generate a file of all occurrences when a concentration value equals or exceeds a user-specified threshold.

### **Scenario A:**

#### Source and Receiving Flock Size

The source infected premises (IP) in this scenario is a 25,000-bird HPAI-infected broiler house, and the receiving flock is a 20,000-bird broiler house. The weight of receiving broilers was assumed to be 6 lb in this scenario where the air intake per bird would be 0.88 m<sup>3</sup>/day using equations presented in Lasiewski and Calder (1971).<sup>352</sup>

#### Meteorological Parameters

Meteorological data such as wind speed, relative humidity, temperature, etc. were obtained from Tupelo, Mississippi, for the year 2011.

#### Aerosol Source Emission Rate

The emission rate was directly estimated from the total suspended particle emission rate for a broiler house in the literature. Burns et al. (2008) estimated a mean particle emission rate of 2.78 ± 1.87 kg/day-house for a broiler house with average placement of 25,000 chickens.<sup>353</sup> Assuming that 50 percent of the suspended particles were contaminated at an HPAI virus concentration of 10<sup>5</sup> EID<sub>50</sub>/g and a particle emission rate of 4.65 (2.78+1.87) kg/day-house, the aerosol source emission rate would be 10<sup>3.43</sup> EID<sub>50</sub> per second (s) = 50%\*4.65\*1,000 g/day/(24\*3,600 s/day)\*10<sup>5</sup>EID<sub>50</sub>/g.

#### Particle Size Distribution

Particle pollution, also known as particulate matter or PM, is a mixture of very small particles and liquid droplets in the air. Generally, the smaller the particle, the more likely it could be inhaled and cause health problems. The EPA is most concerned with particles less than 10 micrometers in diameter, as these are generally considered small enough to pass through the nose and throat and potentially enter the lungs.<sup>354</sup> We considered that 41 percent of particles are PM<sub>10</sub> and 27 percent of PM<sub>10</sub> particles are PM<sub>2.5</sub>. The particle size fraction in this scenario was chosen using a conservative approach from the data points in **Appendix 5 Table 1** to have a greater proportion of small particles. For particles greater than 10 μm in size, a diameter of 25 μm was used based on the mean mass diameter estimate from Redwine et al. (2002).<sup>355</sup>



### HPAI 50 Percent Infectious Dose

As mentioned in the literature review, we used an exponential dose-response model parameterized from data presented in Spekrijse et al. (2012). The specific data points from Spekrijse et al. (2012) that we included in our analysis are shown in **Appendix 5 Table 2**. In our analysis, we excluded data points where the concentration of HPAI virus was zero. We used a chicken air intake rate of  $1.2 \text{ m}^3/\text{day}/\text{bird}$ <sup>356</sup> in order to convert the concentration into dose per chicken per day.

The parameter for exponential distribution  $r$  was estimated to be  $2.517 \times 10^{-6}/\text{EID}_{50}$ . There is considerable uncertainty regarding the parameters as well as the shape of the dose-response model for the aerosol route, given the limited data (**Appendix 5 Table 2**). In particular, as aerosol may represent a very low-concentration exposure for a large number of birds, the risk estimate would be considerably lower if there existed a threshold dose below which the probability of an exposed bird becoming infected were zero. The exponential dose-response model is a “single hit” model without a threshold dose.

**Appendix 5 Table 1.** Fraction of particles from poultry operations with size less than or equal to  $10\mu\text{m}$  ( $\text{PM}_{10}$ ) and  $2.5\mu\text{m}$  ( $\text{PM}_{2.5}$ ).

Study	$\text{PM}_{10}$ fraction of total suspended particles (%)	$\text{PM}_{2.5}$ fraction of PM 10 (%)
Burns et al. (2008) <sup>353</sup> (background PM controlled)	41	
Roumeliotis (2010) <sup>357</sup> (background PM controlled)		15.6
Wathes et al. (1997) <sup>358</sup>	10	
Li et al. (2008), <sup>359</sup> turkey toms		11
Redwine (2002) <sup>355</sup> (background PM not controlled)	5.9 (2.7-8.4)	
Takai et al. (1998), <sup>360</sup> inhalable vs. respirable dust (background PM not controlled )	13	
Li et al. (2009), <sup>361</sup> layers (background PM not controlled)		10.5
Roumeliotis (2007) <sup>362</sup> (background PM not controlled)	77	27
Unpublished industry data	50-70	50

**Appendix 5 Table 2.** Data points from Spekrijse et al. (2012) used to parameterize an exponential dose-response model through maximum likelihood methods.<sup>356</sup>


<b>Trial</b>	<b>Concentration EID<sub>50</sub>/m<sup>3</sup></b>	<b>Dose (EID<sub>50</sub>)</b>	<b>Number infected out of 14 birds</b>
Trial 2 day 4	2.3	2.38	0
Trial 2 day 5	2	2.08	0
Trial 2 day 6	1.5	1.58	1
Trial 2 day 9	2.2	2.28	0
Trial 2 day 10	2.8	2.88	0
Trial 4 day 2	2.5	2.58	0
Trial 4 day 3	2.9	2.98	0
Trial 4 day 4	3.5	3.58	0
Trial 4 day 5	3.2	3.28	0
Trial 4 day 6	4.1	4.18	0
Trial 4 day 7	3.9	3.98	0
Trial 4 day 8	3.5	3.58	0
Trial 4 day 9	4	4.08	0
Trial 4 day 10	3.7	3.78	0
Trial 4 day 13	3.5	3.58	2
Trial 4 day 14	3.7	3.78	0
Trial 1 day 2	2.1	2.18	0
Trial 1 day 3	2.9	2.98	0
Trial 1 day 4	3.1	3.18	0
Trial 1 day 5	2.6	2.68	0
Trial 1 day 6	2.8	2.88	0
Trial 1 day 7	2.9	2.98	0
Trial 1 day 8	2.7	2.78	0
Trial 3 day 2	2.4	2.48	0
Trial 3 day 4	3.05	3.12	0
Trial 3 day 5	2.7	2.78	0
Trial 3 day 6	3.1	3.18	0
Trial 3 day 8	3.51	3.59	0

**Appendix 5 Table 3.** Data points from Spekrijse et al. (2012) used to parameterize an exponential dose-response model through maximum likelihood methods.<sup>356</sup> cont.

<b>Trial, cont.</b>	<b>Concentration EID<sub>50</sub>/m<sup>3</sup></b>	<b>Dose (EID<sub>50</sub>)</b>	<b>Number infected out of 14 birds</b>
Trial 3 day 9	2.75	2.83	0
Trial 3 day 10	3.28	3.36	0
Trial 3 day 13	2.91	2.99	0
Trial 3 day 14	3.50	3.58	0

**Scenario A results:**


Scenario A results are shown in **Appendix 5 Figures 1-2** and **Appendix 5 Table 3**. We can see from **Appendix 5 Figure 1** that the predicted aerosol concentration is highest in the downwind direction and decreases sharply with distance. The predicted concentration is low farther than 2.5 km from the source broiler IP.



**Appendix 5 Figure 1** AERMOD model-predicted HPAI virus concentrations at various points downwind of a source broiler IP (Scenario A)

**Appendix 5 Table 3.** AERMOD model-predicted HPAI virus concentrations and daily exposure probability at various points downwind of a source broiler IP (Scenario A).

Outcome Variable	Distance from source (known infected broiler premises)				
	0.5 km	1 km	1.5 km	3 km	5 km
Predicted HPAI concentration (EID <sub>50</sub> /m <sup>3</sup> )	0.55	0.19	0.1	0.04	0.02
Predicted probability of exposure in a day	0.0242	0.0085	0.0046	0.0016	0.000738



**Appendix 5 Figure 2.** AERMOD model-predicted daily probability of infection for a susceptible broiler flock located at various points downwind of a source broiler IP (Scenario A).

## **Scenario B:**

### Source and Receiving Flock Size

The source farm in this scenario is a 14,000-bird HPAI-infected turkey flock, and the receiving flock is a 20,000-bird broiler house. The weight of receiving broilers was assumed to be 6 lb in this scenario in which case the air intake per bird would be  $0.88 \text{ m}^3/\text{day}$ , obtained using the equations presented in Lasiewski and Calder 1971.<sup>352</sup>

### Meteorological Parameters

Meteorological data such as wind speed, relative humidity, and cloud cover, etc., were obtained from Olivia, Minnesota, for the year 2011.

### Aerosol Source Emission Rate

We used a  $\text{PM}_{10}$  (particulate matter with diameter less than  $10 \mu\text{m}$ ) emission rate for 14-week-old tom turkeys ( $19 \text{ g}/500 \text{ kg}$  Animal Unit/day) with a bird weight of  $10.7 \text{ kg}$  from Li et al., (2008).<sup>363</sup> We assumed 50 percent of the suspended particles were contaminated at an HPAI virus titer of  $10^{5.5} \text{ EID}_{50}/\text{g}$ . Cambra-Lopez et al. (2011) suggest that 60 to 70 percent of  $\text{PM}_{10}$  may originate from organic matter from birds, such as feathers or manure.<sup>364</sup> The aerosol source emission rate for  $\text{PM}_{10}$  was calculated as  $10^{4.02} \text{ EID}_{50}/\text{s} = 14,000 (\text{birds}) * 10.7 (\text{kg}/\text{bird}) * 1/500 (\text{AU}/\text{kg}) * 19 (\text{g}/\text{day}/\text{AU}) * 10^{5.5} \text{ EID}_{50}/\text{g} * 1/(24 * 3600) (\text{day}/\text{s}) * 0.5$ . The overall suspended particle emission rate of  $10^{4.62} \text{ EID}_{50}/\text{s}$  was calculated by assuming 25 percent of all particles are  $\text{PM}_{10}$  ( $10^{4.02}/0.25$ ) from broiler house data.<sup>365</sup>

### Particle Size Distribution


We assumed 25 percent of suspended particles in the air from a turkey house are  $\text{PM}_{10}$  and 11 percent of  $\text{PM}_{10}$  particles are  $\text{PM}_{2.5}$ .<sup>353,358,359</sup> We assigned a diameter of  $25 \mu\text{m}$  to particles larger than  $10 \mu\text{m}$ , based on the mean mass diameter estimate from Redwine et al. (2002).<sup>355</sup> Diameters of  $6.25 \mu\text{m}$  and  $1.5 \mu\text{m}$  were assigned to  $\text{PM}_{10}$  and  $\text{PM}_{2.5}$  size fractions, respectively.

### HPAI 50 percent Embryo Infectious Dose

The infectious dose for this scenario is the same as in Scenario A.

## **Scenario B results:**

Scenario B results are shown in **Appendix 5 Figures 3** and **Appendix 5 Table 4**.



**Appendix 5 Figure 3.** AERMOD model-predicted daily probability of infection for a susceptible broiler flock located at various points downwind of a source turkey IP (Scenario B).

**Appendix 5 Table 4** AERMOD model-predicted HPAI virus concentrations and daily exposure probability at various points downwind of a source turkey IP (Scenario B).


Outcome variable	Distance from source (turkey premises)				
	0.5 km	1 km	1.5 km	3 km	5 km
Predicted HPAI concentration (EID <sub>50</sub> /m <sup>3</sup> )	2.82	0.89	0.46	0.16	0.07
Predicted probability of exposure in a day	0.1176	0.0388	0.0205	0.0069	0.0031

**Scenario C:**

This scenario is aimed at showing the impact of the uncertainty in the aerosol infectious dose. In this scenario, the 50 percent HPAI virus infectious dose of  $10^4$  EID<sub>50</sub> was used instead of the  $10^{5.44}$  used in Scenario A. The rest of the inputs were not changed from Scenario A.

**Scenario C results:**

Scenario C results are shown in **Appendix 5 Figures 4** and **Appendix 5 Table 5**. We can see that the AERMOD model-predicted transmission likelihoods are much higher and at longer distances when using a lower infectious dose.



**Appendix 5 Figure 4.** AEROMOD model-predicted daily probability of infection for a susceptible broiler flock located at various points downwind of a source broiler IP (Scenario C)

**Appendix 5 Table 5.** AERMOD model-predicted HPAI virus concentrations and daily exposure probability at various points downwind of a source broiler IP (Scenario C).

Outcome variable	Distance from source (known infected broiler premises)				
	0.5 km	1 km	1.5 km	3 km	5 km
Predicted HPAI concentration (EID <sub>50</sub> /m <sup>3</sup> )	0.55	0.19	0.1	0.04	0.02
Predicted probability of exposure in a day	0.49	0.20	0.12	0.04	0.02



## Appendix 6: Expert Polling on Aerosol Transmission Route

A panel of eight experts in the turkey and broiler industries with field experience managing AI was surveyed between November 2013 and January 2014 on risk of HPAI transmission via multiple routes of infection. Surveys were administered through the online polling service SurveyMonkey.<sup>h</sup> Experts were asked to provide their opinion, based on previous experience, of perceived risk for given scenarios. Qualitative risk rating definitions were provided and match those used in this risk assessment (with the exception that the survey did not include a “very low risk” option) (see section 5, Overview of Data Analysis Approaches, for risk rating definitions). Below is the subset of questions that pertain to spread by aerosol transmission under two scenarios: with and without depopulation activities happening at source farm. Associated expert responses to these questions are shown in **Appendix 6 Tables 1-2** and **Appendix 6 Figures 1-2**.


*Q1: Please qualitatively rate the likelihood of AI transmission from a known infected flock to an uninfected turkey flock located at distances specified in the table. In this scenario, **there are no** depopulation activities happening at source flock. Please complete the following table, selecting a risk rating for each scenario as negligible, low, moderate, high, or extremely high, based on your expert opinion.*

*Q2: Please qualitatively rate the likelihood of AI transmission from a known infected flock to an uninfected turkey flock located at distances specified in the table. In this scenario, **there are** depopulation activities happening at source flock. Please complete the following table, selecting a risk rating for each scenario as negligible, low, moderate, high, or extremely high, based on your expert opinion.*

**Appendix 6 Table 1.** Expert responses (n=8) to the question of likelihood of AI transmission from a known infected flock to an uninfected turkey flock at specified distances when no depopulation activities are happening at source flock (Question 1).

Distance from source flock	Likelihood Rating				
	Negligible	Low	Moderate	High	Extremely high
1 km	0	4	4	0	0
1.5 km	2	5	1	0	0
2 km	7	0	1	0	0
3 km	7	1	0	0	0


<sup>h</sup> Survey Monkey, Inc., Palo Alto, CA, [www.surveymonkey.com](http://www.surveymonkey.com)



**Appendix 6 Figure 1.** Expert responses (n=8) to the question of likelihood of AI transmission from a known infected flock to an uninfected turkey flock at specified distances when no depopulation activities are happening at source flock (Question 1).

**Appendix 6 Table 2.** Expert responses (n=8) to the question of likelihood of AI transmission from a known infected flock to an uninfected turkey flock at specified distances where depopulation activities are happening at source flock (Question 2).

Distance from source flock	Likelihood rating				
	Negligible	Low	Moderate	High	Extremely high
1 km	1	1	2	4	0
1.5 km	1	2	4	1	0
2 km	4	3	1	0	0
3 km	5	3	0	0	0



**Appendix 6 Figure 2.** Expert responses (n=8) to the question of likelihood of AI transmission from a known infected flock to an uninfected turkey flock at specified distances where depopulation activities are happening at source flock (Question 2).

## Appendix 7: Cross-Commodity Pre-Movement Isolation Period (PMIP)

**TO MOVE BIRDS DURING AN HPAI OUTBREAK, PRODUCERS NEED TO AGREE TO A PMIP FOR A SET NUMBER OF DAYS PRIOR TO MOVEMENT. DURING THE PMIP:**

1. **No live or dead poultry will be moved onto or off the premises.**
2. **Only critical operational visits to the premises will continue.**
3. **Manure, litter, and garbage will not be removed from the premises; the producer is responsible for managing the risks associated with any on-site movement that must occur.**
4. **Enhanced biosecurity for people and vehicles; no off-site equipment will be pre-staged.**

**GOAL:** for producers to actively and effectively implement enhanced biosecurity procedures in the critical time period before live poultry is moved, thus reducing the risk of lateral HPAI transmission.

### What is PMIP?

- The Pre-Movement Isolation Period (PMIP) is a critical biosecurity component of the process to obtain a continuity of business permit that involves a defined period of greatly intensified biosecurity for an entire premises prior to permitted movement of live poultry.
- The PMIP is a component of the Secure Poultry Supply Plan, which provides guidelines for poultry premises that seek to move poultry products or live poultry within, into, or out of a regulatory Control Area during an HPAI outbreak.

### What poultry movements require a PMIP?

- The PMIP is required for these live poultry movements:
  - **Terminal movements** (e.g., broilers to processing, turkeys to processing).
    - All out movements, and
    - Movements that do not remove all birds from a premises (e.g., multi-age premises).
  - **Transfer movements** (i.e., live bird movements between farms)
    - All out movements, and
    - Movements that do not remove all birds from a premises.
- These live poultry movements have a moderate to high risk of causing lateral disease transmission if infected but undetected poultry are moved.

- Movement of poultry that are infected but undetected will have subsequent epidemiologic, regulatory, and economic consequences.

### **What poultry movements do not require a PMIP?**

- The PMIP does not apply to day-old chicks or poults (however, a post-move quarantine does apply to these birds).
- The PMIP does not apply to poultry by-products (except as specifically related to live bird movement).
- The PMIP does not apply to eggs or egg products.

### **When does the PMIP start and end?**

- The PMIP starts the specified number of days prior to the scheduled movement date and ends when load-out begins (i.e., the hours or days of load-out are not considered part of the PMIP).
- The load-out period begins when the first crew, vehicle, or equipment arrives on the premises and ends when the last load of birds departs the premises. Pre-staging of equipment during the PMIP is prohibited.

### **How long is the PMIP?**

- The PMIP takes place for a defined number of days immediately prior to the permitted movement of poultry. This period is as follows (determined by some combination of 95% probability of detection for the type of poultry, the type of housing [contact rate], and the characteristics of HPAI viruses [mean death time], as well as by the ultimate consequences of moving an infected, undetected specific type of flock):
  - For 8 days prior to movement of turkeys to processing
  - For 5 days prior to movement of broilers to processing
- Movements of other types of live poultry may not be advised or may require additional post-movement quarantine.

**What critical operational visits to a premises may continue during the PMIP (when specific biosecurity measures are in place)? *Please defer to designated regulatory officials to determine what types of movements require a permit in the Control Area; while these critical operational visits need to continue during the PMIP, a permit may or may not be required.***

1. Feed delivery in a dedicated truck directly from a stand-alone feed mill. Trucks delivering individual feed ingredients that are stored on poultry premises will require a permit.
2. Emergency repair of critical mechanical equipment.
3. Service visits to address changes in bird health.

### **What is the specific biosecurity for these critical operational visits?**

- People who have contact with other poultry must shower and change clothes before entering the premises and also wear necessary protective clothing and footwear as described in appropriate biosecurity protocols.
- All vehicles and equipment will be cleaned and disinfected prior to entering premises.

### **What activities are prohibited during PMIP?**

Activities that pose a risk for lateral transmission of HPAI virus are prohibited. Specific prohibited activities are:

1. Off-farm disposal of mortality is prohibited. Risks associated with dead birds on-site must be managed.
2. Off-farm removal of manure or litter is prohibited. Risks associated with manure or litter movement on-site must be managed.
3. Off-farm garbage disposal is prohibited. Risks associated with garbage storage on-site must be managed.
4. Visiting other poultry farms is prohibited for people who work on poultry farms. People should have contact only with their assigned flock.
5. All non-critical visitors are prohibited from entering farms. All non-critical, routine, or operational visits must be replaced by telephone communication or must be scheduled outside of the PMIP. Non-critical visitors who work with or have contact with another commercial poultry operation (farm, hatchery, processing plant, etc.) or have contact with a noncommercial poultry flock (backyard birds, hobby farms with birds, or game birds) are prohibited from entering farms.
6. Entering a poultry house is prohibited unless the person is wearing clothing dedicated to the farm and footwear dedicated to the house.
7. Noncritical equipment from off-site is prohibited from being moved on-site.
8. Moving live poultry onto or off the premises is prohibited.

### **Why is the PMIP critical?**

- The PMIP biosecurity requirements will minimize the likelihood of a flock being exposed to HPAI close to its scheduled movement date.
- The PMIP will increase the likelihood of detection of an infected flock prior to movement of birds.
- The PMIP will decrease the likelihood of moving infected but undetected poultry, thus reducing the risk of lateral transmission of HPAI virus from specific categories of live poultry.
- The PMIP assists regulatory officials evaluating movement permit requests, since producers seeking permits will actively document, for the record, that they have achieved specific biosecurity requirements.

### **How does the PMIP work?**

- Signs of disease take time to develop following exposure to and infection with HPAI virus.
- If a flock is exposed to HPAI virus close to its scheduled movement date, signs of disease (or significant disease spread and mortality) within the flock may not be evident yet, and thus the probability of virus detection is relatively lower.
- If the flock is exposed to and becomes infected with HPAI virus prior to the implementation of the PMIP, disease signs of HPAI are likely to be detected by the day of scheduled movement (i.e., by the end of the PMIP); thus, movement can be halted and the flock is unlikely to pose a movement-associated risk.

### **What can be done for birds scheduled to move before a PMIP can be completed?**

- When a Control Area is first established or is expanded, some flocks newly in that Control Area may already be scheduled to move before a full PMIP can be completed. Additionally, rapid marketing to reduce the susceptible poultry population in a Control Area may be beneficial for HPAI outbreak control, provided that birds can be moved in a biosecure manner. In such situations, PMIP biosecurity measures should be implemented as soon as possible, and additional information must be provided to the designated regulatory officials to request movement prior to a full PMIP.
- Additional information that must be provided to the designated regulatory officials includes: mortality and morbidity data, test results, destination premises location, farm visitor and farm activity records from the days prior to the Control Area designation, additional equipment requirements, route to be used for transport, defined load-out duration and situation, and defined biosecurity and downtime protocols for load-out crews.

## Appendix 8: Modeling Technical Details

This appendix provides the technical details for the methods applied in estimating the detection probabilities evaluated in section 9.4, Likelihood of Detecting HPAI in an Infected Broiler House. The probability of detection before the start of load-out and the probability of detection prior to movement to processing are estimated from simulation models consisting of a stochastic disease transmission model and active surveillance model. A description of the transmission and surveillance model algorithms can be found in Weaver et al. (2016).<sup>328</sup> The models from Weaver et al. (2016) were reparametrized according to broilers for use in the analyses presented in this risk assessment.<sup>328</sup> The derivation of the broiler-specific parameters is detailed in the section following the introduction.

The probability of detection prior to the start of load-out as estimated from the simulation models is a critical component in estimating the overall likelihood of not detecting HPAI in a flock prior to the start of load-out. The overall likelihood combines the probability of two events: First, the probability a susceptible flock is infected given it is some distance from an infectious premises; and second, the probability the infection is not detected in the flock prior to the start of load-out. As previously mentioned, the second probability is estimated using the simulation models. The first probability, that a susceptible premises a given distance from an infectious premises is itself infected, is estimated using a spatial transmission kernel, which estimates the hazard rate posed by an infectious premises to a susceptible premises at a given distance. The two probabilities are combined into the overall likelihood following a method outlined in Weaver et al. (2016).<sup>328</sup> Two different transmission kernels were used to estimate the overall likelihood: a transmission kernel derived from data on the 2003 HPAI H7N7 outbreak in the Netherlands by Boender et al. (2007) and a transmission kernel estimated from data on the 2015 HPAI H5N2 outbreak in Minnesota.<sup>133</sup> Details on the estimation of the Minnesota transmission kernel are given following an explanation of the estimation of the broiler-specific disease transmission model parameters used in the simulation.

### ***Estimation of Transmission Model Parameters***

#### ***Adequate Contact Rate***

The distribution for the contact rate was chosen based on a literature review of estimates from experimental and outbreak data for chickens. Contact rates estimated from outbreak data were given greater weight, particularly estimates derived by Bos et al. (2009) and in a white paper on contact rate estimation.<sup>366,367</sup> Both studies used mortality data from flocks infected with HPAI H7N7 during the 2003 outbreak in the Netherlands. Considerable uncertainty exists in the parameter estimates: (1) Contact rate estimates are highly sensitive to assumptions about the length of the latent period, and (2) estimates based on experimental data were generally lower.

In order to capture this uncertainty, a PERT distribution was chosen under the assumption that extreme contact rate values are less likely. The parameters of the distribution are a minimum of 2.50, mode of 4.77, maximum of 9.00, and  $\lambda$  equal to 4.00. Because of the uncertainty in the parameter estimates, a second scenario was evaluated assuming a slow contact rate, which is more representative of the rates observed in experimental settings. The distribution used in this scenario is uniform with a minimum of 1.0 and maximum of 5.0.



### Latent and Infectious Period Distributions

Due to a scarcity of adequate experimental data for Pennsylvania HPAI H5N2, the latent period distribution for this strain was estimated from a mixture of Pennsylvania HPAI H5N2 and HPAI H5N1 (multiple clades) experiments using a maximum likelihood method. HPAI H5N1 data from Bouma et al. (2009),<sup>123</sup> Das et al. (2007),<sup>63</sup> and Poetri et al. (2011),<sup>327</sup> and Pennsylvania HPAI H5N2 data from Spackman et al. (2014)<sup>368</sup> and van der Goot et al. (2003)<sup>121</sup> were used in estimating the latent period for the Pennsylvania HPAI H5N2 strain. Except for Das et al. (2007), the experiments tested for HPAI in each inoculated bird at regular time intervals (e.g., daily or half-day intervals).<sup>63</sup> The latent period is assumed to begin once the bird has been inoculated and to end sometime between the last negative test and first positive test for that particular bird. Let  $t_a$  be the time of the last negative test and let  $t_b$  be the time of first positive test, so the transition from the latent to the infectious period occurs in  $(t_a, t_b]$ . The probability of observing the transition in this time period is given by  $F(t_b) - F(t_a)$ , where  $F$  is the distribution of the latent period, here assumed to be gamma distributed.

In Das et al. (2007), pools of inoculated birds were periodically sampled without replacement and tested for HPAI. Let  $t_c$  be the sampling time. The probability that the transition from the latent to the infectious period occurred prior to  $t_c$  in birds for whom the test is positive is  $F(t_c)$ , while the probability the transition occurs after  $t_c$  in birds testing negative is  $1 - F(t_c)$ . Parameters for the gamma distribution were estimated by maximizing the cumulative likelihood of the observed transition from the latent to the infectious period in each inoculated chicken in each of the cited experiments. The likelihood was maximized using the “nlminb” algorithm, a bounds-constrained quasi-Newton method in R’s “optimx” function.<sup>369-371</sup> The shape parameter was estimated to be 0.82 and the scale parameter was estimated to be 0.44.

A similar method was used to estimate an infectious period distribution, assumed to be Weibull distributed. The estimated shape parameter is 1.965 and estimated scale parameter is 4.238, on the basis of Pennsylvania HPAI H5N2 data from van der Goot et al. (2003) and Swayne et al. (2012).<sup>65,166</sup> For more details, see An Assessment of the Risk Associated with the Movement of Turkey Hatching Eggs into, within, and out of a Control Area during a Highly Pathogenic Avian Influenza Outbreak.<sup>119</sup>

As of this writing, limited data are available on the EA/AM HPAI H5N2 strain due to the relative recency of the outbreak. In an experiment performed at the USDA Southeast Poultry Research Laboratory (SEPR), two mean times to death were identified in chickens, 3.2 days observed in 8-week-old broilers and adult breeders, and 4.8 days observed in 5-week-old broilers (D. Swayne, personal communication, Oct 2016). However, bird-specific inoculation data, as were used in estimating the latent and infectious period distribution for the Pennsylvania HPAI H5N2 strain, are not available. Since only the mean time to death is known, distributions from other strains were used to approximate EA/AM HPAI H5N2. The latent period distribution used to model EA/AM HPAI H5N2 for both mean times to death was estimated from HPAI H5N1 data available in Bouma et al. (2009), Das et al. (2007), Poetri et al. (2011), Spekrijse et al. (2011a), and Spekrijse et al. (2011b) by the same method used to estimate the Pennsylvania HPAI H5N2 latent period distribution.<sup>123,327 106,170</sup> The latent period distribution, assumed to be gamma distributed, has an estimated shape parameter of 0.89 and scale parameter of 1.18.

The infectious period distribution used differs depending on the mean time to death of the EA/AM HPAI H5N2 strain. To model the variability in the infectious period, the scale parameter

of the Weibull distribution estimated for the Pennsylvania HPAI H5N2 infectious period was adjusted until the sum of the expected values of the latent and infectious period distributions matched the desired mean time to death. The shape parameter was kept the same. The resulting parameters of the Weibull infectious period distributions are a shape of 1.965 and scale of 2.90 for the EA/AM HPAI H5N2 strain with mean time to death of 3.2 days, and a shape of 1.965 and scale of 4.70 for the EA/AM HPAI H5N2 strain with mean time to death of 4.8 days.


Number of Broilers per House

The flock size is randomly generated from a lognormal distribution truncated at 13,000 and 50,000, estimated from industry data. The mean flock size is 24,257 birds with a standard deviation of 9,787 birds.

**Estimation of Active Surveillance Model Parameters**

Daily Mortality

Daily normal mortality data for 79 broiler houses and weekly mortality data for an additional 146 houses was provided by industry representatives. Normal mortality is simulated by first selecting 14 consecutive days of daily mortality from one of the 79 broiler houses. The end point for the 14 days of daily mortality is randomly selected to be 1, 2, or 3 days prior to the movement day. The daily mortality is then rescaled by a random number generated from a lognormal distribution with a mean of -5.4850 and standard deviation of 0.7551 estimated from the weekly mortality data, thereby introducing variability into the daily mortality. A histogram of the simulated normal mortality is given in **Appendix 8 Figure 1**. The simulated normal mortality has a mean of 0.8 birds per 1,000 with a 5<sup>th</sup> percentile of 0.10 and 95<sup>th</sup> percentile of 2.41. Disease mortality is selected from the disease transmission model output.



**Appendix 8 Figure 1.** Histogram of simulated daily mortality in a broiler house in the last two weeks prior to movement

Diagnostic Test Sensitivity


The sensitivity of the rRT-PCR test is estimated to be 86.5 percent, meaning there is a 13.5 percent chance the infection will not be detected even when the pooled sample contains an

HPAI-positive swab.<sup>372</sup> AI experts noted this sensitivity estimate is conservative considering recent enhancements to test protocols.<sup>119</sup>

Supplementary antigen capture immune assays using lateral flow devices are utilized in additional testing protocols evaluated in Appendix 10: Supplementary Testing Protocols. These tests require high virus concentrations to detect AI virus (detection limit is between  $10^4$  and  $10^6$  EID<sub>50</sub>).<sup>373-375</sup> The diagnostic sensitivity of these tests therefore depends on the clinical status of the infectious birds, which impacts the level of virus shedding.

A study performed at the USDA SEPRL was undertaken to provide data on AC (antigen capture) test performance in dead birds infected with HPAI viruses. AC test sensitivity was estimated for two strains separately using a Bayesian approach from swabs taken from 14 and 46 dead chickens following exposure to HPAI H7N3 Jalisco and Pennsylvania HPAI H5N2, respectively. In addition, the AC test sensitivity was estimated for HPAI H5N1 (several clades) from a literature review. The resulting posterior distributions are given in **Appendix 8 Figure 2**. The estimated means and 95% credibility intervals for the AC test sensitivities are 57% (33-80%) for the HPAI H7N3 Jalisco strain, 86% (80-91%) for the HPAI H5N1 strain, and 97.9% (92-99.9%) for the HPAI H5N2 strain.<sup>368</sup> The wider credibility interval in the case of HPAI H7N3 is due to the smaller sample size and correspondingly greater uncertainty.

The estimated AC test sensitivities suggest that there is considerable between-strain variation, which is likely due to the variation in virus shedding between different strains, which affects detection because of the low analytic sensitivity of the AC test. AC test sensitivities for LPAI as identified through a literature review were generally lower than the estimates for HPAI, with an average of about 50%. Given the uncertainty and variance surrounding the estimates for AC test sensitivity, a conservative estimate of 50% is chosen for this analysis.



**Appendix 8 Figure 2.** Statistical distributions for the diagnostic sensitivity of antigen capture immunoassays for different HPAI strains.

## **Estimation of the 2015 HPAI H5N2 Minnesota Outbreak Spatial Transmission Kernel**

### Spatial Transmission Kernel Model Comparison

Four candidate parameterizations of the spatial transmission kernel were assessed for best fit based on the 2015 HPAI H5N2 Minnesota outbreak data using Akaike's Information Criterion (AIC).<sup>376</sup> The likelihood function used in the evaluation of AIC takes the form given in Boender et al. (2007).<sup>133</sup> The resulting AIC values from the comparison of the four spatial transmission kernel parameterizations were all within 2 of each other, meaning none of the parameterizations is a definitive best fit. The same parametrization as was used in Boender et al. (2007) was chosen for use in the Minnesota outbreak analysis in order to make the two spatial transmission kernels be more comparable.<sup>133</sup> The spatial transmission kernel is given below as a function of distance between susceptible premises  $i$  and infectious premises  $j$ :

$$h(d_{ij}) = \frac{h_0}{1 + \left(\frac{d_{ij}}{r_0}\right)^\alpha}$$

$h_0$ ,  $r_0$ , and  $\alpha$  are constants to be estimated from outbreak data, where  $h_0$  is the maximum daily hazard rate (occurring when the inter-premises distance is zero), and  $r_0$  and  $\alpha$  determine the decline in the hazard rate as inter-premises distance increases from zero.

Due to phylogenetic evidence of primary introductions occurring concurrently with lateral spread, an additional parameter was added to the force-of-infection equation from Boender et al. (2007).<sup>38,133</sup> The force of infection describes the overall hazard faced by susceptible premises  $i$  at time  $t$ , and in Boender et al. (2007) it depends solely on the number of infectious premises.<sup>133</sup> The force of infection from Boender et al. (2007) is given below as a function of  $t$ :

$$\lambda_i(t) = \sum_{i \neq j} h(d_{ij}) 1\{j \text{ is infectious}\}$$

This equation is modified for use in the Minnesota outbreak through the addition of a parameter,  $k$ , allowing for infection to occur independently of the number of infectious premises:

$$\lambda_i(t) = \left(\sum_{i \neq j} h(d_{ij}) 1\{j \text{ is infectious}\}\right) + k$$

Note that  $k$  is constant and distance-independent in addition to not being reliant on the number of infectious premises at time  $t$ . Therefore,  $k$  largely expresses the risk posed by distance-independent environmental factors such as wild birds. However, as some of the risk from long-distance movements of people and equipment may be captured by  $k$ , a third force-of-infection equation was evaluated with a constant, distance-independent parameter,  $\delta$ , that varies with the number of infectious premises:

$$\lambda_i(t) = \left(\sum_{i \neq j} (h(d_{ij}) + \delta) 1\{j \text{ is infectious}\}\right) + k$$

Infection risk related to distance-independent movements of people and equipment and distance-independent environmental factors is therefore partitioned between  $\delta$  and  $k$ , respectively. The three different force-of-infection equations were compared using AIC. Adding  $k$  to the force of infection significantly improved the model fit based on the resulting AIC values. The addition of

$\delta$ , on the other hand, resulted in a larger AIC (AIC = 1393.393 with  $k$ , compared with 1395.332), which suggests that the inclusion of  $\delta$  is trivial, and that long-distance movements of people and equipment provide only marginal contributions to the risk represented by  $k$ . Thus, the force of infection with  $k$  only was chosen for use in the analysis of the Minnesota outbreak.

Estimation of the Spatial Transmission Kernel Parameters

The four parameters,  $h_0$ ,  $r_0$ ,  $\alpha$ , and  $k$ , were estimated following the maximum likelihood method approach described in Boender *et al.* (2007).<sup>133</sup> The method depends only on inter-premises distance and premises-level infection status. As the exact days on which the infectious period of a case premises started and ended are unknown, a number of simplifying assumptions must be made. For the Minnesota outbreak, case premises are assumed to be infected eight days prior to the detection date. The infectious period is assumed to begin three days later, five days prior to the detection date. The infectious period lasts up to and including the day on which disposal of the depopulated poultry carcasses begins. The mean parameter estimates and 95% confidence intervals under these assumptions regarding infection status are given in **Appendix 8 Table 1**, along with the parameters estimated from the Netherlands outbreak by Boender *et al.* (2007).<sup>133</sup> Infection risk in Minnesota—with its higher mean hazard rate, significantly higher  $r_0$ , which suggests higher infection risk persisted over significantly longer distances, and additional parameter  $k$  representing distance-independent environmental risk factors—would be expected to be considerably higher and less responsive to changes in distance, thereby posing an overall greater threat of HPAI spread.

**Appendix 8 Table 1.** Mean estimates and 95% confidence intervals of spatial transmission kernel model parameters estimated from HPAI outbreaks in Minnesota and the Netherlands.

Description	$h_0$	$r_0$	$\alpha$	$k(10^{-4})$
Minnesota 2015 HPAI H5N2: Case premises are infected 8 days prior to detection; infectious period starts 5 days prior to detection and lasts up to and including compost start date.	0.0061 (0.0025, 0.0137)	7.02 (3.07, 16.16)	2.46 (1.80, 4.38)	3.2 (1.6, 5.2)
Netherlands 2003 HPAI H7N7: Case premises are infected 6 days prior to the first rise in mortality; infectious period starts 2 days later and lasts until depopulation. Estimates from Boender <i>et al.</i> (2007). <sup>133</sup>	0.0020 (0.0012, 0.0039)	1.9 (1.1, 2.9)	2.1 (1.8, 2.4)	NA

Estimation of the Probability of Infection

The spatial transmission kernel is used to estimate the probability that a susceptible premises becomes infected given it is some distance from an infectious premises through the force of

infection. The probability that a susceptible premises  $i$  becomes infected on day  $t$ ,  $q_i(t)$ , is given below:

$$q_i(t) = 1 - e^{-\lambda_i(t)}$$

The mean parameter estimates from the Minnesota and Netherlands outbreaks are used to estimate the probability of infection applied in the estimation of the overall probability. Note that as the force of infection increases, the probability of infection increases. As the mean hazard rate is uniformly higher over distance, and with the addition of  $k$ , the force of infection is higher under the Minnesota outbreak model. Thus, the Minnesota outbreak transmission kernel poses a higher probability of infection, which then results in a higher estimated overall probability of not detecting HPAI in a flock prior to the start of load-out.

## Appendix 9: Poultry Industry Survey on Garbage Management Practices

A convenience sample of veterinarians and other managers in the turkey (n=15), broiler (n=8), and layer (n=40) industries was surveyed between June and August 2016 on standard practices for garbage management on farms that they manage or supervise. Surveys were administered by the University of Minnesota HPAI Team using the online polling service Qualtrics.<sup>i</sup>


A convenience sample of participants with significant experience in the poultry industry was solicited; however, this survey was limited by small sample size. No additional analyses were conducted for these data beyond descriptive statistics. Still, the results are informative for the purpose of the risk assessment, and serve to illustrate the variations in industry practice and potential differences between poultry sectors that may operate in the same geographic area. As such, readers should note that absence of an affirmative response to a high-risk activity does not definitively indicate it is not occurring. The results of the survey are shown in **Appendix 9 Figures 1- 19**. Of note, some survey questions and answer choices were modified to better match the industry to which the survey was distributed. Additionally, some participants declined to answer all questions within the survey.

---

<sup>i</sup> Qualtrics© 2015 Provo, UT, USA. <http://www.qualtrics.com>

**Types of Items Disposed in the Garbage on Poultry Premises**


Appendix 9 Figures 1-3 show items disposed in garbage which may be potentially infectious or contaminated with HPAI in the event of an outbreak. Respondents answered the question: Which of the following are disposed of in trash/garbage on your premises (check all that apply)?<sup>j</sup>




**Appendix 9 Figure 1.** Responses of broiler industry representatives (n=8) to types of items disposed in the garbage on broiler premises. Respondents in the broiler industry wrote in additional items such as boxes, buckets, jugs from disinfectants, litter treatments, disposable chick feeder lids, cans, and bottles.

<sup>j</sup> Item abbreviated for graphic display (\*). Full text prompt as used in survey: “Used needles/syringes/diagnostic supplies that have contacted birds;” “Household garbage from farm manager residence or any other residence”






**Appendix 9 Figure 2.** Responses of layer industry representatives (n=39) to types of items disposed in the garbage on layer premises.




**Appendix 9. Figure 3.** Responses of turkey industry representatives (n=15) to types of items disposed in the garbage on turkey premises. Respondents in the turkey industry wrote in additional items such as trash from farm manager residence, empty medication containers, poult box papers, and supply containers.

**Disposal of Potentially HPAI Infectious or Contaminated Items in the Garbage on Poultry Premises**


Appendix 9 Figures 4-7 show percentages of poultry industry respondents surveyed that reported disposing in the garbage potentially high-risk items, such as wild bird carcasses, poultry carcasses, and other items that may act as fomites.



**Appendix 9 Figure 4.** Responses of poultry industry representatives to disposal of wild bird carcasses in the garbage (layer sector: n=40; broiler sector: n=8; turkey sector: n=15). The answer choice “not sure” was not available in the survey of layer industry representatives.




**Appendix 9 Figure 5.** Responses of poultry industry representatives regarding the frequency of disposing poultry carcasses in the garbage (layer sector: n=39; broiler sector: n=8; turkey sector: n=15). The carcass type (layer/pullet, broiler, or turkey) cited in the question matched the industry sector to which the survey was distributed.



**Appendix 9 Figure 6.** Responses of poultry industry representatives regarding disposal of poultry carcasses in the garbage in the event of an outbreak (layer sector: n=39; broiler sector:


n=8; turkey sector: n=15). The carcass type (layer/pullet, broiler, or turkey) cited in the question matched the industry sector to which the survey was distributed.




**Appendix 9 Figure 7.** Responses of poultry industry representatives regarding disposal of other items that may act as fomites (layer sector: n=39; broiler sector: n=8; turkey sector: n=15). The answer choice “maybe” was not available in the survey of layer industry representatives.

**Transportation of Garbage to Landfill**


Industry representatives indicated that most commercial poultry operations utilize an off-site landfill for garbage disposal (BWG, TWG, personal communication, May 2016). **Appendix 9 Figures 8–10** show survey results related to garbage transportation to landfill, such as use of a contracted service and the garbage truck route.



**Appendix 9 Figure 8.** Responses of poultry industry representatives regarding methods of transporting garbage to landfill (layer sector: n=39; broiler sector: n=7; turkey sector: n=15). Respondents wrote in that some premises may use a combination of grower transport and contracted hauling services.



**Appendix 9 Figure 9.** Responses of poultry industry representatives regarding whether their garbage hauler collects waste from multiple poultry premises (layer sector: n=36; broiler sector: n=7; turkey sector: n=15).




**Appendix 9 Figure 10.** Responses of broiler and turkey industry representatives concerning their awareness of other farms on the garbage truck route (broiler sector: n=7; turkey sector: n=15). This question was not asked of layer industry representatives.


**Location of Garbage Collection Area on Commercial Poultry Premises**

Premises often have dumpsters or a designated location where trash is collected for transportation to a landfill. **Appendix 9 Figures 11–15** show survey results related to location of the garbage collection area on the premises relative to other features on the farm, such as poultry barns, other premises, and rendering collection point.


Note that in **Appendix 9 Figure 14**, the option “no rendering used” was available to respondents in the layer sector. The low number of responses from the broiler and turkey sectors may suggest that some individuals declined to answer this question because it was not applicable to their premises. An additional question about biosecurity practices at the dumpster site was posed only to layer industry representatives (**Appendix 9 Figure 15**).



**Appendix 9 Figure 11.** Responses of broiler and turkey industry representatives regarding the location of the dumpster or trash collection point on their premises (broiler sector: n=7; turkey sector: n=15).




**Appendix 9 Figure 12.** Responses of layer industry representatives (n=39) regarding the location of the dumpster or trash collection point on the premises.



**Appendix 9 Figure 13.** Responses of poultry industry representatives regarding the distance of the dumpster or trash collection point from the nearest poultry barn (layer sector: n=36; broiler sector: n=7; turkey sector: n=14). In the survey of layer industry representatives, it was specified that the nearest poultry barn may be on the same premises or neighboring premises.






**Appendix 9 Figure 14.** Responses of poultry industry representatives regarding the distance of garbage collection from the rendering area (layer sector: n=36; broiler sector: n=3; turkey sector: n=7). A majority (75%) of layer industry respondents indicated no rendering is used (not shown).



**Appendix 9 Figure 15.** Responses of layer industry representatives (n=34) concerning maintenance of the trash collection equipment/dumpster area.

**Feasibility of Suspending Garbage Collection during PMIP**


Premises that wish to move product out of a Control Area during an HPAI outbreak likely will need to observe a Pre-Movement Isolation Period (PMIP), during which no non-critical operations (including off-site disposal of poultry mortality and garbage) are allowed. **Appendix 9 Figure 16** shows survey results related to the feasibility of suspending garbage collection during PMIP. The duration of PMIP may vary by industry sector and type of movement requested. In addition to the results shown, respondents indicated their answers may depend on the size of the farm and the duration of PMIP. For suspension of garbage services to be feasible, some respondents noted they would need more on-site trash storage.




**Appendix 9 Figure 16.** Responses of poultry industry representatives concerning the feasibility of suspending garbage collection on the farm during PMIP (layer sector: n=36; broiler sector: n=7; turkey sector: n=15).

**Frequency of Garbage Pickup and Responsibility of Contracting Garbage Service**


The person responsible for contracting third-party garbage hauling services and the frequency of garbage collection on a poultry premises may vary by farm size and type of operation and are not uniform across the poultry industry. **Appendix 9 Figures 17-19** show survey results related to frequency of garbage pickup and responsibility of contracting garbage service.



**Appendix 9 Figure 17.** Responses of poultry industry representatives regarding the frequency of garbage pickup (layer sector: n=36; broiler sector: n=7; turkey sector: n=15). Turkey industry respondents wrote in that frequency of garbage pickup may depend on the size of the farm and on some premises it is not pre-scheduled but occurs as needed.



**Appendix 9 Figure 18.** Responses of broiler and turkey industry representatives concerning the responsibility for contracting garbage pickup/disposal service (broiler sector: n=7; turkey sector: n=15).



**Appendix 9 Figure 19.** Responses of layer industry representatives regarding the responsibility of contracting garbage pickup/disposal service (layer sector: n=36). Answer choices presented in the layer sector survey were modified from those presented to broiler and turkey representatives to better align with industry practices.

## Appendix 10: Supplementary Testing Protocols

### ***Likelihood of Detecting HPAI in a Broiler Flock Prior to Movement: Protocols for Premises with rRT-PCR Test Result Turnarounds Greater than 12 Hours***

The protocol evaluated in the main risk assessment document involving samples of 11 swabs taken for rRT-PCR the day of and day before the start of load-out assumes the turnaround time needed to obtain results from NAHLN labs is less than 12 hours. As a same-day turnaround is not always feasible, an additional protocol is evaluated in **Appendix 10 Table 1** consisting of one sample of 11 swabs taken for rRT-PCR testing one and two days prior to the start of load-out to allow more time for sending and receiving the test results. Earlier sampling for rRT-PCR generally results in lower detection probabilities because the infection has less time to spread through the flock. Supplementary AC testing performed immediately prior to load-out can be employed to offset the loss in detection probability from earlier rRT-PCR sampling.

To estimate the effect of AC testing on the detection probability, a third protocol was evaluated consisting of one sample of 11 swabs taken for rRT-PCR one and two days prior to movement with the addition of two samples of five swabs each taken for AC testing at the same time immediately prior to the start of load-out. The detection probabilities, and mean with the 5<sup>th</sup> and 95<sup>th</sup> percentile of the number of infectious birds present in an undetected flock at the time of movement, are given in **Appendix 10 Table 1** under the assumption that exposure occurred between 5 and 10 days prior to movement due to a 100% effective five-day PMIP. The estimates are obtained from 6,000 iterations of the simulation model.

As expected, the protocol with the earlier rRT-PCR sampling times is estimated to have lower detection probabilities than the baseline protocol of one rRT-PCR sample of 11 swabs taken the day of and day before movement. Furthermore, the mean number of infectious and undetected birds is lower under the baseline protocol, making it clearly preferable to the early rRT-PCR sampling protocol. The results show that supplementary AC testing in addition to the rRT-PCR samples taken one and two days prior to movement can counteract some of the drawbacks of earlier testing, because detection probabilities are at least as high as under the baseline active surveillance protocol in each strain and contact rate scenario. Since the mean number of infectious and undetected birds is generally lower under the baseline active surveillance protocol, the baseline remains the optimal testing strategy. However, the earlier rRT-PCR sampling with supplementary AC testing represents a highly effective approach when the turnaround time for rRT-PCR results exceeds 12 hours.

**Appendix 10 Table 1.** Likelihood of AI detection and mean number of infectious undetected birds for three different active surveillance protocols and four scenarios varying by the strain or contact rate. A 100% effective 5-day PMIP is assumed to have been implemented.

Strain/contact rate scenario	Active surveillance protocol <sup>a</sup>		
	One sample rRT-PCR taken day of and day before movement	One sample rRT-PCR taken one and two days before movement	One sample rRT-PCR taken one and two days before movement with supplementary AC testing <sup>b</sup>
	Predicted detection probability <sup>c</sup> Mean number of infected undetected birds (5th, 95th percentile)		
PA HPAI H5N2	0.99 176 (0, 1,157)	0.99 1,075 (0, 3478)	0.99 587 (0, 2,748)
EA/AM HPAI H5N2: MTD 3.2	0.99 79 (0, 383)	0.97 224 (0, 1,108)	0.99 81 (0, 368)
EA/AM HPAI H5N2: MTD 4.8; baseline contact rate	0.98 266 (0, 1,260)	0.96 616 (0, 2,536)	0.98 487 (0, 2,386)
EA/AM HPAI H5N2: MTD 4.8; slow contact rate	0.90 51 (0, 199)	0.82 82 (0, 334)	0.90 47 (0, 212)

<sup>a</sup> Samples taken for rRT-PCR testing consist of 11 swabs.

<sup>b</sup> The supplementary AC testing consists of two pools each with five swabs taken at the same time immediately prior to the start of load-out.

<sup>c</sup> Probabilities are estimated from 6,000 simulation iterations.

***Likelihood of Moving Infectious and Undetected Broilers Following Exposure during Load-out: Evaluating the Effect of Supplementary AC Testing***

Exposures occurring during the load-out process are difficult to detect as they typically occur close to the time of movement. Supplementary AC testing can be used to improve the likelihood of detection since the samples can be taken immediately prior to movement, allowing greater time for HPAI to move through the flock. **Appendix 10 Table 2** compares the detection probabilities for the baseline protocol of daily rRT-PCR testing of 11 swabs with a protocol consisting of two samples of five swabs taken for AC testing immediately prior to movement in addition to the daily rRT-PCR sampling. The probabilities are estimated for a single broiler flock from 6,000 simulation iterations using the Pennsylvania HPAI H5N2 strain and considering different numbers of days post-exposure to HPAI and levels of contamination on the load-out crew and equipment. As detection probabilities are already generally quite high under the

baseline testing option, the addition of supplementary AC testing only marginally improves the probability of detection in most cases. However, considerable gains in detection are estimated 2 to 3 days post-exposure in the case of one bird initially infected and 2 days post-exposure when 10 birds are initially infected. As these are the days with the highest risk of releasing large numbers of infectious but undetected birds (see Appendix 10 Table 3), the benefits of the higher detection probabilities due to supplementary AC testing are immense. Supplementary AC testing would furthermore be expected to reduce the mean number of infectious but undetected birds in all cases. Thus, the baseline protocol with the addition of AC testing represents the optimal strategy for minimizing the risk related to exposures during load-out.

**Appendix 10 Table 2.** The likelihood of AI detection for 2 to 6 days following exposure.

Initial number of birds infected <sup>a</sup>	Protocol <sup>b</sup>	Days post-exposure				
		2	3	4	5	6
		Predicted detection probability <sup>c</sup>				
1	rRT-PCR testing only	0.27	0.73	0.95	0.99	1.00
	rRT-PCR with supplementary AC testing <sup>d</sup>	0.51	0.88	0.98	1.00	1.00
10	rRT-PCR testing only	0.84	0.99	1.00	1.00	1.00
	rRT-PCR with supplementary AC testing	0.96	1.00	1.00	1.00	1.00
100	rRT-PCR testing only	0.99	1.00	1.00	1.00	1.00
	rRT-PCR with supplementary AC testing	1.00	1.00	1.00	1.00	1.00

<sup>a</sup> The initial number of infected birds is meant to represent the level of contamination on the load-out personnel and equipment

<sup>b</sup> The rRT-PCR testing consists of daily samples of 11 swabs.

<sup>c</sup> Probabilities estimated from 6,000 simulation iterations using the Pennsylvania HPAI H5N2 strain with a baseline contact rate.

<sup>d</sup> The AC testing consists of two pooled samples of five swabs taken immediately prior to movement.

***Likelihood of at least 100, 500, or 1,000 Infectious but Undetected Broilers in a Flock at the Time of Movement following Exposure during Load-out***

The likelihood of at least 300 infectious but undetected broilers in a flock at the time of movement following exposure to HPAI during load-out was used in Section 9.4.3 (Likelihood of Moving Infectious but Undetected Broilers Following Exposure during Load-out) to demonstrate that despite lower probabilities of detection, the likelihood of releasing large numbers of infectious but undetected birds is predicted to be low. Similar probabilities are provided in **Appendix 10 Tables 3-5** for the cases in which at least 100, 500, and 1,000 infectious but undetected broilers are present in a flock at the time of movement. **Appendix 10 Table 5** provides evidence that the likelihood of releasing 1,000 or more infectious broilers is negligible. Similarly, **Appendix 10 Table 4** suggests the likelihood of releasing 500 or more infectious broilers would be quite low. The likelihood of moving at least 100 infectious but undetected broilers to processing, on the other hand, is nontrivially high in a couple of the cases given in **Appendix 10 Table 3**, specifically the likelihood 3 days post-exposure with one bird initially infected and the likelihood 2 days post-exposure with 10 birds initially infected. The results in

**Appendix 10 Table 3** reiterate the possible risk related to exposure during load-out and the importance of practicing the suggested biosecurity measures during this period. Supplementary AC testing can also be utilized to reduce the likelihood of moving at least 100 infectious but undetected birds to processing.

**Appendix 10 Table 3.** The estimated percent probability of not detecting HPAI in a flock following exposure during load-out where the number of infectious but undetected broilers at the time of movement exceeds 100 birds.\*

Initial no. of birds infected	Days post-exposure				
	2	3	4	5	6
	<b>Predicted percent probability of at least 100 infected but undetected broilers at the time of movement</b>				
1	5.87%	12.63%	3.58%	0.52%	0.03%
10	15.63%	1.00%	0.00%	0.00%	0.00%
100	0.55%	0.00%	0.00%	0.00%	0.00%

\*The initial number of birds infected represents the level of contamination present on the load-out equipment crew and equipment. Percent probabilities are estimated from 6,000 simulation using the Pennsylvania HPAI H5N2 strain and an active surveillance protocol of one sample of 11 swabs taken daily for rRT-PCR testing

**Appendix 10 Table 4.** The estimated percent probability of not detecting HPAI in a flock following exposure during load-out where the number of infectious but undetected broilers at the time of movement exceeds 500 birds.

Initial no. of birds infected	Days post-exposure				
	2	3	4	5	6
	<b>Predicted percent probability of at least 500 infected but undetected broilers at the time of movement</b>				
1	0.02%	2.45%	1.35%	0.25%	0.00%
10	4.90%	0.97%	0.00%	0.00%	0.00%
100	0.55%	0.00%	0.00%	0.00%	0.00%

\*The initial number of birds infected represents the level of contamination present on the load-out equipment crew and equipment. Percent probabilities are estimated from 6,000 simulation using the Pennsylvania HPAI H5N2 strain and an active surveillance protocol of one sample of 11 swabs taken daily for rRT-PCR testing



**Appendix 10 Table 5.** The estimated percent probability of not detecting HPAI in a flock following exposure during load-out where the number of infectious but undetected broilers at the time of movement exceeds 1,000 birds.

Initial no. of birds infected	Days post-exposure				
	2	3	4	5	6
	<b>Predicted percent probability of at least 1,000 infected but undetected broilers at the time of movement</b>				
1	0.00%	0.85%	0.5%	0.12%	0.00%
10	0.90%	0.62%	0.00%	0.00%	0.00%
100	0.55%	0.00%	0.00%	0.00%	0.00%

\*The initial number of birds infected represents the level of contamination present on the load-out equipment crew and equipment. Percent probabilities are estimated from 6,000 simulation using the Pennsylvania HPAI H5N2 strain and an active surveillance protocol of one sample of 11 swabs taken daily for rRT-PCR testing

## Appendix 11: Load-out Mitigation Measures

### **Background and Rationale**


According to the Broiler Working Group, under normal (non-outbreak) conditions, the cumulative length of time required to complete load-out, transport, and any yard (wait) time before slaughter of broilers is  $\leq 12$  hours per barn. This timeline is optimized and likely cannot be shortened during an outbreak situation.

During an outbreak, according to the working group, the total load-out, transport, and yard time will be  $\leq 12$  hours per barn (there is existing motivation for this time to always be less than 12 hours). For a commercial broiler premises practicing an “all-in, all-out” strategy, it is estimated that during an outbreak the cumulative time from beginning of load-out until the last load of birds from a premises is slaughtered (i.e., the total load-out, transport, and yard time) would be  $\leq 24$  hours for at least 80% of farms and would be  $\leq 48$  hours for 90% of farms.

Due to the increased risk associated with extended load-out durations (demonstrated by simulation modeling), additional biosecurity measures are required for longer load-out, transit, and yard time at the premises level. Additional biosecurity and mitigation measures are summarized by premises-wide load-out duration in the following description and in **Appendix 11 Table 1**:

1. For premises with load-out, transport, and yard time  $\leq 24$  hours (estimated to be at least 80% of premises in an outbreak), existing SBS biosecurity measures are adequate. These are:
  - The load-out crew should shower and wear freshly laundered clothing before going on to other activities after load-out.
  - The route for live-haul vehicles and equipment should be selected in consultation with a poultry veterinarian or production manager to minimize contact with and proximity to live poultry and poultry products.
  - When moving equipment from the Infected Zone to the Buffer Zone, C&D of live-haul equipment (power wash) with at least 24-hour downtime is required.
2. For premises with load-out, transport, and yard time of 25 hours to 48 hours (~10% of premises), biosecurity measures in addition to those listed above are required, as follows:
  - Load-out equipment and crews:
    - Must have been adequately cleaned and disinfected using one of the following methods:
      - Sanitization by a disinfectant approved to inactivate HPAI virus
      - Adequate downtime before use
      - Heat treatment to inactivate HPAI virus
    - OR
    - Must have been used only on Monitored Premises during an outbreak. (Equipment and crews that have moved birds of unknown status or positive birds cannot be used.)
  - In all cases, additional barn-to-barn biosecurity must be implemented:

- Pre-staging of equipment in a barn prior to beginning load-out in that barn is not allowed.
  - Other than load-out crew, all persons entering a barn must use *barn-specific* footwear and *farm-specific* clothing.
  - Handwashing is required after working with any barn or birds currently or already loaded out and before entering any other occupied barns on the premises.
3. For premises with load-out, transport, and yard time >48 hours (~10% of premises), biosecurity measures in addition to those listed above (both 1 and 2) are required, as follows:
- Load-out equipment and crews:
    - Must have been adequately cleaned and disinfected using one of the following methods:
      - Sanitization by a disinfectant approved to inactivate HPAI
      - Adequate downtime before use
      - Heat treatment to inactivate HPAI
    - OR
    - Must have been used only on specific Monitored Premises during an outbreak as outlined below:
      - Used exclusively to load out premises where PCR testing occurred within 24 hours of movement AND entire premises was depopulated in under 24 hours.
      - Have not moved birds of unknown or positive status.
  - In all cases, load-out crew must shower and put on clean clothes before entering the premises to be loaded out.
  - In all cases, for all barns loading out after initial 48 hours on a premises, daily testing is required.
    - Negative results of tests taken within 24 hours of scheduled movement (at barn level) must be documented before birds depart premises. (This is diagrammed in **Appendix 11 Figure 1** below.)
    - Any positive results on supplementary (i.e., optional) AC testing must be reported to Incident Command before movement.



**Appendix 11 Figure 1.** Daily testing is required of birds in all barns that are not loaded out within 48 hours of the start of premises-level load out. In this example, there are eight barns on a single-age premises and complete depopulation of the premises takes 96 hours.

**Appendix 11 Table 1.** Biosecurity and mitigation measures by premises-wide load-out duration (including load-out, transport, and any yard [wait] time before slaughter).

Duration of load-out (premises-wide)	≤24 hours	25–48 hours	>48 hours
<p>Existing SBS Biosecurity measures:</p> <ul style="list-style-type: none"> <li>• Load-out crew should shower and wear freshly laundered clothing before going on to other activities after load-out.</li> <li>• Route for live-haul vehicles and equipment should be selected in consultation with a poultry veterinarian or production manager to minimize contact with and proximity to live poultry and poultry products.</li> <li>• When moving equipment from the Infected Zone to the Buffer Zone, C&amp;D of live-haul equipment (power wash) with at least 24-hour downtime is required.</li> </ul>	X	X	X
<p>Load-out equipment and crews:</p> <ul style="list-style-type: none"> <li>• Must have been adequately cleaned and disinfected using one of the following methods: <ul style="list-style-type: none"> <li>○ Sanitization by a disinfectant approved to inactivate HPAI virus</li> <li>○ Adequate downtime before use</li> <li>○ Heat treatment to inactivate HPAI virus</li> </ul> </li> <li>OR</li> <li>• Must have been used only on Monitored Premises during an outbreak. (Equipment and crews that have moved birds of unknown status or positive birds cannot be used.)</li> </ul>		X	
<p>Additional barn-to-barn biosecurity measures:</p> <ul style="list-style-type: none"> <li>• Pre-staging of equipment in a barn prior to beginning load-out in that barn is not allowed.</li> <li>• Other than load-out crew, all persons entering a barn must utilize barn-specific footwear and farm-specific clothing.</li> <li>• Handwashing is required after working with any barn or birds currently or already loaded out and before entering any other occupied barns on the premises.</li> </ul>		X	X

**Appendix 11 Table 1.** Biosecurity and mitigation measures by premises-wide load-out duration (including load-out, transport, and any yard [wait] time before slaughter), cont.

Duration of load-out (premises-wide), cont.	≤24 hours	25–48 hours	>48 hours
<p>Load-out equipment and crews (enhanced):</p> <ul style="list-style-type: none"> <li>• Load-out crew <u>must</u> shower and put on clean clothes before entering the premises to be loaded out AND equipment must have been <u>adequately cleaned and disinfected using one of the following methods</u>:                             <ul style="list-style-type: none"> <li>○ Sanitization by a disinfectant approved to inactivate HPAI</li> <li>○ Adequate downtime before use</li> <li>○ Heat treatment to inactivate HPAI</li> </ul> </li> </ul> <p>OR</p> <ul style="list-style-type: none"> <li>• Load-out crew <u>must</u> shower and put on clean clothes before entering the premises to be loaded out AND equipment must have been <u>used only on specific Monitored Premises during an outbreak as outlined below</u>:                             <ul style="list-style-type: none"> <li>○ Used exclusively to load out premises where PCR testing occurred within 24 hours of movement AND entire premises was depopulated in under 24 hours</li> <li>○ Have not moved birds of unknown or positive status</li> </ul> </li> </ul>			X
<p>Load-out PCR testing:</p> <ul style="list-style-type: none"> <li>• For all barns loading out after initial 48 hours on a premises, daily testing is required.                             <ul style="list-style-type: none"> <li>○ Negative test results taken within 24 hours of scheduled movement (at barn level) must be documented before birds depart premises.</li> <li>○ Any positive results on supplementary AC testing must be reported to Incident Command before movement.</li> </ul> </li> </ul>			X

## References

1. Ritz CW, Worley JW. Poultry mortality composting management guide In: Extension UoGC, ed. Athens, GA, 2012.
2. The National Poultry Improvement Plan. Report of Voting Results on 9- CFR Proposed Changes. NPIP 43rd NPIP Biennial Conference 2016;89-92.
3. USDA: APHIS: VS. Standard E - Biosecurity Principles, National Poultry Improvement Plan (NPIP) Program Standards, 2017;60-64.
4. World Organization for Animal Health (OIE). *Handbook on Import Risk Analysis for Animals and Animal Products*. 2 ed. Paris, France: The World Organization for Animal Health (OIE), 2010.
5. USDA: APHIS: VS: CEAH. An Assessment of the Risk Associated with the Movement of Broiler Hatching Eggs Into, Within, and Out of a Control Area During a Highly Pathogenic Avian Influenza Outbreak. Oct 2012, Egg Sector Working Group, the University of Minnesota, Center for Animal Health and Food Safety, and USDA-APHIS-VS-CEAH. Fort Collins, CO, 2012.
6. USDA:APHIS:VS:NAHMS. Poultry 2010: Structure of the U.S. Poultry Industry, 2011.
7. USDA: Economic Research Service. Food Availability (Per Capita) Data System. <https://www.ers.usda.gov/data-products/food-availability-per-capita-data-system/>, 2017.
8. Bell DD, Weaver WD. *Commercial Chicken Meat and Egg Production*. 5th ed. Norwell, Massachusetts: Kluwer Academic Publishers, 2002.
9. National Chicken Council. U. S. Broiler Production. <http://www.nationalchickencouncil.org/about-the-industry/statistics/u-s-broiler-production/> 2017.
10. Hubbard, LLC. Management guide: Broiler.
11. Annual Report to the United States Animal Health Association, Fiscal Year 2002, National Center for Import and Export, Animal and Plant Health Inspection Service, Veterinary Services. Proceedings of the 106th Annual Meeting of the United States Animal Health Association 2002-2003.
12. USDA APHIS. Poultry Industry Manual In: PReP F, ed, 2013.
13. USDA :APHIS :VS: CEAH: NAHMS. Poultry 2010: Reference of Health and Management Practices on Breeder Chicken Farms in the United States, 2010. Fort Collins, CO, 2011.
14. Moore Jr. PA, Daniel TC, Sharpley AN, et al. Poultry Manure Management In: Wright RJ, Kemper, W.D., Millner, P.D., Power, J.F., Korcak, R.F., ed. *Agricultural Uses of Municipal, Animal, and Industrial Byproducts*. 44 ed: United States Department of Agricultural: Agricultural Research Service, 1998.
15. Cobb-Vantress I. Broiler Management Guide, 2012.
16. Northcutt JKK, Brian. UGA Reference Guide for Solving Poultry Processing Problems (B 1156): UGA extension, 2012.
17. Aviagen. Ross Broiler Management Manuel, 2009.
18. Maryland Department of Agriculture. Poultry Operation Record Keeping Guide & Quick Reference Booklet, 2014.
19. National Chicken Council. Avian Influenza Biosecurity Principles. Washington, D.C., 2015.
20. Segal Y. Prevention and Control of poultry diseases for better farm profitability.: Food and Agriculture Organization.

21. USDA: APHIS. Biosecurity Guide for Poultry and Bird Owners. *Program Aid No 1885*, 2014.
22. Council NC. Poultry biosecurity officer information manual, 2015.
23. USDA. Checklist for Self-Assessment of Enhanced biosecurity.
24. Murray N, MacDiarmid S, Wooldridge M, et al. Handbook on Import Risk Analysis for Animals and Animal Products:.. 2 ed. Paris, France: World Organization for Animal Health (OIE), 2010.
25. Swayne DE. *Avian Influenza*. First ed. Ames, Iowa: Blackwell Publishing, 2008.
26. Perdue ML, Suarez DL, Swayne DE. Avian Influenza in the 1990s. *Avian and Poultry Biology Reviews* 2000;11:1-20.
27. Tong S, Zhu X, Li Y, et al. New world bats harbor diverse influenza A viruses. *PLoS Pathog* 2013;9:e1003657.
28. World Organisation for Animal Health. Update on Highly Pathogenic Avian Influenza In Animals (Type H5 and H7), 2016.
29. Choi YK, Nguyen TD, Ozaki H, et al. Studies of H5N1 Influenza Virus Infection of Pigs by Using Viruses Isolated in Vietnam and Thailand in 2004. *Journal of Virology* 2005;79:10821-10825.
30. Clifford JR. Veterinary Services Memorandum No. 565.14; Reporting Confirmed Findings of Low Pathogenic Notifiable Avian Influenza (LPNAI) (H5 and H7 Subtypes) to the World Organization for Animal Health (OIE) and to Trading Partners In: VS Management Team DVS, ed, 2006.
31. Gaidet N, Cattoli G, Hammoumi S, et al. Evidence of Infection by H5N2 Highly Pathogenic Avian Influenza Viruses in Healthy Wild Waterfowl. *PLoS Pathog* 2008;4:e1000127.
32. Lee DH, Torchetti MK, Winker K, et al. Intercontinental Spread of Asian-Origin H5N8 to North America through Beringia by Migratory Birds. *J Virol* 2015;89:6521-6524.
33. Torchetti MK, Killian ML, Dusek RJ, et al. Novel H5 Clade 2.3.4.4 Reassortant (H5N1) Virus from a Green-Winged Teal in Washington, USA. *Genome Announc* 2015;3.
34. Zhou L-C, Liu J, Pei E-L, et al. Novel Avian Influenza A(H5N8) Viruses in Migratory Birds, China, 2013–2014. *Emerging Infectious Disease journal* 2016;22:1121.
35. Lee D, Bahl J, Torchetti M, et al. Highly pathogenic avian influenza viruses and generation of novel reassortants, United States, 2014–2015. *Emerg Infect Dis* 2016;22.
36. Sleeman JM. Detection of Novel Highly Pathogenic Avian Influenza Viruses in Wild Birds In: Center NWH, ed. 2015-01 ed: USGS, 2015.
37. Ip HS, Torchetti MK, Crespo R, et al. Novel Eurasian highly pathogenic influenza A H5 viruses in wild birds, Washington, USA, 2014. *Emerg Infect Dis* 2015;21.
38. USDA APHIS VS. Epidemiologic and Other Analyses of HPAI-Affected Poultry Flocks: September 9, 2015 Report. 2015.
39. USDA-APHIS. December 2014 – June 2015 Wild Bird Highly Pathogenic Avian Influenza Cases in the United States, 2015.
40. Dejesus E, Costa-Hurtado M, Smith D, et al. Changes in adaptation of H5N2 highly pathogenic avian influenza H5 clade 2.3. 4.4 viruses in chickens and mallards. *Virology* 2016;499:52-64.
41. Senne DA, Suarez DL, Stallnecht DE, et al. Ecology and epidemiology of avian influenza in North and South America. *Developments in biologicals* 2006;124:37-44.



42. USDA:APHIS:VS:STAS:CEAH. Epidemiologic and Other Analyses of Indiana HPAI/LPAI-Affected Poultry Flocks: DRAFT March 4, 2016 Report. In: Health UAVSCfEaA, ed. Fort Collins, CO, 2016;64.
43. Swayne DE. Understanding the complex pathobiology of high pathogenicity avian influenza viruses in birds. *Avian Dis* 2007;51:242-249.
44. Boyce WM, Sandrock C, Kreuder-Johnson C, et al. Avian influenza viruses in wild birds: a moving target. *Comp Immunol Microbiol Infect Dis* 2009;32:275-286.
45. Alexander DJ. An overview of the epidemiology of avian influenza. *Vaccine* 2007;25:5637-5644.
46. Stallknecht DE, Brown JD. Wild birds and the epidemiology of avian influenza. *Journal of Wildlife Diseases* 2007;43:S15-S20.
47. Zhou L-C, Liu J, Pei E-L, et al. Novel Avian Influenza A (H5N8) Viruses in Migratory Birds, China, 2013–2014. *Emerging infectious diseases* 2016;22:1121.
48. Pantin-Jackwood M, Kapczynski D, Spackman E, et al. Pathogenicity and transmission of Eurasian HPAI H5 clade 2.3.4.4 viruses in avian species, 2015.
49. Hinshaw V, Webster R, Easterday B, et al. Replication of avian influenza A viruses in mammals. *Infection and Immunity* 1981;34:354-361.
50. Englund L, Klingeborn B, Mejerland T. Avian Influenza-a Virus Causing an Outbreak of Contagious Interstitial Pneumonia in Mink. *Acta Veterinaria Scandinavica* 1986;27:497-&.
51. Hall JS, Bentler KT, Landolt G, et al. Influenza infection in wild raccoons. *Emerging infectious diseases* 2008;14:1842.
52. Cardona CJ, Xing Z, Sandrock CE, et al. Avian influenza in birds and mammals. *Comparative Immunology, Microbiology and Infectious Diseases* 2009;32:255-273.
53. Root JJ, Shriner SA, Ellis JW, et al. When fur and feather occur together: interclass transmission of avian influenza A virus from mammals to birds through common resources. *Scientific Reports* 2015;5:14354.
54. Bui CM GL, MacIntyre CR. Highly pathogenic avian influenza virus, midwestern United States [letter]. *Emerg Infect Dis* 2016.
55. Arzey G. The Role of Wild Aquatic Birds in the Epidemiology of Avian Influenza in Australia. *Australian Veterinary Journal* 2004;82:377-378.
56. Bean B, Moore BM, Sterner B, et al. Survival of influenza viruses on environmental surfaces. *J Infect Dis* 1982;146:47-51.
57. Shortridge KF, Zhou NN, Guan Y, et al. Characterization of Avian H5N1 Influenza Viruses From Poultry in Hong Kong. *Virology* 1998;252:331-342.
58. Brahmakshatriya V, Lupiani B, Brinlee J, et al. Preliminary study for evaluation of avian influenza virus inactivation in contaminated poultry products using electron beam irradiation. *Avian Pathology* 2009;38:245-250.
59. Beato MS, Mancin M, Bertoli E, et al. Infectivity of H7 LP and HP influenza viruses at different temperatures and pH and persistence of H7 HP virus in poultry meat at refrigeration temperature. *Virology* 2012;433:522-527.
60. Chmielewski R, Swayne DE. Avian Influenza: Public Health and Food Safety Concerns. *Annual Review of Food Science and Technology* 2011;2:21.
61. Spackman E, Gelb J, Preskenis LA, et al. The pathogenesis of low pathogenicity H7 avian influenza viruses in chickens, ducks and turkeys. *Virology journal* 2010;7:1.

62. Toffan A, Serena Beato M, De Nardi R, et al. Conventional inactivated bivalent H5/H7 vaccine prevents viral localization in muscles of turkeys infected experimentally with low pathogenic avian influenza and highly pathogenic avian influenza H7N1 isolates. *Avian Pathology* 2008;37:407-412.
63. Das A, Spackman E, Thomas C, et al. Detection of H5N1 high-pathogenicity avian influenza virus in meat and tracheal samples from experimentally infected chickens. *Avian Dis* 2008;52:40-48.
64. Bertran K, Swayne DE, Pantin-Jackwood MJ, et al. Lack of chicken adaptation of newly emergent Eurasian H5N8 and reassortant H5N2 high pathogenicity avian influenza viruses in the U.S. is consistent with restricted poultry outbreaks in the Pacific flyway during 2014-2015. *Virology* 2016;494:190-197.
65. Swayne DE, Eggert D, Beck JR. Reduction of high pathogenicity avian influenza virus in eggs from chickens once or twice vaccinated with an oil-emulsified inactivated H5 avian influenza vaccine. *Vaccine* 2012;30:4964-4970.
66. Moses H, Brandly C, Jones EE, et al. The isolation and identification of fowl plague virus. *Am J Vet Res* 1948;9:314-328.
67. Starick E, Werner O. Detection of H7 Avian Influenza Virus Directly From Poultry Specimens. *Avian Diseases* 2003;47:1187-1189.
68. Pillai S, Saif Y, Lee C. Detection of influenza A viruses in eggs laid by infected turkeys. *Avian Diseases* 2010;54:830-833.
69. Suarez D, Woolcock P, Bermudez A, et al. Isolation from turkey breeder hens of a reassortant H1N2 influenza virus with swine, human, and avian lineage genes. *Avian Diseases* 2002;46:111-121.
70. Mohan R, Saif Y, Erickson G, et al. Serologic and epidemiologic evidence of infection in turkeys with an agent related to the swine influenza virus. *Avian Diseases* 1981;25:11-16.
71. Ficken M, Guy J, Gonder E. An outbreak of influenza (H1N1) in turkey breeder hens. *Avian Diseases* 1989;33:370-374.
72. Akey BL. Low-pathogenicity H7N2 avian influenza outbreak in Virginia during 2002. *Avian Diseases* 2003;47:1099-1103.
73. Narayan O, Lang G, Rouse B. A new influenza A virus infection in turkeys. *Archives of Virology* 1969;26:149-165.
74. Birnbaum NG, O'Brien B. Methods for Inactivation of Avian Influenza Virus in the Environment. 1st ed. Ames, Iowa: Wiley-Blackwell, 2008;391-405.
75. De Benedictis P, Beato MS, Capua I. Inactivation of Avian Influenza Viruses by Chemical Agents and Physical Conditions: A Review. *Zoonoses and Public Health* 2007;54:51-68.
76. Lombardi ME, Ladman BS, Alphin RL, et al. Inactivation of avian influenza virus using common detergents and chemicals. *Avian Dis* 2008;52:118-123.
77. Fichtner GJ. The Pennsylvania/Virginia experience in eradication of avian influenza (H5N2). *Avian Diseases* 2003;33-38.
78. Beard CW, Brugh M, Johnson DC. Laboratory studies with the Pennsylvania avian influenza viruses (H5N2) 1984;462-473.
79. Songserm T, Jam-On R, Sae-Heng N, et al. Survival and stability of HPAI H5N1 in different environments and susceptibility to disinfectants. *Developments in biologicals* 2006;124:254.

80. Alexander DJ. The Epidemiology and Control of Avian Influenza and Newcastle-Disease. *Journal of Comparative Pathology* 1995;112:105-126.
81. Ssematimba A, Hagensars TJ, de Jong MCM. Modelling the Wind-Borne Spread of Highly Pathogenic Avian Influenza Virus between Farms. *PLoS ONE* 2012;7:e31114.
82. Ypma RJF, Jonges M, Bataille A, et al. Genetic data provide evidence for wind-mediated transmission of highly pathogenic avian influenza. *Journal of Infectious Diseases* 2012.
83. McQuiston JH, Garber LP, Porter-Spalding BA, et al. Evaluation of Risk Factors for the Spread of Low Pathogenicity H7N2 Avian Influenza Virus Among Commercial Poultry Farms. *JAVMA-Journal of the American Veterinary Medical Association* 2005;226:767-772.
84. Beato MS, Capua I, Alexander DJ. Avian influenza viruses in poultry products: a review. *Avian Pathology* 2009;38:193-200.
85. Kreager K. Avian Influenza Control Philosophies in the Layer and Layer Breeder industries. *Avian Diseases* 2003;344-348.
86. Ssematimba A, Hagensars T, de Wit J, et al. Avian influenza transmission risks: Analysis of biosecurity measures and contact structure in Dutch poultry farming. *Preventive Veterinary Medicine* 2013;109:106-115.
87. Stegeman JA, Bouma A. Epidemiology and Control of Avian Influenza. 11th International Conference of the Associations for Tropical Veterinary Medicine and 16th Veterinary Association Malaysia Congress 2004;141-143.
88. Samadieh B, Bankowski RA. Transmissibility of Avian Influenza-A Viruses. *American Journal of Veterinary Research* 1971;32:939-945.
89. Canadian Food Inspection Agency. Comprehensive Report on the 2004 Outbreak of High Pathogenicity Avian Influenza (H7N3) in the Fraser Valley of British Columbia, Canada., 2004.
90. Beard CW, Brugh M. Laboratory Studies on the Pennsylvania Isolates of Avian Influenza (H5N2) in Specific Pathogen-Free Chickens. *Journal of the American Veterinary Medical Association* 1984;185:340.
91. Swayne DE, Suarez DL, Sims LD. Influenza. *Diseases of Poultry*. 13th ed. Ames, IA: Wiley-Blackwell, 2013;181-218.
92. McCapes RH, Bankowski R, West GB. Avian Influenza in California: the nature of the clinical disease 1964-1985. *Avian Diseases* 2003;47:118-132.
93. Smithies L, Emerson F, Robertson S, et al. Two different type A influenza virus infections in turkeys in Wisconsin II. 1968 outbreak. *Avian Diseases* 1969;13:606-610.
94. Pedroni E, Muñoz X, Sotomayor V, et al. Outbreak of human A (H1N1) influenza in turkeys of a commercial poultry farm, Valparaiso, Chile: August 2009. *Revista chilena de infectologia: organo oficial de la Sociedad Chilena de Infectologia* 2012;29:420-426.
95. Kleven SH, Nelson RC, Deshmukh DR, et al. Epidemiologic and field observations on avian influenza in Minnesota turkeys. *Avian Diseases* 1970:153-166.
96. Halvorson DA. The control of H5 or H7 mildly pathogenic avian influenza: a role for inactivated vaccine. *Avian Pathology* 2002;31:5-12.
97. Halvorson DA. Personal Communication: AI virus isolation from turkey semen. email ed, 2012.
98. Samadieh B, Bankowski R. Effect of avian influenza-A viruses upon egg production and fertility of turkeys. *Avian Diseases* 1970:715-722.

99. Pantin-Jackwood M, Wasilenko JL, Spackman E, et al. Susceptibility of turkeys to pandemic-H1N1 virus by reproductive tract insemination. *Virology Journal* 2010;7:27.
100. Swayne DE. Personal Communication: AI virus isolation in turkey semen. email ed, 2012.
101. Ali A, Yassine H, Awe OO, et al. Replication of Swine and Human Influenza Viruses in Juvenile and Layer Turkey Hens. *Veterinary Microbiology* 2013;163:71-78.
102. Aldous EW, Seekings JM, McNally A, et al. Infection dynamics of highly pathogenic avian influenza and virulent avian paramyxovirus type 1 viruses in chickens, turkeys and ducks. *Avian Pathology* 2010;39:265-273.
103. Nili H, McNally A, Aldous E, et al. Pathological changes in turkeys experimentally infected with different doses of A/ostrich/Italy/984/2000 H7N1 avian influenza virus. *Iranian Journal of Veterinary Research* 2008;9:330-335.
104. Homme PJ, Easterday BC. Avian Influenza Virus Infections .III. Antibody Response by Turkeys to Influenza-a/Turkey/Wisconsin/1966 Virus. *Avian Diseases* 1970;14:277-284.
105. Swayne DE, Slemons RD. Using mean infectious dose of high- and low-pathogenicity avian influenza viruses originating from wild duck and poultry as one measure of infectivity and adaptation to poultry. *Avian Dis* 2008;52:455-460.
106. Spekrijse D, Bouma A, Stegeman JA, et al. The effect of inoculation dose of a highly pathogenic avian influenza virus strain H5N1 on the infectiousness of chickens. *Veterinary Microbiology* 2011;147:59-66.
107. Kitajima M, Huang Y, Watanabe T, et al. Dose–response time modelling for highly pathogenic avian influenza A (H5N1) virus infection. *Letters in applied microbiology* 2011;53:438-444.
108. Schijven F, Teunis PFM. Quantitative Risk Assessment of Avian Influenza Virus Infection via Water. 2006.
109. Kwon Y, Swayne D. Different routes of inoculation impact infectivity and pathogenesis of H5N1 high pathogenicity avian influenza virus infection in chickens and domestic ducks. *Avian Diseases* 2010;54:1260-1269.
110. Swayne DE, Beck JR. Experimental Study to Determine if Low-Pathogenicity and High-Pathogenicity Avian Influenza Viruses Can Be Present in Chicken Breast and Thigh Meat Following Intranasal Virus Inoculation. *Avian Diseases* 2005;49:81-85.
111. Purchase H. Experiments on the viability of the virus of fowl-plague under trade conditions. *Vet Rec* 1931;11:644-648.
112. Sergeev AA, Demina O, Pyankov O, et al. Infection of chickens caused by avian influenza virus A/H5N1 delivered by aerosol and other routes. *Transboundary and Emerging Diseases* 2013;60:159-165.
113. Spekrijse D, Bouma A, Koch G, et al. Quantification of dust-borne transmission of highly pathogenic avian influenza virus between chickens. *Influenza and other respiratory viruses* 2013;7:132-138.
114. Yee KS, Carpenter TE, Farver TB, et al. An evaluation of transmission routes for low pathogenicity avian influenza virus among chickens sold in live bird markets. *Virology* 2009;394:19-27.
115. USDA: APHIS: VS: CEAH. An Assessment of the Risk Associated with the Movement of Eggshells and Inedible Egg Product Into, Within, and Out of a Control Area During a Highly Pathogenic Avian Influenza Outbreak. March, 2013, Collaboration with

University of Minnesota, Center for Animal Health and Food Safety, Egg Sector Working Group. Fort Collins, CO, 2013.

116. Elbers ARW, Fabri THF, De Vries TS, et al. The Highly Pathogenic Avian Influenza a (H7N7) Virus Epidemic in the Netherlands in 2003 - Lessons Learned From the First Five Outbreaks. *Avian Diseases* 2004;48:691-705.

117. Saenz RA, Essen SC, Brookes SM, et al. Quantifying Transmission of Highly Pathogenic and Low Pathogenicity H7N1 Avian Influenza in Turkeys. *PLoS ONE* 2012;7:e45059.

118. Kilany WH, Abdelwhab E, Arafa AS, et al. Protective efficacy of H5 inactivated vaccines in meat turkey poults after challenge with Egyptian variant highly pathogenic avian influenza H5N1 virus. *Veterinary Microbiology* 2011;150:28-34.

119. USDA: APHIS: VS: CEAH, University of Minnesota Center for Animal Health and Food Safety, Turkey Sector Working Group,. Draft Assessment of the Risk Associated with the Movement of Turkey Hatching Eggs Into, Within, and Out of a Control Area During a Highly Pathogenic Avian Influenza Outbreak, Last reviewed: Jan 2015. Fort Collins, CO, 2015.

120. van der Goot JA, Koch G, De Jong MCM, et al. Quantification of the Effect of Vaccination on Transmission of Avian Influenza (H7N7) in Chickens. *Proceedings of the National Academy of Sciences of the United States of America* 2005;102:18141-18146.

121. van der Goot JA, De Jong MCM, Koch G, et al. Comparison of the Transmission Characteristics of Low and High Pathogenicity Avian Influenza A Virus (H5N2). *Epidemiology and Infection* 2003;131:1003-1013.

122. Pfeiffer J, Pantin-Jackwood M, To TL, et al. Phylogenetic and biological characterization of highly pathogenic H5N1 avian influenza viruses (Vietnam 2005) in chickens and ducks. *Virus Res* 2009;142:108-120.

123. Bouma A, Claassen I, Natih K, et al. Estimation of transmission parameters of H5N1 avian influenza virus in chickens. *PLoS Pathog* 2009;5:e1000281.

124. Bertran K, Lee D-H, Balzli C, et al. Age is not a determinant factor in susceptibility of broilers to H5N2 clade 2.3.4.4 high pathogenicity avian influenza virus. *Veterinary Research* 2016;47:116.

125. Swayne DE, Pantin-Jackwood M. Pathobiology of Avian Influenza Virus Infections in Birds and Mammals In: Swayne DE, ed. *Avian Influenza*. 1st ed. Ames, IA: Blackwell Publishing, 2008;87-122.

126. Irvine R, Banks J, Londt B, et al. Outbreak of highly pathogenic avian influenza caused by Asian lineage H5N1 virus in turkeys in Great Britain in January 2007. *Veterinary record* 2007;161:100-101.

127. Swayne DE, Suarez DL. Highly Pathogenic Avian Influenza. *Revue Scientifique et Technique de l'Office International des Epizooties* 2000;19:463-482.

128. Swayne DE, Halvorson DA. Influenza In: Saif YM, Fadly AM, Glisson JR, et al., eds. *Diseases of Poultry*. Ames, IA: Blackwell Publishing, 2008;168.

129. Mutinelli E, Capua I, Terregino C, et al. Clinical, Gross, and Microscopic Findings in Different Avian Species Naturally Infected During the H7N1 Low- and High-Pathogenicity Avian Influenza Epidemics in Italy During 1999 and 2000. *Avian Diseases* 2003;47:844-848.

130. USDA: APHIS: VS. Highly pathogenic avian influenza response plan, The Red Book; Foreign Animal Disease Preparedness & Response Plan FAD PReP. 2015.

131. USDA: APHIS: VS. Avian Influenza Diagnostics and Testing In: Service USDoAAaPHI, ed, 2013.
132. Dorigatti I, Mulatti P, Rosà R, et al. Modelling the spatial spread of H7N1 avian influenza virus among poultry farms in Italy. *Epidemics* 2010;2:29-35.
133. Boender GJ, Hagenaaars TJ, Bouma A, et al. Risk maps for the spread of highly pathogenic avian influenza in poultry. *PLoS Comput Biol* 2007;3:e71.
134. Rorres C, Pelletier STK, Bruhn MC, et al. Ongoing Estimation of the Epidemic Parameters of a Stochastic, Spatial, Discrete-Time Model for a 1983–84 Avian Influenza Epidemic. *Avian diseases* 2011;55:35-42.
135. Busani L, Valsecchi MG, Rossi E, et al. Risk factors for highly pathogenic H7N1 avian influenza virus infection in poultry during the 1999–2000 epidemic in Italy. *The Veterinary Journal* 2009;181:171-177.
136. Sharkey KJ, Bowers RG, Morgan KL, et al. Epidemiological consequences of an incursion of highly pathogenic H5N1 avian influenza into the British poultry flock. *Proceedings of the Royal Society B: Biological Sciences* 2008;275:19-28.
137. Brugh M, Johnson DC. Epidemiology of Avian Influenza in Domestic Poultry. *Avian Diseases* 1986;47:177-186.
138. Wilson D, Schmidtman E, Richard R, et al. Isolation of avian influenza from insects. Arbovirus Research in Australia-Proceedings 4th Symposium 1986.
139. Axtell RC. Poultry integrated pest management: status and future. *Integrated Pest Management Reviews* 1999;4:53-73.
140. Halvorson DA. Avian Influenza: a Minnesota cooperative control program. *Avian Diseases* 2003;47:327-336.
141. Cardona C. Low-Pathogenicity Avian Influenza virus outbreak in commercial poultry in California. The threat of pandemic influenza: are we ready. *National Academy Press, Washington* 2005:243-253.
142. USDA: APHIS: VS: CEAH, University of Minnesota Center for Animal Health and Food Safety, Broiler Sector Working Group. An Assessment of the Risk Associated with the Movement of Broiler Day Old Chicks, Within, and Out of a Control Area during a Highly Pathogenic Avian Influenza Outbreak. Fort Collins, CO, 2013.
143. Sawabe K, Hoshino K, Isawa H, et al. Detection and isolation of highly pathogenic H5N1 avian influenza A viruses from blow flies collected in the vicinity of an infected poultry farm in Kyoto, Japan, 2004. *Am J Trop Med Hyg* 2006;75:327-332.
144. Sawabe K, Hoshino K, Isawa H, et al. Blow Flies Were One of the Possible Candidates for Transmission of Highly Pathogenic H5N1 Avian Influenza Virus during the 2004 Outbreaks in Japan. *Influenza Research and Treatment* 2011;2011:8.
145. Nielsen AA, Skovgard H, Stockmarr A, et al. Persistence of low-pathogenic avian influenza H5N7 and H7N1 subtypes in house flies (Diptera: Muscidae). *J Med Entomol*;48:608-614.
146. Wanaratana S, Panyim S, Pakpinyo S. The potential of house flies to act as a vector of avian influenza subtype H5N1 under experimental conditions. *Med Vet Entomol* 2011;25:58-63.
147. Sawabe K, Tanabayashi K, Hotta A, et al. Survival of avian H5N1 influenza A viruses in *Calliphora nigribarbis* (Diptera: Calliphoridae). *J Med Entomol* 2009;46:852-855.
148. Wanaratana S, Amonsin A, Chaisingh A, et al. Experimental assessment of house flies as vectors in avian influenza subtype H5N1 transmission in chickens. *Avian Diseases* 2013.

149. Tsuda Y, Hayashi T, Higa Y, et al. Dispersal of a blow fly, *Calliphora nigribarbis*, in relation to the dissemination of highly pathogenic avian influenza virus. *Japanese Journal of Infectious Diseases* 2009;62:294-297.
150. Tsuda Y. Personal Communication: HPAI transmission through flies, 2012.
151. Stafford KC. *Fly management handbook A guide to biology, dispersal, and management of the house fly and related flies for farmers, municipalities, and public health officials*: The Connecticut Agricultural Experiment Station, New Haven, 2008.
152. James M, Harwood R. The house fly and its relatives. *Herms's Medical Entomology McMillan Company 6th edition, London, England* 1969:249-265.
153. Campbell J. G89-954 A Guide for Managing Poultry Insects (Revised April 1996) In: Extension. HMfUoN-L, ed. *Paper 1147*, 1989.
154. Crippen TL, Sheffield CL, Esquivel SV, et al. The acquisition and internalization of Salmonella by the lesser mealworm, *Alphitobius diaperinus* (Coleoptera: Tenebrionidae). *Vector Borne Zoonotic Dis* 2009;9:65-72.
155. Hosen M, Khan AR, Hossain M. Growth and Development of the Lesser Mealworm, *Alphitobius diaperinus* (Panzer) (Coleoptera: Tenebrionidae) on Cereal Flours. *Pakistan Journal of Biological Sciences* 2004.
156. Winpisinger KA, Ferketich AK, Berry RL, et al. Spread of *Musca domestica* (Diptera: Muscidae), from two caged layer facilities to neighboring residences in rural Ohio. *Journal of Medical Entomology* 2005;42:732-738.
157. Sergeev AA, Demina O, Pyankov O, et al. Infection of chickens caused by avian influenza virus A/H5N1 delivered by aerosol and other routes. *Transboundary and emerging diseases* 2013;60:159-165.
158. Guan J, Fu Q, Chan M, et al. Aerosol transmission of an avian influenza H9N2 virus with a tropism for the respiratory tract of chickens. *Avian diseases* 2013;57:645-649.
159. Selleck PW, Arzey G, Kirkland PD, et al. An Outbreak of Highly Pathogenic Avian Influenza in Australia in 1997 Caused by an H7N4 Virus. *Avian Diseases* 2003;47:806-811.
160. Inter American Institute for Cooperation on Agriculture. Canada's experiences with avian influenza (AI). A compilation of documents on AI and the response of the Canadian Government and poultry sector to the 2004 AI outbreak in British Columbia. 2005.
161. USDA APHIS VS. Epidemiologic and Other Analyses of HPAI-Affected Poultry Flocks: September 9, 2015 Report, D. Halvorson, Personal Communication. 2015.
162. Henzler DJ, Kradel DC, Davison S, et al. Epidemiology, Production Losses, and Control Measures Associated With an Outbreak of Avian Influenza Subtype H7N2 in Pennsylvania (1996-98). *Avian Diseases* 2003;47:1022-1036.
163. Schofield L, Ho J, Kournikakis B, et al. Avian Influenza Aerosol Sampling Campaign in the British Columbia Fraser Valley, 9-19 April 2004: Defense Research and Development Canada, 2005.
164. USDA-APHIS. Epidemiologic and Other Analyses of HPAI-Affected Poultry Flocks: July 15, 2015 Report, 2015.
165. Forman AJ, Parsonson IM, Doughty WJ. The Pathogenicity of an Avian Influenza-Virus Isolated in Victoria. *Australian Veterinary Journal* 1986;63:294-296.
166. van der Goot JA, Koch G, de Jong MC, et al. Transmission dynamics of low- and high-pathogenicity A/Chicken/Pennsylvania/83 avian influenza viruses. *Avian Dis* 2003;47:939-941.

167. Homme PJ, Easterday BC, Anderson DP. Avian Influenza Virus Infections II. Experimental Epizootiology of Influenza a/Turkey/Wisconsin/1966 Virus in Turkeys. *Avian Diseases* 1970;14:240-247.
168. Zhang P, Tang Y, Liu X, et al. Characterization of H9N2 influenza viruses isolated from vaccinated flocks in an integrated broiler chicken operation in eastern China during a 5 year period (1998-2002). *J Gen Virol* 2008;89:3102-3112.
169. Tsukamoto K, Imada T, Tanimura N, et al. Impact of different husbandry conditions on contact and airborne transmission of H5N1 highly pathogenic avian influenza virus to chickens. *Avian Dis* 2007;51:129-132.
170. Spekrijse D, Bouma A, Koch G, et al. Airborne transmission of a highly pathogenic avian influenza virus strain H5N1 between groups of chickens quantified in an experimental setting. *Veterinary Microbiology* 2011;152:88-95.
171. Zhong L, Wang X, Li Q, et al. Molecular mechanism of the airborne transmissibility of H9N2 avian influenza A viruses in chickens. *Journal of virology* 2014;88:9568-9578.
172. Weber TP, Stilianakis NI. Inactivation of influenza A viruses in the environment and modes of transmission: a critical review. *J Infect* 2008;57:361-373.
173. Tellier R. Review of aerosol transmission of influenza A virus. *Emerg Infect Dis* 2006;12:1657-1662.
174. McDevitt JJ, Rudnick SN, Radonovich LJ. Aerosol susceptibility of influenza virus to UV-C light. *Appl Environ Microbiol* 2012;78:1666-1669.
175. Krauss S, Walker D, Pryor SP, et al. Influenza A viruses of migrating wild aquatic birds in North America. *Vector Borne Zoonotic Dis* 2004;4:177-189.
176. De Marco MA, Foni E, Campitelli L, et al. Long-Term Monitoring for Avian Influenza Viruses in Wild Bird Species in Italy. *Veterinary Research Communications* 2003;27:107-114.
177. Brown JD, Stallknecht DE, Beck JR, et al. Susceptibility of North American ducks and gulls to H5N1 highly pathogenic avian influenza viruses. *Emerg Infect Dis* 2006;12:1663-1670.
178. Arnal A, Vittecoq M, Pearce-Duvel J, et al. Laridae: A neglected reservoir that could play a major role in avian influenza virus epidemiological dynamics. *Critical reviews in microbiology* 2015;41:508-519.
179. Mathieu C, Moreno V, Pedersen J, et al. Avian Influenza in wild birds from Chile, 2007-2009. *Virus Res* 2015;199:42-45.
180. Alexander DJ, Brown IH. Recent Zoonoses Caused by Influenza a Viruses. *Revue Scientifique Et Technique De L Office International Des Epizooties* 2000;19:197-225.
181. Capua I, Alexander DJ. Avian influenza infections in birds—a moving target. *Influenza and other respiratory viruses* 2007;1:11-18.
182. Kou Z, Li Y, Yin Z, et al. The Survey of H5N1 Flu Virus in Wild Birds in 14 Provinces of China from 2004 to 2007. *PLoS ONE* 2009;4:e6926.
183. Gilbert M, Jambal L, Karesh WB, et al. Highly Pathogenic Avian Influenza Virus among Wild Birds in Mongolia. *PLoS ONE* 2012;7:e44097.
184. World Organisation for Animal Health (OIE). Highly pathogenic avian influenza, United States of America 16/12/2014, 2014.
185. World Organisation for Animal Health (OIE). Highly pathogenic avian influenza, United States of America 20/01/2015, 2015a.



186. Verhagen JH, van der Jeugd HP, Nolet BA, et al. Wild bird surveillance around outbreaks of highly pathogenic avian influenza A(H5N8) virus in the Netherlands, 2014, within the context of global flyways. *Euro Surveill* 2015;20.
187. Krauss S, Stallknecht DE, Slemons RD, et al. The enigma of the apparent disappearance of Eurasian highly pathogenic H5 clade 2.3. 4.4 influenza A viruses in North American waterfowl. *Proceedings of the National Academy of Sciences* 2016;113:9033-9038.
188. Sa e Silva M, Mathieu-Benson C, Kwon YK, et al. Experimental infection with low and high pathogenicity H7N3 Chilean avian influenza viruses in Chiloe wigeon (*Anas sibilatrix*) and cinnamon teal (*Anas cyanoptera*). *Avian Dis* 2011;55:459-461.
189. Alexander DJ, Parsons G, Manvell RJ. Experimental Assessment of the Pathogenicity of 8 Avian Influenza a Viruses of H-5 Subtype for Chickens, Turkeys, Ducks and Quail. *Avian Pathology* 1986;15:647-662.
190. Wood J, Webster R, Nettles V. Host range of A/Chicken/Pennsylvania/83 (H5N2) influenza virus. *Avian Diseases* 1985:198-207.
191. Pantin-Jackwood MJ, Costa-Hurtado M, Shepherd E, et al. Experimental infection of mallard ducks with different subtype H5 and H7 highly pathogenic avian influenza viruses. *AAAP*, 2014.
192. Garber L, Bjork KE, Patyk KA, et al. Factors associated with highly pathogenic avian influenza H5N2 infection on table egg layer farms in the Midwest, United States, 2015. *Avian Diseases* 2016.
193. Burns TE, Ribble C, Stephen C, et al. Use of observed wild bird activity on poultry farms and a literature review to target species as high priority for avian influenza testing in 2 regions of Canada. *Can Vet J* 2012;53:158-166.
194. CFIA. Avian influenza investigation in British Columbia - 2014/2015, 2014-2015.
195. Pasick J, Handel K, Robinson J, et al. Relationship between H5N2 avian influenza viruses isolated from wild and domestic ducks in British Columbia, Canada. *Avian Dis* 2007;51:429-431.
196. USDA. Update on avian influenza findings in the Pacific Flyway, 2015.
197. Bevins SN, Dusek RJ, White CL, et al. Widespread detection of highly pathogenic H5 influenza viruses in wild birds from the Pacific Flyway of the United States. *Scientific Reports* 2016;6:28980.
198. World Organisation for Animal Health (OIE). Highly pathogenic avian influenza, United States of America 25/01/2015, 2015b.
199. Christopher SJ, Michelle C, Erik CH, et al. Surveillance for Highly Pathogenic Avian Influenza Virus in Wild Birds during Outbreaks in Domestic Poultry, Minnesota, 2015. *Emerging Infectious Disease journal* 2016;22.
200. Alexander DJ. A Review of Avian Influenza in Different Bird Species. *Veterinary Microbiology* 2000;74:3-13.
201. Cunningham DL, Fairchild BD. Biosecurity basics for poultry growers In: Extension UoGC, ed, 2012;4.
202. Morishita TY. Biosecurity for poultry In: Extension TOSU, ed, 2001.
203. Carey JB, Prochaska F, Jeffrey JS. Poultry Facility Biosecurity In: Texas Agricultural Extension Service TTAMUS, ed. College Station, TX.
204. Utah Department of Agriculture and Food. High Pathogenic Avian Flu, 2015.
205. Becker WB. The isolation and classification of Tern virus: influenza A-Tern South Africa--1961. *J Hyg (Lond)* 1966;64:309-320.

206. Hesterberg U, Harris K, Stroud D, et al. Avian influenza surveillance in wild birds in the European Union in 2006. *Influenza and Other Respiratory Viruses* 2009;3:1-14.
207. Shriner SA, Root JJ, Mooers NL, et al. Susceptibility of rock doves to low-pathogenic avian influenza A viruses. *Archives of virology* 2016;161:715-720.
208. Guarino JL. Bird movements in relation to control. 1968.
209. Craven S, Stern N, Line E, et al. Determination of the incidence of Salmonella spp., Campylobacter jejuni, and Clostridium perfringens in wild birds near broiler chicken houses by sampling intestinal droppings. *Avian diseases* 2000:715-720.
210. Kalthoff D, Breithaupt A, Helm B, et al. Migratory Status Is Not Related to the Susceptibility to HPAIV H5N1 in an Insectivorous Passerine Species. *PLoS ONE* 2009;4:e6170.
211. Hinkle NC, Hickle LA. California caged layer pest management evaluation. *The Journal of Applied Poultry Research* 1999;8:327-338.
212. USDA: APHIS: VS: STAS: CEAH. Risk that Poultry Feed made with Corn—Potentially Contaminated with Eurasian - North American Lineage H5N2 HPAI Virus from Wild Migratory Birds — Results in Exposure of Susceptible Commercial Poultry, Sept 2015 In: USDA, ed. Fort Collins. CO, 2015.
213. Nestorowicz A, Kawaoka Y, Bean WJ, et al. Molecular analysis of the hemagglutinin genes of Australian H7N7 influenza viruses: role of passerine birds in maintenance or transmission? *Virology* 1987;160:411-418.
214. Villareal C, Flores A. The Mexican avian influenza (H5N2) outbreak. *Avian Diseases* 2003;47:18-22.
215. Feare CJ. Role of wild birds in the spread of highly pathogenic avian influenza virus H5N1 and implications for global surveillance. *Avian Dis* 2010;54:201-212.
216. Kapczynski DR, Pantin-Jackwood M, Guzman SG, et al. Characterization of the 2012 highly pathogenic avian influenza H7N3 virus isolated from poultry in an outbreak in Mexico: pathobiology and vaccine protection. *J Virol* 2013;87:9086-9096.
217. Minnesota Department of Natural Resources. Second confirmed case of avian influenza reported in wild birds, July 10, 2015, 2015.
218. Stallknecht DE, Shane SM. Host Range of Avian Influenza-Virus in Free-Living Birds. *Veterinary Research Communications* 1988;12:125-141.
219. Brown JD, Luttrell MP, Berghaus RD, et al. Prevalence of antibodies to type A influenza virus in wild avian species using two serologic assays. *Journal of Wildlife Diseases* 2010;46:896-911.
220. Schnebel B, Dierschke V, Rautenschlein S, et al. No Detection of Avian Influenza a Viruses of the Subtypes H5 and H7 and Isolation of Lentogenic Avian Paramyxovirus Serotype 1 in Passerine Birds During Stopover in the Year 2001 on the Island Helgoland (North Sea). *Deutsche Tierärztliche Wochenschrift* 2005;112:456-460.
221. Račnik J, Slavec B, Trilar T, et al. Evidence of avian influenza virus and paramyxovirus subtype 2 in wild-living passerine birds in Slovenia. *European Journal of Wildlife Research* 2008;54:529-532.
222. Gronesova P, Kabat P, Trnka A, et al. Using nested RT-PCR analyses to determine the prevalence of avian influenza viruses in passerines in western Slovakia, during summer 2007. *Scand J Infect Dis* 2008;40:954-957.
223. Han Y, Hou G, Jiang W, et al. A Survey of Avian Influenza in Tree Sparrows in China in 2011. *PLoS ONE* 2012;7:e33092.

224. Boon AC, Sandbulte MR, Seiler P, et al. Role of terrestrial wild birds in ecology of influenza A virus (H5N1). *Emerg Infect Dis* 2007;13:1720-1724.
225. Brown JD, Stallknecht DE, Berghaus RD, et al. Infectious and lethal doses of H5N1 highly pathogenic avian influenza virus for house sparrows (*Passer domesticus*) and rock pigeons (*Columbia livia*). *J Vet Diagn Invest* 2009;21:437-445.
226. Perkins LEL, Swayne DE. Comparative Susceptibility of Selected Avian and Mammalian Species to a Hong Kong-Origin H5n1 High-Pathogenicity Avian Influenza Virus. *Avian Diseases* 2003;47:956-967.
227. Perkins LEL, Swayne DE. Varied Pathogenicity of a Hong Kong-Origin H5n1 Avian Influenza Virus in Four Passerine Species and Budgerigars. *Veterinary Pathology* 2003;40:14-24.
228. Forrest HL, Kim JK, Webster RG. Virus shedding and potential for interspecies waterborne transmission of highly pathogenic H5N1 influenza virus in sparrows and chickens. *J Virol* 2010;84:3718-3720.
229. Yamamoto Y, Nakamura K, Yamada M, et al. Pathogenesis in Eurasian tree sparrows inoculated with H5N1 highly pathogenic avian influenza virus and experimental virus transmission from tree sparrows to chickens. *Avian Dis* 2013;57:205-213.
230. Alfonso CP, Cowen BS, Vancampen H. Influenza-a Viruses Isolated From Waterfowl in 2 Wildlife Management Areas of Pennsylvania. *Journal of Wildlife Diseases* 1995;31:179-185.
231. van den Brand JMA, Krone O, Wolf PU, et al. Host-specific exposure and fatal neurologic disease in wild raptors from highly pathogenic avian influenza virus H5N1 during the 2006 outbreak in Germany. *Veterinary Research* 2015;46:24.
232. Lierz M, Hafez HM, Klopffleisch R, et al. Protection and virus shedding of falcons vaccinated against highly pathogenic avian influenza A virus (H5N1). *Emerg Infect Dis* 2007;13:1667-1674.
233. Marinova-Petkova A, Georgiev G, Seiler P, et al. Spread of influenza virus A (H5N1) clade 2.3. 2.1 to Bulgaria in common buzzards. *Emerging infectious diseases* 2012;18:1596.
234. USDA. Update on Avian Influenza Findings. Poultry Findings Confirmed by USDA's National Veterinary Services Laboratories, 2015.
235. Naguib MM, Kinne J, Chen H, et al. Outbreaks of highly pathogenic avian influenza H5N1 clade 2.3.2.1c in hunting falcons and kept wild birds in Dubai implicate intercontinental virus spread. *J Gen Virol* 2015.
236. Redig PT, Goyal SM. Serologic evidence of exposure of raptors to influenza A virus. *Avian Dis* 2012;56:411-413.
237. Gunnarsson G, Jourdain E, Waldenstrom J, et al. Zero prevalence of influenza A virus in two raptor species by standard screening. *Vector Borne Zoonotic Dis* 2010;10:387-390.
238. Peterson MJ, Aguirre R, Ferro PJ, et al. Infectious Disease Survey of Rio Grande Wild Turkeys in the Edwards Plateau of Texas. *Journal of Wildlife Diseases* 2002;38:826-833.
239. Ferro PJ, Khan O, Vuong C, et al. Avian influenza virus investigation in wild bobwhite quail from Texas. *Avian Dis* 2012;56:858-860.
240. Hall JS, Ip HS, Franson JC, et al. Experimental infection of a North American raptor, American Kestrel (*Falco sparverius*), with highly pathogenic avian influenza virus (H5N1). *PLoS One* 2009;4:e7555.

241. Zabel CJ, McKelvey K, Ward Jr JP. Influence of primary prey on home-range size and habitat-use patterns of northern spotted owls (*Strix occidentalis caurina*). *Canadian Journal of Zoology* 1995;73:433-439.
242. Chiang S, Bloom P, Bartuszevige A, et al. Home range and habitat use of Cooper's hawks in urban and natural areas. In: Christopher A. Lepczyk and Paige S. Warren, ed. *Urban bird ecology and conservation Studies in Avian Biology (no 45)*: University of California Press, Berkeley, CA., 2012.
243. Whitacre D, Burnham W. Home Range Size, Nesting Density, and Body Size in Raptors. In *Neotropical birds of prey: Biology and ecology of a forest raptor community.*: Cornell University Press., 2013;349- 350.
244. Nuradji H, Bingham J, Lowther S, et al. A comparative evaluation of feathers, oropharyngeal swabs, and cloacal swabs for the detection of H5N1 highly pathogenic avian influenza virus infection in experimentally infected chickens and ducks. *Journal of Veterinary Diagnostic Investigation* 2015;27:704-715.
245. Busquets N, Abad FX, Alba A, et al. Persistence of highly pathogenic avian influenza virus (H7N1) in infected chickens: feather as a suitable sample for diagnosis. *Journal of General Virology* 2010;91:2307-2313.
246. Yamamoto Y, Nakamura K, Yamada M, et al. Persistence of avian influenza virus (H5N1) in feathers detached from bodies of infected domestic ducks. *Applied and Environmental Microbiology* 2010;76:5496-5499.
247. Yamamoto Y, Nakamura K, Yamada M, et al. Comparative pathology of chickens and domestic ducks experimentally infected with highly pathogenic avian influenza viruses (H5N1) isolated in Japan in 2007 and 2008. *Japan Agricultural Research Quarterly: JARQ* 2010;44:73-80.
248. Aiello R, Beato MS, Mancin M, et al. Differences in the detection of highly pathogenic avian influenza H5N1 virus in feather samples from 4-week-old and 24-week-old infected Pekin ducks (*Anas platyrhynchos* var. *domestica*). *Veterinary microbiology* 2013;165:443-447.
249. Spackman E, Pantin-Jackwood MJ, Kapczynski DR, et al. H5N2 Highly Pathogenic Avian Influenza Viruses from the US 2014-2015 outbreak have an unusually long pre-clinical period in turkeys. *BMC Veterinary Research* 2016;12:260.
250. Dee S, Deen J, Rossow K, et al. Mechanical transmission of porcine reproductive and respiratory syndrome virus throughout a coordinated sequence of events during cold weather. *Canadian Journal of Veterinary Research* 2002;66:232-239.
251. USDA: APHIS: VS: CEAH. Appendix 5 of An Assessment of the Risk Associated with the Movement of Broiler Hatching Eggs Into, Within, and Out of a Control Area During a Highly Pathogenic Avian Influenza Outbreak. Oct 2012. Fort Collins, CO, 2012.
252. Dee S, Deen J, Rossow K, et al. Mechanical transmission of porcine reproductive and respiratory syndrome virus throughout a coordinated sequence of events during warm weather. *Canadian journal of veterinary research* 2003;67:12.
253. Davison S, Dufour-Zavala L, Garcia M, et al. Vaccinal laryngotracheitis—overview in the United States. Proc 109th Annual Meeting of the United States Animal Health Association, Hershey, PA 2005;580.
254. Dufour-Zavala L. Epizootiology of infectious laryngotracheitis and presentation of an industry control program. *Avian diseases* 2008;52:1-7.

255. Volkova V, Thornton D, Hubbard SA, et al. Factors Associated with Introduction of Infectious Laryngotracheitis Virus on Broiler Farms During a Localized Outbreak. *Avian diseases* 2012;56:521-528.
256. Nishiguchi A, Kobayashi S, Yamamoto T, et al. Risk Factors for the Introduction of Avian Influenza Virus into Commercial Layer Chicken Farms During the Outbreaks Caused by a Low-Pathogenic H5N2 Virus in Japan in 2005. *Zoonoses and Public Health* 2007;54:337-343.
257. Busani L, Valsecchi MG, Rossi E, et al. Risk factors for highly pathogenic H7N1 avian influenza virus infection in poultry during the 1999–2000 epidemic in Italy. *The Veterinary Journal* 2009;181:171-177.
258. Weaver JT, Malladi S, Goldsmith TJ, et al. Impact of Virus Strain Characteristics on Early Detection of Highly Pathogenic Avian Influenza Infection in Commercial Table-Egg Layer Flocks and Implications for Outbreak Control. *Avian diseases* 2012;56:905-912.
259. Vieira AR, Hofacre CL, Smith JA, et al. Human contacts and potential pathways of disease introduction on Georgia poultry farms. *Avian Diseases* 2009;53:55-62.
260. USDA:APHIS:VS:STAS:CEAH. Epidemiologic and Other Analyses of Indiana HPAI/LPAI- Affected Poultry Flocks: March 18, 2016 Report. 2016:56.
261. USDA: APHIS: VS: UMN CAHFS. Highly pathogenic avian influenza Secure broiler supply plan, Foreign Animal Disease Preparedness & Response Plan FAD PRoP, National Animal Health Emergency Management System. 2015.
262. USDA APHIS VS CEAH U. Highly Pathogenic Avian Influenza Secure Turkey Supply Plan, Turkey Sector Working Group. 2015.
263. Halvorson DA, Hueston WD. The development of an exposure risk index as a rational guide for biosecurity programs. *Avian diseases* 2006;50:516-519.
264. Leibler JH, Carone M, Silbergeld EK. Contribution of company affiliation and social contacts to risk estimates of between-farm transmission of avian influenza. *PLoS ONE* 2010;5:e9888.
265. Dorea F, Vieira A, Hofacre C, et al. Stochastic model of the potential spread of highly pathogenic avian influenza from an infected commercial broiler operation in Georgia. *Avian Diseases* 2010;54:713-719.
266. USDA: APHIS: VS. Poultry Industry Manual FAD PRoP Foreign Animal Disease Preparedness & Response Plan. 2013.
267. Halvorson DA. Prevention and management of avian influenza outbreaks: experiences from the United States of America. *Rev Sci Tech* 2009;28:359-369.
268. Dunn PA, Wallner-Pendleton EA, Lu H, et al. Summary of the 2001-02 Pennsylvania H7N2 Low Pathogenicity Avian Influenza Outbreak in Meat Type Chickens. *Avian Diseases* 2003;47:812-816.
269. Wood JP, Choi YW, Chappie DJ, et al. Environmental persistence of a highly pathogenic avian influenza (H5N1) virus. *Environmental Science & Technology* 2010;44:7515-7520.
270. Tiwari A, Patnayak DP, Chander Y, et al. Survival of two avian respiratory viruses on porous and nonporous surfaces. *Avian Dis* 2006;50:284-287.
271. Glanville Wd, Idris S, Costard S, et al. A quantitative risk assessment for the onward transmission of highly pathogenic avian influenza H5N1 from an infected small-scale broiler farm in Bogor, West Java, Indonesia. *Pro-Poor HPAI Risk Reduction*, 2010.

272. Ansari SA, Springthorpe VS, Sattar SA, et al. Potential role of hands in the spread of respiratory viral infections: studies with human parainfluenza virus 3 and rhinovirus 14. *J Clin Microbiol* 1991;29:2115-2119.
273. Glass SE, Naqi SA, Grumbles LC. Isolation of Avian Influenza-Virus in Texas. *Avian Diseases* 1981;25:545-549.
274. Utterback W. Update on avian influenza through February 21, 1984 in Pennsylvania and Virginia. Proceedings-Western Poultry Disease Conference (USA) 1984.
275. Dorea F, Berghaus R, Hofacre C, et al. Survey of biosecurity protocols and practices adopted by growers on commercial poultry farms in Georgia, US A. *Avian diseases* 2010;54:1007-1015.
276. Beard C, Brugh M, Johnson D. Laboratory studies with the Pennsylvania avian influenza viruses (H5N2). Proceedings annual meeting-United States Animal Health Association (USA) 1984.
277. Kurmi B, Murugkar H, Nagarajan S, et al. Survivability of highly pathogenic avian influenza H5N1 virus in poultry faeces at different temperatures. *Indian Journal of Virology* 2013;24:272-277.
278. Burns T, Guerin M, Kelton D, et al. On-farm Study of Human Contact Networks to Document Potential Pathways for Avian Influenza Transmission between Commercial Poultry Farms in Ontario, Canada. *Transboundary and emerging diseases* 2011;58:510-518.
279. Malone B. Catastrophic mortality management. 2005 Pennsylvania Poultry Sales and Service Conference 2005.
280. Sander JE, Warbington MC, Myers LM. Selected methods of animal carcass disposal. *Journal of the American veterinary medical association* 2002;220:1003-1005.
281. Senne DA, Panigrahy B, Morgan RL. Effect of composting poultry carcasses on survival of exotic avian viruses: highly pathogenic avian influenza (HPAI) virus and adenovirus of egg drop syndrome-76. *Avian diseases* 1994;38:733-737.
282. Gear DA, Dusek RJ, Walsh DP, et al. No Evidence of Infection or Exposure to Highly Pathogenic Avian Influenzas in Peridomestic Wildlife on an Affected Poultry Facility. *Journal of Wildlife Diseases* 2016.
283. Blake JP, Donald JO. Alternatives for the disposal of poultry carcasses. *Poultry Science* 1992;71:1130-1135.
284. Blake J, Donald J. Rendering - A disposal method for dead birds In: System ACE, ed, 1995.
285. Alabama Department of Agriculture and Industries, Alabama Department of Environmental Management. Authorized Methods of Poultry Mortality Management, 2010.
286. Elving J, Emmoth E, Albihn A, et al. Composting for avian influenza virus elimination. *Applied and environmental microbiology* 2012;78:3280-3285.
287. Ahmed ZA, Hussin H, Rohaim M, et al. Efficacy of composting dead poultry and farm wastes infected with avian influenza virus H5N1. *Am Eurasian J Agric Environ Sci* 2012;12:588-596.
288. Guan J, Chan M, Grenier C, et al. Survival of avian influenza and Newcastle disease viruses in compost and at ambient temperatures based on virus isolation and real-time reverse transcriptase PCR. *Avian Diseases* 2009;53:26-33.
289. Tablante NL, Malone GW. Controlling avian influenza through in-house composting of depopulated flocks: Sharing Delmarva's experience. Proceedings of 2006 National Symposium on Carcass Disposal 2006.

290. Lipatov AS, Kwon YK, Pantin-Jackwood MJ, et al. Pathogenesis of H5N1 influenza virus infections in mice and ferret models differs according to respiratory tract or digestive system exposure. *Journal of Infectious Diseases* 2009;199:717-725.
291. Reperant LA, Van Amerongen G, van de Bildt MW, et al. Highly pathogenic avian influenza virus (H5N1) infection in red foxes fed infected bird carcasses. *Emerging infectious diseases* 2008;14:1835.
292. Vahlenkamp TW, Teifke JP, Harder TC, et al. Systemic influenza virus H5N1 infection in cats after gastrointestinal exposure. *Influenza and Other Respiratory Viruses* 2010;4:379-386.
293. Songserm T, Amonsin A, Jam-on R, et al. Fatal avian influenza A H5N1 in a dog. *Emerging infectious diseases* 2006;12:1744.
294. Root JJ, Bentler KT, Shriner SA, et al. Ecological routes of avian influenza virus transmission to a common mesopredator: an experimental evaluation of alternatives. *PLoS one* 2014;9:e102964.
295. Root JJ, Shriner SA, Ellis JW, et al. When fur and feather occur together: interclass transmission of avian influenza A virus from mammals to birds through common resources. *Sci Rep* 2015;5:14354.
296. Root JJ, Shriner SA, Bentler KT, et al. Extended Viral Shedding of a Low Pathogenic Avian Influenza Virus by Striped Skunks (*Mephitis mephitis*). *PLoS one* 2014;9:e70639.
297. Root JJ, Bosco-Lauth AM, Bielefeldt-Ohmann H, et al. Experimental infection of peridomestic mammals with emergent H7N9 (A/Anhui/1/2013) influenza A virus: Implications for biosecurity and wet markets. *Virology* 2016;487:242-248.
298. Reperant L, Rimmelzwaan G, Kuiken T. Avian influenza viruses in mammals. *Revue scientifique et technique* 2009;28:137.
299. Tesky JL. *Vulpes vulpes*. *Fire Effects Information System*: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory, 1995.
300. Kern Jr. WH. Northern Raccoon: University of Florida IFAS Extension, 2012.
301. Georgia Department of Natural Resources WRD. Opossum Fact Sheet, 2006.
302. Kiiskila J. *Mephitis mephitis* In: University of Michigan MoZ, ed. *Animal Diversity Web*, 2014.
303. Conservation ODoW. Turkey Vulture, 2011.
304. USDA APHIS UoM, CAHFS. Highly Pathogenic Avian Influenza, Secure Broiler Supply Plan, 2015.
305. Wilkinson K. The biosecurity of on-farm mortality composting. *Journal of Applied Microbiology* 2007;102:609-618.
306. Halvorson DA. Risk reduction when building or remodeling a poultry farm In: Owen RL, Barger K, eds. *A practical guide for managing risk in poultry production*: American Association of Avian Pathologists, 2011;26-31.
307. Zhang C, Xuan Y, Shan H, et al. Avian influenza virus H9N2 infections in farmed minks. *Virology Journal* 2015;12:1-8.
308. Hookham M. Caring for mink in third generation: Agri-View, 2015.
309. Graitcer P. Plucky Former Poultry Farmer Goes Wild For Gators: National Public Radio, 2012.

310. Keawcharoen J, Oraveerakul K, Kuiken T, et al. Avian Influenza H5n1 in Tigers and Leopards. *Emerging Infectious Diseases* 2004;10:2189-2191.
311. Fur Commission USA. Super Duper Recyclers | Fur Commission USA, 1999.
312. UMN:CAHFS. Epidemiologic Study of Highly Pathogenic Avian Influenza H5N2 among Turkey Farms: University of Minnesota, Center for Animal Health and Food Safety, 2015.
313. Halvorson D. Prevention and management of avian influenza outbreaks: experiences from the United States of America. *Revue scientifique et technique* 2009;28:R O'Connor, Personal Communication.
314. Thiry E, Zicola A, Addie D, et al. Highly pathogenic avian influenza H5N1 virus in cats and other carnivores. *Veterinary Microbiology* 2007;122:25-31.
315. Davis LM, Spackman E. Do crocodilians get the flu? Looking for influenza A in captive crocodilians. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* 2008;309A:571-580.
316. USDA: APHIS: VS: CEAH, University of Minnesota Center for Animal Health and Food Safety, Egg Sector Working Group. An Assessment of the Risk Associated with the Movement of Nest Run Eggs Into, Within, and Outside of a Control Area During a Highly Pathogenic Avian Influenza Outbreak. Fort Collins, CO, 2010.
317. US Fox Shippers Council. Feeding: US Fox Shippers Council.
318. Canadian Food Inspection Agency. Section 3: Operational Management - National Farm-Level Mink Biosecurity Standard - Animals - Canadian Food Inspection Agency: Canadian Food Inspection Agency, 2013.
319. USDA-NASS-Agricultural Statistics Board. Mink 07/06/2012 - USDA\_Mink-Production-Report-July062012-2011.pdf In: Service USD oANAS, ed, 2012.
320. Yagyu K, Yanagawa R, Matsuura Y, et al. Contact infection of mink with influenza A viruses of avian and mammalian origin. *Arch Virol* 1981;68.
321. Garber L, Voelker L, Hill G, et al. Description of live poultry markets in the United States and factors associated with repeated presence of H5/H7 low-pathogenicity avian influenza virus. *Avian diseases* 2007;51:417-420.
322. Sheta BM, Fuller TL, Larison B, et al. Putative human and avian risk factors for avian influenza virus infections in backyard poultry in Egypt. *Veterinary microbiology* 2014;168:208-213.
323. Code of Federal Regulations. Title 40, Protection of Environment, 40CFR1.258, 2005;Criteria for municipal solid waste landfills, Subpart C
324. Poss PE, Friendshuh KA, Ausherman LT. The control of avian influenza. *Avian Diseases* 2003;47, Special Issue:318-326.
325. Halvorson DA, Frame DD, Friendshuh KA, et al. Outbreaks of low pathogenicity avian influenza in USA. *Avian diseases* 2003;36-46.
326. Van Buskirk MA. Control of Avian Influenza from the Perspective of State Government. *Avian Diseases* 2003;47:347-357.
327. Poetri O, Bouma A, Claassen I, et al. A single vaccination of commercial broilers does not reduce transmission of H5N1 highly pathogenic avian influenza. *Veterinary research* 2011;42:1.
328. Weaver JT, Malladi S, Bonney PJ, et al. A Simulation Based Evaluation of Pre-movement Active Surveillance Protocol Options for the Managed Movement of Turkeys to



Slaughter during an Outbreak of Highly Pathogenic Avian Influenza in the United States. *Avian Diseases* 2015.

329. Graiver DA, Topliff CL, Kelling CL, et al. Survival of the avian influenza virus (H6N2) after land disposal. *Environmental Science & Technology* 2009;43:4063-4067.
330. Shahid MA, Abubakar M, Hameed S, et al. Avian influenza virus (H5N1); effects of physico-chemical factors on its survival. *Virology* 2009;6:38.
331. Brown JD, Swayne DE, Cooper RJ, et al. Persistence of H5 and H7 avian influenza viruses in water. *Avian Dis* 2007;51:285-289.
332. World Organization of Animal Health (OIE). Terrestrial Animal Health Code, Chapter 10.4 Infection With Avian Influenza Viruses, 2016.
333. USDA. FY2016 HPAI Response Using Heat Treatment for Virus Elimination In: Agriculture USDo, ed, 2016.
334. Nazir J, Haumacher R, Ike AC, et al. Persistence of avian influenza viruses in lake sediment, duck feces, and duck meat. *Applied and environmental microbiology* 2011;77:4981-4985.
335. Lu H, Castro AE, Pennick K, et al. Survival of Avian Influenza Virus H7N2 in SPF Chickens and Their Environments. *Avian Diseases* 2003;47:1015-1021.
336. Webster RG, Yakhno M, Hinshaw VS, et al. Intestinal influenza: replication and characterization of influenza viruses in ducks. *Virology* 1978;84:268-278.
337. Nazir J, Haumacher R, Ike A, et al. Long-term study on tenacity of avian influenza viruses in water (distilled water, normal saline, and surface water) at different temperatures. *Avian Diseases* 2010;54:720-724.
338. Zarkov IS. Survival of avian influenza viruses in filtered and natural surface waters of different physical and chemical parameters. *Revue de médecine vétérinaire* 2006;157:471.
339. Thomas C, Swayne DE. Thermal inactivation of H5N1 high pathogenicity avian influenza virus in naturally infected chicken meat. *Journal of Food Protection®* 2007;70:674-680.
340. Wanaratana S, Tantilertcharoen R, Sasipreeyajan J, et al. The inactivation of avian influenza virus subtype H5N1 isolated from chickens in Thailand by chemical and physical treatments. *Vet Microbiol* 2010;140:43-48.
341. Zou S, Guo J, Gao R, et al. Inactivation of the novel avian influenza A (H7N9) virus under physical conditions or chemical agents treatment. *Virology journal* 2013;10:1.
342. Greatorex JS, Digard P, Curran MD, et al. Survival of Influenza A (H1N1) on Materials Found in Households: Implications for Infection Control. *PLoS ONE* 2011;6:e27932.
343. Noyce JO, Michels H, Keevil CW. Inactivation of influenza A virus on copper versus stainless steel surfaces. *Appl Environ Microbiol* 2007;73:2748-2750.
344. Sakaguchi H, Wada K, Kajioka J, et al. Maintenance of influenza virus infectivity on the surfaces of personal protective equipment and clothing used in healthcare settings. *Environmental Health and Preventive Medicine* 2010;15:344-349.
345. McDevitt J, Rudnick S, First M, et al. Role of absolute humidity in the inactivation of influenza viruses on stainless steel surfaces at elevated temperatures. *Applied and Environmental microbiology* 2010;76:3943-3947.
346. Pelzel AM, McCluskey BJ, Scott AE. Review of the highly pathogenic avian influenza outbreak in Texas, 2004. *Journal of the American Veterinary Medical Association* 2006;228:1869-1875.

347. Senne D, Holt T, Akey B. An overview of the 2002 outbreak of low-pathogenic H7N2 avian influenza in Virginia, West Virginia and North Carolina. *Frontis* 2005;8:41-47.
348. Mannelli A, Ferrè N, Marangon S. Analysis of the 1999–2000 highly pathogenic avian influenza (H7N1) epidemic in the main poultry-production area in northern Italy. *Preventive Veterinary Medicine* 2006;73:273-285.
349. Halvorson DA, Karunakaran D, Newman JA. Avian Influenza in Caged Laying Chickens. *Avian Diseases* 1980;24:288-294.
350. Cimorelli AJ, Perry SG, Venkatram A, et al. AERMOD: A dispersion model for industrial source applications. Part I: General model formulation and boundary layer characterization. *Journal of applied meteorology* 2005;44:682-693.
351. US Environmental Protection Agency (EPA). Support Center for Regulatory Atmospheric Modeling (SCRAM) Preferred/Recommended Models. [https://www3.epa.gov/scram001/dispersion\\_prefrec.htm](https://www3.epa.gov/scram001/dispersion_prefrec.htm), 2016.
352. Lasiewski RC, Calder WA. A preliminary allometric analysis of respiratory variables in resting birds. *Respiration physiology* 1971;11:152-166.
353. Burns R, Li H, Moody L, et al. Quantification of particulate emissions from broiler houses in the southeastern United States. Bonn: International Commission of Agricultural Engineering (CIGR), Institut fur Landtechnik, 2008;unpaginated.
354. EPA USEPA. Particulate Matter (PM), 2016.
355. Redwine J, Lacey R, Mukhtar S, et al. Concentration and emissions of ammonia and particulate matter in tunnel-ventilated broiler houses under summer conditions in Texas. *Transactions of the ASAE* 2002;45:1101-1109.
356. Spekrijse D, Bouma A, Koch G, et al. Quantification of dust-borne transmission of highly pathogenic avian influenza virus between chickens. *Influenza and Other Respiratory Viruses* 2012:no-no.
357. Roumeliotis TS, Dixon BJ, Van Heyst BJ. Characterization of gaseous pollutant and particulate matter emission rates from a commercial broiler operation part I: Observed trends in emissions. *Atmospheric Environment* 2010;44:3770-3777.
358. Wathes CM, Holden MR, Sneath RW, et al. Concentrations and emission rates of aerial ammonia, nitrous oxide, methane, carbon dioxide, dust and endotoxin in UK broiler and layer houses. *British Poultry Science* 1997;38:14-28.
359. Li H, Xin, H., Burns, R., Hoff, S., Harmon, J., Jacobson, L., Noll, S., Koziel, J. Ammonia and PM Emissions from a tom turkey barn in Iowa. *2008 ASABE Annual International Meeting*. Providence, Rhode Island, 2008.
360. Takai H, Pedersen S, Johnsen JO, et al. Concentrations and Emissions of Airborne Dust in Livestock Buildings in Northern Europe. *Journal of Agricultural Engineering Research* 1998;70:59-77.
361. Li H, Burns R. Particulate Matter Emissions from a High-rise Layer House in Iowa. *2009 ASABE Annual International Meeting*. Reno, Nevada, 2009.
362. Roumeliotis TS, Van Heyst BJ. Size fractionated particulate matter emissions from a broiler house in Southern Ontario, Canada. *Science of the Total Environment* 2007;383:174-182.
363. Li H, Xin H, Burns RT, et al. Effects of bird activity, ventilation rate and humidity on PM10 concentration and emission rate of a turkey barn 2008.

364. Cambra-López M, Hermosilla T, Lai H, et al. Particulate matter emitted from poultry and pig houses: source identification and quantification. *Transactions of the ASABE* 2011;54:629-642.
365. USDA: APHIS: VS: STAS: CEAH. An Assessment of the Risk Associated with the Movement of Day-Old Turkey Poults into, within, and out of a Control Area During a Highly Pathogenic Avian Influenza Outbreak. March, 2014, Collaboration with University of Minnesota, Center for Animal Health and Food Safety and Turkey Sector Working Group, 2014.
366. Bos ME, Nielen M, Koch G, et al. Back-calculation method shows that within-flock transmission of highly pathogenic avian influenza (H7N7) virus in the Netherlands is not influenced by housing risk factors. *Prev Vet Med* 2009;88:278-285.
367. USDA:APHIS:VS:CEAH. Estimating transmission rate parameters: new and old approaches applied to synthetic and outbreak data. Fort Collins, CO, 2014.
368. Spackman E, Weaver J.T., Malladi. S. Detection of H5 and H7 highly pathogenic avian influenza virus with lateral flow devices: Performance with healthy, sick, and dead chickens. Oral presentation. American Association of Veterinary Laboratory Diagnosticians, 57th annual meeting; October 16-22, 2014 2014.
369. R Development Core Team. R: a language environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, 2010.
370. Nash JC. On best practice optimization methods in R. *Journal of Statistical Software* 2014;60:1-14.
371. Nash JC, Varadhan R. Unifying optimization algorithms to aid software system users: optimx for R. *Journal of Statistical Software* 2011;43:1-14.
372. Spackman E, Senne D, Bulaga L, et al. Development of real-time RT-PCR for the detection of avian influenza virus. *Avian diseases* 2003;47:1079-1082.
373. Marché S, Van Den Berg T. Evaluation of rapid antigen detection kits for the diagnosis of highly pathogenic avian influenza H5N1 infection. *Avian diseases* 2010;54:650-654.
374. Soliman M, Selim A, Coward VJ, et al. Evaluation of two commercial lateral flow devices (LFDs) used for flockside testing of H5N1 highly-pathogenic avian influenza infections in backyard gallinaceous poultry in Egypt. *Journal of molecular and genetic medicine: an international journal of biomedical research* 2010;4:247.
375. Slomka MJ, To TL, Tong HH, et al. Evaluation of lateral flow devices for identification of infected poultry by testing swab and feather specimens during H5N1 highly pathogenic avian influenza outbreaks in Vietnam. *Influenza and other respiratory viruses* 2012;6:318-327.
376. Akaike H. Akaike's Information Criterion. *International Encyclopedia of Statistical Science*: Springer, 2011;25-25.