

THIS ARTICLE IS SPONSORED BY THE  
MINNESOTA DAIRY HEALTH CONFERENCE.



UNIVERSITY OF MINNESOTA

---

College of Veterinary Medicine

VETERINARY CONTINUING EDUCATION



ST. PAUL, MINNESOTA  
UNITED STATES OF MINNESOTA

# **The Role of Selenium and Vitamin E in Mastitis and Reproduction of Dairy Cattle**

**Jerry D. Olson, DVM, MS  
Clinical and Population Sciences  
College of Veterinary Medicine  
University of Minnesota**

## **Role of Selenium and Vitamin E**

Selenium was first recognized as an essential trace element in 1957 and identified as an essential component of the enzyme glutathione peroxidase (GSH-Px) in 1972. The highest concentration of selenium in the body of animals fed selenium-adequate diets is found in the kidney followed by the liver. Erythrocytes and leukocytes contains approximately three fourths of whole blood selenium. However, almost all of the selenium associated GSH-Px activity in whole blood is associated with erythrocytes (Scholz and Hutchinson, 1979). Glutathione peroxidase catalyzes the conversion of hydrogen peroxide to water and converts lipid peroxides to lipid alcohols. Glutathione peroxidase functions in the cytosol of the cell. In addition to the role of GSH-Px as antioxidant, vitamin E also functions as an intercellular and intracellular antioxidant. Vitamin E protects polyunsaturated fatty acids, enzymes, and transport proteins in membranes from free radical attack. When acting as an antioxidant, vitamin E supplies become depleted.

## **MASTITIS**

Experiments have demonstrated that the selenium and/or vitamin E status of dairy cattle can affect the incidence, duration, and severity of clinical mastitis. Smith, et al., 1984, assigned 80 non-lactating Holstein cows to one of four treatment groups: 1) Control, 2) Selenium (Se), 3) Vitamin E (E), and 4) Selenium and vitamin E (Se + E). Cows receiving selenium were injected with 0.1 mg selenium/kg of body weight i.m. 21 days prior to expected calving. Cows receiving vitamin E were supplemented with .74 g vitamin E per day in the ration. The incidence of clinical cases of mastitis per lactating quarter was reduced 12 - 37% in treated groups. The duration of clinical mastitis was reduced 13 - 41%. The predominance of clinical mastitis in this trial was associated with environmental pathogens. The plasma concentration of vitamin E during the prepartum period in the unsupplemented cows averaged 1.75 ug/ml and 2.74 ug/ml vitamin E supplemented cows. The serum concentration of selenium at calving for the control cows 0.0306 ug/ml and 0.0570 ug/ml for cows injected with selenium. Evidence from this trial would suggest that deficiencies selenium and vitamin E may increase the incidence and duration of mastitis resulting from environmental pathogens. In an epidemiological study, Erskine, et al., 1987, reported the blood selenium concentrations in dairy herds with high and low somatic cell counts. Two groups, each consisting of 16 dairy herds which were enrolled in Dairy

Herd Improvement Association, were studied to evaluate the relationship between mean blood selenium concentrations and the prevalence of mastitis in the herd. The low somatic cell (SCC) herds had 12 month mean SCC of <150,000 cells/ml and the high SCC herds had a 12 month mean herd SCC >700,000 cells/ml. The herds with low SCC had significantly higher whole blood concentrations of selenium (0.133 ug/ml) and blood GSH-Px than herds with high SCC (0.074 ug/ml). The serum concentration of vitamin A, vitamin E, and beta-carotene were not significantly different between low and high SCC herds. The herds with high SCC had higher prevalence of Streptococcus agalactia and Staphylococcus aureus. Erskine, et al., 1989, experimentally challenged quarters of 20 Holstein heifers with coliform organisms in the 14th week of lactation. The heifers were divided into two groups; one group of 10 Holstein heifers was fed a selenium-deficient diet (0.04 mg of Se/kg of total ration on a dry matter basis) beginning 3 months before calving and continuing through the first lactation and one group of 10 heifers fed a diet supplemented with 2 mg selenium per day. The peak bacterial concentration in milk was higher and the duration of infection was longer for selenium-deficient cows. In addition, selenium-supplementation significantly reduced the number of agalactic and atrophied quarters compared to selenium-deficient heifers. Erskine, et al., 1990, experimentally challenged quarters of 10 selenium-deficient (0.04 ppm Se in ration on a dry matter basis) cows and 10 selenium-supplemented (0.14 ppm Se in ration on a dry matter) cows with Staphylococcus aureus. Infections were established in 14 of the 16 quarters from selenium-deficient cows and in 16 of the 19 quarters from selenium-supplemented cows. Peak bacterial concentrations occurred earlier and were higher and in the selenium deficient cows. Similarly, somatic cell counts peaked earlier in selenium deficient cows. However, peak somatic cell counts did not differ between selenium-deficient and -supplemented groups. With respect to experimental staphylococcal mastitis, selenium status did not affect the percentage of challenge exposures resulting in infection, the duration of infection, or the severity of experimentally induced staphylococcal mastitis. From these experiments, the following conclusion can be drawn: 1) Selenium-supplementation of selenium-deficient cows is more effective in reducing the severity of environmental mastitis from coliform mastitis than staphylococcal mastitis. 2) Selenium-supplementation of selenium-deficient cows can reduce the severity of clinical signs including duration of febrile response, duration of clinical signs, and glandular atrophy and agalactia. 3) The mean whole blood selenium concentrations were significantly higher in herds with low somatic cell counts than in herds with high somatic cell counts. This suggests that low blood selenium concentrations are a risk factor for a higher prevalence of mastitis in the herd.

## **REPRODUCTION**

### **Retained Fetal Membranes**

In a review of literature that summarized greater than 60,000 calvings, the average incidence of retained fetal membranes (RFM) was 10.3%. The causes of RFM can be divided into nutritional and non-nutritional categories. Selenium and vitamin E deficiencies are important nutritional causes of RFM. Trinder, et al., first reported on the possible role of selenium in RFM in 1969. The incidence of RFM was higher at 42%

(19/45) in untreated herd mates compared to 0% (0/20) in cows treated one month before calving with an injection of 15 mg selenium and 680 IU vitamin E i.m.. In a later report (Trinder, et al., 1973), the incidence of RFM for untreated cows was 44% (35/79), 5% (7/61) for cows treated with 15 mg Se and 680 IU vitamin E i.m., and 29% (9/31) for cows treated with only 15 mg Se i.m.. The concentration of selenium in the diet of these cows was between 0.026 and 0.039 ppm of dry matter. The results of these trials suggest the incidence of RFM may further decrease when vitamin E supplementation is added. Gwazdauskas, et al., 1979, randomly assigned 351 dairy cows to be either controls or to receive an injection of 21.9 mg Se and 500 mg vitamin E at 28 to 30 days prior to projected calving. The incidence of RFM in control cows was 10% (19/190) and 13.1% (21/160) for Se-vitamin E treated cows. This study did not report either the dietary intake of selenium or the concentration of selenium in plasma, serum or blood in control or treated cows. This study suggests there is no advantage to selenium-supplementation in a herd with a low incidence of RFM. Segerson, et al., 1981, reported the results of a study where prepartum Holstein cows in four North Carolina herds were injected with 50 mg Se in combination with 680 IU vitamin E. The incidence of RFM was reduced in cows having borderline deficiencies of selenium prior to supplementation, but the incidence was not reduced in cows without deficiencies or in cows extremely deficient prior to selenium supplementation. Eger, et al., 1985, reported on series of five experiments involving selenium and vitamin E injected i.m. 3 weeks prepartum to Israeli-Holstein cows. The mean incidence of RFM for 348 control multiparous cows was 27.6% and 16.5% for control primiparous cows. Selenium with or without vitamin E significantly reduced the incidence of RFM to 14.7% in 428 multiparous cows and 7.0% in 186 primiparous cows. Selenium in the prepartum diet ranged from 0.035 to 0.109 ppm in the prepartum diet. Selenium alone was at least as effective as a combination of selenium and vitamin E in this study. Hidioglou, et al., evaluated the incidence of RFM in 627 parturitions. The herd was divided into 3 groups: 1) Control (n=217 cows), 2) Cows injected with 45 mg Se and 2040 IU of vitamin E (n=190), 3) Cows treated Se-containing intraruminal boluses (n=220). The incidence of RFM (22.1%) was not reduced either selenium supplement. The plasma selenium concentration at parturition in control cows was 0.0684 ug/ml and averaged 0.085 ug/ml in selenium-supplemented cows. Plasma selenium concentration were adequate in control cows in this study, thus explaining the failure of selenium-supplementation to reduce the incidence of RFM. The following conclusions can be drawn from these studies relative to the benefit of selenium supplementation: 1) Prepartum diets must be selenium deficient (<0.06 ppm of dry matter) if cattle are to benefit from selenium supplementation. 2) A benefit of selenium-supplementation on decreasing the incidence of RFM will not likely occur unless the incidence of RFM is greater than 20%. 3) The incidence of RFM will decrease with selenium-supplementation only if serum or whole blood selenium concentration are deficient before supplementation and Se-adequate post-supplementation. 4) Dietary supplementation of vitamin E of cows that deficient may reduce the incidence of RFM, although the evidence of the benefit is not as strong as with Se. 5) The amount of selenium supplemented by injection to dry cows has ranged from 2.3 to 50 mg Se. 6) To effectively evaluate the benefit of selenium and vitamin E supplementation, the selenium and vitamin E status of the animal must be determined both pre- and post-supplementation.

## Cystic Ovarian Disease

Harrison, et al., observed that cumulative incidence of cystic ovarian disease through the 14th post-partum week was 19% for cows supplemented with Se and 47% for cows not treated with selenium. Vitamin E status of these cows did not affect the incidence of cystic ovarian disease.

## Fertility

There is limited information on the effect of selenium and vitamin E supplementation on fertility.

## EVALUATION OF SELENIUM STATUS OF CATTLE

The clinician can evaluate the selenium status of animals either directly through laboratory determination of selenium concentrations in serum, plasma, or whole blood or glutathione peroxidase activity in whole blood, or indirectly through feed or ration analysis. The concentration of selenium in whole blood is about three times the concentration selenium in serum or plasma. Since the concentration of selenium in erythrocytes is higher than in serum or plasma, hemolysis will result in an increase in serum or plasma selenium concentrations. Current recommendations suggest that whole blood be collected in either EDTA or heparin and used to estimate the Se status in cows (Maas, et al., 1992). Since the life span of the bovine erythrocyte is 160 days and circulating erythrocytes do not have ability to incorporate selenium into Se-dependent GSH-Px, whole blood selenium is an excellent indicator of long-term selenium status in cattle.

**Table 1., Evaluation of Selenium Status**

<b>Selenium Status</b>	<b>Serum ug/ml</b>	<b>Whole Bld Se ug/ml</b>
Deficient	<0.04	<0.11
Marginal	0.04 - 0.07	0.11 - 0.2
Adequate	>0.07	>0.2

Determining serum vitamin E concentrations may be the most efficient method of determining the vitamin E status of cattle.

**Table 2. Evaluation of Vitamin E Status**

Vitamin E Status	Serum ug/ml
Deficient	< 1.5
Marginal	1.5 - 4
Adequate	> 4

### SELENIUM SUPPLEMENTATION

The literature would suggest the selenium requirement for most species of livestock is between 0.05-0.3 ppm (mg/kg) of dietary dry matter. The current requirement for all classes of dairy cattle is 0.30 ppm (NRC, 1989). Selenium requirements are affected by dietary vitamin E intake, by disease process being prevented, and by the concentration of dietary antagonists including sulfur, iron, copper, and calcium.

#### Selected References:

Eger, S., D. Drori, I. Kadoori, N. Miller, and H. Schindler. 1985. Effects of selenium and vitamin E on incidence of retained placenta. *J Dairy Sci* 68:2119-2122.

Erskine, R.J., R.J. Eberhart, L.J. Hutchinson, and R.W. Scholz. 1987. Blood selenium concentrations and glutathione peroxidase activities in dairy herds with high and low somatic cell counts. *JAVMA* 190:1417-1421.

Erskine, R.J., R.J. Eberhart, L.J. Hutchinson, and S.B. Spencer. 1987. Herd management and prevalence of mastitis in dairy herds with high and low somatic cell counts. *JAVMA* 190:1411-1416.

Erskine, R.J., R.J. Eberhart, P.J. Grasso, and R.W. Scholz. 1989. Induction of *Escherichia coli* mastitis in cows fed selenium-deficient or selenium-supplemented diets. *Am J Vet Res* 50:2093-2100.

Erskine, R.J., R.J. Eberhart, and R.W. Scholz. 1990. Experimentally induced *Staphylococcus aureus* mastitis in selenium-deficient and selenium-supplemented dairy cows. *Am J Vet Res* 51:1107-1111.

Harrison, J.H., D.D. Hancock, and H.R. Conrad. 1984. Vitamin E and selenium for reproduction of the dairy cow. *J Dairy Sci* 67:123-132.

Hidiroglou, M., A.J. McAllister, and C.J. Williams. 1987. Prepartum supplementation of selenium and vitamin E to dairy cows: Assessment of selenium status and reproductive performance. *J Dairy Sci* 70:1281-1288.

Maas, J., 1983. Diagnosis and management of selenium-responsive diseases in cattle. *Comp Cont Ed* 7:S393-S400.

Maas, J., F.D. Galey, J.R. Peauroi, J.T. Case, E.S. Littlefield, C.C. Gay, L.D. Koller, R.O. Crisman, D.W. Weber, D.W. Warner, and M.L. Tracy. 1992. The correlation between serum selenium and blood selenium in cattle. *J Vet Invest* 4:48-52

Olson, J.D., 1992. Health and reproductive aspects of the peripartum cow. *Feedin and Managing for Improved Profitability, Four States Dairy Extension Program, Dec 8-11, 1992.*

Scholz, R., and L.J. Hutchinson. 1979. Distribution of glutathione peroxidase activity and selenium in the blood of dairy cows. *Am J Vet Res* 40:245-249.

Segerson, E.C., G.J. Riviere, H.L. Dalton, and M.D. Whitacre. 1981. Retained placenta of Holstein cows treated with selenium and vitamin E. *J Dairy Sci* 64:1833-1836.

Smith, K.L., J.H. Harrison, D.D. Hancock, D.A. Todhunter, and H.R. Hunter. 1984. Effect of vitamin E and selenium supplementation on incidence of clinical mastitis and duration of clinical symptoms. *J Dairy Sci* 67:1293-1300.

Smith, K.L., J.S. Hogan, and H.R. Conrad. 1988. Selenium in dairy cattle: Its role in disease resistance. *Vet Med* 83:72-78.

Trinder, N., R.J. Hall, and C.P. Renton. 1973. The relationship between intake of selenium and vitamin E on the incidence of retained placentae in dairy cows. *Vet Rec* 93:641-644.

Trinder, N., C.D. Woodhouse, and C.P. Renton. 1969. The effect of vitamin E and selenium on the incidence of retained placentae in dairy cows. *Vet Rec* 85:550-553.

Van Saun, R.J. 1990. Rational approach to selenium supplementation essential. *Feedstuffs* 62:15-17