

E-supplement to: Efficacy of antibiotic and non-antibiotic interventions in preventing and treating necrotic enteritis in broiler chickens: a systematic review

1. Ahiwe EU, Chang'a EP, Abdallh ME, Al-Qahtani M, Kheravii SK, Wu S, Graham H, Iji PA. Dietary hydrolysed yeast cell wall extract is comparable to antibiotics in the control of subclinical necrotic enteritis in broiler chickens. *British Poultry Science*. 60:757-765; 2019. <https://doi.org/10.1080/00071668.2019.1664727>.
2. Al-Baadani HH, Abudabos AM, Al-Mufarrej SI, Alzawqari M. Effects of dietary inclusion of probiotics, prebiotics and synbiotics on intestinal histological changes in challenged broiler chickens. *South African Journal of Animal Sciences*. 46:157-165; 2016. <https://doi.org/10.4314/sajas.v46i2.6>.
3. Alizadeh M, Rogiewicz A, McMillan E, Rodriguez-Lecompte JC, Patterson R, Slominski BA. Effect of yeast-derived products and distillers dried grains with solubles (DDGS) on growth performance and local innate immune response of broiler chickens challenged with *Clostridium perfringens*. *Avian Pathology*. 45:334-345; 2016. <https://doi.org/10.1080/03079457.2016.1155693>.
4. Ao Z, Kocher A, Choct M. Effects of dietary additives and early feeding on performance, gut development and immune status of broiler chickens challenged with *Clostridium perfringens*. *Asian-Australasian Journal of Animal Sciences*. 25:541-551; 2012. <https://doi.org/10.5713/ajas.2011.11378>.
5. Charal JW, Bidner TD, Southern LL, Janes ME, Gutierrez ME, Lavergne TA. Anise oil dosage and its effect on growth performance and jejunal lesions during a *Clostridium perfringens* challenge in battery trials, and growth performance in a floor pen trial. *Journal of Applied Poultry Research*. 26:240-252; 2017. <https://doi.org/10.3382/japr/pfw068>.
6. Cheng Y, Zhang N, Han J, Chang C, Hsiao S, Yu Y. Optimization of surfactin production from *Bacillus subtilis* in fermentation and its effects on *Clostridium perfringens*-induced necrotic enteritis and growth performance in broilers. *Journal of Animal Physiology and Animal Nutrition*. 102:1232-1244; 2018. <https://doi.org/10.1111/jpn.12937>.
7. Collier CT, Van der Klis JD, Deplancke B, Anderson DB, Gaskins HR. Effects of tylosin on bacterial mucolysis, *Clostridium perfringens* colonization, and intestinal barrier function in a chick model of necrotic enteritis. *Antimicrobial Agents and Chemotherapy*. 47:3311-3317; 2003. <https://doi.org/10.1128/AAC.47.10.3311-3317.2003>.

8. Daneshmand A, Kermanshahi H, Sekhavati MH, Javadmanesh A, Ahmadian M, Alizadeh M, Aldawoodi A. Effects of cLF chimera peptide on intestinal morphology, integrity, microbiota, and immune cells in broiler chickens challenged with necrotic enteritis. *Scientific Reports*. 10; 2020. <https://doi.org/10.1038/s41598-020-74754-x>.
9. Das Q, Islam MR, Lepp D, Tang J, Yin X, Mats L, Liu H, Ross K, Kennes YM, Yacini H, et al. Gut microbiota, blood metabolites, and spleen immunity in broiler chickens fed berry pomaces and phenolic-enriched extractives. *Frontiers in Veterinary Science*. 7; 2020. <https://doi.org/10.3389/fvets.2020.00150>.
10. Emami NK, Calik A, White MB, Young M, Dalloul RA. Necrotic enteritis in broiler chickens: The role of tight junctions and mucosal immune responses in alleviating the effect of the disease. *Microorganisms*. 7; 2019. <https://doi.org/10.3390/microorganisms7080231>.
11. Emami NK, Calik A, White MB, Kimminau EA, Dalloul RA. Effect of probiotics and multi-component feed additives on microbiota, gut barrier and immune responses in broiler chickens during subclinical necrotic enteritis. *Frontiers in Veterinary Science*. 7; 2020. <https://doi.org/10.3389/fvets.2020.572142>.
12. Geier MS, Mikkelsen LL, Torok VA, Allison GE, Olhood CG, Boulianne M, Hughes RJ, Choct M. Comparison of alternatives to in-feed antimicrobials for the prevention of clinical necrotic enteritis. *Journal of Applied Microbiology*. 109:1329-1338; 2010. <https://doi.org/10.1111/j.1365-2672.2010.04758.x>.
13. Gharib-Naseri K, Kheravii SK, Li L, Wu S. Buffered formic acid and a monoglyceride blend coordinately alleviate subclinical necrotic enteritis impact in broiler chickens. *Poultry Science*. 100; 2021. <https://doi.org/10.1016/j.psj.2021.101214>.
14. Hofacre CL, Froyman R, George B, Goodwin MA, Brown J. Use of aviguard, virginiamycin, or bacitracin MD against *Clostridium perfringens*-associated necrotizing enteritis. *Journal of Applied Poultry Research*. 7:412-418; 1998. <https://doi.org/10.2307/1592685>.
15. Hofacre CL, Beacorn T, Collett S, Mathis G. Using competitive exclusion, mannan-oligosaccharide and other intestinal products to control necrotic enteritis. *Journal of Applied Poultry Research*. 12:60-64; 2003. <https://doi.org/10.1093/japr/12.1.60>.

16. Hofacre CL, Reynolds DJ, Mathis GF, Lumpkins BS, Ollis N, Smith JA, Demey V. Effect of a competitive exclusion culture in a necrotic enteritis challenge model in broilers. *Journal of Applied Poultry Research*. 28:350-355; 2019. <https://doi.org/10.3382/japr/pfy078>.
17. Hofacre CL, Mathis GF, Lumpkins BS, Sygall R, Vaessen S, Hofacre CS, Smith JA, Clanton E. Efficacy of butyric and valeric acid esters in a necrotic enteritis challenge model. *Avian Diseases*. 64:407-414; 2020. <https://doi.org/10.1637/aviandiseases-D-19-00124>.
18. Hussein EOS, Ahmed SH, Abudabos AM, Suliman GM, Abd El-Hack ME, Swelum AA, Alowaimer AN. Ameliorative effects of antibiotic-, probiotic-and phytobiotic-supplemented diets on the performance, intestinal health, carcass traits, and meat quality of *Clostridium perfringens*-infected broilers. *Animals*. 10; 2020. <https://doi.org/10.3390/ani10040669>.
19. Jayaraman S, Das PP, Saini PC, Roy B, Chatterjee PN. Use of *Bacillus subtilis* PB6 as a potential antibiotic growth promoter replacement in improving performance of broiler birds. *Poultry Science*. 96:2614-2622; 2017. <https://doi.org/10.3382/ps/pex079>.
20. Kan L, Guo F, Liu Y, Pham VH, Guo Y, Wang Z. Probiotics *Bacillus licheniformis* improves intestinal health of subclinical necrotic enteritis-challenged broilers. *Frontiers in Microbiology*. 12; 2021. <https://doi.org/10.3389/fmicb.2021.623739>.
21. Kang M, Oh JY, Cha SY, Kim WI, Cho HS, Jang HK. Efficacy of polymers from spontaneous carotenoid oxidation in reducing necrotic enteritis in broilers. *Poultry Science*. 97:3058-3062; 2018. <https://doi.org/10.3382/ps/pey180>.
22. Keerqin C, Rhayat L, Zhang ZH, Gharib-Naseri K, Kheravii SK, Devillard E, Crowley TM, Wu SB. Probiotic *Bacillus subtilis* 29,784 improved weight gain and enhanced gut health status of broilers under necrotic enteritis condition. *Poultry Science*. 100; 2021. <https://doi.org/10.1016/j.psj.2021.01.004>.
23. Knap I, Lund B, Kehlet AB, Hofacre C, Mathis G. *Bacillus licheniformis* prevents necrotic enteritis in broiler chickens. *Avian Diseases*. 54:931-935; 2010. <https://doi.org/10.1637/9106-101509-ResNote.1>.
24. Krueger LA, Spangler DA, Vandermyde DR, Sims MD, Ayangbile GA. Avi-Lution[®] supplemented at 1.0 or 2.0 g/kg in feed improves the growth performance of broiler chickens during challenge with

bacitracin-resistant *Clostridium perfringens*. *Poultry Science*. 96:2595-2600; 2017.
<https://doi.org/10.3382/ps/pex074>.

25. Kumar A, Kheravii SK, Li L, Wu SB. Monoglyceride blend reduces mortality, improves nutrient digestibility, and intestinal health in broilers subjected to clinical necrotic enteritis challenge. *Animals*. 11; 2021a. <https://doi.org/10.3390/ani11051432>.

26. Kumar A, Toghiani M, Kheravii SK, Pineda L, Han Y, Swick RA, Wu SB. Potential of blended organic acids to improve performance and health of broilers infected with necrotic enteritis. *Animal Nutrition*. 7:440-449; 2021b. <https://doi.org/10.1016/j.aninu.2020.11.006>.

27. Li Z, Wang W, Lv Z, Liu D, Guo Y. *Bacillus subtilis* and yeast cell wall improve the intestinal health of broilers challenged by *Clostridium perfringens*. *British Poultry Science*. 58:635-643; 2017.
<https://doi.org/10.1080/00071668.2017.1370697>.

28. Lillehoj HS, Brett L, Chris C, Fang C, Greg FM, Misun J, Ron LC, Soon SP, Sung HL, Yeaseul L. Calcium montmorillonite-based dietary supplement attenuates necrotic enteritis induced by *Eimeria maxima* and *Clostridium perfringens* in broilers. *Journal of Poultry Science*. 53:329-340; 2016.
<https://doi.org/10.2141/jpsa.0150182>.

29. McDougald LR, Hofacre C, Mathis G, Fuller L, Hargrove JL, Greenspan P, Hartle DK. Enhancement of resistance to coccidiosis and necrotic enteritis in broiler chickens by dietary muscadine pomace. *Avian Diseases*. 52:646-51; 2008. <https://doi.org/10.1637/8306-041508-Reg.1>.

30. McKnight LL, Page G, Han Y. Effect of replacing in-feed antibiotics with synergistic organic acids, with or without trace mineral and/or water acidification, on growth performance and health of broiler chickens under a *Clostridium perfringens* Type A challenge. *Avian Diseases*. 64:374-378; 2020.
<https://doi.org/10.1637/aviandiseases-D-19-00115>.

31. M'Sadeq SA, Wu S, Choct M, Forder R, Swick RA. Use of yeast cell wall extract as a tool to reduce the impact of necrotic enteritis in broilers. *Poultry Science*. 94:898-905; 2015.
<https://doi.org/10.3382/ps/pev035>.

32. Musa BB, Duan Y, Khawar H, Sun Q, Ren Z, Mohamed MAE, Abbasi IHR, Yang X. *Bacillus subtilis* B21 and *Bacillus licheniformis* B26 improve intestinal health and performance of broiler chickens with

Clostridium perfringens-induced necrotic enteritis. Journal of Animal Physiology and Animal Nutrition. 103:1039-1049; 2019. <https://doi.org/10.1111/jpn.13082>.

33. Oladokun S, Koehler A, MacIsaac J, Ibeagha-Awemu EM, Adewole DI. Bacillus subtilis delivery route: effect on growth performance, intestinal morphology, cecal short-chain fatty acid concentration, and cecal microbiota in broiler chickens. Poultry Science. 100; 2021. <https://doi.org/10.1016/j.psj.2020.10.063>.

34. Sayed ML, Hanan MI, Zeinab MS, Makarita AM. Effect of zinc bacitracin and oligosaccharids on chickens vaccinated with chicken necrotic enteritis vaccine. Veterinary Medical Journal Giza. 57:449-456; 2009.

35. Sokale AO, Menconi A, Mathis GF, Lumpkins B, Sims MD, Whelan RA, Doranalli K. Effect of Bacillus subtilis DSM 32315 on the intestinal structural integrity and growth performance of broiler chickens under necrotic enteritis challenge. Poultry Science. 98:5392-5400; 2019. <https://doi.org/10.3382/ps/pez368>.

36. Tactacan GB, Schmidt JK, Miille MJ, Jimenez DR. A Bacillus subtilis (QST 713) spore-based probiotic for necrotic enteritis control in broiler chickens. Journal of Applied Poultry Research. 22:825-831; 2013. <https://doi.org/10.3382/japr.2013-00730>.

37. Vidanarachchi JK, Mikkelsen LL, Constantinoiu CC, Choct M, Iji PA. Natural plant extracts and prebiotic compounds as alternatives to antibiotics in broiler chicken diets in a necrotic enteritis challenge model. Animal Production Science. 53:1247-1259; 2013.

38. Xue G, Wu S, Choct M, Swick RA. Effects of yeast cell wall on growth performance, immune responses and intestinal short chain fatty acid concentrations of broilers in an experimental necrotic enteritis model. Animal Nutrition. 3:399-405; 2017. <https://doi.org/10.1016/j.aninu.2017.08.002>.

39. Yang Y, Wang Q, Diarra MS, Yu H, Hua Y, Gong J. Functional assessment of encapsulated citral for controlling necrotic enteritis in broiler chickens. Poultry Science. 95:780-789; 2016. <https://doi.org/10.3382/ps/pev375>.

40. Zhang G, Mathis GF, Hofacre CL, Yaghmaee P, Holley RA, Duranc TD. Effect of a radiant energy-treated lysozyme antimicrobial blend on the control of clostridial necrotic enteritis in broiler chickens. *Avian Diseases*. 54:1298-300; 2010. <https://doi.org/10.1637/9370-041410-ResNote.1>