

MN 2000
EB 392
Rev 79
C.2

Insecticide Suggestions to Control Greenhouse and Floriculture Pests in 1979

UNIVERSITY OF MINNESOTA
DOCUMENTS
MAR 26 1979
ST. PAUL CAMPUS LIBRARIES



Extension Bulletin 392 — Revised 1979
AGRICULTURAL EXTENSION SERVICE
UNIVERSITY OF MINNESOTA

This archival publication may not reflect current scientific knowledge or recommendations.
Current information available from University of Minnesota Extension: <http://www.extension.umn.edu>

Insecticide Suggestions to Control Greenhouse and Floriculture Pests in 1979

Do not use after 1979

The major crops and their major pests are:

African violets	Cyclamen mites, also spider mites, Pritchard mealybugs, occasionally aphids
Azaleas	Spider mites, cyclamen mites, whiteflies, greenhouse thrips, leaf miners, leaf rollers
Begonias	Cyclamen mites probably most serious, thrips
Cacti (succulent plants)	Mealybugs, scale insects
Camellias	Aphids, thrips, leaf miners, spider mites
Carnations	Aphids, thrips, spider mites
Chrysanthemums	Aphids, leaf miners (<i>Liriomyza sp.</i>)
Cyclamens	Cyclamen mites, spider mites, aphids
Foliage plants	Root aphids, foliar aphids, scale insects, mealybugs, spider mites, whiteflies
Geraniums	Aphids, whiteflies, plume moths (<i>Platyptilia pica</i>)
Lilies	Aphids
Orchids	Orchid flies (<i>Eurytoma orchidearum</i>), slugs
Poinsettias	Mealybugs, whiteflies, spider mites, scale insects, root aphids
Roses	Spider mites, thrips, aphids, midges, leaf rollers
Snapdragons	Spider mites, aphids, cyclamen mites, whiteflies

The following abbreviations appear throughout this publication:

W.P.—Wettable Powder

E.C.—Emulsifiable Concentrate

S.P.—Soluble Powder

Other publications in this series:

Extension Folder 412—Insecticide Suggestions to Control Household Insects.

Extension Folder 413—Insecticide Suggestions to Control Livestock and Poultry Pests.

Extension Folder 414—Insecticide Suggestions to Control Tree, Shrub, Lawn, and Turf Insects.

Extension Bulletin 387—Insecticides.

Extension Bulletin 388—Insecticide Suggestions to Control Insect Pests of Field Crops.

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Ants	SPRAYS OR DUSTS diazinon	1 to 2 lb. 50% W.P.	Apply 1 lb. of dust or 1 gal. spray per 1,000 sq. ft. soil along walks. Do not apply to foliage.

Authors of this publication are Mark E. Ascerno, assistant professor and extension entomologist; D. M. Noetzel, assistant professor and extension entomologist; L. K. Cutkomp, professor, Department of Entomology, Fisheries and Wildlife.

Certain pesticides may be marketed under brand names other than those listed and are acceptable as long as generic content is the same.

The information given in this publication is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Minnesota Agricultural Extension Service is implied.

Issued in furtherance of cooperative extension work in agriculture and home economics, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Roland H. Abraham, Director of Agricultural Extension Service, University of Minnesota, St. Paul, Minnesota 55108. The University of Minnesota, including the Agricultural Extension Service, is committed to the policy that all persons shall have equal access to its programs, facilities, and employment without regard to race, creed, color, sex, national origin, or handicap.

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks	
Aphids	AEROSOLS and VAPORS		Be sure foliage is dry at time of treatment.	
	dichlorvos (Vapona)	6 oz. smoke generator for 10,000 cu. ft.; 1 oz. 81% E.C. per 10,000 cu. ft. on steam pipes.	Can be used on cucumbers, tomatoes, and lettuce in greenhouses. Close greenhouse vents for 2 hrs. after treatment. Do not use on mum varieties Pink Champagne and Nightingale or on Shasta lily. May be phytotoxic to tomatoes.	
	naled (Dibrom)	1 fl. oz. 8 lb. E.C. per 10,000 cu. ft. on steam pipes.	Can be used on cucumbers and tomatoes in greenhouse. Close greenhouse vents for at least 2 hrs. after treatment. May injure rose varieties White Butterfly and Golden Rapture, Pink Champagne mums, Green Wandering Jew, poinsettias, and Dutchman's Pipe.	
	parathion	10% aerosol using 1 lb. per 50,000 cu. ft.	Causes leathery leaves on gardenias; not recommended on asters and white cyclamen. Can be used on vegetables in the greenhouse. For use by commercial growers only.	
	sulfotepp (Dithio)	Smoke or 5% aerosol bomb using 1 lb. per 50,000 cu. ft.		
	nicotine	Smoke generator. Check label for amount to use and additional precautions.	Can be used on vegetables in the greenhouse. May damage tender plants or tender portions of plants. Do not use on mum varieties Iceberg and Albatross. For use by commercial growers only.	
	SPRAYS			
	endosulfan (Thiodan)	1 lb. 50% W.P. or 1 qt. 24% E.C.	Also may use 3% dusts. Can be used on tomatoes in the greenhouse. May injure chrysanthemum varieties Bonnafon Deluxe, Fred Shoesmith, and White Knight, and also some geranium varieties.	
malathion	2 lb. 25% W.P. or 1½ pt. 57% E.C.	Dosage doubled for scale insects. May injure some crassula, cucurbit, fern,		

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Aphids (continued)	malathion (continued)		petunia, orchid, sweet pea, saint paulia, violet, gloxinia, red carnation, and rose varieties. Phytotoxicity greatest with E.C. Dusts also may be applied. Do not use for green peach aphid. Can be used on vegetables in the greenhouse.
	acephate (Orthene)	$\frac{1}{3}$ - $\frac{2}{3}$ lb. 75% S.P.	For use on roses, carnations, and chrysanthemums. Do not apply more than once every 28 days to carnations and chrysanthemums. Has injured chrysanthemum varieties Show Off, Iceberg, Wild Honey, Mt. Snow, Albatross, Bonnie Jean, Always Pink, Knob Hill, and Statesman. Phytotoxicity also reported on Schefflera, carnations, poinsettias, and ruffled fern at the $\frac{2}{3}$ lb. rate. Phytotoxicity greatly reduced at the lower rates. Phytotoxicity may occur on other mum varieties. Spray a few plants and observe for 2 weeks before treating entire crop.
	Enstar 5E	10 oz. 65% E.C. when aphids first noted or 20 oz. for heavy populations.	Manufacture discontinued; stocks available until depleted. Should be applied in pre-bloom stage. May injure bracts of some poinsettia varieties.
	pirimicarb (Pirimor)	4 to 8 oz. 50% W.P.	Approved on chrysanthemums. Do not apply more than four times.
	demeton (Systox)	1 to 1½ pt. 25% E.C.	May injure Croft lilies. Effective on foliar nematodes. For use by commercial growers only.
	oxydemetonmethyl (Meta-Systox-R)	1 to 1½ pt. 25% E.C.	May injure mum varieties Hurricane, Iceberg, Whitetop, and Pennant; also Easter lilies.
	SOIL TREATMENT oxydemetonmethyl (Meta-Systox-R)	1½ pt. 25% E.C. Apply ½ cup of the mixture per 6- to 8-in. pot, or use 1 Tbsp. in 3 gal. for 100 6-in. pots.	Causes leaf scorch on Easter Lily; injury noted on mum varieties Hurricane, Iceberg, Whitetop, and Pennant.
	aldicarb (Temik)	30 to 40 oz. Temik 10 G/1,000 sq. ft. on soil surface.	Should give 6 weeks' protection; approved on mums, gerberas, roses, carnations, Easter lilies, and snapdragons. For use by commercial growers only.

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Aphids (continued)	dimethoate (Cygon)	6 fl. oz. 30.5% E.C. diluted in enough water to treat 1,000 sq. ft. of bench.	For use on carnations.
	demeton (Systox)	1 to 1½ pt. 25% E.C. Apply 1 pt. of the mixture per sq. ft. of bench or ¼ pt. per 6-in. pot.	May injure African violets, Cibotium ferns, and Croft lilies. For use by commercial growers only.
Caterpillars (corn borer, leaf roller, loopers, army worms, etc.)	SPRAYS		
	carbaryl (Sevin)	2 lb. 50% W.P. or 1¼ lb. 80% sprayable.	Except for older stages of cabbage looper. May injure Boston ivy and Maidenhair fern.
	methoxychlor	2 lb. 50% W.P.	Preferred for sensitive foliage. Can be used on lettuce, cucumbers, tomatoes, and radishes in the greenhouse.
	trichlorfon (Dylox, Proxol)	20 to 30 oz. 80% S.P. or 2 to 3 pt. 40.5% E.C.	Some injury to certain varieties of carnation, zinnia, and hydrangia.
	<i>Bacillus thuringiensis</i> (Dipel, Biotrol, Thuricide)	½ to 1 lb. W.P.	May take 2 to 3 days before mortality results. Can be used on vegetables in the greenhouse.
	AEROSOLS and VAPORS		
naled (Dibrom)	1 fl. oz. 8 lb. E.C. per 10,000 cu. ft. on steam pipes.	For use against leaf rollers. Can be used on cucumbers and tomatoes in greenhouse. Close greenhouse vents for at least 2 hrs. after treatment. May injure rose varieties White Butterfly and Golden Rapture, Pink Champagne mums, Green Wandering Jew, poinsettias, and Dutchman's Pipe.	
dichlorvos (Vapona)	6 oz. smoke generator for 10,000 cu. ft.; 1 oz. 81% E.C. per 10,000 cu. ft. on steam pipes.	Can be used on cucumbers, tomatoes, and lettuce in greenhouse. Close greenhouse vents for 2 hrs. after treatment. Do not use on mum varieties Pink Champagne and Nightingale or on Shasta lily. May be phytotoxic to tomatoes.	

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Centipedes, symphylids	lindane	10 oz. 25% W.P. per 1,000 sq. ft.	Work into soil. Steam sterilization also is advised.
Cutworms	carbaryl (Sevin)	2 lb. 50% W.P., 1¼ lb. 80% W.P., or 5% bait.	Apply to soil and base of plant. May injure Boston ivy and Maidenhair fern.
	trichlorfon (Dylox, Proxol)	20 to 30 oz. 80% S.P. or 2 to 3 pt. 40.5% E.C.	Some injury to certain varieties of carnation, zinnia, and hydrangia. Apply to soil and base of plant.
Cyclamen mites	dicofol (Kelthane)	1⅓ lb. 35% W.P. or 1 pt. 18.5% E.C.	Incompatible with sulfur. Make 2 or 3 applications at 2-week intervals. Use sanitation and isolation during clean-up period.
	endosulfan (Thiodan)	1 lb. 50% W.P. or 1 qt. 24% E.C.	Can be used on tomatoes in the greenhouse. May injure chrysanthemum varieties Bonnafon Deluxe, Fred Shoemith, and White Knight; also some geranium varieties. Make 2 or 3 applications at 2-week intervals. Use sanitation and isolation during clean-up period.
Fungus gnats	malathion	4% dust or 1½ to 2 pt. 57% E.C.	Look under "Aphids" for malathion cautions. Malathion can be used on vegetables in the greenhouse.
	lindane	10 oz. 25% W.P. per 1,000 sq. ft.	Lightly drench soil for best larval control. Treat soil weekly until infestation is controlled. Surface dusting controls adults. Dichlorvos (Vapona) for aphids also will control fungus gnat adults.
	resmethrin	1% bomb using 1 lb. for 3,000 sq. ft.	For fungus gnat adults. Effectiveness reduced in sunlight and at temperatures above 72° F.
Grasshoppers	carbaryl (Sevin)	2 lb. 80% W.P.	May injure Boston ivy and Maidenhair fern.

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Grasshoppers (continued)	malathion	2 pt. 57% E.C.	Look under "Aphids" for cautions. Can be used on vegetables in the greenhouse.
Leafhoppers	carbaryl (Sevin)	2 lb. 50% W.P. or 1¼ lb. 80% sprayable.	May injure Boston ivy and Maidenhair fern.
Leaf miners	SPRAYS malathion	2 lb. 25% W.P. or 1½ pt. 57% E.C.	Look under "Aphids" for cautions. Can be used on vegetables in the greenhouse.
	oxydemetonmethyl (Meta-Systox-R)	1½ pt. 25% E.C.	Soil drench may be used as given under "Aphids." May injure mum varieties Hurricane, Iceberg, Whitetop, and Pennant; also Easter lilies.
	parathion	1 lb. of 25% W.P. or 1 pt. 25% E.C.	Avoid excessive treatment of asters, gardenias, and white cyclamen. Can be used on vegetables in the greenhouse. For use by commercial growers only.
	resmethrin	1 pt. to 1 qt. 24.3% E.C.	For <i>Liriomyza</i> leaf miner on mum.
	trichlorfon (Dylox, Proxol)	20 to 30 oz. 80% S.P. or 2 to 3 pt. 40.5% E.C.	Some injury to certain varieties of carnation, zinnia, and hydrangia.
	SOIL TREATMENT aldicarb (Temik)	30 to 40 oz. Temik 10 G/1,000 sq. ft. on soil surface (¼ tsp./6-in. pot).	For use by commercial growers only. Approved on mums and gerberas.
	Leaf rollers	acephate (Orthene)	⅓-¾ lb. 75% S.P. Also may use any insecticides listed under "Caterpillars," at dosages recommended.
Mealybugs	SPRAYS lindane	½ lb. 25% W.P. or 1½ pt. 20% E.C.	
	malathion	3 lb. 25% W.P. or 1½ pt. 57% E.C.	Look under "Aphids" for cautions. Do not use on cacti. Can be used on vegetables in the greenhouse.
	parathion	1 lb. 25% W.P.	Repeat in 3 to 4 weeks. For use by commercial growers only. Can be used on vegetables in the greenhouse. Avoid excessive treatment of asters, gardenias, and white cyclamen.

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Mealybugs (continued)	demeton (Systox)	1 to 1½ pt. 25% E.C.	May injure Croft lilies. Effective on foliar nematodes. For use by commercial growers only.
	AEROSOLS and VAPORS		
	dichlorvos (Vapona)	6 oz. smoke generator for 10,000 cu. ft. 1 oz. 81% E.C. per 10,000 cu. ft. on steam pipes.	Can be used on cucumbers, tomatoes, and lettuce in the greenhouse. Look under "Aphids" for cautions.
	malathion	15% bomb using 1 lb. per 50,000 cu. ft.	Can be used on vegetables in the greenhouse. Look under "Aphids" for cautions.
	naled (Dibrom)	1 fl. oz. 60% E.C. per 10,000 cu. ft. on steam pipes.	Can be used on cucumbers and tomatoes in the greenhouse. Look under "Aphids" for cautions.
	sulfotepp (Dithio)	15% bomb using 1 lb. per 50,000 cu. ft.	
	SOIL TREATMENT		
	aldicarb (Temik)	30 to 40 oz. Temik 10 G/1,000 sq. ft. on soil surface (¼ tsp./6-in. pot).	For use by commercial growers only. Approved on poinsettias.
Plant bugs	carbaryl (Sevin)	2 lb. 50% W.P.	May injure Boston ivy and Maidenhair fern.
	trichlorfon (Dylox, Proxol)	20 to 30 oz. 80% S.P. or 2 to 3 pt. 40.5% E.C.	Some injury to certain varieties of carnation, zinnia, and hydrangia.
Plume moth	carbaryl (Sevin)	2 lb. 50% W.P. or 1¼ lb. 80% sprayable.	Principal geranium pest. Sprays preferable to aerosols, especially on older larvae. Vapona can be used on cucumbers, tomatoes, and lettuce in the greenhouse, but may be phytotoxic to tomatoes.
	dichlorvos (Vapona)	As recommended for mealybugs.	
Roaches	diazinon	1 to 2 lb. 50% W.P.	Apply 1 lb. dust or 1 gal. of spray per 1,000 sq. ft. soil along walks. Do not apply to foliage.
Scale insects	dichlorvos (Vapona) malathion parathion sulfotepp (Dithio) demetox (Systox)	Same as for mealybugs.	3 or 4 applications, but repeat as necessary. Best control when crawlers are present. Do not use parathion for soft scales.

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Slugs, snails	Mesurool metaldehyde, or proprietary baits containing metaldehyde	1 lb. 2% bait per 1,000 sq. ft., 2 oz. 15% dust per 100 sq. ft., or 1 oz. 20% E.C. to cover 100 sq. ft.	Apply once every 2 weeks to soil, using fresh material.
Sowbugs (pillbugs)	malathion	4% dust or 1½ to 2 pt. 57% E.C.	Look under "Aphids" for cautions. Can be used on vegetables in the greenhouse.
	lindane	1 lb. 25% W.P.	
Spider mites	SPRAYS carbophenothion (Trithion)	1½ to 2 pt. 45.9% E.C.	For use by commercial growers only. May injure crassula, some ferns, and some roses. Use caution on tender plants.
	dicofol (Kelthane)	1⅓ lb. 35% W.P. or 1 pt. 18.5% E.C.	Incompatible with sulfur. May injure roses. Some mite populations may be resistant to dicofol.
	malathion	1½ pt. 57% E.C.	May injure some crassula, cucurbit, fern, petunia, orchid, sweet pea, saint paulia, violet, gloxinia, red carnation, and rose varieties. Can be used on vegetables in the greenhouse. Some mite populations may be resistant to malathion.
	oxythioquinox (Morestan)	½ to 1 lb. 25% W.P.	Certain rose varieties damaged.
	propargite (Omite)	½ to 1 lb. 30% W.P.	May injure new growth on roses.
	oxydemetonmethyl (Meta-Systox-R)	1½ pt. 25% E.C.	Soil drench may be used as given under "Aphids." May injure mum varieties Hurricane, Iceberg, Whitetop, and Pennant; also Easter lilies.
	Pentac	½ lb. 50% W.P.	May cause damage to chrysanthemum foliage. Effectiveness reduced at higher temperatures.
	Plictran	4 oz. 50% W.P.	May cause damage to mum blossoms, poinsettia bracts, and rose foliage. May be

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Spider mites (continued)	Plictran (continued)		toxic to other crops at recommended rates. Check on a few plants before treating entire crop. Slow-acting with long residual. Effectiveness reduced at higher temperatures and humidities.
	tetradifon (Tedion)	½ lb. 50% W.P. or 1 qt. 12.3% E.C.	Avoid treatment of White Butterfly and Cinderella roses. Other rose varieties also may be injured. Can be used on vegetables in greenhouse.
	demeton (Systox)	1 to 1½ pt. 25% E.C.	May injure Croft lilies. Effective on foliar nematodes. For use by commercial growers only.
	Vendex	8 to 16 oz. 50% W.P.	Do not repeat more than once every 28 days. Do not apply to chrysanthemums in bloom or poinsettias with bracts. Activity severely reduced below 55°F.
	AEROSOLS dichlorvos (Vapona)	6 oz. for 10,000 cu. ft. or paint 1 fl. oz. 81% E.C. per 10,000 cu. ft. on steam pipes.	Dichlorvos (Vapona) can be used on cucumbers, tomatoes, and lettuce in the greenhouse but may be phytotoxic to tomatoes. Look under "Aphids" for cautions.
	malathion	15% bomb using 1 lb. for 50,000 cu. ft.	
	parathion	10% aerosol using 1 lb. for 50,000 cu. ft.	Malathion and parathion can be used on vegetables in the greenhouse. Look under "Aphids" for cautions.
	sulfotepp (Dithio)	5% aerosol using 1 lb. for 50,000 cu. ft.	
	VAPORS OR SMOKES naled (Dibrom)	Paint 1 fl. oz. 60% E.C. per 10,000 cu. ft. on steam pipes.	Corrosive to metals. Look under "Aphids" for cautions. Can be used on cucumbers and tomatoes in the greenhouse.
	sulfotepp (Dithio)	As directed on label.	
	tetradifon (Tedion)	Paint 10 oz. 50% W.P. as slurry on steam pipes for	Avoid treatment of White Butterfly and Cinderella

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Spider mites (continued)	tetradifon (continued)	200,000 cu. ft.	roses. Other rose varieties also may be injured. Can be used on vegetables in the greenhouse.
	tetradifon 15% plus sulfotepp 12.5%	Smoke generator 6 oz. for 10,000 cu. ft.	Check label for cautions.
	SOIL TREATMENT aldicarb (Temik)	30 to 40 oz. Temik 10 G/1,000 sq. ft. on soil surface (1/8 tsp./6-in. pot).	Should control mites for 40 to 60 days. Approved on mums, poinsettias, gerberas, roses, carnations, orchids, and snapdragons. For use by commercial growers only.
	dimethoate (Cygon)	6 oz. 30.5% E.C. in enough water to cover 1,000 sq. ft. of bench.	Carnations only. Can also be used as a spray at 1 1/2 pt. 30.5% E.C. in 100 gal. water.
	demeton (Systox)	1 to 1 1/2 pts. 25% E.C. Apply 1 pt. of the mixture per sq. ft. of bench or 1/4 pt. per 6-in pot.	May injure African violets, Cibotium ferns, and Croft lilies. For use by commercial growers only.
Spittle bugs	lindane	1/2 lb. 25% W.P. or 1 1/2 pt. 20% E.C.	
Springtails	SOIL TREATMENT malathion	As recommended for fungus gnats.	Can be used on vegetables in the greenhouse.
Symphylids	lindane	10 oz. 25% W.P. per 1,000 sq. ft.	Work into soil. Steam sterilization also is advised.
Thrips	SPRAYS malathion	1 1/2 pt. 57% E.C.	Look under "Aphids" for cautions. Can be used on vegetables in the greenhouse.
	dimethoate (Cygon)	1 1/2 pt. 30.5% E.C.	Carnations only.
	methoxychlor	2 lb. 50% W.P.	Preferred for sensitive foliage. Can be used on lettuce, cucumbers, tomatoes, and radishes in the greenhouse.
	parathion	1 lb. 25% W.P. or 1 pt. 25% E.C.	Avoid excessive treatment of asters, gardenias, and white cyclamen. Can be used on vegetables in the greenhouse. For use by commercial growers only.

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Thrips (continued)	AEROSOLS and VAPORS		
	malathion	15% bomb using 1 lb. for 50,000 cu. ft.	Look under "Aphids" for cautions. Can be used on vegetables in the greenhouse.
	parathion, dichlorvos (Vapona), sulfotepp (Dithio), or tetradifon 15% plus sulfotepp 12.5%	As recommended for spider mites.	Parathion can be used on vegetables in the greenhouse. Dichlorvos (Vapona) can be used on cucumbers, tomatoes, and lettuce in the greenhouse.
	resmethrin	1% bomb using 1 lb. for 3,000 sq. ft.	For exposed flower thrips on mums. Effectiveness reduced in sunlight and at temperatures above 72°F.
	SOIL TREATMENT		
	aldicarb (Temik)	30 to 40 oz. Temik 10 G/1,000 sq. ft. soil surface (1/8 tsp./6-in. pot)	For use by commercial growers only. Aldicarb recommended only for thrips on chrysanthemums.
	dimethoate (Cygon)	6 oz. 30.5% E.C. in enough water to cover 1,000 sq. ft. of bench.	Carnations only.
Whiteflies	SPRAYS		
	endosulfan (Thiodan)	1 lb. 50% W.P. or 1 qt. 24% E.C.	Can be used on tomatoes in the greenhouse. Also 3% dust may be used. May injure mum varieties Bonnafon Deluxe, Fred Shoemith, and White Knight; also some geranium varieties.
	malathion	1½ pt. 57% E.C. or 2½ lb. 25% W.P.	Look under "Aphids" for cautions. Can be used on vegetables in the greenhouse.
	parathion	1 lb. 25% W.P. or 1 pt. 25% E.C.	Avoid excessive treatment of asters, gardenias, and white cyclamen. Can be used on vegetables in the greenhouse. For use by commercial growers only.
	endosulfan plus malathion or parathion	1 lb. of 50% W.P. plus 2½ lb. of 25% W.P., or 1 lb. of 25% W.P.	
	resmethrin	1 pt. to 1 qt. 24.3% E.C.	
	Enstar 5E	10 oz. 65% E.C. when pest first noted or 20 oz. for heavy populations.	Manufacture discontinued. Stocks available until depleted. Should be applied in pre-bloom stage. May injure bracts of some poinsettia varieties.

Pest	Pesticide	Dosage per 100 gal. spray unless otherwise stated	Remarks
Whiteflies (continued)	oxydemetonmethyl (Meta-Systox-R)	1½ pt. 25% E.C.	May injure mum varieties Hurricane, Iceberg, Whitetop, and Pennant; also Easter lilies.
	demeton (Systox)	1 to 1½ pt. 25% E.C.	May injure Croft lilies. Effective on foliar nematodes. For use by commercial growers only.
AEROSOLS and VAPORS			
	dichlorvos (Vapona)	6 oz. for 10,000 cu. ft., or paint 1 fl. oz. 81% E.C. per 10,000 cu. ft. on steam pipes.	Close greenhouse vents for 2 hrs. after application. Look under "Aphids" for cautions. Can be used on cucumbers, tomatoes, and lettuce in the greenhouse. May be phytotoxic to tomatoes.
	endosulfan (Thiodan)	1 lb. 10% aerosol per 50,000 cu. ft.	Check label for cautions. Can be used on tomatoes in the greenhouse.
	resmethrin	1% bomb using 1 lb. for 3,000 sq. ft.	Effectiveness reduced in sunlight at temperatures above 72°F.
	sulfotepp (Dithio)	5% bomb using 1 lb. per 50,000 cu. ft.	
	malathion	15% bomb using 1 lb. for 50,000 cu. ft.	Look under "Aphids" for cautions. Can be used on vegetables in the greenhouse.
	naled (Dibrom)	Paint 1 fl. oz. 60% E.C. per 10,000 cu. ft. on steam pipes.	Corrosive to metals. Look under "Aphids" for cautions. Can be used on cucumbers and tomatoes in the greenhouse.
SOIL TREATMENT			
	aldicarb (Temik)	30 to 40 oz. Temik 10G/1,000 sq. ft. soil surface (¼ tsp./6-in. pot).	For use by commercial growers only. Should control whiteflies for 3 to 4 weeks. Approved on mums, poinsettias, and gerbera.
	demeton (Systox)	1 to 1½ pts. 25% E.C. Apply 1 pt. of the mixture per sq. ft. of bench or ¼ pt. per 6-in. pot.	May injure African violets, Cibotium ferns, and Croft lilies. For use by commercial growers only.

Resistance of Certain Aphids (Particularly Green Peach Aphids) and Red Spider Mites

There is no easy way to minimize the development of resistance except to minimize the use of chemicals; however, the following guidelines may be helpful:

- (1) Prevent introduction of new strains from other sources by careful inspection of the plants upon receipt; preferably, isolate for a time. If an infestation is present, give a thorough treatment before permitting the plant to be mixed with the others.
- (2) Keep conditions as cool and unfavorable for the aphids and mites as possible without interfering with the growing conditions of the plants. Aphids do best at temperatures of 72-75° F and a relative humidity of about 75%. Mites do best at temperatures of 75° F and above and low relative humidity. Therefore, you can lower temperature for aphid and mite control and raise humidity for mite control.
- (3) Make each treatment thorough, but do not overtreat.
- (4) Alternate insecticides if resistance is suspected. Be sure to select the replacement from a different chemical group. Chemicals and their groups are given below:

Aphids: (1) Chlorinated hydrocarbons — endosulfan (Thiodan); (2) Organophosphates — demeton (Systox), dichlorvos (Vapona), dimethoate (Cygon), malathion, naled (Dibrom), oxydemetonmethyl (Meta-Systox-R), parathion, sulfotepp (Dithio), acephate (Orthene); (3) Carbamate — aldicarb (Temik), pirimicarb (Pirimor); (4) Botanical — nicotine; (5) Insect growth regulator — Enstar 5E.

Spider Mites: (1) Chlorinated hydrocarbons — dicofol (Kelthane), Pentac; (2) Organophosphates — carbophenothion (Trithion), demeton (Systox), dichlorvos (Vapona), dimethoate (Cygon), naled (Dibrom), oxydemetonmethyl (Meta-Systox-R), parathion, malathion, sulfotepp (Dithio); (3) Carbamate — aldicarb (Temik); (4) An organic carbonate — Morestan; (5) Sulfur compounds (not phosphates) — tetradifon (Tedion), Omite; (6) Organotin compound — Plictran, Vendex

Toxicity and Hazard

The hazard from insecticides is not only dependent upon the toxicity of the chemical but upon how it is used and what form of the insecticide might be contacted.

For human safety, when using aldicarb (Temik 10 G) use rubber gloves and a respirator when in an enclosure; never use treated plants for food or feed purposes; do not market potted plants within 4 weeks after treatment; wait 100 days before planting food crops in treated soil. Other chemicals also require special care. ALWAYS read the label for information on proper precautions.

The following table illustrates relative differences due to the physical state (+ + + + = very hazardous, + = least hazardous).

Physical State	Hazard		
	To Lungs	Through Skin	By Mouth
Droplets of:			
Emulsions	+ +	+ + +	+
Wettable or flowables	+ +	+ +	+
Solutions	+ +	+ + +	+
Smokes	+ + +	+	+
Vapors (aerosols)	+ + + +	+	+
Dusts	+ + +	+ +	+
Granules	+	+ +	+ +
The <i>highly</i> toxic (less than 5 grams or 1/8 oz. active ingredient for a man):			
demeton (Systox)	carbophenothion (Trithion)	sulfotepp (Dithio)	
parathion	aldicarb (Temik)	nicotine	
The <i>moderately</i> toxic (1/8 oz. to 2 oz. for a man):			
	lindane	dichlorvos (Vapona)	
	endosulfan (Thiodan)	diazinon (oral)	
	Mesuroil	naled (Dibrom)	
		oxydemetonmethyl (Meta-Systox-R) (oral)	
		pirimicarb (Pirimor)	



Extension Bulletin 392-1979
AGRICULTURAL EXTENSION SERVICE
UNIVERSITY OF MINNESOTA