Technical Report

Department of Computer Science
and Engineering
University of Minnesota
4-192 EECS Building
200 Union Street SE
Minneapolis, MN 55455-0159 USA

TR 97-043

Low Rank Approximation of a Hankel Matrix by Structured Total Least Norm

by: Haesun Park, Lei Zhang, and
J. Ben Rosen

Low Rank Approximation of a Hankel Matrix by Structured Total Least Norm

Haesun Park, Lei Zhang * and J. Ben Rosen†

Abstract

The structure preserving rank reduction problem arises in many important applications. The singular value decomposition (SVD), while giving the best low rank approximation to a given matrix, may not be appropriate for these applications since it does not preserve the given structure.

We present a new method for structure preserving low rank approximation of a matrix, which is based on Structured Total Least Norm (STLN). The STLN is an efficient method for obtaining an approximate solution to the overdetermined linear system $AX \approx B$ preserving the given linear structure in A or $[A \mid B]$, where errors can occur in both the right hand side matrix B and the matrix A. The approximate solution can be obtained to minimize the error in the L_p norm, where $p = 1, 2$, or ∞. An algorithm is described for Hankel structure preserving low rank approximation using STLN with L_p norm. Computational results are presented, which compare performances of the STLN based method for L_1 and L_2 norms and other existing methods for reduced rank approximation for Hankel matrices.

Keywords: overdetermined linear systems, rank deduction, Hankel structure, Toeplitz structure, structured total least norm, total least squares, 1-norm, 2-norm, singular value decomposition

1 Introduction

An important practical tool in many modern engineering and science is the singular value decomposition (SVD) [12]. For any matrix $Z \in \mathbf{C}^{m \times n}$, there exist unitary matrices

$$U = [u_1, \ldots, u_m] \in \mathbf{C}^{m \times m} \quad \text{and} \quad V = [v_1, \ldots, v_n] \in \mathbf{C}^{n \times n}$$

*The work of all three authors was supported in part by the National Science Foundation grant CCR-9509085. Computer Science Dept., Univ. of Minnesota, Minneapolis, MN 55455, U.S.A., e-mail:*hpark@cs.umn.edu, lzhang@cs.umn.edu.*

†Computer Science Dept., Univ. of Minnesota, Minneapolis, MN 55455 and Dept. of Computer Science and Engineering, Univ. of California, San Diego, La Jolla, CA 92093, U.S.A. B. Rosen is supported in part by an ARPA/NIST grant and the Minnesota Supercomputer Institute.

such that

$$Z = U\Sigma V^H \tag{1.1}$$

where $\Sigma = diag(\sigma_1, \ldots, \sigma_p) \in \mathbf{R}^{m\times p}$, where $p = min(m, n)$, is a diagonal matrix and $\sigma_1 \geq \cdots \geq \sigma_n \geq 0$. Throughout this paper, we will consider only the case when $m \geq n$, i.e., $p = n$. This SVD can also be represented as

$$Z = \sum_{i=1}^{n} \sigma_i u_i v_i^H. \tag{1.2}$$

It is well known that the SVD gives the best low rank approximation $\hat{Z}$, to the given matrix Z, which satisfies

$$\min \|Z - \hat{Z}\|_p, \quad rank(\hat{Z}) \leq k, \tag{1.3}$$

for any number $k \leq n$, where $p = 2$ or F. The solution is given by taking only the first k terms in (1.2) as

$$\hat{Z} = \sum_{i=1}^{k} \sigma_i u_i v_i^H. \tag{1.4}$$

A problem which is closely related to (1.3) is the Total Least Squares (TLS) problems [22]. For an overdetermined system

$$Ax \approx b \tag{1.5}$$

where $A \in \mathbf{C}^{m\times l}$ with $m > l$ and $b \in \mathbf{C}^{m\times 1}$, the TLS gives the solution that satisfies

$$\min_{E,x} \|[E\ r]\|_F \qquad \text{such that} \qquad (A+E)x = b + r. \tag{1.6}$$

The matrix $[A + E \quad b + r]$ is rank deficient since $b + r \in Range(A + E)$, where $(x \quad -1)^T$ is the right singular vector corresponding to the smallest singular value of $[A + E \quad b + r]$. It is well known that this minimum perturbation $[E\ r]$ for $[A\ b]$ can be represented as

$$[E\ r] = -[A\ b]v_{l+1}v_{l+1}^H = -\sigma_{l+1}u_{l+1}v_{l+1}^H$$

when $Z \equiv [A\ b]$ has its SVD

$$Z = \sum_{i=1}^{l+1} \sigma_i u_i v_i^H.$$

This shows the relation between low rank approximation and TLS solution since the rank deficient matrix

$$[A + E \quad b + r] = \sum_{i=1}^{l} \sigma_i u_i v_i^H \in \mathbf{C}^{m\times(l+1)}$$

is same as the best low rank approximant $\hat{Z}$ in (1.3), with $k = l$, for the matrix $[A\ b]$. Note that

$$\|[E\ r]\|_F = \sigma_{l+1}. \tag{1.7}$$

In many applications the data matrix Z has a special structure, such as Toeplitz, Hankel or sparse. For these applications, the SVD based solution for low rank approximation may not be appropriate, since it does not preserve the given structure. In fact, the solution (1.4) based on the SVD will typically be dense with no special structure, even when A is sparse or has a structure.

Recently, a new problem formulation, called Structured Total Least Norm (STLN), and algorithm for computing the STLN solution have been developed [12, 13]. The STLN method preserves any given linear structure in E or $[E\ r]$ and can minimize these structured perturbations in the discrete L_p norm, where $p = 1, 2\ or\ \infty$ [13]. A theoretical justification and computational testing of the STLN algorithm confirm that it is an efficient method for solving structured problems or when errors can occur only in some of the elements of A or $[A\ \ b]$ [13].

In this paper, we present a new STLN based method for structure preserving rank reduction, formulating the structure preserving low rank approximation problem as an optimization problem. The new method takes advantage of the relation between the low rank approximation via SVD and the TLS solution for overdetermined systems, which is discussed above.

Other structure preserving methods for solving overdetermined systems such as STLS [8] and CTLS [1] can also be applied to the structure preserving rank reduction problem. However, these methods can produce solutions with $\mathbf{p} = 2$ only (also see [5]) and reduce the rank of the given matrix by one only. Another method which was developed specifically for structure preserving low rank approximation is due to Cadzow [4]. The performance of Cadzow's method is compared to that of the STLN based method for a certain rank reduction problem in Section 4.

A classical problem where the new STLN based algorithm can have important impact is the model reduction problem with Hankel structure. This problem arises in may areas of engineering, such as speech encoding, filter design and high-order linear state-space models of dynamic systems, where reduced-order models can give significant improvement in solution techniques [23, 14]. In model reduction, given a Hankel matrix H, a Hankel matrix $\hat{H}$ of given rank k is to be found such that $\|H - \hat{H}\|_p$, for some matrix norm, is minimized. The STLN method for preserving Hankel structure in $[A\ \ b]$ can be easily modified to solve this problem for various norms. Moreover, STLN can be further extended to compute approximant of any given rank k. In this paper, we will concentrate on the Hankel structure. It is straightforward to apply the results on Hankel matrices to Toeplitz matrices since the Toeplitz structure can be obtained simply by permuting the rows or columns of the Hankel matrix. Our recent results on nonlinear extension of STLN [17] can also be applied to obtain nonlinear structured low rank approximation as well.

The organization of the paper is as follows. In Section 2, a formulation and algorithm of STLN for solving an overdetermined system (1.5) with Hankel structured $[A\ b]$ and its extension to multiple right hand side vectors is presented. For L_1 norm minimization, we show how the formulation can be modified so that problems with complex data can be handled as linear program as well [16, 17]. Then in Section 3, it is shown that the Hankel structure preserving method presented in Section 2 can be used to find a lower rank Hankel structured approximant. Computational test results, which compare the performance of the new STLN based method

Cadzow's method, are in Section 4. Finally, difficulties involved with structure preserving low rank approximation are discussed.

2 STLN for Overdetermined Systems with Hankel Structure

In STLN for solving the overdetermined system

$$AX \approx B \tag{2.8}$$

where $A \in \mathbf{C}^{m\times l}$ and $B \in \mathbf{C}^{m\times d}$, and $[A\ B]$ has Hankel structure, we are to find the minimum Hankel structured perturbation $[E\ F]$ so that $B + F \in Range(A + E)$.

First, we show that when all the antidiagonals in $[A\ B]$ are allowed to be perturbed, it is always possible to find a Hankel perturbation $[E\ F]$, so that $B + F \in Range(A + E)$, therefore, the solution to Eqn. (2.8) exists.

Theorem 2.1 *For any given dimensions m, l, d, with $m \geq l + d \geq 1$, and for an integer j, $1 \leq j \leq l + 1$, there exists a Hankel matrix $H \in \mathbf{C}^{m\times(l+d)}$ such that $rank(H) = l$, and from the jth to the $(j + d - 1)$th columns of H are linearly dependent on the rest l columns of H.*

Proof. The following simple example proves the theorem. Suppose the Hankel matrix $H \in \mathbf{C}^{m\times(l+d)}$ is represented as

$$H = \begin{pmatrix} \eta_1 & \cdots & \eta_l & \eta_{l+1} & \cdots & \eta_{l+d} \\ \eta_2 & \cdots & \eta_{l+1} & \eta_{l+2} & \cdots & \eta_{l+d+1} \\ \vdots & \ldots & \vdots & \vdots & \ldots & \vdots \\ \eta_m & \cdots & \eta_{l+m-1} & \eta_{l+m} & \cdots & \eta_{l+d+m-1} \end{pmatrix},$$

Let

$$\eta_i = \begin{cases} 1 & \text{if } i = l + k(l - j + 1),\ k \geq 0, \\ 0 & \text{otherwise.} \end{cases} \tag{2.9}$$

Then the following properties hold:

P1. The matrix P consisting of the first l rows and l columns of H is of the form

$$P = \begin{pmatrix} 0 & 0 & \ldots & 0 & 1 \\ 0 & 0 & \ldots & 1 & * \\ \vdots & \vdots & \ldots & \vdots & \vdots \\ 0 & 1 & \ldots & * & * \\ 1 & * & \ldots & * & * \end{pmatrix}.$$

P2. For $j \leq l$, the $(j+i-1)$th column and $(l+i)$ column of H are identical, $i = 1, ..., d$. If $j = l+1$, (2.9) becomes $\eta_l = 1$ and $\eta_k = 0$, for all $k \neq l$. This implies that the last d columns of H are all zeros.

By P1 and P2, the result follows.

■

Corollary 2.2 *If every anti-diagonal in a Hankel matrix* $[A \ B] \in \mathbf{C}^{m\times(l+d)}$, *with* $m \geq l+d$, *is allowed to be perturbed, then it is always possible to find a Hankel perturbation matrix* $[E \ F] \in \mathbf{C}^{m\times(l+d)}$ *so that* $B + F \in Range(A+E)$.

■

Remark 2.3 *If some anti-diagonals in* H *should be fixed, the above results may not be true. Consider the example*

$$H = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Suppose the top three anti-diagonals are not to be perturbed, then it is not possible to perturb the rest of H *such that the third column lies in the range space of the first two columns. This example shows that the existence of the solution depends on the structure and perturbation pattern.*

■

The following discussion on the formulation on STLN explains how we determine the minimum Hankel perturbation matrix $[E \ F]$. Also we show how the formulation changes when we use L_1 norm to handle complex valued data, so that linear programming techniques can still be used for L_1 norm minimization with complex data. One of the advantages of STLN formulation is that structured problems can be solved using different norms. In particular, when L_1 norm is used, then the method is more robust in the presence of outliers [16, 17]. For the general discussion of Structured Total Least Norm (STLN) that preserves the given structure in E or $[E \ F]$, see [16, 23].

2.1 Single Right Hand Side Vector

In this section, we present the STLN method for solving the overdetermined system

$$Ax \approx b \tag{2.10}$$

where $A \in \mathbf{C}^{m\times l}$ and $b \in \mathbf{C}^{m\times 1}$, and $[A \ b]$ has Hankel structure, and all the anti-diagonals in $[A \ b]$ are subject to error. The STLN solution will find the minimum $[E \ f] \in \mathbf{C}^{m\times(l+1)}$, which also has Hankel structure, so that

$$(A+E)x = b + f \tag{2.11}$$

for some vector x, i.e., $b + f \in Range(A + E)$.

In solving the problem (2.8) using STLN, a vector $\eta \in \mathbf{C}^{(m+l)\times 1}$ is used to represent the $m + l$ free elements (the anti-diagonals) in the Hankel perturbation matrix $[E\ f]$. Specifically, when

$$[E\ f] = \begin{pmatrix} \eta_1 & \eta_2 & \cdots & \eta_{l+1} \\ \eta_2 & \eta_3 & \cdots & \eta_{l+2} \\ \vdots & \vdots & \cdots & \vdots \\ \eta_m & \eta_{m+1} & \cdots & \eta_{l+m} \end{pmatrix},$$

the vector η is defined as

$$\eta = (\ \eta_1 \quad \eta_2 \quad \cdots \quad \eta_{l+m}\ )^T.$$

Note that the perturbation E depends only on

$$\alpha \equiv (\ \eta_1 \quad \eta_2 \quad \cdots \quad \eta_{l+m-1}\ )^T \tag{2.12}$$

and the perturbation on the right hand side is

$$f = (\ \eta_{l+1} \quad \eta_{l+2} \quad \cdots \quad \eta_{l+m}\ )^T.$$

The vectors α and f can be represented as $\alpha = P_0\eta$ and $f = P_1\eta$, where

$$P_0 = (\ I_{(l+m-1)\times(l+m-1)} \quad 0_{(l+m-1)\times 1}\ ) \in \mathbf{R}^{(l+m-1)\times(l+m)}, \qquad P_1 = (\ 0_{m\times l} \quad I_{m\times m}\ ) \in \mathbf{R}^{m\times(l+m)}.$$

The vector α and the Hankel matrix E are equivalent in the sense that given α, E is known, and vice versa.

Now, the *structured residual* $\hat{r}$ for the perturbation $[E\ f]$ is defined as

$$\hat{r} = \hat{r}(\eta, x) = b + f - (A + E)x.$$

As shown in Theorem 2.1, when $[A\ b]$ has Hankel structure, it is always possible to find the perturbation $[E\ f]$ with Hankel structure such that $b + f \in Range(A + E)$. Therefore, we can force $\hat{r}$ to be zero to obtain Hankel structured solution $[A + E\ \ b + f]$. Accordingly, the problem is formulated as the following constraint minimization problem [2, 24],

$$\begin{aligned} \min_{\eta,\, x} \quad & \|D\eta\|, \\ s.t.\ \hat{r} = 0 & \end{aligned} \tag{2.13}$$

where D is a diagonal matrix of positive weights that accounts for the repetition of each η_i. There are many ways to solve this minimization problem. We solve it by using the weighting method [2], which transforms (2.13) into

$$\min_{\eta,x} \left\| \begin{pmatrix} \omega\hat{r}(\eta, x) \\ D\eta \end{pmatrix} \right\|_p, \tag{2.14}$$

where ω is a large penalty value.

Then it is solved by using a linear approximation to $\hat{r}(\eta, x)$. Let $\Delta\eta$ and Δx represent a small change in η and x, respectively, and ΔE represent the corresponding change in E. Then the first order approximation to $\hat{r}(\eta + \Delta\eta, x + \Delta x)$ is

$$\begin{aligned} \hat{r}(\eta + \Delta\eta, x + \Delta x) &= b + P_1(\eta + \Delta\eta) - (A + E + \Delta E)(x + \Delta x) \\ &\approx b + P_1\eta - (A + E)x + P_1\Delta\eta - (A + E)\Delta x - \Delta E x. \end{aligned} \tag{2.15}$$

Introducing a matrix $Y \in \mathbf{C}^{m\times(m+l-1)}$ that satisfies

$$Ex = Y\alpha = YP_0\eta, \tag{2.16}$$

(2.14) becomes

$$\min_{\Delta x, \Delta\eta} \left\| \begin{pmatrix} \omega(YP_0 - P_1) & \omega(A + E) \\ D & 0 \end{pmatrix} \begin{pmatrix} \Delta\eta \\ \Delta x \end{pmatrix} + \begin{pmatrix} -\omega\hat{r} \\ D\eta \end{pmatrix} \right\|_p . \tag{2.17}$$

For details on how the matrix Y is formed, see [16]. The (1,1) block of the matrix

$$M \equiv \begin{pmatrix} YP_0 - P_1 & A + E \\ D & 0 \end{pmatrix} \in \mathbf{C}^{(2m+l)\times(2l+m)}$$

has Toeplitz structure as shown in the following example.

Example 2.4 *Suppose*

$$E = \begin{pmatrix} \eta_1 & \eta_2 & \eta_3 \\ \eta_2 & \eta_3 & \eta_4 \\ \eta_3 & \eta_4 & \eta_5 \\ \eta_4 & \eta_5 & \eta_6 \end{pmatrix}, \quad \textit{and} \quad f = \begin{pmatrix} \eta_4 \\ \eta_5 \\ \eta_6 \\ \eta_7 \end{pmatrix}.$$

Then

$$D^2 = diag(1, 2, 3, 4, 3, 2, 1), \qquad Y = \begin{pmatrix} x_1 & x_2 & x_3 & 0 & 0 & 0 \\ 0 & x_1 & x_2 & x_3 & 0 & 0 \\ 0 & 0 & x_1 & x_2 & x_3 & 0 \\ 0 & 0 & 0 & x_1 & x_2 & x_3 \end{pmatrix}$$

and $YP_0 = (Y\ 0) \in \mathbf{C}^{m\times(m+l)}$ *is also Toeplitz.*

■

The above discussions are summarized in Algorithm H-STLN for Hankel structure preserving solution for (2.8). For a fast algorithm that takes advantage of the structure of the matrix M, see [15].

For $p = 1$ and $p = \infty$, Eqn (2.17) can be solved as a linear program when the matrix A and the vector b are real [16. 17]. However, when the data has complex values, then we cannot use a

linear program since the absolute value of a complex number involves computation of squares. Accordingly, we modify the formulation (2.14) into

$$\min_{\eta_R,\eta_I,x_R,x_I} \left\| \begin{pmatrix} \omega \hat{r}_R(\eta, x) \\ \omega \hat{r}_I(\eta, x) \\ D\eta_R \\ D\eta_I \end{pmatrix} \right\|_p, \tag{2.18}$$

where the subscripts R and I denote real and pure imaginary parts, respectively, e.g., $x = x_R + \sqrt{-1}x_I$. In fact, for $p = 2$, (2.18) is the same as (2.14) since

$$\left\| \begin{pmatrix} \omega(\hat{r}_R + \sqrt{-1}\hat{r}_I) \\ D(\eta_R + \sqrt{-1}\eta_I) \end{pmatrix} \right\|_2 = \left\| \begin{pmatrix} \omega \hat{r}_R \\ \omega \hat{r}_I \\ D\eta_R \\ D\eta_I \end{pmatrix} \right\|_2. \tag{2.19}$$

After linearization, (2.18) becomes:

$$\min_{\substack{\Delta x_R,\ \Delta x_I, \\ \Delta\eta_R,\ \Delta\eta_I}} \left\| \begin{pmatrix} \omega(Y_R P_0 - P_1) & \omega(-Y_I P_0) & \omega(A_R + E_R) & -\omega(A_I + E_I) \\ \omega(Y_I P_0) & \omega(Y_R P_0 - P_1) & \omega(A_I + E_I) & \omega(A_R + E_R) \\ D & 0 & 0 & 0 \\ 0 & D & 0 & 0 \end{pmatrix} \begin{pmatrix} \Delta\eta_R \\ \Delta\eta_I \\ \Delta x_R \\ \Delta x_I \end{pmatrix} + \begin{pmatrix} -\omega \hat{r}_R \\ -\omega \hat{r}_I \\ D\eta_R \\ D\eta_I \end{pmatrix} \right\|_p. \tag{2.20}$$

The matrices Y_R and Y_I can be chosen as in (2.16) to satisfy

$$E_R x_R = Y_R \alpha_R,\ E_I x_R = Y_R \alpha_I,\ E_R x_I = Y_I \alpha_R,\ E_I x_I = Y_I \alpha_I.$$

With this formulation, since every entry is real, $p = 1$ and $p = \infty$ cases can be solved as linear programs. For problems with real data, (2.20) becomes (2.17).

2.2 Multiple Right Hand Side Vectors

The STLN method presented in Section 2.1 for a single right hand side vector can be extended to handle multiple right hand side (RHS) vectors [23]. Suppose the following overdetermined system with d RHS is given:

$$AX \approx B, \tag{2.21}$$

where $[A \quad B]$ has Hankel structure, $A \in \mathbf{C}^{m \times l}$, and denoting the ith column of B as $B_{\cdot i}$, $B = [B_{\cdot 1} \ \ldots \ B_{\cdot d}] \in \mathbf{C}^{m \times d}$. The problem is to find the minimum $[E \quad F] \in \mathbf{C}^{m \times (l+d)}$, which also has Hankel structure, so that $(A + E)X = B + F$ for some matrix $X = [X_{\cdot 1} \ \ldots \ X_{\cdot d}] \in \mathbf{C}^{n \times d}$.

Algorithm H-STLN

Input – A vector b and a matrix A with Hankel structure in $[A \;\; b]$, and tolerance tol. $E = 0, f = 0$.
Output – Hankel matrix $[E \;\; f]$, structured residual $\hat{r}$, and vector x such that $(A + E)x = E + f + \hat{r}$. In ideal situation, $\hat{r} = 0$.

1. Compute x from $\min \|Ax - b\|_p$. Form Y from x. Set $\hat{r} = b - Ax$.

2. **repeat**

   (a) $\underset{\Delta x, \Delta\eta}{\text{minimize}} \left\| \begin{pmatrix} \omega(YP_0 - P_1) & \omega(A+E) \\ D & 0 \end{pmatrix} \begin{pmatrix} \Delta\eta \\ \Delta x \end{pmatrix} + \begin{pmatrix} -\omega\hat{r} \\ D\eta \end{pmatrix} \right\|_p .$

   (b) Set $x := x + \Delta x$, $\eta := \eta + \Delta\eta$.

   (c) Form E and f from η, form Y from x, $\hat{r} = b + f - (A + E)x$.

   until ($\|\Delta x\| \leq tol$ and $\|\Delta\eta\| \leq tol$)

Assume the Hankel matrix $[E \;\; F]$ is represented as

$$[E\ F] = \begin{pmatrix} \eta_1 & \cdots & \eta_l & \eta_{l+1} & \cdots & \eta_{l+d} \\ \eta_2 & \cdots & \eta_{l+1} & \eta_{l+2} & \cdots & \eta_{l+d+1} \\ \vdots & \cdots & \vdots & \vdots & \cdots & \vdots \\ \eta_m & \cdots & \eta_{l+m-1} & \eta_{l+m} & \cdots & \eta_{l+d+m-1} \end{pmatrix}.$$

As in the case of $d = 1$, a vector

$$\eta = (\, \eta_1 \quad \cdots \quad \eta_{m+l+d-1} \,)^T \in \mathbf{C}^{(m+l+d-1)\times 1}$$

is used to represent $m + l + d - 1$ free elements in the Hankel perturbation matrix $[E \;\; F]$. Then the perturbation on E depends only on α as defined in (2.12). In addition, the ith column $F_{\cdot i}$ of $F = [F_{\cdot 1} \; \cdots \; F_{\cdot d}]$ is

$$F_{\cdot i} = (\, \eta_{l+i} \quad \eta_{l+i+1} \cdots \quad \eta_{l+m+i-1} \,)^T .$$

Therefore,

$$\alpha = P_0\eta \qquad \text{where} \qquad P_0 = (\, I_{(l+m-1)\times(l+m-1)} \quad 0_{(l+m-1)\times d} \,) \in \mathbf{R}^{(l+m-1)\times(l+m+d-1)}$$

and each vectors $F_{\cdot i}$, $1 \leq i \leq d$ can be represented as

$$F_{\cdot i} = P_i\eta \qquad \text{where} \qquad P_i = (\, 0_{m\times(l+i-1)} \quad I_{m\times m} \quad 0_{m\times(d-i)} \,) \in \mathbf{R}^{m\times(l+m+d-1)}.$$

The structured residual matrix $\hat{R}$ is defined as

$$\hat{R}(\eta, X) = B + F - (A + E)X.$$

For any matrix $X = [X_{.1}\ X_{.2}\ \cdots\ X_{.d}]$, define

$$vec(X) = \begin{pmatrix} X_{.1} \\ X_{.2} \\ \vdots \\ X_{.d} \end{pmatrix}$$

by stacking the columns of X into a long vector. The key idea in formulating the multiple RHS problem as STLN lies in the fact that (2.21) can be written as the following equivalent structured overdetermined system with a single right hand side vector,

$$diag(A, \ldots, A)vec(X) \equiv \begin{pmatrix} A & 0 & \ldots & 0 \\ 0 & A & \ldots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \ldots & 0 & A \end{pmatrix}_{md \times nd} \begin{pmatrix} X_{.1} \\ X_{.2} \\ \vdots \\ X_{.d} \end{pmatrix} \approx \begin{pmatrix} B_{.1} \\ B_{.2} \\ \vdots \\ B_{.d} \end{pmatrix} \equiv vec(B). \quad (2.22)$$

When A is perturbed to $A + E$ and B is perturbed to $B + F$ we want $(A + E\ \ B + F)$ to be Hankel. Also the diagonal blocks in (2.22) can only be perturbed to be the same matrix. When (2.22) is solved, the solution matrix X will be found in the vectorized form $x = vec(X)$. As in the case of $d = 1$, which is shown in Section 2.1, the problem is now formulated as the following minimization problem,

$$min \| \begin{pmatrix} \omega vec(\hat{R}(\eta, x)) \\ D\eta \end{pmatrix} \|_p. \quad (2.23)$$

In order to compute the STLN solution $x = vec(X)$, we need to construct an $md \times (m+l-1)$ matrix $Y \in \mathbf{R}^{md \times (m+l-1)}$ such that

$$diag(E, \ldots, E)x = Y\alpha. \quad (2.24)$$

Letting $Y = [Y_1^T\ Y_2^T\ \ldots Y_d^T]^T$, where each $Y_i \in \mathbf{C}^{m \times (m+l-1)}$, (2.24) gives

$$EX_{.i} = Y_i\alpha,$$

for $1 \leq i \leq d$. Because Y is completely determined by the structure of E, and E is independent of the column index i of $X_{.i}$, all Y_i's have the same structure. The only difference is the values of the matrix elements in Y_i: Y_i consists of the n elements of $X_{.i}$, $1 \leq i \leq d$. For example, consider Example 2.4 in Section 2.1 and assume that there are two right hand side vectors. Then, we

have

$$Y = \begin{pmatrix} Y_1 \\ Y_2 \end{pmatrix} = \begin{pmatrix} x_{11} & x_{21} & x_{31} & 0 & 0 & 0 \\ 0 & x_{11} & x_{21} & x_{31} & 0 & 0 \\ 0 & 0 & x_{11} & x_{21} & x_{31} & 0 \\ 0 & 0 & 0 & x_{11} & x_{21} & x_{31} \\ x_{12} & x_{22} & x_{32} & 0 & 0 & 0 \\ 0 & x_{12} & x_{22} & x_{32} & 0 & 0 \\ 0 & 0 & x_{12} & x_{22} & x_{32} & 0 \\ 0 & 0 & 0 & x_{12} & x_{22} & x_{32} \end{pmatrix} \quad \text{where } X = [X_{.1}\ X_{.2}] = \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \\ x_{31} & x_{32} \end{pmatrix}. \tag{2.25}$$

Now, the vectorized form of the structured residual $\hat{R}$ is

$$\hat{r}(\eta, X) = vec(\hat{R}(\eta, X)) = vec(B + F) - diag(A + E, \ldots, A + E)vec(X).$$

For small changes $\Delta\eta$ and $vec(\Delta X)$ in η and X, respectively, the first order approximation to $\hat{r}(\eta, X)$ is

$$\begin{aligned} \hat{r}(\eta + \Delta\eta, X + \Delta X) &= vec(B) + \begin{pmatrix} P_1 \\ \vdots \\ P_d \end{pmatrix}(\eta + \Delta\eta) \\ &\quad -diag(A + E + \Delta E, \ldots, A + E + \Delta E)vec(X + \Delta X) \\ &\approx \hat{r}(\eta, X) + \begin{pmatrix} P_1 \\ \vdots \\ P_d \end{pmatrix}\Delta\eta - diag(A + E, \ldots, A + E)vec(X). \end{aligned}$$

Since

$$diag(\Delta E, \ldots, \Delta E)vec(X) = Y\Delta\alpha = YP_0\Delta\eta,$$

with the linearization, the problem (2.23) becomes

$$\underset{\Delta x, \Delta\eta}{\text{minimize}} \left\| \begin{pmatrix} \omega diag(A + E, \ldots, A + E) & \omega(YP_0 - \begin{pmatrix} P_1 \\ \vdots \\ P_d \end{pmatrix}) \\ 0 & D \end{pmatrix} \begin{pmatrix} \Delta x \\ \Delta\eta \end{pmatrix} + \begin{pmatrix} \omega\hat{r} \\ D\eta \end{pmatrix} \right\|_p. \tag{2.26}$$

As shown in Section 2.1 for $d = 1$, the formulation (2.23) needs to be modified if we want to solve it as a linear program when $[A \quad B]$ is complex. We skip the details here since the reformulation is analogous to what is shown in section 2.1.

3 Hankel Structure Preserving Rank Reduction

In this section, we show how the results presented in Section 2 can be applied to find the Hankel structure preserving low rank approximation.

Assume that a Hankel matrix $H \in \mathbf{C}^{m\times n}$, $m > n$, with $rank(H) = n$, is given and we want to find another Hankel matrix $\hat{H} \in \mathbf{C}^{m\times n}$ such that

$$min\|H - \hat{H}\|_p \quad and \quad rank(\hat{H}) \leq k < n. \tag{3.27}$$

It is easy to show that when the optimal solutions for problem (3.27) are $\hat{H}_k$ and $\hat{H}_{k-1}$ with the requirements $rank(\hat{H}_k) \leq k$ and $rank(\hat{H}_{k-1}) \leq k-1$, respectively, the inequality

$$\|H - \hat{H}_k\|_p \leq \|H - \hat{H}_{k-1}\|_p$$

should hold. Therefore, the solution $\hat{H}$ for (3.27) will satisfy $rank(\hat{H}) = k$ in general. For simplicity of discussion, we first assume that $k = n-1$. Suppose the matrix H is partitioned as

$$H = [A\ b] \tag{3.28}$$

where A consists of the first $n-1$ columns of H and b is the last column of H. If we solve the overdetermined system

$$Ax \approx b \tag{3.29}$$

using STLN method for Hankel structured $[A\ b]$ presented in Section 2.1, then we will obtain a minimum Hankel perturbation $[E\ f]$ such that

$$b + f \in Range(A + E).$$

Therefore, $[A+E\ \ b+f]$ is a solution that has Hankel structure and the property that $rank([A+E\ \ b+f]) \leq n-1$.

When $rank(H) = n$, any of the n columns of H can be chosen to be the right hand side vector b in obtaining the reduced rank approximant $\hat{H}$ by solving an overdetermined system with a structure preserving method such as STLN. Therefore, one main difference between STLN for solving the overdetermined system $Ax \approx b$ where $[A\ \ b]$ is Hankel, and using it for structure preserving low rank approximation is that in the latter case, there are n choices for the right hand side vector b. If the structure is not imposed on $\hat{H}$, i.e., when the problem is solved as a TLS problem with the partitioning shown as (3.28), it is clear that the choice of column from H to be the right hand side vector does not affect the solution. This is due to the fact that the SVD of $[A_1\ b\ A_2]$ gives the same expression (1.4) regardless of the value of j.

For the structure preserving rank reduction, the choice of the column to be the right hand side also should not make difference. Assume that the jth column of H is chosen to be the right hand side vector b, i.e.,

$$H = [A_1\ b\ A_2]$$

where $A_1 \in \mathbf{C}^{m\times(j-1)}$, $b \in \mathbf{C}^{m\times 1}$ and $A_2 \in \mathbf{C}^{m\times(n-j)}$, $1 \leq j \leq n$. Also assume that the actual solution for the problem (3.27) is $\hat{H}$. Then the perturbation $[E_1\ f\ E_2]$ with the Hankel structure, which we obtain from solving the overdetermined system

$$[A_1\ \ A_2]x \approx b, \tag{3.30}$$

should satisfy $[A_1 + E_1 \;\; b + f \;\; A_2 + E_2] = \hat{H}$ when the solution $\hat{H}$ is unique, regardless of the location of b. Even if the solution is not unique, the solution of (3.30) should satisfy

$$\|H - \hat{H}\|_p = \|H - [A_1 + E_1 \;\; b + f \;\; A_2 + E_2]\|_p.$$

However, the computed solution obtained from the STLN (2.14) may be different depending on the choice of the right hand side vector b since this amounts to different initial value for the perturbation E and f.

For multiple rank reduction, say rank reduction by $n-k > 1$, we can apply similar procedure as for the case of rank 1 reduction ($k = n - 1$), which is described above. Suppose the matrix H is partitioned as

$$H = [A \;\; B]$$

where A consists of the first k columns of H and B of the last $n - k$ columns of H. If we solve the overdetermined system

$$AX \approx B$$

using STLN method for Hankel structured $[A \;\; B]$ presented in Section 2.2, then we will obtain a minimum Hankel perturbation $[E \;\; F]$ such that

$$B + F \in Range(A + E).$$

Therefore, $[A + E \; B + F]$ is a solution that has Hankel structure and $rank([A + E \; B + F]) \leq k$ where $k < n - 1$. As in the case of $k = n - 1$, the choice of the right hand side will affect the solution obtained by the STLN based method. For more discussion on the choice of the columns of A to be the right hand side vectors, see Section 4.3.

4 Computational Results

The STLN based Hankel structure preserving rank reduction algorithm has been implemented in MATLAB in order to investigate its computational performance. Specifically, we present the computational results that illustrate the behavior of STLN algorithm with L_1 and L_2 norms for producing the rank reduced Hankel matrix and compare the results to those of Cadzow's algorithm [4]. The test is performed for single rank reduction as well as rank reduction by 2.

For each of our tests, we constructed an $m \times n$ Hankel matrix H_c with $rank(H_c) = k < n$ for a specific value of k. Then the matrix H_c was perturbed to another Hankel matrix H so that $rank(H) = n$, which is used as the given data. In Section 4.1, we present two methods that can be used for constructing the Hankel matrix H_c with any given rank. The computational results are presented in Section 4.2. Finally, we present difficulties in structure preserving rank reduction, as well as in Cadzow's and STLN based algorithms, and discuss some ways to improve them.

4.1 Generating a Hankel Matrix of Given Rank

Unlike the unstructured cases, constructing a Hankel structured matrix of the given dimension and rank is not trivial. In this section, we present two methods for generating a Hankel matrix of any dimension $m \times n$ with $m \geq n$ and of a given rank $k \leq n$. These results are used in generating the test matrices in our computational tests.

The first method for generating a Hankel matrix of the given rank is well known and it is based on problems that occur in linear prediction [20, 19]. Suppose each of $m+n-1$ data samples η_j can be represented as a sum of $k \leq n$ terms as

$$\eta_j = \sum_{l=1}^{k} \beta_l z_l^j, \quad \text{for } j = 1, 2, ..., m+n-1, \tag{4.31}$$

where $\beta_l \neq 0$, $z_l \neq 0$, $l = 1, ..., k$, and $z_p \neq z_q$ for $p \neq q$. If we arrange these data in a Hankel matrix $H \in \mathbf{C}^{m \times n}$ as

$$H = \begin{pmatrix} \eta_1 & \eta_2 & \cdots & \eta_n \\ \eta_2 & \eta_3 & \cdots & \eta_{n+1} \\ \vdots & \vdots & \cdots & \vdots \\ \eta_m & \eta_{m+1} & \cdots & \eta_{m+n-1} \end{pmatrix},$$

then $rank(H) = k$. Also the jth column of H is a linear combination of the $(j-1)$th, $(j-2)$th, ..., $(j-k)$th columns of H, $j = k+1, ..., n$. [20, 19]. Notice that the rank of H is the same as the number of terms in the summation in (4.31). By using this method, we can construct either complex or real Hankel matrix of a given rank.

The second method is new and it is based on certain properties of circulant matrices [6]. Consider the $t \times t$ discrete Fourier transformation (DFT) matrix F [25],

$$F = [f_1 \ ... \ f_t] = \frac{1}{\sqrt{t}} \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & \omega & \omega^2 & \cdots & \omega^{t-1} \\ 1 & \omega^2 & \omega^4 & \cdots & \omega^{2(t-1)} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & \omega^{t-1} & \omega^{2(t-1)} & \cdots & \omega^{(t-1)(t-1)} \end{pmatrix}$$

where $\omega = exp(2\pi i/t)$. It is well known that the matrix F can diagonalize any circulant matrix of order t. In other words, any circulant matrix $C \in \mathbf{C}^{t \times t}$ can be represented as

$$C = F \Lambda F^* = \sum_{i=1}^{t} \lambda_i f_i f_i^* \tag{4.32}$$

where λ_i, $1 \leq i \leq t$ are the eigenvalues of C and $\Lambda = diag(\lambda_1 \ \lambda_2 \cdots \lambda_t)$.

On the other hand, any matrix $\tilde{C}$ that can be represented as

$$\tilde{C} = F\tilde{\Lambda}F^* = \sum_{i=1}^{t} \alpha_i f_i f_i^* \in \mathbf{C}^{t\times t} \tag{4.33}$$

for any $\alpha_i \in \mathbf{C}$, for $1 \le i \le t$, where $\tilde{\Lambda} = diag(\alpha_1\ \alpha_2 \cdots \alpha_t)$ is a circulant matrix. If exactly k of α_i's are non-zero, then $\tilde{C}$ has rank k. Moreover, the first k columns of $\tilde{C}$ are linearly independent. To show this, let P be the permutation matrix such that

$$P^*\tilde{\Lambda}P = \begin{pmatrix} \tilde{\Lambda}_1 & 0 \\ 0 & 0 \end{pmatrix},$$

where $\tilde{\Lambda}_1 \in \mathbf{C}^{k\times k}$ and $rank(\tilde{\Lambda}_1) = k$. Let $\tilde{F} = FP$. Then Eqn. (4.33) becomes

$$\tilde{C} = FPP^*\tilde{\Lambda}PP^*F^* = \tilde{F}\begin{pmatrix} \tilde{\Lambda}_1 & 0 \\ 0 & 0 \end{pmatrix}\tilde{F}^*.$$

It's sufficient to show that the submatrix G consisting of the first k rows and k columns of $\tilde{F}$ has full rank. The result follows since

$$\tilde{F}(1:k,1:k) = G = \begin{pmatrix} 1 & 1 & \dots & 1 \\ \omega^{i_1} & \omega^{i_2} & \dots & \omega^{i_k} \\ \omega^{2i_1} & \omega^{2i_2} & \dots & \omega^{2i_k} \\ \vdots & \vdots & & \vdots \\ \omega^{i_1(k-1)} & \omega^{i_2(k-1)} & \dots & \omega^{i_k(k-1)} \end{pmatrix}$$

where $0 \le i_j \le t-1$, $j = 1, ..., k$ and $i_p \ne i_q$ when $p \ne q$. Therefore, when $t = m + n - 1$ for any positive integers m and n with $m \ge n$ and $k \le n$, the $m \times n$ submatrix T of $\tilde{C}$, which is defined as

$$T = \sum_{i=1}^{k} \alpha_i f_i(1:m) f_i^*(1:n) \in \mathbf{C}^{m\times n}$$

is Toeplitz and $rank(T) = k$. Moreover, the first k column of T are linearly independent. By permuting the rows of T by applying the anti-diagonal matrix I_m^A of order m from its left, we obtain the Hankel matrix

$$H = I_m^A T \tag{4.34}$$

with the property that $H \in \mathbf{C}^{m\times n}$, $rank(H) = rank(H(:,1:k)) = k \le n$.

In general, the Hankel matrix generated by the above method will have complex elements. However, using some properties of circulant matrices, we can generate a *real* Hankel matrix of a given rank. This is based on the fact that if α_i's in (4.33) are chosen such that

$$\alpha_1 \in \mathbf{R}, \quad \alpha_{i+1} = \overline{\alpha_{t-i+1}}, \quad i = 1, ..., t-1, \tag{4.35}$$

then $\tilde{C}$ is real. In fact, the (i,j)th element of $\tilde{C}$ is

$$\frac{1}{t}\sum_{l=0}^{t-1} \omega^{(i-j)l}\alpha_{l+1}.$$

Note that $\omega^{(i-j)l}$ and $\omega^{(i-j)(t-l)}$ are conjugate to each other and when (4.35) holds, the (i,j)th element of $\tilde{C}$ is real. Consequently, the Hankel matrix generated by the above method is also real.

4.2 Performance of the STLN based Rank Reduction Method

The STLN based rank reduction algorithm has been implemented in MATLAB in order to investigate its computational performance. We also compared our STLN based algorithm to the Cadzow's algorithm. To our knowledge, the Cadzow's method and our STLN based method are the only two existing methods that can be used for reducing rank of a matrix by any given number while preserving the structure. We denote by STLN1 and STLN2, the STLN based algorithms with $p = 1$ and $p = 2$, respectively. In general, the STLN based rank reduction algorithms will converge to the desired Hankel structured low rank approximation if there is an exact solution close by, i.e., the given Hankel data matrix H of full rank is already close to the optimal Hankel matrix of desired lower rank. However, the practical question of how large the perturbation can be for convergence can only be answered by computational testing. For the STLN based methods, the last column of the matrix H was used as the right hand side vector b, and the penalty value of $\omega = 10^8$ was used.

For each test, a Hankel matrix $H_c \in \mathbf{R}^{7\times 5}$ with $rank(H_c) = k$ where $k = 3$ or $k = 4$ was generated by using either the linear prediction method or the circulant matrix method as described in Section 4.1. Then random perturbation vector $\delta \in \mathbf{R}^{11\times 1}$, where the components of δ are uniformly distributed random variables in [0, 1], was arranged to give a Hankel matrix $\Delta \in \mathbf{R}^{7\times 5}$. The perturbed Hankel matrix was generated as

$$H = H_c + \tau * \Delta,$$

where τ is a constant which determined the level of perturbation in the given data matrix H. Then only the Hankel matrix H is assumed to be known. The objective is to compare the accuracy and speed of the three methods, STLN1 and STLN 2 based methods and Cadzow's method, in recovering the actual solution H_c by reducing the rank of H preserving the Hankel structure.

For the purpose of comparing the speed, all three methods were iterated until the absolute value of the smallest singular value of the computed matrix $\hat{H}$ is less than 10^{-14} for rank 1 reduction. In each table, we show the total number of iterations for the STLN based and Cadzow's methods. Note that the major computational complexity of one iteration in the STLN1 and STLN2 based methods and the Cadzow's method requires a linear program, a QR decomposition, and a SVD, respectively. The maximum iteration of each method was set to be 20 linear programs (LP), 100 QR decompositions (QRD), and 100 SVD's, respectively. The structured perturbation size $\|H - \hat{H}\|_F$ after the termination is also shown.

In Table 1, we present the test results of the following problem. In the test, the data generated from 11 samples η_j, was arranged in an 7×5 Hankel matrix H_c. Each data sample

τ	# of iterations			$\|H - \hat{H}\|_F$		
	STLN1 (LP)	STLN2 (QRD)	Cadzow (SVD)	STLN1	STLN2	Cadzow
10^{-8}	1	3	52	4.002×10^{-8}	4.189×10^{-8}	4.189×10^{-8}
10^{-7}	2	2	78	3.141×10^{-7}	3.103×10^{-7}	3.103×10^{-7}
10^{-6}	2	2	88	3.631×10^{-6}	3.768×10^{-6}	3.768×10^{-6}
10^{-5}	2	3	100*	3.016×10^{-5}	2.670×10^{-5}	2.670×10^{-5}
10^{-4}	3	4	100*	3.557×10^{-4}	3.407×10^{-4}	3.407×10^{-4}
10^{-3}	4	7	100*	2.812×10^{-3}	2.840×10^{-3}	2.840×10^{-3}
10^{-2}	6	18	100*	2.942×10^{-2}	3.000×10^{-2}	3.000×10^{-2}
10^{-1}	4	31	100*	4.213×10^{-1}	4.027×10^{-1}	4.016×10^{-1}

Table 1: Rank 1 reduction of a Hankel matrix

τ	# of iterations			$\|H - \hat{H}\|_F$		
	STLN1 (LP)	STLN2 (QRD)	Cadzow (SVD)	STLN1	STLN2	Cadzow
10^{-8}	8	100*	95	3.130×10^{-8}	1.806×10^{-2}	1.744×10^{-2}
10^{-7}	8	100*	100*	4.052×10^{-7}	1.806×10^{-2}	1.744×10^{-2}
10^{-6}	8	100*	90	4.864×10^{-6}	1.806×10^{-2}	1.745×10^{-2}
10^{-5}	8	100*	95	3.764×10^{-5}	1.807×10^{-2}	1.745×10^{-2}
10^{-4}	8	100*	99	2.742×10^{-4}	1.819×10^{-2}	1.756×10^{-2}
10^{-3}	6	100*	100*	3.104×10^{-3}	1.952×10^{-2}	1.893×10^{-2}
10^{-2}	20	100*	100*	4.637×10^{-2}	3.597×10^{-2}	3.502×10^{-2}
10^{-1}	10	100*	100*	4.972×10^{-1}	4.180×10^{-1}	3.804×10^{-1}

Table 2: Rank 1 reduction of a Hankel matrix with an outlier 0.01 at the 8th anti-diagonal

η_j is represented as a sum of 4 exponential terms as in (4.31) where

$$\beta = (1.0,\ 2.0,\ 0.5\ 1.5) \qquad \text{and} \qquad z = (e^{-0.1},\ e^{-0.2},\ e^{-0.3},\ e^{-0.35}).$$

Therefore, $rank(H_c) = 4$. Note that for all values of τ, the smallest singular value became smaller than 10^{-14} in only several iterations in the STLN based algorithms when τ is less than 10^{-3}. The STLN1 based algorithm converged particularly fast. For $\tau \geq 10^{-5}$, the Cadzow's method did not satisfy the stopping criterion even after 100 iterations. However, the smallest singular values were 2.267×10^{-14} and 1.925×10^{-14}, for $\tau = 10^{-5}$ and $\tau = 10^{-4}$, respectively, at the termination after 100 iterations. In Cadzow's algorithm, the smallest singular value became less than the value of τ after about 5 iterations. However, the convergence speed after the first several iterations was extremely slow, reducing the smallest singular value by a factor of 10^{-1} in every 10 to 15 more iterations.

The results presented in Table 2 are obtained with the identical Hankel matrix $H_c \in \mathbf{R}^{7\times 5}$ as in the previous test. Then H_c was perturbed by a random Hankel matrix Δ, which is also generated as before, except that now an outlier of size 0.01 was added to η_8 in each case. It

is seen that for the STLN2 based algorithm and Cadzow's algorithm, the outlier dominated the convergence speed and the error in recovering H_c. In contrast, the STLN1 based algorithm produced a solution, which ignores the outlier and the accuracy of solution was influenced only by the size of τ. In particular, the error $\|H - \hat{H}\|_F$ with STLN1 is essentially proportional to τ only. This shows that the STLN1 based algorithm can be very robust in the presence of a large error in the data.

In the second test, we used the Hankel matrix H_c generated by the circulant matrix based method as described in Section 4.1 and solve the problem to reduce the rank by 2. The elements of the circulant matrix were randomly chosen from interval [0, 1]. The Hankel matrix $H_c \in \mathbf{R}^{7\times 5}$, with $rank(H_c) = 3$, was then perturbed similarly to generate the full rank data matrix H as in the first test. Again three methods were compared in terms of accuracy and speed. The convergence criterion was that the 2 norm of the vector consisting of the smallest 2 singular values of the computed matrix $\hat{H}$ should be less than 10^{-14}. The results are shown in Table 3 and Table 4. The results presented in Table 3 show that all three methods produced good answers in this test. However, even in the best case of Cadzow's with $\tau = 10^{-8}$, it required 28 SVD's, which is prohibitively expensive. Table 4 shows the results obtained with the same Hankel matrix H_c as those in Table 3 except that now a large error of size 0.01 was added to η_9. The solutions produced by all three methods were apparently influenced by the largest perturbation, which is the outlier. The STLN1 based method converged extremely fast especially when the noise level τ is low. However, the solutions produced by STLN1 are also affected by the outlier unlike the results presented in Table 2. This can be explained using our theoretical analysis: L_1 norm minimization is robust in the presence of outliers, but the solution may be affected by the outliers depending on the *locations* of the outliers [18].

We have conducted many other tests using different data matrix of various sizes, and the similar results were obtained. Based on these implementation results, we conclude that the STLN based method is very efficient for obtaining low rank approximation with Hankel structure. The Cadzow's method often has difficulties with convergence and its computational complexity is high even when it converges. In all our implementations, E and f were initially set to be zero. For improving the speed and convergence of the STLN based methods, it will be important to find a good initial guess for them. The results from Cadzow's method can, in fact, give an initial guess. However, using the Cadzow's result after several iterations as the initial guess for the STLN based method did not necessarily improve the solution. Also Cadzow's method is too expensive even for providing initial guess since each iteration involves on SVD.

4.3 Discussions

In this section, we discuss some difficulties involved with the problem of finding a structured low rank approximant and also involved with the algorithms we discussed in this paper for solving the problem. As shown in Section 2, the structured low rank approximation problem may not even have a solution. The existence of the solution depends on the structure which is imposed on the solution. For the Hankel structured problem where every anti-diagonal of the matrix is allowed to be perturbed, a solution always exists as shown in Theorem 2.1. On the other hand,

	# of iterations			$\|H-\hat{H}\|_F$		
τ	STLN1 (LP)	STLN2 (QRD)	Cadzow (SVD)	STLN1	STLN2	Cadzow
10^{-8}	1	1	26	2.845×10^{-8}	2.366×10^{-8}	2.318×10^{-8}
10^{-7}	3	3	33	4.126×10^{-7}	3.904×10^{-7}	3.904×10^{-7}
10^{-6}	3	3	35	3.645×10^{-6}	3.955×10^{-6}	3.954×10^{-6}
10^{-5}	5	5	37	3.091×10^{-5}	2.994×10^{-5}	2.994×10^{-5}
10^{-4}	6	6	49	2.494×10^{-4}	2.498×10^{-4}	2.947×10^{-4}
10^{-3}	6	6	38	3.026×10^{-3}	3.136×10^{-3}	3.136×10^{-3}
10^{-2}	14	10	43	3.469×10^{-2}	3.568×10^{-2}	3.568×10^{-2}
10^{-1}	15	66	61	4.714×10^{-1}	5.598×10^{-1}	3.645×10^{-1}

Table 3: Rank 2 reduction of a Hankel matrix

	# of iterations			$\|H-\hat{H}\|_F$		
τ	STLN1 (LP)	STLN2 (QRD)	Cadzow (SVD)	STLN1	STLN2	Cadzow
10^{-8}	3	12	91	2.127×10^{-2}	1.236×10^{-2}	1.234×10^{-2}
10^{-7}	3	12	91	2.128×10^{-2}	1.236×10^{-2}	1.234×10^{-2}
10^{-6}	3	11	91	2.132×10^{-2}	1.236×10^{-2}	1.234×10^{-2}
10^{-5}	3	14	91	2.073×10^{-2}	1.236×10^{-2}	1.235×10^{-2}
10^{-4}	9	13	91	4.615×10^{-4}	1.248×10^{-2}	1.247×10^{-2}
10^{-3}	9	19	91	4.110×10^{-3}	1.293×10^{-2}	1.292×10^{-2}
10^{-2}	20*	52	90	8.121×10^{-1}	3.904×10^{-2}	3.903×10^{-2}

Table 4: Rank 2 reduction of a Hankel matrix with an outlier 0.01 at the 9th anti-diagonal

the difficulty with Hankel structured problem involves the fact that there may be infinitely many global solutions as illustrated in the following simple 2×2 example.

Consider the 2×2 anti-diagonal matrix

$$I_2^A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

and the problem

$$min\|I_2^A - T\|_F, \quad T \in \mathbf{R}^{2\times 2}, \quad rank(T) = 1 \tag{4.36}$$

where the solution T is required to have Toeplitz structure. It can be shown that (4.36) has infinitely many global solutions: the matrix

$$T = \begin{pmatrix} \frac{\mu}{\mu^2+1} & \frac{\mu^2}{\mu^2+1} \\ \frac{1}{\mu^2+1} & \frac{\mu}{\mu^2+1} \end{pmatrix}$$

is a solution for (4.36) for any $\mu \in \mathbf{R}$ and $\|T - I_2^A\|_F = 1$.

The next example shows that the choice of column which is to be the right hand side vector in the STLN based algorithm is not trivial. The example also illustrates that the permutation affects the solution in Cadzow's algorithm. Suppose the optimal Hankel approximant H_c with $rank(H_c) \leq k$, for a Hankel matrix H has the minimum distance $\beta = \|H - H_c\|_F$. Then the distance between the permuted Hankel matrix $I^A H I^A$ and its best structure preserving reduced rank approximant should also have the same distance β since

$$\|I^A H I^A - I^A H_c I^A\|_F = \beta.$$

However, the following example shows that this is not necessarily true in both STLN based algorithm and Cadzow's algorithm. Consider the rank 1 reduction problem for the Hankel matrix

$$H = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & \mathbf{1} \end{pmatrix} \tag{4.37}$$

and its optimal Hankel approximant H_c with $rank(H_c) \leq 2$. The Cadzow's method gives the solution

$$\hat{H} = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

after one iteration and further iteration does not change this solution. The STLN2 based method also gives the same solution in one iteration. It is easy to see that the computed solution $\hat{H}$ is an optimal solution since $\hat{H}$ is also the solution obtained by SVD without preserving the structure.

However, when these methods are applied to the permuted Hankel matrix

$$I_3^A H I_3^A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \tag{4.38}$$

they produce poor results. In Cadzow's method, after the ith iteration of SVD and averaging along antidiagonals, we obtain

$$\hat{H}^{(i)} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 2^{-i} \\ 0 & 2^{-i} & 0 \end{pmatrix}.$$

Therefore, the computed answer converges to the matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{4.39}$$

reducing the rank of H by 2 instead of 1, with $\|H - \hat{H}\|_F = \sqrt{2}$. The same problem is observed when our STLN based method was used: the computed solution was very close to the matrix (4.39) from the first iteration. In case of STLN based algorithms, the permutation in (4.38) amounts to choosing the first column, instead of the last column, of H in (4.37) to be the right

hand side vector b. However, all three columns of H are orthogonal to each other and it is not clear how to determine the column which would be the best candidate to be the right hand sice vector b. Moreover, the situation is reversed if we want rank 2 reduction. For rank 2 reduction, all three methods produced the correct answer $\hat{H}$ of (4.39) in one step when we start with the permuted matrix (4.38), while all methods had difficulty with the matrix H of Eqn. (4.37), producing a zero matrix, which is the result of rank 3 reduction. However, in case of STLN based method, we have some preliminary results which indicate that incorporating regularization can solve the above difficulty.

Although the STLN based algorithm seems far superior to other existing methods as shown in this paper, in order to more fully develop the STLN based algorithm for structured low rank approximation problems, a number of areas need further study. Future work will include finding a good initial guess for STLN, choice of the column to be the right hand side vector, and applying regularization.

References

[1] T.J. Abatzoglou, J.M. Mendel, G.A. Harada. The constrained total least squares technique and its application to harmonic superresolution. *IEEE Transactions on Signal Processing*, 39:1070–1087, 1991.

[2] A.A. Anda and H. Park. Self-scaling fast rotations for stiff least squares problems, *Linear Algebra and its Applications*, 234:137-161, 1996.

[3] H. Chen, S. Van Huffel and J. Vandewalle. Structured total least norm and its application to exponential data modeling. Proceedings PRORISC/IEEE Benelux workshop on Circuits, Systems and Signal Processing, Mierlo, The Netherlands, March 23-24, 1995, to appear.

[4] J. Cadzow. Signal enhancement: a composite property mapping algorithm. *IEEE Trans. on Acoustics, Speech, and Signal Processing*, ASSP–36:49–62, 1988.

[5] M.T. Chu, R.E. Funderlic, and R.J. Plemmons. Least squares approximation of structured lower rank matrices, draft manuscript, 1997.

[6] P.J. Davis. Circulant Matrices, Wiley-Interscience, 1979.

[7] B. De Moor. Structured total least squares and L_2 approximation problems. *Linear Algebra and its Applications*, special issue on Numerical Linear Algebra Methods in Control, Signals and Systems, Eds. Van Dooren, et. al., 188-189:163-207, 1993.

[8] B. De Moor. Total least squares for affinely structured matrices and the noisy realization problem. *IEEE Transactions on Signal Processing*, SP–42:3104–3113, 1994.

[9] R. Fletcher. *Practical Methods of Optimization, Vol. 1, Unconstrained Optimization.* John Wiley, New York, 1980.

[10] K. Glover. All optimal Hankel-norm approximations of linear multivariable systems and their l_∞-error bounds. *Int. J. Control*, 1115-1193, 1984.

[11] G. H. Golub and C. F. Van Loan. An analysis of the total least squares problem. *SIAM Journal on Numerical Analysis*, 17:883–893, 1980.

[12] G. H. Golub and C. F. Van Loan. *Matrix Computations*, second edition. Johns Hopkins University Press, Baltimore, 1989.

[13] S. Haykin. *Adaptive Filter Theory*, second edition. Prentice Hall, Englewood Cliffs, NJ, 1991.

[14] P. Lemmerling, I. Dologlou, and S. Van Huffel. Speech compression based on exact modeling and structured total least norm optimization. Technical Report, ESAP-SISTA/TR 1997-31, Katholieke Universiteit Leuven, Belgium, 1997.

[15] J.B. Rosen, H. Park, and J. Glick. The total least norm problem: formulation and algorithms, Army High Performance Computing Research Center preprint 94-041, University of Minnesota, November, 1993, revised July, 1994.

[16] J.B. Rosen, H. Park, and J. Glick. Total least norm formulation and solution for structured problems. *SIAM Journal on Matrix Anal. Appl.*. 17-1:110-128, 1996.

[17] J.B. Rosen, H. Park, and J. Glick. Structured total least norm for nonlinear problems, to appear in *SIAM Journal on Matrix Anal. Appl.*.

[18] J. B. Rosen, H. Park, and L. Zhang, On robustness of L_1 norm minimization in the presence of outliers, Technical report, Computer Science Department, University of Minnesota, 1997, in preparation.

[19] E. Sjöström. Singular value computations for Toeplitz matrices. Linköping studies in Science and Technology. Theses No. 554. Department of Mathematics. Linköping Univesrity, Sweden, 1996.

[20] D.S. Stephenson. Linear prediction and maximum entropy methods in NMR spectroscopy. *Progress in NMR spectroscopy*, 20:515-626, 1988.

[21] S. Van Huffel and J. Vandewalle. Analysis and properties of the generalized total least squares problem $AX \approx B$ when some or all columns of A are subject to errors. *SIAM Journal on Matrix Anal. Appl.*, 10:294–315, 1989.

[22] S. Van Huffel and J. Vandewalle. *The Total Least Squares Problem, Computational Aspects and Analysis*. SIAM, Philadelphia, 1991.

[23] S. Van Huffel, H. Park, and J.B. Rosen. Total least norm solution for structured problems in parameter estimation, *IEEE Transactions on Signal Processing*, 44-10:2464-2474,, 1996.

[24] C.F. Van Loan. On the method of weighting for equality constrained least squares problems. *SIAM J. Numer. Anal.*, 22:851–864, 1985.

[25] C.F. Van Loan. Computational Frameworks for the Fast Fourier Transform. SIAM, 1992.